

**2013 Annual Management Report  
Norton Sound-Port Clarence Area and Arctic-  
Kotzebue Area**

by

**Jim Menard,**

**Joyce Soong,**

**Scott Kent,**

**Lauri Harlan,**

**and**

**Ashley Brown**

March 2015

---

Alaska Department of Fish and Game

Divisions of Sport Fish and Commercial Fisheries



## Symbols and Abbreviations

The following symbols and abbreviations, and others approved for the Système International d'Unités (SI), are used without definition in the following reports by the Divisions of Sport Fish and of Commercial Fisheries: Fishery Manuscripts, Fishery Data Series Reports, Fishery Management Reports, and Special Publications. All others, including deviations from definitions listed below, are noted in the text at first mention, as well as in the titles or footnotes of tables, and in figure or figure captions.

Weights and measures (metric)		General		Mathematics, statistics		
centimeter	cm	Alaska Administrative Code	AAC	all standard mathematical signs, symbols and abbreviations		
deciliter	dL	all commonly accepted abbreviations	e.g., Mr., Mrs., AM, PM, etc.	alternate hypothesis	H <sub>A</sub>	
gram	g	all commonly accepted professional titles	e.g., Dr., Ph.D., R.N., etc.	base of natural logarithm	e	
hectare	ha			catch per unit effort	CPUE	
kilogram	kg	at	@	coefficient of variation	CV	
kilometer	km			common test statistics	(F, t, $\chi^2$ , etc.)	
liter	L	compass directions:		confidence interval	CI	
meter	m	east	E	correlation coefficient (multiple)	R	
milliliter	mL	north	N	correlation coefficient (simple)	r	
millimeter	mm	south	S	covariance	cov	
Weights and measures (English)		west	W	degree (angular )	°	
	cubic feet per second	ft <sup>3</sup> /s	copyright	degrees of freedom	df	
	foot	ft	corporate suffixes:	expected value	E	
	gallon	gal	Company	greater than	>	
	inch	in	Corporation	greater than or equal to	≥	
	mile	mi	Incorporated	harvest per unit effort	HPUE	
	nautical mile	nmi	Limited	less than	<	
	ounce	oz	District of Columbia	less than or equal to	≤	
	pound	lb	et alii (and others)	logarithm (natural)	ln	
	quart	qt	et cetera (and so forth)	logarithm (base 10)	log	
yard	yd	exempli gratia		logarithm (specify base)	log <sub>2</sub> , etc.	
Time and temperature		(for example)	e.g.	minute (angular)	'	
	day	d	Federal Information Code	not significant	NS	
	degrees Celsius	°C	id est (that is)	null hypothesis	H <sub>0</sub>	
	degrees Fahrenheit	°F	latitude or longitude	percent	%	
	degrees kelvin	K	monetary symbols	probability	P	
	hour	h	(U.S.)	probability of a type I error (rejection of the null hypothesis when true)	α	
	minute	min	months (tables and figures): first three letters	probability of a type II error (acceptance of the null hypothesis when false)	β	
	second	s	registered trademark	second (angular)	"	
	Physics and chemistry		trademark	Jan.,...,Dec	standard deviation	SD
		all atomic symbols		®	standard error	SE
alternating current		AC	™	variance		
ampere		A	United States (adjective)	population	Var	
calorie		cal	United States of America (noun)	sample	var	
direct current		DC	U.S.C.			
hertz		Hz	U.S. state			
horsepower		hp				
hydrogen ion activity (negative log of)		pH				
parts per million		ppm				
parts per thousand	ppt, ‰					
volts	V					
watts	W					

***FISHERY MANAGEMENT REPORT NO. 15-09***

**2013 ANNUAL MANAGEMENT REPORT  
NORTON SOUND-PORT CLARENCE AREA AND ARCTIC-KOTZEBUE  
AREA**

By

Jim Menard, Joyce Soong, Scott Kent, Lauri Harlan, and Ashley Brown  
Alaska Department of Fish and Game, Division of Commercial Fisheries, Nome

Alaska Department of Fish and Game  
Division of Sport Fish, Research and Technical Services  
333 Raspberry Road, Anchorage, Alaska, 99518-1565

March 2015

The Fishery Management Reports series was established in 1989 by the Division of Sport Fish for the publication of an overview of management activities and goals in a specific geographic area, and became a joint divisional series in 2004 with the Division of Commercial Fisheries. Fishery Management Reports are intended for fishery and other technical professionals, as well as lay persons. Fishery Management Reports are available through the Alaska State Library and on the Internet: <http://www.adfg.alaska.gov/sf/publications/>. This publication has undergone regional peer review.

*Jim Menard, Joyce Soong, Scott Kent, and Ashley Brown  
Alaska Department of Fish and Game, Division of Commercial Fisheries,  
P.O. Box 1148, Nome, AK 99762, USA*

*This document should be cited as:*

*Menard, J., J. Soong, S. Kent, L. Harlan, and A. Brown. 2015. 2013 Annual management report Norton Sound-Port Clarence Area and Arctic-Kotzebue. Alaska Department of Fish and Game, Fishery Management Report No. 15-09, Anchorage.*

The Alaska Department of Fish and Game (ADF&G) administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act (ADA) of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

**If you believe you have been discriminated against in any program, activity, or facility please write:**

ADF&G ADA Coordinator, P.O. Box 115526, Juneau, AK 99811-5526

U.S. Fish and Wildlife Service, 4401 N. Fairfax Drive, MS 2042, Arlington, VA 22203

Office of Equal Opportunity, U.S. Department of the Interior, 1849 C Street NW MS 5230, Washington DC 20240

**The department's ADA Coordinator can be reached via phone at the following numbers:**

(VOICE) 907-465-6077, (Statewide Telecommunication Device for the Deaf) 1-800-478-3648,

(Juneau TDD) 907-465-3646, or (FAX) 907-465-6078

**For information on alternative formats and questions on this publication, please contact:**

ADF&G, Division of Sport Fish, Research and Technical Services, 333 Raspberry Rd, Anchorage AK 99518 (907) 267-2375

# TABLE OF CONTENTS

	Page
LIST OF TABLES.....	v
LIST OF FIGURES.....	v
LIST OF APPENDICES .....	v
ABSTRACT .....	1
INTRODUCTION.....	1
<u>SECTION 1: MANAGEMENT AREA OVERVIEWS</u> .....	<u>2</u>
BOUNDARIES .....	2
SALMON OVERVIEW .....	2
Commercial Salmon Fishery .....	3
Subsistence Salmon Fishery .....	3
Salmon Management .....	4
NORTON SOUND SALMON OVERVIEW.....	5
District Boundaries .....	5
Historical Fishery Use .....	6
Commercial Fishery Overview.....	7
Commercial Fishery Management.....	8
Subsistence Fishery Overview.....	9
Historical Regulatory Actions in Norton Sound Subdistricts .....	10
PORT CLARENCE SALMON OVERVIEW .....	13
District Boundaries .....	13
Commercial Fishery Overview.....	13
Subsistence Fishery Overview.....	14
KOTZEBUE SALMON OVERVIEW .....	15
District Boundaries .....	15
Commercial Fishery Overview.....	16
Subsistence Fishery Overview.....	18
ARCTIC SALMON OVERVIEW .....	18
District Boundaries .....	18
Subsistence Fishery Overview.....	18
PACIFIC HERRING OVERVIEW .....	19
District Boundaries .....	19
Spawning Areas and Timing .....	19
NORTON SOUND PACIFIC HERRING OVERVIEW.....	21
Commercial Fishery Overview.....	21
Sac Roe.....	21
Spawn-on-Kelp.....	22
Food and Bait Fishery.....	22
Commercial Fishery Management.....	23

## TABLE OF CONTENTS (Continued)

	<b>Page</b>
Subsistence Fishery Use .....	24
PORT CLARENCE AND KOTZEBUE PACIFIC HERRING OVERVIEW .....	24
Commercial Fishery Overview .....	24
Historical Resource Investigations .....	24
KING CRAB OVERVIEW .....	25
Norton Sound King Crab Overview .....	25
District Boundaries .....	25
Abundance .....	25
Commercial Fishery Overview .....	27
CDQ Fishery .....	28
Commercial Catch Sampling .....	28
Subsistence Fishery Overview .....	28
St. Lawrence Island And Kotzebue King Crab Overview .....	29
District Boundaries .....	29
Abundance .....	29
Commercial Fishery Overview .....	30
MISCELLANEOUS FISH OVERVIEW .....	30
Inconnu (Sheefish).....	30
Spawning Areas and Timing.....	30
Historical Fishery Use .....	31
Subsistence Fishery .....	32
Commercial Fishery.....	32
Sport Fishery.....	32
Historical Escapement .....	33
Dolly Varden .....	33
Spawning Areas and Timing.....	34
Subsistence Fishery .....	34
Commercial Fishery.....	34
Sport Fishery.....	35
Historical Escapement .....	35
Whitefish .....	35
Spawning Areas and Timing.....	35
Commercial Fishery.....	36
Subsistence Fishery .....	36
Historical Escapement .....	37
Saffron Cod .....	37
Miscellaneous Finfish Species .....	37
Subsistence Fishery .....	37
Commercial Fishery.....	37
Sport Fishery.....	37
<b>SECTION 2: SALMON FISHERIES</b> .....	<b>38</b>
2013 Norton Sound Salmon Fishery.....	38
2013 Norton Sound Fisheries Outlook .....	38
Commercial Fishery Season Summary .....	38
Subsistence Fishery Season Summary .....	40

## TABLE OF CONTENTS (Continued)

	<b>Page</b>
Season Summary by Subdistrict .....	40
Nome–Norton Sound Subdistrict 1 .....	40
Golovin–Norton Sound Subdistrict 2 .....	41
Elim–Norton Sound Subdistrict 3 .....	42
Norton Bay–Norton Sound Subdistrict 4 .....	43
Shaktoolik and Unalakleet–Norton Sound Subdistricts 5 and 6.....	44
Escapement.....	45
Chinook Salmon .....	45
Coho Salmon .....	46
Pink Salmon .....	47
Sockeye Salmon .....	47
Enforcement.....	48
2014 Norton Sound Salmon Outlook.....	48
2013 Port Clarence Salmon Fishery .....	49
Commercial Fishery Season Summary .....	49
Subsistence Fishery Season Summary .....	49
Escapement .....	50
Enforcement.....	50
2014 Port Clarence Salmon Outlook .....	50
2013 Kotzebue Sound Salmon Fishery.....	50
Commercial Fishery Season Summary .....	50
Subsistence Fishery Season Summary .....	51
Escapement .....	52
Enforcement.....	52
2014 Kotzebue Salmon Outlook.....	52
<b><u>SECTION 3: PACIFIC HERRING FISHERIES</u></b>	<b><u>53</u></b>
2013 Norton Sound Pacific Herring Fishery .....	53
Sac Roe.....	53
Spawn-on-Kelp.....	53
Bait Fishery.....	53
Commercial Fishery Management .....	53
Catch Reporting and Enforcement.....	54
Biomass Determination .....	54
2014 Norton Sound Pacific Herring Outlook .....	54
<b><u>SECTION 4: KING CRAB FISHERIES</u></b>	<b><u>55</u></b>
Norton Sound Crab Fishery .....	55
Abundance .....	55
Summer Open Access Commercial Fishery .....	55
CDQ Fishery.....	56
Harvest Areas and Commercial Catch Sampling.....	56
Enforcement.....	57
Winter Commercial Fishery.....	57
Subsistence Fishery .....	57
Sport Fishery.....	58
Future Resource Investigations.....	58
St. Lawrence Island Crab Fishery.....	58
Commercial Fishery.....	58
<b><u>SECTION 5: MISCELLANEOUS SPECIES</u></b>	<b><u>59</u></b>

## TABLE OF CONTENTS (Continued)

	<b>Page</b>
Inconnu (Sheefish).....	59
Commercial Fishery.....	59
Subsistence and Sport Fishery .....	59
Escapement .....	59
Dolly Varden .....	59
Commercial Fishery.....	59
Subsistence and Sport Fishery .....	59
Escapement.....	60
Whitefish .....	60
Commercial Fishery.....	60
Subsistence and Sport Fishery .....	60
Saffron Cod .....	60
Commercial Fishery.....	60
Subsistence and Sport Fishery .....	60
Capelin.....	61
Commercial Fishery.....	61
Subsistence .....	61
ACKNOWLEDGEMENTS.....	61
REFERENCES CITED .....	62
TABLES .....	65
APPENDIX A: NORTON SOUND FISHERIES.....	81
APPENDIX B: PORT CLARENCE FISHERIES .....	117
APPENDIX C: KOTZEBUE FISHERIES .....	123
APPENDIX D: HERRING FISHERIES .....	135
APPENDIX E: KING CRAB FISHERIES .....	149
APPENDIX F: MISCELLANEOUS FISHERIES .....	173
APPENDIX G: OVERVIEW OF 2013 .....	183
APPENDIX H: ARCTIC FISHERIES .....	227



## LIST OF FIGURES

Figure	Page
1 Norton Sound, Port Clarence, Kotzebue Sound, and Arctic management districts.....	2
2 Norton Sound commercial salmon fishing subdistricts. ....	5
3 Port Clarence District. ....	13
4 Seward Peninsula with road accessible waters.....	15
5 Kotzebue Sound District, villages and subsistence fishing area.....	16
6 Kotzebue Sound commercial salmon fishing subdistricts and statistical areas. ....	17
7 Arctic management district. ....	19
8 Commercial herring districts and statistical areas of Norton Sound, Port Clarence, and Kotzebue Sound.....	20
9 King crab fishing districts and sections of Statistical Area Q. ....	26
10 Kotzebue and Kobuk River Valley villages and their spatial relationship with inconnu spawning and overwintering areas. ....	31

## LIST OF TABLES

Table	Page
1 Norton Sound commercial salmon harvest summary by subdistrict, 2013.....	66
2 Subsistence salmon harvest for northern Norton Sound, 2013.....	67
3 Salmon counts of rivers and associated salmon escapement goal ranges, Norton Sound and Port Clarence, 2013.....	68
4 Commercial salmon set gillnet catches from Golovin, Subdistrict 2, Norton Sound, 2013. ....	70
5 Commercial salmon set gillnet catches from Elim, Subdistrict 3, Norton Sound, 2013.....	70
6 Commercial salmon set gillnet catches from Norton Bay, Subdistrict 4, Norton Sound, 2013.....	71
7 Commercial salmon set gillnet catches from Shaktoolik, Subdistrict 5, Norton Sound, 2013. ....	72
8 Commercial salmon set gillnet catches from Unalakleet, Subdistrict 6, Norton Sound, 2013. ....	73
9 Kotzebue District commercial chum salmon catch and average weight by date, 2013.....	74
10 Historical chum salmon catch for Kobuk River drift test fishery, 1993–2013. ....	75
11 Commercial herring sac roe harvest summary by subdistrict, Norton Sound District, 2013.....	76
12 Daily catch for the open access and CDQ summer commercial king crab harvests, Norton Sound Section, Eastern Bering Sea, July 3–September 14, 2013. ....	77
13 Commercial harvest of red king crab from Norton Sound Section by statistical area, Norton Sound District, 2013.....	80

## LIST OF APPENDICES

Appendix	Page
A1 Commercial salmon catch by species, Norton Sound District, 1990–2013.....	82
A2 Number of commercial salmon permits fished, Norton Sound, 1990–2013. ....	83
A3 Round weight and value of commercially caught salmon by species, Norton Sound District, 1990–2013.....	84
A4 Estimated mean prices paid to commercial salmon fishermen in dollars, Norton Sound District, 1990–2013.....	85
A5 Mean commercial salmon harvest weights, Norton Sound District, 1990–2013.....	86
A6 Commercial and subsistence salmon catch by species, by year in Nome Subdistrict, Norton Sound District, 1990–2013. ....	87
A7 Commercial and subsistence salmon catch by species, by year in Golovin Subdistrict, Norton Sound District, 1990–2013. ....	88
A8 Commercial and subsistence salmon catch by species, by year in Elim Subdistrict, Norton Sound District, 1990–2013. ....	89

## LIST OF APPENDICES (Continued)

Appendix	Page
A9 Commercial and subsistence salmon catch by species, by year in Norton Bay Subdistrict, Norton Sound District, 1990–2013.....	90
A10 Commercial and subsistence salmon catch by species, by year in Shaktoolik Subdistrict, Norton Sound District, 1990–2013.....	91
A11 Commercial and subsistence salmon catch by species, by year in Unalakleet Subdistrict, Norton Sound District, 1990–2013.....	92
A12 Subsistence salmon catch by species and year for St. Michael and Stebbins in Norton Sound District, 1994–2013.....	93
A13 Commercial, subsistence, and sport salmon catch by species, by year for Subdistricts 1–6 in Norton Sound District, 1990–2013.....	94
A14 Sport salmon harvest by species, by year for the Unalakleet River, 1990–2013.....	95
A15 Sport salmon harvest by species, by year for the Fish/Niukluk rivers, 1990–2013.....	96
A16 Sport salmon harvest by species, by year for the Nome River, 1990–2013.....	97
A17 Comparative salmon aerial survey escapement indices of Norton Sound streams unless noted otherwise, 1990–2013.....	98
A18 Combined aerial survey numbers of chum, pink, coho, and Chinook salmon for Norton Sound, 1990–2013.....	102
A19 Total escapement for chum, pink, coho, and Chinook salmon from weir and tower projects at Kwiniuk, Niukluk, Nome, and Snake rivers, North River, Eldorado River.....	103
A20 Total escapement and catch for chum, pink, coho, and Chinook salmon for Norton Sound District, 1995–2013.....	104
A21 Nome Subdistrict chum salmon estimated escapement, 1999–2013.....	105
A22 Historical escapement of salmon and Dolly Varden at Eldorado River counting tower, 1997–2002 and weir, 2003–2013.....	107
A23 Historical escapement of salmon and Dolly Varden at Snake River counting tower, 1995–2002 and weir 2003–2013.....	108
A24 Historical salmon escapement at Kwiniuk River counting tower, 1990–2013.....	109
A25 Historical salmon escapement at Niukluk River counting tower, 1995–2012.....	110
A26 Historical salmon escapement at Nome River counting tower, 1993–1995, and weir, 1996–2013.....	110
A27 Salmon escapement at Solomon River weir, 2013.....	111
A28 Historical sockeye salmon escapement at Glacial Lake weir, 2000–2013.....	111
A29 Historical salmon escapement at Inglutalik River counting tower, 2011–2013.....	112
A30 Historical salmon escapement at North River counting tower, 1996–2013.....	112
A31 Historical salmon escapement at Unalakleet River weir, 2010–2013.....	112
A32 Chum salmon escapement by river, Nome Subdistrict, 1993–2013.....	113
A33 Pink salmon escapement by year and river, Nome Subdistrict, 1993–2013.....	114
A34 Number of customary trade permits issued, Norton Sound District and Port Clarence District, 2007–2013.....	115
B1 Comparative sockeye salmon aerial survey indices, Port Clarence District, 1990–2013.....	118
B2 Historical escapement of salmon and Dolly Varden at Pilgrim River counting tower, 1997–2002 and weir 2003–2013.....	119
B3 Estimated number of subsistence fishing families and harvest in Port Clarence District, 1994–2013.....	120
B4 Application of 20-05-00 liquid blend of phosphorous and nitrogen fertilizer to Salmon Lake, 1997–2013.....	121
C1 Kotzebue District chum salmon catch statistics, 1990–2013.....	124
C2 Kotzebue District chum salmon type of processing and weights, 1990–2013.....	125
C3 Kotzebue District mean prices paid per pound in dollars to salmon fishermen by species, 1990–2013.....	126
C4 Kotzebue District commercial fishery dollar value estimates, 1990–2013.....	127
C5 Kotzebue District commercial and subsistence salmon catches, 1990–2004 and 2012–2013.....	128
C6 Kotzebue District subsistence chum salmon catches by village, 1990–2004 and 2012–2013.....	129

## LIST OF APPENDICES (Continued)

Appendix	Page
C7 Kotzebue District average subsistence chum salmon harvest per household by village, 1990–2004 and 2012–2013.....	130
C8 Kotzebue District chum salmon aerial survey counts, 1990–2009.....	131
C9 Kobuk River chum salmon drift test fish cumulative catch per unit effort, 2007–2013.....	133
D1 Norton Sound herring and spawn-on-kelp harvests by U.S. commercial fishermen, 1990–2013. ....	136
D2 Commercial herring fishery summary information, Norton Sound District, 1990–2013.....	137
D3 Norton Sound commercial herring harvest by subdistrict, by year, 1990–2013.....	138
D4 Port Clarence District commercial herring fishery, 1986–1996. ....	139
D5 Norton Sound herring age class composition by percentage of commercial catch, commercial gear combined, 1987–1993. ....	140
D6 Norton Sound herring age class composition by percentage of commercial catch, commercial gear combined, 1994–1999. ....	141
D7 Norton Sound herring age class composition by percentage of commercial catch, commercial gear combined, 2000–2006. ....	142
D8 Norton Sound herring age class composition by percentage of commercial catch, commercial gear, 2013.....	143
D9 Norton Sound herring age class composition by percentage of total catch, variable mesh gillnets, 1987–1992.....	144
D10 Norton Sound herring age class composition by percentage of total catch, variable mesh gillnets, 1993–1998.....	145
D11 Norton Sound herring age class composition by percentage of total catch, variable mesh gillnets, 1999–2004.....	146
D12 Norton Sound herring age class composition by percentage of total catch, variable mesh gillnets, 2005–2012.....	147
D13 Norton Sound Pacific herring age composition comparison of the 2013 variable mesh gear and the projected age composition of the 2014 return. ....	148
E1 Historical commercial summer harvest of red king crab from Norton Sound Section, Eastern Bering Sea, by statistical areas, 1990–2013 .....	150
E2 The results of the population assessment surveys conducted for red king crab in Norton Sound since 1976.....	153
E3 Historical summer commercial red king crab fishery economic performance, Norton Sound Section, Eastern Bering Sea, 1990–2013. ....	154
E4 Average length and percentage of recruit and postrecruit male red king crab from summer commercial fishery catch samples in Norton Sound Section, Eastern Bering Sea, 1990–2013.....	156
E5 Winter commercial and subsistence red king crab harvests, Norton Sound, Eastern Bering Sea, 1990–2013.....	157
E6 Summer subsistence red king crab harvests, Norton Sound, Eastern Bering Sea, 2004–2013.....	158
E7 Number of crab pots lost during the subsistence and commercial winter crab fisheries, and ADF&G winter studies, 2006–2013.....	159
E8 Size composition by percent of red king crab from winter research pots near Nome, Norton Sound, Eastern Bering Sea, 1990–2012. ....	160
E9 Current and historical catch performance for the Norton Sound summer commercial crab fishery, 2008–2013.....	161
E10 Norton Sound crab exvessel value and fishery price per pound, 1994–2013. ....	162
E11 Closed water regulations in effect for the Norton Sound summer commercial crab fishery, with dotted line showing temporary boundaries for 2013 fishery only. ....	163
E12 The percent of crab harvested during the Norton Sound summer commercial red king crab fishery east of 164° west longitude, 1990–2013. ....	164
E13 Norton Sound male red king crab size distribution from trawl assessment surveys conducted by the National Marine Fisheries Service in 1991, and by ADF&G in 1996, 1999, and 2002. ....	165
E14 Norton Sound male red king crab size distribution from trawl assessment surveys conducted by ADF&G in 2006, 2008, and 2011. ....	166

## LIST OF APPENDICES (Continued)

<b>Appendix</b>	<b>Page</b>
E15 Length composition of Norton Sound red king crab summer commercial harvests, 1990–1994.....	167
E16 Length composition of Norton Sound red king crab summer commercial harvests, 1995–1998.....	168
E17 Length composition of Norton Sound red king crab summer commercial harvests, 1999–2002.....	169
E18 Length composition of Norton Sound red king crab summer commercial harvests, 2003–2006.....	170
E19 Length composition of Norton Sound red king crab summer commercial harvests, 2007–2010.....	171
E20 Length composition of Norton Sound red king crab summer commercial harvests, 2011–2013.....	172
F1 Kotzebue District winter commercial sheefish harvest statistics, 1990–2013.....	174
F2 Kotzebue District reported subsistence harvests of sheefish, 1991–2004 and 2012–2013.....	175
F3 Non-salmon sport fish harvests in Norton Sound and Kotzebue/Chukchi Sea, 1990–2013.....	176
F4 Kotzebue District incidentally caught and sold Dolly Varden during the commercial salmon fishery, 1990–2013.....	177
F5 Subsistence harvests of Dolly Varden from the villages of Kivalina and Noatak, 1991–2013.....	178
F6 Dolly Varden sport fish harvests in Norton Sound, by river, 1990–2013.....	179
F7 Aerial survey counts of overwintering and spawning Dolly Varden in the Kotzebue District, 1990– 2013.....	180
F8 Subsistence whitefish catch and effort in the Kotzebue District, 1991–1993, 1997–2004, and 2012– 2013.....	181
F9 Norton Sound District winter commercial whitefish harvest statistics, 2007–2013.....	182
F10 Norton Sound District winter commercial saffron cod harvest statistics, 1994–1995 and 2010–2013.....	182
G1 List of common and scientific names of finfish species of the Norton Sound, Port Clarence, Kotzebue and Arctic Districts.....	184
G2 Alaska Department of Fish and Game and associated cooperative studies conducted within the Norton Sound, Port Clarence, Kotzebue, and Arctic Districts, 2013.....	185
G3 Commercial processors and buyers operating in Norton Sound and Kotzebue Sound, 2013.....	188
G4 Unalakleet subsistence salmon harvest survey form, 2013.....	189
G5 Shaktoolik subsistence salmon harvest survey form, 2013.....	190
G6 Koyuk Subdistrict subsistence salmon harvest survey form, 2013.....	191
G7 Emergency Orders issued during 2013.....	192
H1 Commercial freshwater finfish harvest and sales, Colville River, Arctic Area, 1990–2007.....	228

## ABSTRACT

This report provides information about the 2013 commercial and subsistence fisheries of Norton Sound-Port Clarence and Arctic-Kotzebue management areas of the Arctic-Yukon-Kuskokwim Region of the Alaska Department of Fish and Game, Division of Commercial Fisheries. The management areas consist of all waters from Point Romanof north of the Yukon River and west of 141 degrees W longitude and those waters draining into the Bering Sea north of Yukon River; the Chukchi Sea, Beaufort Sea and Arctic Ocean. Commercial and subsistence fisheries target 5 species of salmon (Chinook *Oncorhynchus tshawytscha*, sockeye *O. nerka*, chum *O. keta*, coho *O. kisutch*, and pink *O. gorbuscha* salmon), Pacific herring *Clupea pallasii*, red king crab *Paralithodes camtschaticus*, and miscellaneous species such as inconnu (sheefish) *Stenodus leucichthys*, whitefish *Coregonus laurettae*, Dolly Varden *Salvelinus malma*, saffron cod *Eleginus gracilis*, and capelin *Mallotus villosus*.

Key words: Norton Sound, Port Clarence, Kotzebue Sound, Arctic, subsistence, commercial fishery, management, escapement, salmon, Chinook salmon *Oncorhynchus tshawytscha*, chum salmon *Oncorhynchus keta*, coho salmon *Oncorhynchus kisutch*, pink salmon *Oncorhynchus gorbuscha*, sockeye (red) salmon *Oncorhynchus nerka*, red king crab *Paralithodes camtschaticus*, Pacific herring *Clupea pallasii*, inconnu sheefish *Stenodus leucichthys*, whitefish *Coregonus laurettae*, *Coregonus pidschian*, *Coregonus sardinella*, *Coregonus nasus*, Dolly Varden *Salvelinus malma*, saffron cod *Eleginus gracilis*, Annual Management Report (AMR), Fishery Management Report (FMR).

## INTRODUCTION

This report summarizes the 2013 season and historical information concerning management of the commercial and subsistence fisheries of Norton Sound-Port Clarence, Arctic-Kotzebue management areas of the Arctic-Yukon-Kuskokwim Region. Data from select management and research projects are included in this report. A more complete documentation of project results is presented in separate reports. Historical harvest and escapement information in this report goes back to 1990. For information prior to 1990 see Menard et al. 2013.

Data presented in this report supersede information found in previous management reports. An attempt has been made to correct errors present in earlier reports. Previously unreported data were included and are indicated by appropriate footnotes. Current-year catch data presented were derived from seasonal field data.

This report is organized into the following major sections:

- 1) Management Area Overviews
- 2) Salmon Fisheries
- 3) Pacific Herring Fisheries
- 4) King Crab Fisheries
- 5) Miscellaneous Species

Tabular data have been separated into 2 categories to facilitate use of this report: 1) Tables 1–13 present annual data, and 2) Appendices generally present historical comparisons. Not all appendices are cited in the text, and those that are cited are not necessarily cited in order.

## SECTION 1: MANAGEMENT AREA OVERVIEWS

### BOUNDARIES

Norton Sound-Port Clarence Area and Arctic-Kotzebue Area include all waters from Point Romanof in southern Norton Sound and St. Lawrence Island and west of 141 degrees W longitude, to the U.S.-Canada border (Figure 1). This area encompasses over 100,000 mi<sup>2</sup>, and has a coastline exceeding that of California, Oregon, and Washington combined. For crab management the southern boundary is Cape Romanzof.




Figure 1.—Norton Sound, Port Clarence, Kotzebue Sound, and Arctic management districts.

### SALMON OVERVIEW

There are 5 species of Pacific salmon indigenous to the area; however, chum *Oncorhynchus keta* and pink salmon *O. gorbuscha* historically are the most abundant. Chum and Chinook (king) salmon *O. tshawytscha* are found as far north as Barrow, but they are less common north of the Kotzebue Sound drainages. The northernmost large concentrations of chum salmon are found

within Kotzebue Sound drainages, but large numbers of Chinook and coho *O. kisutch* salmon are not found north of Norton Sound. Small sockeye (red) salmon *O. nerka* populations exist within a few Southern Seward Peninsula drainages. Pink salmon have been observed by aerial survey in increasing numbers in rivers north of Point Hope to Barrow. Small numbers of chum, pink, sockeye, and Chinook salmon have been reported by subsistence fishermen along the Arctic coast.

## **COMMERCIAL SALMON FISHERY**

In 1959 and 1960, Alaska Department of Fish and Game (ADF&G) biologists conducted resource inventories that indicated harvestable surpluses of salmon were available in several river systems of Norton Sound, Port Clarence, and Kotzebue districts. Historically, ADF&G has supported liberalizing various regulations by encouraging processors to explore and develop new fishing grounds since statehood. As a result, commercial salmon fishing activity grew significantly in the region and enabled some local residents to obtain cash income.

Currently, most commercial fishermen and many buying station workers are resident Native Alaskans (Yupik, Inupiat, and Siberian Yupik). Commercial fishermen operate set gillnets from outboard powered skiffs, and all commercial caught salmon are harvested in coastal marine waters.

There is no commercial salmon fishery in the Arctic District.

## **SUBSISTENCE SALMON FISHERY**

There are approximately 23,000 people in the area, the majority of whom are Native Alaskans residing in more than 40 small villages scattered along the coast and major river systems. Nearly all local residents are dependent to varying degrees on fish and game resources for their livelihoods.

Subsistence fishermen operate gillnets or seines in the main rivers and to a lesser extent in coastal marine waters to harvest salmon. Beach seines are used to catch schooling or spawning salmon and other species of fish. The major portion of fish taken during summer months is air-dried or smoked for later consumption by residents or occasionally their dogs.

Historical subsistence harvest information is discontinuous. Prior to 1960, subsistence data are either incomplete or entirely lacking. From the early 1960s until 1982, ADF&G conducted annual household surveys in communities with major salmon fisheries. In 1983, budgetary restrictions made it impossible to conduct surveys in each Norton Sound village, so surveys in many areas were suspended until 1994, when ADF&G initiated a new annual postseason household subsistence salmon harvest survey program. This program was also cut after the 2003 season in Norton Sound and after 2004 in Kotzebue Sound due to budget constraints. However, expansion of subsistence salmon permits in 2004 to Port Clarence District (affecting the communities of Teller and Brevig Mission), and Norton Sound Subdistricts 2 and 3 (affecting the communities of Council, White Mountain, Golovin, and Moses Point/Elim) has resulted in fewer household surveys because subsistence harvests for those communities are now reported through subsistence permits.

Also, in 2004, the Division of Commercial Fisheries began doing subsistence salmon household surveys annually in Shaktoolik and Unalakleet (and in Koyuk starting in 2008) and in other southern Norton Sound villages periodically. Surveyors attempt to contact all households.

ADF&G staff members use a community household list and each year update any new households and delete those no longer there. Salmon survey data are expanded to include those households that usually fish but ADF&G was unable to contact.

Prior to the fishing season, ADF&G personnel usually make one visit to each village to issue subsistence salmon fishing permits. Villagers can also call the Nome office toll free, and a permit will be mailed or faxed when possible. Village residents are able to mail completed permits to the Nome office postage free. Attempts are made to contact, by phone or letter, all permit holders who did not return their household permit. Also, trips to villages are made postseason by ADF&G personnel to collect permits and discuss the fishing season.

In 2008, a cooperative project was initiated and is ongoing (ADF&G Divisions of Commercial Fisheries, Habitat, and Subsistence; and North Slope Borough Department Wildlife Management and Planning) to assess Pacific salmon resources in the Arctic District. Components of the project include 1) documenting subsistence salmon fishing patterns such as species targeted, fishing gear and methods, harvest timing, local salmon abundance and run timing, historical knowledge, and observations of spawning locations; 2) conducting aerial surveys to document adult salmon distribution in river systems and determine which rivers could be used as index areas for future monitoring; and 3) acquiring age, sex, length and genetic samples for salmon.

## **SALMON MANAGEMENT**

The Division of Commercial Fisheries of ADF&G is responsible for management of commercial and subsistence fisheries in this vast area. Permanent full-time staff assigned to this area during 2013 consisted of an Area Management Biologist, an Assistant Area Management Biologist, a Research Biologist, and a Fish and Game Program Technician stationed in the Nome office. In addition, seasonal assistance in conducting various management and research activities was provided by approximately 20 seasonal biologists and technicians in Norton Sound, Port Clarence, and Kotzebue Sound. Biologists from regional staff provided additional assistance. In 2013, interns funded by Norton Sound Economic Development Corporation (NSEDG) were utilized as fisheries technicians at some projects. There are 5 cooperative projects staffed by NSEDG and 2 projects jointly operated by NSEDG and ADF&G in Norton Sound that supplemented salmon escapement monitoring activities of area staff.

The main objective of ADF&G's program is to manage commercial and subsistence salmon fisheries on a sustained yield basis. Field projects are conducted to provide information on salmon abundance, migration, and stock composition. Summaries of ADF&G and NSEDG projects are presented in Appendix G2.

Management of salmon fisheries is complicated by insufficient comparative catch and return information and difficulties in obtaining accurate escapement data. Management difficulties are compounded by the need to provide not only for adequate escapements, but also for the needs of several different user groups. Alaska law requires subsistence users to receive priority over other users of fish and wildlife resources. If subsistence harvest increases, commercial fishing and sport fishing may be restricted.

The cornerstone regulation that governs commercial salmon harvest in all districts is the scheduled weekly fishing period. Commercial salmon fishing regulations allow for variable fishing periods per week during the open season depending on area and season differences. ADF&G attempts to distribute fishing effort throughout the entire return to avoid harvesting only



particular segments of the run. Occasionally, fishing time is increased or decreased by emergency order. Emergency orders issued in 2013 are listed in Appendix G9. Managers issue these orders depending upon fishing conditions and strength of runs or spawning escapements, as determined by evaluation of available run timing and abundance indicators. Weekly fishery reports with fishery status and schedules are broadcast during the fishing season over radio stations KICY and KNOM in Nome, and fishery news articles are published in the *Nome Nugget* and *Arctic Sounder*.

## NORTON SOUND SALMON OVERVIEW

### DISTRICT BOUNDARIES

Norton Sound Salmon District consists of all waters between Cape Douglas in the north and Point Romanof in the south. The district is divided into 6 subdistricts and corresponding statistical areas: Subdistrict 1, Nome (333-10); Subdistrict 2, Golovin (333-20); Subdistrict 3, Elim (333-31, 32, 33); Subdistrict 4, Norton Bay (333-40); Subdistrict 5, Shaktoolik (333-50); and Subdistrict 6, Unalakleet (333-60). The subdistrict and statistical area boundaries were established to facilitate management of individual salmon stocks, and each subdistrict contains at least 1 major salmon-producing stream (Figure 2).



Figure 2.—Norton Sound commercial salmon fishing subdistricts.

All commercial salmon fishing in the district is by set gillnets in marine waters; however, fishing effort is usually concentrated near river mouths. Commercial fishing typically begins in June and targets Chinook salmon if sufficient run strength exists. Emphasis switches to chum salmon in July and the coho salmon fishery begins the fourth week of July and closes in September. Pink salmon are much more abundant in even-numbered year returns. A pink salmon directed fishery may coincide with or may be scheduled to alternate periods with the historical chum salmon directed fishery.

Salmon management had changed significantly since the mid-1990s because of limited market conditions and marginal returns of several salmon stocks within the district; however, rebounding salmon returns in the mid-2000s resulted in renewed buyer interest. There had been no commercial interest in pink salmon from 2000 to 2006, but beginning in 2007 there was some commercial fishing to harvest a small portion of the pink salmon run. Also, since 2007 there has been renewed buyer interest in Golovin and Elim Subdistricts and since 2008 in Norton Bay Subdistrict. Commercial fishery managers use estimates of run strength from escapement counting projects, test fishing, aerial surveys, and commercial fishing indexes. Nome Subdistrict is managed intensively for subsistence use: Tier II chum salmon subsistence permits, registration permits, closed waters, setting fishing period length, limiting gear, and harvest limits are all tools that can be employed throughout the season to provide for escapement needs and to maximize subsistence opportunity.

## **HISTORICAL FISHERY USE**

Archeological evidence dating back 2,000 years indicates fishing has been a part of life for Norton Sound residents for many centuries (Bockstoe 1979). The largest precontact settlements on the Bering Strait Islands and the western Seward Peninsula were located where marine mammals were the primary subsistence resource. The rest of the region's population lived in small groups scattered along the coast, often moving seasonally to access fish and wildlife resources (Thomas 1982). During summer months, residents would usually disperse in groups composed of 1 or 2 families and set up camps near the mouths of streams. Harvest levels of fish on any 1 stream were relatively small because of low concentrations of people who caught only what their families and 1 or 2 dogs needed through the winter (Thomas 1982).

A large-scale fur trade was developed by the Russians in the late 1800s and continued after the American purchase (Magdanz and Punguk 1981). These activities and support for hundreds of commercial whalers and trading ships caused trading to increase in the region around 1848 (Ray 1975). Increased competition for walrus, caribou, and other species from outsiders may have increased the importance of salmon to area residents (Magdanz and Punguk 1981). In the late 1890s, gold was discovered on the Seward Peninsula and boom towns sprang up with thousands of new immigrants flocking to the region. Commerce and the establishment of missions drew people to central year-round communities.

Mining affected fish populations significantly. Nearly every stream on the Seward Peninsula has had some sort of mining operation, ranging from simple gold panning or sluice boxes to hydraulic giants or bucket-line dredges. One example of extensive impact is the Solomon River, which is only 30 miles long but had 13 dredges working at one time. Another obvious impact was the large number of people who came to live in the region between 1900 and 1930. Communities like Nome, which had a population of 30,000, and Council, which had 10,000 residents, did not exist before gold was discovered.

In the late nineteenth century the size of dog teams increased from 2 or 3 to as many as 10 to 20. At about the same time, wooden boats began to replace kayaks (Thomas 1982). Consequently, the demand for dried fish to feed the dog teams increased with the development of better means to harvest fish. Winter transportation throughout the region was hired dog teams and drivers who carried mail or freight along the coast and across the state to the ice-free port at Seward. Dried fish, primarily chum and pink salmon, became a major barter item in response to the increased demand for dog food (Thomas 1982).

Local residents spent most of their summers catching and drying large amounts of salmon, some of which they kept for themselves; the rest they bartered or sold to mining camps, roadhouses, and trading posts or stores. For example, the Haycock mining camp on the Koyuk River bought about 2 tons of dried fish each year. Roadhouses were located at Golovin, Walla Walla, Moses Point, Isaac's Point, Ungalik, Robertvale, Foothills (south of Shaktoolik), Egavik, and other locations. Dried fish was bought in units of bundles (50 dried fish tied together) at a typical price of \$0.10 per lb from the fishermen. One elder in the area thought fishermen retained more fish for their own use, which may have averaged 5 to 10 bundles per household, compared to the amount sold (Thomas 1982).

The number of people gradually decreased over the next 20 years after the gold rush and the gold deposits were worked out. The number of dog teams diminished by the mid-1930s when mail planes and mechanical tractors were introduced, and the last dog-team mail contract ended in 1962 at Savoonga. However, local stores continued to trade and barter in dry fish at Shaktoolik, St. Michael, Unalakleet, and Golovin. An example of quantity was the 8x20x40-foot cache at the Shaktoolik store filled to the top with dry fish. One elder said the stores would buy the fish for \$0.06 per lb and then sell them for \$0.10 per lb or their equivalent in groceries and supplies (Thomas 1982). By the early 1960s, commercial salmon fishing developed into a source of summer cash and snow machines were replacing the need for dog teams. The use of dry fish to feed dogs decreased and cash became more available for exchange at stores.

## **COMMERCIAL FISHERY OVERVIEW**

Commercial salmon fishing in Norton Sound District began in Shaktoolik and Unalakleet Subdistricts in 1961. Most early interest involved Chinook and coho salmon flown in dressed condition to Anchorage for further processing. A single U.S. freezer ship purchased and processed chum and pink salmon during 1961. In 1962, two floating cannery ships operated in the district and commercial fishing was extended into Norton Bay, Moses Point, and Golovin. The peak in salmon canning operations occurred in 1963.

Since then, markets have been sporadic and some subdistricts have often been unable to attract buyers for entire seasons. A joint venture between KEG (Koyuk–Elim–Golovin) Fisheries and NPL Alaska, Inc. operated from 1984 until midseason in 1988. Two Japanese freezer ships were permitted to buy directly from domestic fishermen limited to salmon caught in the internal waters of Golovnin and Norton Bays. The most consistent markets are at Shaktoolik and Unalakleet, and onshore processing occurs at Unalakleet. Appendix G3 provides a list of commercial processors and buyers that operated in Norton Sound and Kotzebue Sound in 2013.

The commercial salmon fishing season usually opens by emergency order between June 8 and July 1 but depends on run timing within each subdistrict. The season closes by regulation on August 31 in Subdistricts 1, 2, and 3, and on September 7 in Subdistricts 4, 5, and 6, but processors often terminated their operations before regulatory closure dates in the past. However, during recent years Norton Sound Seafood Products (NSSP) has remained operational until the regulatory fishing season closure. Commercial fishing periods are set by emergency order. No commercial salmon fishing periods occurred in the Nome Subdistrict 1997–2012 because of regulatory restrictions on chum salmon, lack of buyer interest, or weak runs. In 2013, limited commercial fishing occurred for chum and pink salmon (Appendix A6).

Commercial fishing gear is restricted to gillnets. A maximum aggregate length of 100 fathoms is allowed for each fisherman and there are no depth restrictions. However, mesh size is often

restricted in an attempt to direct harvest toward a specific species of salmon. Fishing periods restricted to 6.0 in and smaller mesh gillnets are used to target chum and coho salmon. Most gillnets fished are approximately 5.875-inch stretched mesh. In Unalakleet and Shaktoolik Subdistricts, 8.25-inch stretched mesh gillnets are commonly used if there are Chinook salmon fishing periods in June through early July. During years when large pink salmon runs occur and there is a buyer, ADF&G establishes fishing periods allowing only 4.5-inch mesh or less to be used. These special small-mesh periods are an attempt to target pink salmon while reducing harvest of larger sized salmon species.

## **COMMERCIAL FISHERY MANAGEMENT**

Norton Sound District is managed on comparative commercial catch data, escapements, and weather conditions. A combination of factors are considered before managers issue emergency orders affecting seasons, fishing periods, allowable mesh size, and fishing areas.

Aerial surveys are used to monitor escapements in most Norton Sound streams. Weather conditions, time of day, type of aircraft, water and bottom conditions, date of survey, and efficiency of surveyor and pilot must be taken into account when making interannual aerial survey comparisons. Counting towers and weirs are a more consistent and accurate method of obtaining migration information and have been utilized on several river systems in Norton Sound. In 2013, there were 3 counting towers and 7 weirs in operation. One sonar project was operated on the Shaktoolik River, but the project was still in development and was not used for inseason management.

Early management emphasis is on Chinook salmon switching to chum salmon around July 1, and then gradually shifting to coho salmon during the fourth week in July. Pink salmon are abundant during even-numbered years, but often no buyer is available for this species except as incidentally caught fish when there are other salmon directed fisheries. Coho salmon catches have remained fairly stable in recent years, although they have dropped from the record levels seen in Norton Sound in the mid-2000s. Chum salmon catches have been rebounding in recent years. Management actions have consisted of a series of emergency orders that open and close fishing seasons and periods and establish gillnet mesh size specifications.

Commercial fisheries in Golovin and Elim Subdistricts have targeted chum salmon and during even-numbered years pink salmon in June and July, and coho salmon in late July and August. Commercial chum salmon harvests have dropped dramatically since the mid-1980s. Poor chum salmon runs resulted in restrictive management actions during the late 1990s and early 2000s, but in the mid-2000s there was little market interest even as runs began to rebound. However, continued improving chum salmon runs in the late 2000s in Norton Sound has sparked renewed buyer interest in the northern subdistricts.

Little or no commercial salmon harvest had occurred in Nome and Norton Bay Subdistricts since the early 1980s. Nome Subdistrict has had very depressed chum salmon stocks that, until the mid-2000s, had required closure or severe restrictions of the subsistence fishery. Although salmon runs have improved greatly with record runs of pink and coho salmon in recent years and the best chum salmon runs since the 1980s, Nome Subdistrict had been unable to attract a buyer for pink and coho salmon until recently and was closed to commercial chum salmon fishing by regulation until 2013. The Norton Bay Subdistrict often has healthy stocks, but it had been unable to attract markets willing to operate in this remote area until recently. Since 2008, improving market conditions resulted in NSSP bringing more tenders to the subdistrict, and

commercial salmon fishing has resumed in Norton Bay. Commercial salmon harvest for Norton Sound in 2013 by subdistrict is listed in Table 1.

## **SUBSISTENCE FISHERY OVERVIEW**

Norton Sound District subsistence salmon harvest surveys have been conducted sporadically since statehood. From 1994 through 2003, ADF&G conducted an annual subsistence postseason salmon harvest assessment effort in northwest Alaska to provide more extensive, complete, and reliable salmon harvest estimates than had previously existed. These household subsistence harvest surveys were primarily funded by ADF&G Division of Commercial Fisheries and were conducted by the Division of Subsistence during the fall in 8 villages (Brevig Mission, Teller, Golovin, White Mountain, Elim, Koyuk, Shaktoolik, and Unalakleet). In 2004, surveys were replaced by permits in most of northern Norton Sound. Over the last 10 years in Norton Sound Subdistricts 1–6 (2003–2012), the average subsistence harvest was 70,149 salmon, with the majority being pink salmon (Appendix A13). However, from 2004 to 2007, the village of Koyuk was not surveyed, and therefore no harvest data from Norton Bay Subdistrict are included for those years.

Two goals of the postseason household subsistence survey are to collect harvest data to estimate subsistence salmon catch by species and community, and to compile information on gear types, participation rates, sharing, use of salmon for dog food, and household size. A copy of the Norton Sound subsistence salmon harvest survey form is shown by village in Appendices G4–G6.

In 2004, ADF&G’s subsistence salmon harvest assessment program changed substantially when household surveys were discontinued in most communities because the household subsistence permit system was expanded from Nome to include Port Clarence District (affecting the communities of Teller and Brevig Mission) and Norton Sound Subdistricts 2 and 3 (affecting the communities of Council, White Mountain, Golovin, and Elim). Thereafter, subsistence salmon harvest for those communities are reported totals from subsistence permits, so household surveys have not been necessary. Permits issued at the Nome office, and by ADF&G personnel in the field, identify gear restrictions, bag limits, subsistence zones (for Subdistrict 1, Salmon Lake and Pilgrim River only), location and access descriptions, and subsistence regulations for each location or body of water. In addition, the permit contains a catch calendar for household members to record gear type used, area fished, and catch in numbers by species for each day fished. If subsistence fishermen reach their harvest limit in one river, they can fish in other rivers until they reach the limit in those rivers. Subsistence permits are important to management because they identify users, fishing effort, harvests, and limits. Subsistence salmon harvests for the current year in northern Norton Sound are listed in Table 2.

In Subdistrict 1 (Nome), low salmon stock levels combined with a large concentration of users has required subsistence fishing permits since 1975. By regulation, permits with catch calendars are issued to each requesting household listing all Nome Subdistrict fishing locations, catch limits, and gear restrictions. After the fishing season, households are required to return the completed permit to ADF&G, whether or not they actually fished. Due to the subsistence permit program, all subsistence salmon catches from Norton Sound Subdistrict 1 have been determined from returned permits since 1975. However, not all fishermen obtained or returned permits from 1975 to 2003, and the data were not expanded for unreturned permits because the assumption was that those permit holders did not fish. Beginning in 2004, stricter enforcement of regulations

including fines for failure to return a permit resulted in nearly 99% of all permits issued being returned, and since 2010, all subsistence salmon permits issued have been returned or permit holders have reported catches in person, by telephone, or by email.

Norton Bay, Shaktoolik, and Unalakleet Subdistricts have continued to be surveyed postseason by household interviews. Additionally, daily surveys of Unalakleet River and ocean subsistence fishermen have been conducted annually during the Chinook salmon run since 1985. Although total harvests by subsistence fishermen were not documented, effort and catch information were used to judge timing and magnitude of the Chinook salmon return. The commercial fishery is delayed until it becomes apparent subsistence needs are being met and Chinook salmon are beginning their upstream migration as indicated by ADF&G test net in lower Unalakleet River.

## **HISTORICAL REGULATORY ACTIONS IN NORTON SOUND SUBDISTRICTS**

Nome Subdistrict (Subdistrict 1) has been the focus of most regulatory actions within the Norton Sound District since the 1970s. Although pink salmon are usually the most abundant species of salmon in Nome Subdistrict streams, the commercial fishery primarily targeted chum salmon during the 1970s. Relatively large chum salmon catches in this subdistrict in conjunction with weak local abundance implied the fishery may have intercepted nonlocal stocks. A 1978–1979 Norton Sound stock separation study (Gaudet and Schaefer 1982) showed that some salmon tagged near Nome were recaptured in fisheries from Golovin (Subdistrict 2) to Kotzebue. In an attempt to provide for spawning requirements and to provide for an important subsistence fishery that targets local stocks, a commercial harvest guideline of 5,000–15,000 chum salmon was adopted as a regulation.

The Alaska Board of Fisheries (BOF), in response to an advisory committee petition, directed ADF&G to manage the Nome Subdistrict commercial fishery for optimal chum salmon escapement after poor chum salmon escapements during the 1982 and 1983 seasons. During 1984 fall BOF meetings, directives in practice that season became regulation. In response to public and advisory committee proposals, the following commercial fishery restrictions were adopted as regulations:

- 1) Salmon may be taken commercially only from July 1 through August 31.
- 2) Fishing periods were restricted to two 24-hour periods per week.
- 3) Waters west of Cape Nome were closed to commercial salmon fishing to allow for rebuilding of river stocks that supported the historical subsistence effort.

ADF&G was directed to allow a harvest at the lower end of the guideline harvest range of 5,000 to 15,000 chum salmon, as stipulated in regulation 5 AAC 04.360. In addition to these restrictions, a proposal to restrict sport fishery in Nome and Snake rivers was adopted in 1984 that allowed “a bag and possession limit of 15 salmon, other than Chinook salmon, of which only 5 could be chum and coho salmon, in combination.”

Subsistence permit limits in Nome and Snake rivers were restricted to 20 chum and 20 coho salmon. The remainder of the permit limit could be filled with salmon other than chum or coho salmon.

Even with these restrictive regulations in place, chum salmon escapement goals were difficult to attain. The 1987 fishing season experienced poor returns of both chum and pink salmon to Nome Subdistrict streams. Numerous management actions were made to curtail commercial fishing activities, and later, sport, personal use, and subsistence fishing were restricted. Even with such

drastic fishery restrictions, escapement goals for chum salmon were not attained during 1987 in Nome, Eldorado, Flambeau, Bonanza, Snake, and Solomon rivers. In response to this continuing trend of decreasing chum and pink salmon returns to Nome Subdistrict, several new regulations were adopted by BOF in 1987 restricting gillnet length and mesh size in the subsistence fishery. Beach seine use in specific waters in the subsistence fishery was also eliminated.

Beginning in 1991, no chum salmon harvests were allowed until escapement goals were likely to be met or conservative management actions were judged to be no longer effective. Regulation changes in 1992 affected the use of beach seines for subsistence fishing in Nome Subdistrict. Managers were given authority to permit subsistence harvest of chum or pink salmon by beach seine if escapement needs were likely to be met. In the past, beach seines were viewed as an overly effective means to harvest fish. However, since 1999, beach seines were used to harvest abundant species and allow live release of other species experiencing depressed runs.

Through a series of BOF-directed meetings, BOF concluded that the previous management plan did not provide adequate opportunity for all subsistence salmon users to supply their annual needs for chum salmon. Therefore, Nome Subdistrict was designated a Tier II subsistence chum salmon permit fishery during a special BOF meeting held in Nome, March 1999. Tier II permits are dispensed to individuals prioritized by fishing history and dependence and are based on projected harvestable surplus. As a result, ADF&G allowed 20 individuals who scored highest on the Tier II application process in 1999 to subsistence fish. The intent was to allow Tier II permit holders first priority over other subsistence users if only a small harvestable surplus of chum salmon returned. If the run was assessed to be strong, then the subsistence fishery would open to all Alaskan residents who obtain a Tier I permit and individual harvests would be restricted to prescribed bag limits. In addition, BOF established “closed waters” areas where no subsistence salmon fishing would be allowed at any time, to protect chum salmon on the spawning grounds, and placed existing chum salmon aerial survey escapement goals for 6 Nome Subdistrict streams into regulation. In 1999, due to poor chum salmon returns, ADF&G closed even the Tier II fishery, and in 2000, only 10 Tier II permits were issued.

During a BOF work session in September 2000, several Norton Sound District chum salmon stocks were determined to be stocks of concern based on the *Policy for the Management of Sustainable Salmon Fisheries*. Chum salmon in Nome Subdistrict were determined to be a stock of management concern, and chum salmon in Golovin and Elim Subdistricts were determined to be a stock of yield concern.

Based upon the stock of concern determinations, BOF made several changes to regulations for management of Norton Sound salmon. In January 2001, BOF repealed the existing biological escapement goals (BEG) in regulation and adopted optimal escapement goals (OEG) for chum salmon for 5 Norton Sound rivers. In the past, escapement goals were expressed as aerial survey counts of salmon. Aerial surveys do not count all salmon present but serve as an index to compare current and previous surveys. New OEGs are in actual number of fish and based on ADF&G escapement goal analysis (Clark 2001). For rivers where an escapement project (tower or weir) are operated, 4 of 5 OEGs were established. BOF established OEGs, by subdistrict, are as follows:

#### Nome Subdistrict (Subdistrict 1)

Snake River: 1,600–2,500 chum salmon

Nome River: 2,900–4,300 chum salmon

Eldorado River: 6,000–9,200 chum salmon

#### Elim Subdistrict (Subdistrict 3)

Kwiniuk River: 11,500–23,000 chum salmon

Tubutulik River: 9,200–18,400 chum salmon

A chum salmon management plan for Nome Subdistrict (Subdistrict 1) and a salmon management plan for Golovin and Elim Subdistricts (Subdistricts 2 and 3) were adopted by BOF. Commercial chum salmon fishing in Nome Subdistrict was closed and the fishery may not be reopened again until the abundance of chum salmon has a harvestable surplus large enough to meet subsistence needs for 4 consecutive years.

ADF&G was given authority to establish subsistence gillnet mesh size restriction of 4.5 in or less by emergency order when necessary to conserve chum salmon in Subdistricts 1, 2, and 3. Also, the Cripple and Penny rivers were closed to subsistence fishing for chum salmon.

In addition, BOF expanded legal gear for the subsistence fishery to include a line attached to a rod or pole, from Cape Espenburg on the northern Seward Peninsula along the coast to Bald Head (between Elim and Koyuk). Bald Head is the boundary between Subdistricts 3 and 4. Therefore, west of Cape Espenburg in the Kotzebue District, in Port Clarence District, and in Norton Sound District from Cape Douglas to Bald Head, hook and line became legal subsistence gear. Although hook and line can be used for subsistence fishing, sport fish methods and means requirements still apply to harvesting of fish (for example, no snagging of fish is allowed). Sport fish bag and possession limits, by species, as specified in regulation 5 AAC 70.022 also apply, except when fishing through ice or in the Nome Subdistrict subsistence areas designated for each river. However, fishermen cannot combine sport fish bag and possession limits with subsistence harvest permit limits.

In 2001, chum salmon runs began to improve in Nome Subdistrict and additional permits were issued in the Tier II chum salmon fishery. Beginning in 2004, BOF expanded the salmon subsistence permit requirement for the Norton Sound area to include all marine waters, and fresh waters flowing into marine waters from Cape Prince of Wales to Bald Head. This regulation required salmon permits to be issued in Brevig Mission, Teller, White Mountain, Golovin, and Elim in addition to Nome.

Improving chum salmon runs in Nome Subdistrict resulted in Tier II chum salmon fishery restrictions being suspended beginning in 2006. A permit is still required for subsistence salmon fishing, but there is no longer a Tier II fishery that restricts participation in subsistence fishing. In addition, the BOF allowed commercial chum salmon fishing beginning in 2013.

Regulatory actions were also undertaken in other subdistricts. Subdistricts 5 and 6 Chinook salmon were designated a stock of yield concern in 2004, and BOF continued this designation in 2007 and 2010. To increase Chinook salmon escapements, BOF also adopted a more conservative *Subdistricts 5 and 6 King Salmon Management Plan* (5 AAC 04.395) that was first implemented during the 2007 season. Under the new plan, commercial fishing directed at Chinook salmon can only occur if the midpoint of the North River tower SEG range is projected to be reached. Additionally, the plan directs ADF&G to provide escapement windows by



restricting subsistence gillnet fishing for salmon from mid-June to mid-July to two 48-hour fishing periods a week in marine waters, and two 36-hour fishing periods a week in Unalakleet River. Subsistence fishing time can only be liberalized if the department projects that the lower end of the SEG range will be achieved. If North River Chinook salmon passage is projected to fall short of the SEG, ADF&G is directed to close the Chinook salmon fishery.

## PORT CLARENCE SALMON OVERVIEW

### DISTRICT BOUNDARIES

Port Clarence District encompasses all waters from Cape Douglas north to Cape Prince of Wales including Salmon Lake and Pilgrim River drainages (Figure 3). Salmon, saffron cod *Eleginus gracilis*, whitefish, and herring *Clupea pallasii* are the major subsistence species.




Figure 3.—Port Clarence District.

*Note:* Cross-hatched area on map shows location where commercial salmon fishing may be opened.

### COMMERCIAL FISHERY OVERVIEW

In contrast to Norton Sound District, commercial fishing has been limited in Port Clarence District. In 1966, a commercial salmon fishery was established in the Grantley Harbor/Tuksuk Channel area of the Port Clarence District, but the fishery that year yielded less than 2,300 combined chum, pink, and sockeye salmon (Menard et al. 2013). It was closed later that same season, due to small salmon runs and concerns from local residents about impacts to area subsistence salmon fisheries, and had remained closed until relatively recently. In the mid-2000s, there were large increases in sockeye salmon runs as well as positive results from an ADF&G

test fishery in 2006. Consequently, in 2007, the BOF reestablished by regulation a Port Clarence District commercial salmon fishery. The BOF also established an inriver run goal of at least 30,000 sockeye salmon as a trigger point to allow a commercial fishery. The 2007 fishery harvest was 1,152 sockeye salmon, and 3,183 chum salmon, whereas the 2008 fishery harvest was 89 sockeye salmon, 256 chum salmon, and 910 pink salmon (Menard et al. 2010). The 2008 commercial fishery was closed when the inriver goal of 30,000 sockeye salmon for Pilgrim River was projected to fall short. The commercial fishery has remained closed since 2009 because of poor runs of sockeye salmon.

## **SUBSISTENCE FISHERY OVERVIEW**

Salmon Lake, which empties into the Pilgrim River in the Port Clarence District, along with Glacial Lake in the northwestern portion of the Nome Subdistrict, supports the northernmost sockeye salmon populations of significant size in North America. Subsistence harvests of sockeye salmon in the Sinuk River, which drains Glacial Lake, have historically been low due to difficulties navigating this shallow, boulder-laden river. In contrast, sockeye salmon harvests in the Pilgrim River are much higher because it is more easily traveled and several beach seining and set gillnet fishing locations are accessible via the Kougarok Road (Nome–Taylor Highway) emanating from Nome. A traditional subsistence salmon fishery has probably occurred within this district for centuries; however, subsistence fishing has only been reported at Salmon Lake since the 1930s and monitored at the upper Pilgrim River since 1962. Data collected by ADF&G personnel showed most fishermen of Brevig Mission fish northern and northeastern sections of Port Clarence District, and Teller fishermen utilize Grantley Harbor and Tuksuk Channel. Interviews with local residents indicated substantial fishing effort within Agiapuk River.

Beginning in 2007, regulations allowed for cash sales of up to \$200 worth of subsistence-taken finfish per household, per year, in the Norton Sound–Port Clarence Area only, and starting in 2013 the amount allowed was raised to \$500. From 2007 to 2012, 5 or less customary trade finfish permits were issued per year. Sales in most years were confidential because less than 4 permits were issued (Appendix A34).

Village subsistence surveys were conducted annually by the Division of Commercial Fisheries until 1983 (Appendix B3). The Division of Subsistence conducted a partial survey of Brevig Mission in 1989 and conducted full-scale household surveys of both villages from 1994 to 2003. Since expansion of the subsistence salmon permit program in 2004, subsistence salmon harvests for residents of Teller and Brevig Mission have been determined from reported totals on permits.

Salmon Lake and Pilgrim River stocks have been fished by Nome residents in addition to residents of Brevig Mission and Teller for quite some time. To conserve declining sockeye salmon stocks, BOF adopted a regulation in 1972 to close Salmon Lake and its tributaries to subsistence salmon fishing from July 15 through August 31. However, because Pilgrim River is accessible from the road system there has been increased fishing effort from Nome area residents due to increased fishing restrictions in Nome Subdistrict beginning in the 1990s (Figure 4) and more so in the mid-2000s when there were record runs of sockeye salmon to Salmon Lake.




Figure 4.—Seward Peninsula with road-accessible waters.

From 1997 to 2001, ADF&G conducted a fertilization program at Salmon Lake, partially funded by NSEDC and the Bureau of Land Management (BLM) to restore sockeye salmon to historical levels by applying liquid fertilizer. However, ADF&G could not determine whether the method was effective and suspended fertilization in 2001. After impressive 2003 sockeye salmon returns, the project was reevaluated and fertilizer was applied at a reduced rate in 2004, stopped again in 2005 and 2006, restarted in 2007 by NSEDC, and has continued in subsequent years (Appendix B4).

## KOTZEBUE SALMON OVERVIEW

### DISTRICT BOUNDARIES

Kotzebue Sound District encompasses all waters from Point Hope to Cape Prince of Wales, including those waters draining into the Chukchi Sea (Figure 5). Salmon, saffron cod, whitefish, and herring are major subsistence species.




Figure 5.—Kotzebue Sound District, villages and subsistence fishing area.

## COMMERCIAL FISHERY OVERVIEW

Kotzebue Sound District supports the northernmost commercial salmon fishery in Alaska. Kotzebue Sound District is divided into 3 subdistricts. Subdistrict 1 has 6 statistical areas where commercial salmon fishing may occur (Figure 6).

The commercial fishery under state management opened in 1962. Salmon harvests consist primarily of chum salmon, although limited amounts of Dolly Varden and a few Chinook, sockeye, pink, and coho salmon are harvested during the salmon fishery.




Figure 6.—Kotzebue Sound commercial salmon fishing subdistricts and statistical areas.

The earliest documented sales of salmon in Kotzebue District were in 1909 when Lockhart's store purchased 21,906 lb of salmon from local Native Alaskans and resold it at \$0.05/lb. Of those sales, 21,366 lb were sold to gold miners on the Kobuk River drainage and 540 lb were sold to a company in Seattle. A commercial fishery occurred from 1914 to 1918. Salmon were canned and the bulk of the harvest is assumed to have been sold to miners who worked in the upper Kobuk River drainage. The next organized commercial fishery began under state management in 1962 and continues to the present. The current fishery became fully developed in the mid-1970s. The fishery displayed a gradually declining pattern of overall run strength with 4-year cycles of stronger returns followed by weaker returns (Appendix C1). In 1987, the fisheries managers' new program emphasized attaining escapement goals. Before 1987, harvests were proportional to total return. Since 1995, poor market conditions and/or limited buyer capacity have caused harvests to fall short of their potential.

In 1981, a chum salmon hatchery was established at Sikasuilaq Springs, a tributary of Noatak River. The hatchery was closed in 1995 due to lack of funding support. At peak production in 1992, the hatchery incubated 11,100,000 eggs. An estimated peak adult hatchery return of 90,000 chum salmon occurred in 1997. The estimated contribution to the commercial fishery was unknown.

## **SUBSISTENCE FISHERY OVERVIEW**

Subsistence salmon fishing in Kotzebue Sound District continues to be important, but fish abundance and fishing activities vary from community to community. Along the Noatak and Kobuk rivers where chum salmon runs are strong, household subsistence activities in middle and late summer revolve around catching, drying, and storing salmon. In southern Kotzebue Sound, fewer salmon are taken for subsistence because of low availability. Some fishermen base their fishing effort out of their village, while others move seasonally to fish camps where they stay for several days to several weeks. The predominant species in the district is chum salmon, although small numbers of other salmon species are present.

Historical subsistence surveys for the Kotzebue area have been less complete than for Norton Sound and Port Clarence Districts. However, expanded documented surveys from 1995–2004 result in an estimated total subsistence salmon harvest for the Kotzebue Sound area to be 57,977 annually (Appendix C5). During these years, ADF&G Division of Subsistence (DOS) conducted annual household subsistence salmon surveys in select Kotzebue District communities. Due to budget constraints these surveys were discontinued in 2005 but were restarted in 2012, when comprehensive subsistence fish harvest data were again collected from Kotzebue area villages by DOS. The town of Kotzebue was surveyed in 1995–2001 using a mail-in postcard but has not been surveyed since.

## **ARCTIC SALMON OVERVIEW**

### **DISTRICT BOUNDARIES**

The Arctic District includes all waters of Alaska north of the latitude of the western most tip of Point Hope and west of 141 degrees W longitude, including those waters draining into the Chukchi Sea, Beaufort Sea, and Arctic Ocean (Figure 7).

### **SUBSISTENCE FISHERY OVERVIEW**

There are no commercial salmon fisheries in the Arctic District. Small numbers of chum, pink, and Chinook salmon have been reported by subsistence fishermen along the Arctic coast, with pink salmon being the most numerous and then chum salmon (Craig George, North Slope Borough, senior wildlife biologist, personal communication). Salmon are caught in gillnets as an incidental species when subsistence fishermen are targeting other non-salmon finfish. In October 2012, a fisherman caught 2 sockeye salmon in Ikroavik Lake, approximately 5 miles south of Barrow, subsistence fishing with gillnets under the ice targeting least cisco (Geoff Carroll, ADF&G, Barrow; personal communication). There are no reliable reports of coho salmon being caught.




Figure 7.—Arctic management district.

## PACIFIC HERRING OVERVIEW

### DISTRICT BOUNDARIES

Pacific herring *Clupea pallasii* are present in Norton Sound, Port Clarence, Kotzebue Sound, and Arctic Districts. Norton Sound Herring District consists of all state waters between the latitude of the westernmost tip of Cape Douglas and the latitude of Point Romanof (Figure 8). Port Clarence Herring District consists of all Alaska waters between the latitude of Cape Douglas and the latitude of Cape Prince of Wales. Kotzebue Sound Herring District consists of all Alaska waters between the latitude of Cape Prince of Wales and the latitude of Point Hope. The Arctic District does not have herring district boundaries in regulation.

### SPAWNING AREAS AND TIMING

Arrival of herring on the spawning grounds is greatly influenced by climate and oceanic conditions, particularly the extent of the Bering Sea ice pack. Most herring spawning populations appear near the eastern Bering Sea coast immediately after ice breakup between mid-May and mid-June. Spawning progresses in a northerly direction and may continue into July or August along portions of the Seward Peninsula or within the Chukchi Sea.

The largest abundance of herring in the Arctic-Yukon-Kuskokwim Region is in Norton Sound District. Primary spawning areas are from Stuart Island to Tolstoi Point. When sea ice has remained in this area into June, spawning has been more extensive along Cape Denbigh and locations along the northern shore of Norton Sound between Bald Head and Bluff. Additional northerly spawning areas have been more difficult to identify because of small herring stock

sizes and limited investigations. Likely spawning areas include Imuruk Basin in Port Clarence District, and Shishmaref Inlet, Deering–Kivalik coast, and Hotham Inlet in Kotzebue District. Although subsistence herring catches have been reported in the Arctic District near Barrow, there is no information available on spawning areas.




Figure 8.—Commercial herring districts and statistical areas of Norton Sound, Port Clarence, and Kotzebue Sound.



# **NORTON SOUND PACIFIC HERRING OVERVIEW**

## **COMMERCIAL FISHERY OVERVIEW**

### **Sac Roe**

The earliest American commercial effort on Bering Sea herring apparently took place in the early part of the 1900s near Golovin in Norton Sound (Menard et al. 2013). Domestic commercial fishing resumed for “spring herring” in Norton Sound in 1964 near Unalakleet and continued sporadically until 1979. Between 1964 and 1978, the fishery averaged about 10 tons of herring annually for sac roe extraction (Menard et al. 2013). In 1979, a domestic herring fishery for sac roe began on a larger scale in Norton Sound when approximately 1,292 tons of herring were taken by 63 fishermen (13 purse seiners, 50 gillnetters). Purse seiners took 70% of the total catch.

After the 1979 season, BOF adopted a public proposal that made gillnets and beach seines the only legal commercial herring fishing gear within Norton Sound. A purse seine fishery could only be opened if the gillnet fleet could not take the allowable harvest. The regulation attempted to encourage local fishermen to participate in this developing fishery.

During the 1980 season, 294 gillnet fishermen harvested 2,452 tons of herring (Menard et al. 2013). Because gillnet fishermen demonstrated they were capable of taking the available harvest, a regulation was passed in 1981 to prohibit any purse seine gear within Norton Sound District.

Before the 1984 season, harvest by beach seine fishermen was negligible, but in 1984, 10 beach seine fishermen harvested 327 tons. In 1984, BOF set a beach seine gear limit of 100 fathoms and limited harvest to “not exceed 10% of the total herring sac roe harvest projections as published by the ADF&G.” During the fall 1987 BOF meetings, beach seine gear was further restricted to a limit of 75 fathoms. Beach seine harvests from 1985 to 2000 were only about 8% of total reported harvest, and since 1998, little market interest has existed for herring caught with beach seines because of the smaller average size of herring captured.

As with most developing fisheries, fishing effort and harvest increased with each season. In 1984, Norton Sound became a superexclusive herring fishing district to slow growth and bolster local involvement, but it had limited success. The 1987 herring sac roe gillnet harvest was 3,759 tons and had the highest level of fishing effort on record (Menard et al. 2013). This effort was more than twice the average from 1980 through 1986, yet Norton Sound area residents accounted for only about a third of both the effort and total harvest. Then, in 1987 after a public proposal adopted at the fall BOF meeting, the Commercial Fisheries Entry Commission (CFEC) changed Norton Sound Herring District to Limited Entry status with a maximum number of 301 gillnet and 4 beach seine permits. Beginning in 1988, a moratorium was placed on Norton Sound and no new entrants were allowed into the sac roe herring fishery.

No harvest occurred in 1992 due to very late ice breakup, but both the gillnet and beach seine fisheries continued with a total of more than 200 participating fishermen until 1998. The 1995 gillnet harvest of 6,033 tons was the highest on record, and the 1993 beach seine harvest of 742 tons was the largest harvest on record by this gear type. Combined dollar value for both the beach seine and gillnet fisheries peaked in 1996 at \$4.5 million (Appendix D2).

Since 1997, poor market conditions have been the primary influence on the level of commercial harvest. There has been no harvest by beach seine since 2000. Number of fishermen has

decreased from 122 in 1999 to an average of 19 for the past 5 years. From 1999 to present, the number of buyers has steadily declined from 4 to 1, with no buyers present in 2004. Even when there was a buyer, sometimes only bait was purchased, as happened in 2007–2009. One bright spot was the high recovery of over 13% roe in 2010 and 2011. In 2012 there was no sac roe fishery due to late ice breakup.

### **Spawn-on-Kelp**

A small-scale spawn-on-kelp *Fucus* sp. fishery existed in Norton Sound from 1977 to 1984. Harvests during the 1977–1984 periods ranged from less than 1 ton (1977) to approximately 47 tons (1981). During the 1984 season, 1 ton of *Macrocystis* kelp imported into Norton Sound resulted in a harvest of approximately 3 tons of product (Menard et al. 2013). In response to a public proposal, BOF closed all spawn-on-kelp fisheries in Norton Sound before the start of the 1985 season.

The 1998 herring market was known to be poor before the southernmost fisheries opened. An experimental herring spawn-on-*Macrocystis*-kelp fishery was approved by BOF to operate in Norton Sound during the 1998 season. The commissioner approved emergency regulations to allow a herring spawn-on-wild-*Fucus*-kelp fishery shortly before the normal start of the sac roe fishery. The intent of these decisions was to allow as much opportunity as possible to sac roe permit holders, because only a small minority would have an opportunity to participate in the sac roe fishery.

At the January 1999 meeting, BOF instituted a *Macrocystis* kelp open pound fishery and allowed for a wild *Fucus* spawn-on-kelp fishery for sac roe permit holders who had not sold sac roe product. Wild *Fucus* harvest is limited to an area west of Wood Point to Canal Point, including Stuart Island, and the guideline harvest level may not exceed 30 metric tons. The herring pound spawn-on-kelp guideline harvest level may not be more than 90 tons, to include combined weight of herring eggs and kelp.

Since 2001, little (less than 1 ton) or no harvest has occurred from either the *Macrocystis* kelp or wild *Fucus* spawn-on-kelp fisheries (Appendix D2).

### **Food and Bait Fishery**

Early records indicate about 3,200 tons of “fall herring” were processed in Norton Sound from 1916 to 1941 (Menard et al. 2013). This fishery, dependent on salt curing, declined because foreign competition produced poor marketing conditions. Japan began gillnetting in Norton Sound during 1968 with 3 vessels. Effort was concentrated about 12 miles offshore between St. Michael and Golovin. Approximately 40 Japanese vessels reported harvesting a record 1,400 tons of herring during 1969 (Menard et al. 2013). An average annual harvest of approximately 450 tons was reported in Norton Sound by the Japanese during 1968–1974. All foreign fleets were prohibited in 1977 from gillnet fishing in the area.

Since 1977, there has not been a consistent domestic commercial food and bait herring fishery in Norton Sound. The majority of reported food and bait herring harvest estimates were initially harvested as sac roe but bought and processed as food and bait, so they were considered food and bait for the purposes of this report. The largest Norton Sound herring harvest in the past 50 years occurred in 1995 when an estimated 6,763 tons of sac roe herring were delivered, of which only 116 tons were purchased as food and bait. Since 1997, no more than 91 tons of herring were sold annually as food and bait (Appendix D1).

## COMMERCIAL FISHERY MANAGEMENT

The overall statewide management strategy is based upon the *Bering Sea Herring Fishery Management Plan* (5 AAC 27.060) to annually harvest 0–20% of the herring biomass. The upper end of the exploitation range is applied to stocks in good condition. The lower end of the exploitation range is applied to stocks exhibiting a trend of decreasing abundance and poor recruitment. If a minimum biomass threshold level of 7,000 tons for Norton Sound is not achieved, no commercial fishery will be allowed.

Typically, herring are long-lived fish and will usually remain harvestable for at least 5 years after recruiting into the fishery. Harvesting only a percentage of the biomass ensures fish will remain for following years. This type of strategy helps mitigate population fluctuations caused by successive years of poor recruitment, a common occurrence in marine-spawning fish. Before 1983, harvests in Norton Sound were regulated by subdistrict so harvests would be dispersed over the entire fishing grounds (Menard et al. 2013). This strategy prevented harvest efforts from concentrating in one area, on what was then thought to be a distinct stock of fish.

Methods to reliably forecast herring returns are still being developed and estimates of recruitment are not available; therefore, inseason assessments of biomass supersede projected biomass for management of Norton Sound herring. The herring fishery is managed for a 20% exploitation rate at biomass levels twice minimum threshold or greater. If the run does not materialize as projected, the harvest exploitation rate may be reduced to a lower level.

Generally, fisheries management staff has tried to set commercial openings to allow gillnetters to fish flood tides as they crest. Figured heavily in this strategy is the belief that ripe females approach the beach at that time to spawn. Because the Norton Sound fishery covers a large area with varying tides, opening at the optimal time throughout the district is not always possible. The fishing fleet must be flexible to maximize catches and roe quality. However, since 1997 there have been limited markets for herring and the catch has been well below the guideline harvest level. Since 2002, to maximize efficiency for fishermen and buyers, ADF&G has opened the fishery continuously once buyers are ready and then buyers direct the fleet when to set and pull nets.

In the past, duration of beach seine openings was dependent on herring abundance near the beach and favorable weather conditions for spotters and fishing. Beach seiners prefer to work flood tides similar to gillnetters; however, fisheries managers frequently provided less optimal fishing times. Beach seiners are able to harvest their allotment of 10% of the preseason harvest goal in a single 3-hour opening under ideal conditions. By nature of the gear, beach seiners have the potential to wrap up large numbers of fish that could potentially exceed their allocation. In the past, management staff often reduced beach seine efficiency by allowing a gillnet opening to occur before a beach seine opening. This opening breaks up school size and reduces likelihood of excessive harvests. Occasionally, the beach seine fleet has been used to test roe quality of herring newly arrived in nearshore waters before a gillnet opening. The potential for waste would have been great had the entire gillnet fleet fished on poor quality herring.

In the 2000s, the market desired a higher roe percent and larger size fish. These criteria have been difficult to achieve with beach seine gear and therefore no buyer interest has existed for herring harvested from beach seines.

## SUBSISTENCE FISHERY USE

Pacific herring were used for subsistence purposes by coastal residents well before the mid-1800s when their use was first documented by early explorers. Subsistence harvest of herring and herring roe on kelp is not documented but is believed to be relatively small. It is also known that St. Michael and Stebbins residents harvest spawn-on-kelp roe for subsistence use.

## PORT CLARENCE AND KOTZEBUE PACIFIC HERRING OVERVIEW

### COMMERCIAL FISHERY OVERVIEW

Port Clarence and Kotzebue commercial herring fisheries have been in regulation since 1982. In Port Clarence and Kotzebue Districts, regulations state that herring may be taken from April 15 through November 15, except that herring may not be taken during the open commercial salmon fishing season. The 1983 and 1984 regulations set a guideline harvest of 150 metric tons (165 tons) for each subdistrict, which is still in effect. Presently, purse seines, beach seines, and gillnets are legal commercial gear within these districts.

Before 1987, no spring sac roe commercial fisheries had ever occurred within these districts. In 1987 and 1988 a spring sac roe herring fishery was attempted in the Port Clarence District. A fish buyer located in Nome in 1994 and 1995 provided a ready crab bait market and transportation for fish, which facilitated a spring harvest. However, no one has fished for bait since 1996 (Appendix D4).

Regulations allow spawn-on-kelp fisheries in Port Clarence and Kotzebue. Attempts at open pound *Macrocystis* harvest in Port Clarence District in 1991 and 1992 were unsuccessful.

### HISTORICAL RESOURCE INVESTIGATIONS

Resource investigations of Port Clarence and Kotzebue Sound area herring stocks were conducted by ADF&G from March 1976 to September 1978 (Barton 1978). These studies indicated herring populations from Golovnin Bay (Norton Sound) northward differed significantly in size and behavioral characteristics from herring populations occurring in the southern Bering Sea. Differences between populations were summarized as follows (Barton 1978):

Seward Peninsula Populations	Southern Norton Sound to Southern Bering Sea Pelagic Populations
Smaller herring at age with lower vertebral counts.	Larger herring with probable higher vertebral counts.
Lower abundance.	Higher abundance.
Subtidal spawning (3m) in shallow bays, inlets and lagoons.	Intertidal and shallow subtidal spawning along exposed rocky headlands.
<i>Zostera sp.</i> primary spawning substrate.	<i>Fucus sp.</i> primary spawning substrate.
More euryhaline.	Less euryhaline.
Over winter in shallow bays; water is warmed by river discharge under ice cover.	Over winter in deep ocean layers near the Pribilof Islands.
Fall (non-spawning) runs documented.	No fall runs documented.
Larval development in brackish water.	Larval development probable in more saline water.

Data collected from herring populations along the Seward Peninsula strongly indicated that a separate stock of herring occurs in Port Clarence and Kotzebue Sound Districts. These data do not preclude possibility of more southern stocks utilizing this region, such as stocks that winter near the Pribilof Islands and migrate to the western Alaska coast to spawn. Migration to central Bering Sea for wintering herring stocks along the western Seward Peninsula is unlikely; rather they might remain in coastal lagoons, bays, or inlets that are warmed by river discharge under the ice (Barton 1978). Size difference may be explained by warmer water temperatures from river discharge. Water temperatures and feeding conditions in deep ocean waters are probably more favorable for growth than those in herring winter habitats along the Seward Peninsula, where apparently they have become adapted to Arctic conditions (Barton 1978).

Aerial surveys are difficult in Port Clarence District because of organic coloring of waters of Imuruk Basin, Tuksuk Channel, Grantley Harbor, and, to a lesser extent, Port Clarence. Presence of other species of fish caught in test commercial gear sets indicate the need for verifying species composition of any biomass sighted. A further complicating factor within Port Clarence is spring ice conditions. Port Clarence is a sheltered body of water that becomes highly stained over winter and takes time to clear once ice melts. Typically, outside waters are significantly warmer than inside waters, which are covered by ice longer, thereby slowing solar gain and water mixing. Soon after ice begins to shift, herring move into the warm shallow lagoons to spawn. Herring are invisible to aerial observation once they enter stained water. The best aerial survey conditions exist just outside the entrance to Port Clarence, where herring mass just before the ice moves. In each of the past several years one or two surveys were flown, but virtually no herring were observed because the narrow window of time for seeing fish was missed.

## **KING CRAB OVERVIEW**

### **NORTON SOUND KING CRAB OVERVIEW**

#### **District Boundaries**

Norton Sound Section (Q3) consists of all waters in Registration Area Q north of the latitude of Cape Romanzof (61 degrees 49 minutes N latitude), east of the International Dateline, and south of 66 degrees N latitude (Figure 9).

#### **Abundance**

Since 1998, a length-based population model has been used to predict biomass for the red king crab population in Norton Sound (Zheng et al. 1998). Incorporating data from trawl surveys (Appendices E2 and E13–E14), historical winter and summer pot studies, and winter and summer fisheries (Appendices E15–E20), the model is used to project abundance estimates of legal male crab even in years when no trawl survey occurs, allowing abundance-based management of the summer commercial crab fishery. Every time new data are incorporated into the population model, the model both estimates current abundance and revises prior years' abundances. It should be noted that estimates prior to 1996 are currently under review because survey extrapolation methodologies changed after that point, and previous biomass estimates may be revised and incorporated into the model as a result of this assessment.




Figure 9.—King crab fishing districts and sections of Statistical Area Q.

The following estimates are based on the model's results from spring of 2013 with the latest data from the 2011 trawl survey, the 2012 summer fishery, and the 2011–2012 winter study. In 2008, legal biomass estimate for the summer crab fishery was 3.35 million lb, an increase of 13% from the 2.96 million lb estimated for 2007. The legal population estimate increased for the following 3 years: to 3.71 million lb in 2009, 4.24 million lb in 2010, and 4.43 million lb in 2011. From 2011 to 2013, the legal abundance estimate decreased, to 4.22 million lb in 2012 and to 4.13 million lb in 2013 (NPFMC 2013).

The latest winter study data indicate that the legal proportion of the catch increased from 2007 to 2010, changed little from 2010 to 2011, and decreased in 2012 (Appendix E8). However, the record high overall male catch per unit of effort (CPUE) seen in 2012 and relatively stable legal CPUE from 2008 to 2011 suggests that the decline in 2012 in proportion of legal crab was caused more by an increase in prerecruit abundance than a decline in legal crab abundance. Because prerecruit-1 crab require 1 molt to become part of legal population the following year but prerecruit-2 crab require 2 molts, the above-average proportion of prerecruit-2 male crab in 2012 indicates a possible recruitment surge in 2014. No winter study took place in 2013 because ADF&G did an expanded spring and summer tagging study.

## COMMERCIAL FISHERY OVERVIEW

The last year that a large-vessel summer commercial crab fishery existed in Norton Sound Section was in 1990. No summer commercial fishery occurred in 1991 because of ADF&G staff constraints. In 1992, the summer commercial fishery resumed. Appendix E1 shows historical summer commercial harvest by year and statistical area for Norton Sound crab fishery since 1990. For historical information before 1990, please refer to the 2012 Annual Management Report (Menard et al. 2013). Regulation changes adopted during the March 1993 BOF meeting changed participation in the fishery to that of small boats. A superexclusive designation went into effect for the Norton Sound commercial crab fishery June 27, 1994. This designation stated that a vessel registered for the Norton Sound crab fishery may not be used to take king crab in any other registration area during that registration year. Later, a vessel moratorium put into place before the 1996 season was intended to precede a license limitation program. Community Development Quota (CDQ) groups were allocated a portion of the summer harvest beginning in 1998, but no CDQ harvest occurred until the 2000 season. The North Pacific License Limitation Program (LLP) went into effect for the Norton Sound crab fishery January 1, 2000. The program states a vessel which exceeds 32 feet in length overall must hold a valid crab license issued under LLP by National Marine Fisheries Service. Regulation changes and location of buyers resulted in harvest distribution moving eastward in Norton Sound in the mid-1990s (Appendix E12).

During the March 1999 BOF meeting a new management strategy was enacted for the Norton Sound summer red king crab fishery. A threshold level of abundance of legal male red king crab biomass was set at 1.5 million lb. A summer commercial season may only open if the legal crab biomass is estimated to be at least 1.5 million lb, and if the legal biomass falls in the range of 1.5 to 2.5 million lb the harvest rate will be no more than 5% so the stock may rebuild. If legal biomass is 2.5 million lb or more, the harvest rate will be no more than 10%. In March of 2012, this regulation was modified by the BOF so that the new threshold level of abundance of legal male red king crab biomass was set at 1.25 million lb. If the estimated legal crab biomass falls within the range of 1.25 to 2.0 million lb, the harvest rate will be no more than 7% of legal male abundance. From 2.0 to 3.0 million lb, the harvest rate will be no more than 13%. If the estimated legal biomass is more than 3.0 million lb, the harvest rate will be no more than 15%. Improved abundance estimates and the current management strategy will greatly reduce the risks of overfishing the stock.

Since 1981, in order to protect crab utilized by the inshore subsistence fishery from commercial harvest, an area delineated by a line approximately 10 to 15 miles off the shores of southern Seward Peninsula from Port Clarence to St. Michael has been closed to the summer commercial fishery. This closure line has relaxed over the years to its current position adopted by the BOF in 2002 (Appendix E11).

To reduce handling mortality of undersized crab and smaller female crab, the BOF at its March 2008 meeting put a new regulation into effect: a minimum of 4 escapement rings are required per pot with each ring having a minimum inside diameter of 4.5 in located within 1 mesh size from the bottom of the pot, or at least one-half of the vertical surface of a square pot or sloping side-wall surface of a conical or pyramid pot must be composed of no less than 6.5 in stretched mesh. Also starting with the 2008 season, even though the minimum legal size of red king crab is 4.75 inches in carapace width (CW), the local seafood plant did not always buy crab less than 5.0 in CW. The Anchorage buyer, however, has continued to buy crab as long as they are of legal size.

In 2010, due to concern over lack of stock status information, the North Pacific Fisheries Management Council closed the area above Cape Prince of Wales to crabbing. Only state waters (within 3 miles of shore) will be open to crabbing north of the latitude of Cape Prince of Wales (Appendix E11).

### **CDQ Fishery**

NSEDC and Yukon Delta Fisheries Development Association (YDFDA) divide the CDQ allocation. Only fishermen designated by these 2 CDQ groups are allowed to participate in this portion of the king crab fishery. Fishermen were required to have a CDQ fishing permit from CFEC and register their vessel with ADF&G before they made their first delivery. Fishermen operated under authority of the CDQ group and each CDQ group decided how their crab quota was harvested.

During the March 2002 BOF meeting, new regulations were adopted that affected the CDQ crab fishery and relaxed closed-water boundaries in eastern Norton Sound and waters west of Sledge Island. Closed-water boundaries are illustrated in Appendix E11. The Norton Sound CDQ fishery may begin at noon on June 15, or no less than 72 hours after commercial gillnet or beach seine herring fishing is closed, whichever is later, through noon on June 28. After July 1, the commissioner may, by emergency order, open a CDQ fishery for any remaining allocation after closure of the open access fishery. At the March 2008 BOF meeting the regulation requiring the herring fishery to be closed was repealed, and the CDQ fishery was allowed to occur by emergency order before, during, or after the open access fishery. Previously, the open access fishery started on July 1, but BOF passed a regulation allowing ADF&G to open the fishery by emergency order anytime beginning on or after June 15.

### **Commercial Catch Sampling**

The Norton Sound red king crab commercial fishery had the benefit of an onboard observer during the 2000 and 2001 seasons because there was a floating processor on the fishing grounds in those years. In years with no onboard observer, a smaller percentage of crab from the commercial harvest is sampled because fishermen deliver at all times of the day and night. The new seafood processing plant, Norton Sound Seafood Products (NSSP), began operating in Nome in summer 2002, greatly improving the ability of Nome ADF&G staff to sample crab brought to the Nome dock. Crab was either sampled at NSSP or at the small boat harbor where nonresident fishermen or catcher-processors not selling to NSSP offload their catch for delivery to Anchorage. ADF&G will continue to make a concerted effort to coordinate catch sampling with fishermen and buyers to ensure optimal commercial harvest data collection.

## **SUBSISTENCE FISHERY OVERVIEW**

Norton Sound residents utilize red king crab for subsistence, mainly during winter. Fishing occurs through cracks or holes cut in the ice with the use of hand lines and pots. To document trends in subsistence harvest, BOF enacted a regulation in 1977 requiring subsistence fishermen in Norton Sound to obtain a permit before fishing. Fishermen record their daily effort and catch on these permits.

For catch information before 1990, please refer to the 2012 Annual Management Report (Menard et al. 2013). Since 1990, the winter subsistence crab fishery harvest has ranged from a low of 256 crab during the 2000–2001 season to a high of 12,152 crab during the 1989–1990 season (Appendix E5). Lack of success in the winter crab fishery during some years has been attributed to



a declining crab population caused by removal of crab in the summer commercial fishery, together with low recruitment, low effort caused by poor ice conditions, and changes in nearshore winter distribution of crab. All these factors in varying degrees affect success of the winter fishery, as well as increased use of more efficient gear (pots instead of hand lines). Unstable ice conditions and record snowfalls adversely affected: 1992–1993, 1996–1997, 2000–2001, 2003–2004, and 2005–2006 catches. During years of stable ice conditions, approximately 85 fishermen averaged 80 crabs each.

## **ST. LAWRENCE ISLAND AND KOTZEBUE KING CRAB OVERVIEW**

### **District Boundaries**

Formerly, St. Lawrence Island Section was located immediately west and north of Norton Sound Section, but in May of 2006, BOF expanded Norton Sound Section to include the St. Lawrence Island Section south of 66 degrees N latitude and west of 168 degrees W longitude (Figure 9). The former St. Lawrence Island Section north of 66 degrees N latitude is now the Kotzebue Section.

### **Abundance**

Unlike Norton Sound, the area of the Bering Strait that includes St. Lawrence Island has never been surveyed consistently by ADF&G. Even though commercial and subsistence harvests are allowed by regulation, ADF&G does not have abundance estimates for this area. In summer of 2005, an exploratory pot survey was conducted by NSEDC in cooperation with ADF&G to assess the number and distribution of male blue king crab in the vicinity of King Island, Wales, and Port Clarence. The survey was only partially successful due to strong currents that made pot retrieval difficult when set deeper than 10 fathoms. Shallow pot placement resulted in a catch primarily of egg-bearing female blue king crab, and indicated that using standard Norton Sound crab gear would only access a nursery site for gravid blue king crab. When more suitable gear becomes available, further surveys will be necessary to determine the feasibility of a summer fishery. At the March 2008 BOF meeting, legal size requirement for blue king crab was changed from 5.5 to 5.0 in. Preliminary data indicate that blue king crab size at maturity is very similar to Norton Sound red king crab.

In summer of 2006, 2008, and 2011, the Northern Bering Sea trawl survey was conducted by NSEDC in cooperation with ADF&G to assess crab resources in the St Lawrence Island and Bering Strait areas of Norton Sound District. The primary focus was to collect information on blue king crab size, distribution, and abundance. The area surveyed lies west and northwest of the standard ADF&G triennial Norton Sound red king crab trawl survey locations. In 2006, trawls were conducted from near the southwest corner of St Lawrence Island to the Bering Strait area southwest of Cape Prince of Wales. Size information and general distribution of blue king crab was collected. In 2008 prior to the trawl survey, a camera sled was towed a few meters above the seabed to observe crab and other species in the St. Lawrence Island area that had been trawled in 2006. The 2008 and 2011 trawl work was focused on looking at the distribution of blue and red king crab in the area between Port Clarence and King Island. More survey work is necessary to generate an abundance estimate and to better understand the distribution of blue king crab. The 2006, 2008, and 2011 survey data should only be considered a starting point to understanding the Bering Strait and St. Lawrence Island blue king crab stock.

## Commercial Fishery Overview

In 1984, a regulation was adopted to close waters within 10 miles of all inhabited islands within the St. Lawrence Island Section (St. Lawrence Island, Little Diomed, and King Island). This regulation attempts to protect stocks targeted by local fishermen and reduce impacts on marine mammal subsistence harvests. Since 1990, commercial catches in the former St. Lawrence Island Section have only been reported for 4 years. In 1992, 53 lb of blue king crab were landed. In 1995, 7,913 lb of blue king crab were delivered from 3 landings (Bue et al. 1997). In 2005, 316 lb of red king crab were harvested in the Kotzebue area, and in 2006, 340 lb were harvested<sup>1</sup>.

Villagers of Little Diomed and St. Lawrence Island have bartered with and sold winter-caught blue king crab to residents of Nome and other villages for years. ADF&G does not have an accurate estimate of the magnitude of this trade. Remoteness of the villages contributes to lack of catch records. Current regulations allow a commercial harvest and sale of king crab caught near shore during winter. However, local residents have decided not to export any of their winter catch for commercial sale.

## MISCELLANEOUS FISH OVERVIEW

Several species other than salmon, crab, and herring are utilized for commercial and subsistence purposes in Norton Sound, Port Clarence, Kotzebue, and Arctic Districts (Appendix G1). Primary species include inconnu or “sheefish” *Stenodus leucichthys*, Dolly Varden *Salvelinus malma*, whitefish (*Coregonus laurettae*, *C. pidschian*, *C. sardinella*, *C. nasus*, and *Prosopium cylindraceum*), *Coregonus* sp., *Prosopium* sp., and saffron cod *Eleginus gracilis*.

These fish are taken by set gillnets, beach seines, “jigging” through the ice, and rod and reel. Subsistence catches taken during summer months are normally air dried, and winter catches are stored frozen. Fish are utilized for human consumption and for dog food. Fish taken for commercial purposes are mainly sold locally, although some are shipped out of the area.

Subsistence harvest of most species is not limited by regulation. Commercial harvest may be prohibited in some freshwater areas, but limited commercial endeavors are allowed in many areas under terms of a permit.

## INCONNU (SHEEFISH)

### Spawning Areas and Timing

Sheefish are distributed throughout nearshore estuarine areas of Kotzebue Sound, with the largest spawning stocks and harvests in the Kobuk–Selawik River drainages and Hotham Inlet. However, there is a small population in the Sheshalik and Krusenstern areas of northern Kotzebue Sound and in the Koyuk River of Norton Bay in Norton Sound (Figure 10).

---

<sup>1</sup> Statewide electronic fish ticket database. 1<sup>st</sup> edition. Alaska Department of Fish and Game, Division of Commercial Fisheries. 1985 to present. [URL is not publically available because some information is confidential.] Hereafter referenced as “fish ticket database.”



Figure 10.—Kotzebue and Kobuk River Valley villages and their spatial relationship with inconnu spawning and overwintering areas.

Inconnu's spawning and overwintering migration behavior makes them available for harvest by various fisheries throughout their life cycle, yet increases their vulnerability to overharvest. Although inconnu are capable of consecutive spawning, most spawn every 2 to 3 years, and slow maturation rates of 5–7 years for males and 7–11 for females increase the time required to restore depleted populations. Sheefish have high fecundity, and large females can carry over 400,000 eggs. Such populations may be subject to episodic recruitment events depending on environmental conditions. If spawner abundance is maintained above a threshold level, intermittent years of good recruitment can carry the population through years of less favorable ice conditions.

After ice breakup in Kotzebue Sound area, adult sheefish migrate upriver to spawning areas on the Kobuk and Selawik rivers. On the Kobuk River, spawning occurs upstream from the village of Kobuk, with the greatest concentration observed between the Mauneluk and Beaver rivers. Then, when spawning is complete in late September and early October, sheefish disperse downstream to overwintering areas within Hotham Inlet/Selawik Lake.

### Historical Fishery Use

During the 1960s, age, sex, and length data indicated inconnu stocks were overharvested by commercial and subsistence fisheries in Kotzebue District. Consequently, an annual area commercial harvest quota of 25,000 lb was instituted, but subsistence is given priority and has remained unrestricted.

## **Subsistence Fishery**

Inconnu have long been utilized for subsistence purposes throughout Kotzebue basin, especially in Kotzebue, Selawik, and the villages along the Kobuk River. In 2004, an estimated 10,163 sheefish were harvested, surpassing the previous record since 1971 estimated at 9,805 in 1997, and 7,823 in 2003 (Appendix F2). These harvests may include winter, summer, and fall catches. Due to budget constraints, the Division of Subsistence did not survey the villages in Kotzebue Sound District for subsistence sheefish harvests from 1988 to 1990, and from 2004 to 2011. Due to limited survey effort during many years, total catch and effort should be regarded as minimum numbers and are not comparable year to year. Subsistence inconnu harvest information was not always collected for the town of Kotzebue, where a sizable ice fishery occurs for sheefish in late winter and spring. In 2012 and 2013, there were comprehensive subsistence surveys for fish and wildlife harvests of 6 Kotzebue area villages, but data is not yet available for 2013.

Summer and fall subsistence fishing for inconnu occur along Kobuk and Selawik rivers from June through October with gillnets, beach seines, and rod and reel. In spring, residents of Kotzebue, Noorvik, and Selawik harvest inconnu with hand jigs through the ice of Hotham Inlet and Selawik Lake. In early winter, Kotzebue, Noorvik, and Selawik fishermen use gillnets set under the ice in Hotham Inlet and Selawik Lake. No requirement exists for harvest reporting; however, during various years from 1973 to 2004 and starting again in 2012, ADF&G Division of Subsistence conducted household subsistence harvest surveys in various villages in Kotzebue District.

In 1987, BOF adopted a regulation limiting size of gillnets used to take inconnu for subsistence to be not more than 50 fathoms in aggregate length or 12 meshes in depth, nor have a mesh size larger than 7.0 in (5 AAC 01.120). This regulation was intended to conserve the larger, breeding portion of the stock. Except for this gear restriction, ADF&G does not restrict timing, area, or quantity of subsistence inconnu harvest.

## **Commercial Fishery**

Most commercial fishing effort occurs through the ice in Hatham Inlet, near Kotzebue, using gillnets from 5.5 in to 7.0 in stretched mesh. Recorded commercial catches are relatively small, but undocumented catches may be significant. Therefore, harvest totals should be considered minimum estimates. Restricted markets outside northwestern Alaska greatly limit commercial activity; however, most individuals participating in the winter commercial fishery also fish for subsistence purposes. Inconnu incidentally caught in the commercial salmon fisheries are sold in years when there is a market, but only in small amounts. Reported harvest and effort in the commercial fishery have declined in the last 15 years. Since 1998, harvest has not exceeded 1,250 lb, compared to the highest harvest of 8,224 lb in the last 23 years (Appendix F1). There were no reported commercial sheefish catches from 2006–2011 and 2013, but fishermen participated in commercial fishing in 2005 and 2012. Although sheefish were probably harvested and sold in the winter of 2011–2012 by several fishermen, only 1 fish ticket from the Kotzebue Sound District was submitted to ADF&G, making that catch information confidential.

## **Sport Fishery**

Kotzebue District sheefish are considered by many to be among the pinnacle of Alaska freshwater sport fishing due to their large size. Since the start of the ADF&G Trophy Fish

Program in 1967, all but one qualifying sheefish came from the Kobuk River. In spite of this, the level of sport fishing effort is still quite low.

Residents of Kobuk River villages have expressed concern over sport fish practices near spawning grounds on the upper Kobuk River. Catch-and-release fishing is considered by some local residents to be disrespectful and damaging to sheefish. Also, the practice of discarding filleted carcasses in the water is thought to drive other sheefish away from the area. In 1986, the Division of Subsistence investigated these concerns and found the concerns could be addressed if sport anglers were more aware of local customs and culture. An educational brochure is now available to fishermen on upper Kobuk River in the hope that proper handling during catch-and-release can minimize impacts on spawning populations. Although overall harvests are substantial, populations appear to be healthy, spawner abundances are increasing, and sport harvests are relatively low (Scanlon 2009). Sheefish sport harvests in the last 10 years have averaged approximately 850 annually (Appendix F3).

### **Historical Escapement**

Historically, aerial surveys were conducted on key inconnu spawning areas incidental to effort of enumerating salmon. These surveys were primarily conducted along upper Kobuk River in September. Survey conditions historically result in either very few or no inconnu being observed. During these surveys, species identification has been a problem. Surveys were not conducted from 1984 through 1990 because of high and/or turbid water, poor weather conditions, or lack of personnel. Through the early 1990s, incomplete escapement and catch data provided little basis for assessing current population status of inconnu in Kotzebue District, but some local residents were concerned that the inconnu stocks were declining.

Because of these concerns, a cooperative tagging project on inconnu in Kotzebue District began in 1994. This study was conducted by Division of Sport Fish, U.S. Fish & Wildlife Service (USFWS), and National Park Service. Spawning inconnu were tagged in Upper Kobuk River and Selawik River. Roughly 600 sheefish were tagged in Kobuk River by Division of Sport Fish and 150 in Selawik River by USFWS in 1994. During the fall of 1995, roughly 617 inconnu were tagged in Upper Selawik River and approximately 1,386 were tagged in Upper Kobuk River. In 1996, 2,300 were tagged in Upper Kobuk and 500 in Selawik River. The Selawik River project ended in 1996. In 1997, 1,757 inconnu were tagged in Upper Kobuk River. Spawning population estimates of inconnu in Upper Kobuk River were 32,273 in 1995, 43,036 in 1996, and 26,800 in 1997. Inconnu spawn upstream of the village of Kobuk; the greatest observed concentrations were between Meneluk and Beaver rivers. After spawning is complete in late September, fish disperse to downstream overwintering areas. In Selawik River, the spawning population estimate was 5,200 and 5,300 for 1995 and 1996, respectively. Tag recoveries showed that these stocks mixed in Hotham Inlet winter habitats but maintained fidelity to their spawning areas (DeCicco 2001).

### **DOLLY VARDEN**

Dolly Varden are distributed throughout Norton Sound, Port Clarence, Kotzebue, and Arctic Districts. Although taxonomists have disagreed on distinguishing Dolly Varden characteristics and distribution of Arctic char and Dolly Varden, most now agree char in this area are the northern form of Dolly Varden. To eliminate confusion, in this report these fish are referred to as Dolly Varden, the common name for this species complex; however, locally they are called trout.

## **Spawning Areas and Timing**

Dolly Varden in northwest Alaska are primarily nonconsecutive spawners. They spawn throughout late summer and fall in almost all drainages of Norton Sound, some northern Seward Peninsula rivers, and the major drainages of Kotzebue Sound and Chukchi Sea. Fry emerge in spring and migrate to the ocean during early summer after spending from 1 to 6 (generally 2–5) years in freshwater. Movements of Norton Sound Dolly Varden coincide with salmon. In spring, Dolly Varden are likely to remain longer in streams following a large pink salmon run to feed on abundant outmigrating fry. Also, they are sometimes present in streams during summer to feed on salmon eggs, especially during years of high pink salmon abundance.

Because Dolly Varden are a late-maturing fish (generally age 6–7), they are susceptible to overfishing by commercial, subsistence, and/or sport fisheries. Consequently, commercial fisheries have been maintained at low levels or prohibited to both reduce potential overharvest and provide for reproductive needs and subsistence uses.

## **Subsistence Fishery**

Dolly Varden is an important component in the diet of subsistence users in Norton Sound–Kotzebue Sound and Arctic areas. In some communities, they outrank salmon and whitefish in importance to subsistence; however, most villagers in Norton Sound District report Dolly Varden as incidental catches in subsistence salmon nets. Subsistence fishermen harvest Dolly Varden with seines in fall, hook and line through ice in winter, and gillnets in spring. The fall seine fishery contributes the greatest number of fish to annual subsistence Dolly Varden harvest.

In Kotzebue District, fall seine fishing is a group effort with several households making up a fishing group. Catch is stored and allowed to freeze in willow cribs located near the seining site. These fish are used throughout the winter by the fishing group. Most Dolly Varden harvests take place before or just after freeze-up. Fishermen from Noatak usually fish before freeze-up, but residents of Kobuk River villages of Shungnak and Noorvik fish for Dolly Varden throughout the winter. Since 1991, subsistence catch of Dolly Varden in Noatak has ranged from almost 3,000 to over 11,000 fish (Appendix F5), but they should be considered minimal figures because of survey timing. Except for 2007, no Dolly Varden harvest surveys have been conducted of Kivalina residents during the last 22 years. In 2012 and 2013, a comprehensive survey of fish harvests was done in Kobuk River villages and Noatak by the Division of Subsistence, but data are not yet available for 2013.

In Arctic District, fishery harvest studies by ADF&G's Division of Subsistence noted that annual community catches of Dolly Varden in Kaktovik (Pedersen and Linn 2005) and Anaktuvuk Pass (Pedersen and Hugo 2005) produced annual catches of “char” (a mix of Arctic char and Dolly Varden).

## **Commercial Fishery**

Dolly Varden generally appear in commercial catches during the last 3 weeks of August and are taken as a nontarget species in the Kotzebue Sound commercial chum salmon fishery. In 1976, regulations closed the commercial chum salmon fishery on August 31 and thus reduced harvest of Dolly Varden. Spawning and overwintering Dolly Varden typically pass through the area during September but typically begin migration along the northern shore of Kotzebue Sound during the third week of August. Reported Dolly Varden sales are dependent upon available markets. The typical season catch, when buyers are purchasing Dolly Varden throughout August,

is approximately 1,000 to 3,000 fish (Appendix F4). However, limited markets in the 2000s have resulted in less than 200 Dolly Varden reported sold each year in Kotzebue Sound, and zero sold since 2005 because the buyer no longer purchases Dolly Varden. Regardless of sales, Dolly Varden catches are still required to be reported on fish tickets. During the 2011–2012 season, three fishermen caught and sold 1,057 lb of Dolly Varden to the fish plant in Nome as bait. This was the first recorded sale of Dolly Varden in Norton Sound in recent history.

## **Sport Fishery**

Drainages of Kotzebue Sound and the Chukchi Sea are known for the large size of anadromous Dolly Varden, but Kotzebue area residents and non-locals boating on Kobuk and Noatak rivers are the primary participants in this area's Dolly Varden sport fishery. Both Noatak and Kobuk rivers are National Wild and Scenic rivers with headwaters included in Gates of the Arctic National Park. However, the Wulik River is probably the most important Dolly Varden stream in northwestern Alaska. The 90-mile Wulik River is known for the largest and most abundant Dolly Varden populations. Located approximately 90 miles north of Kotzebue, Wulik River flows into the Chukchi Sea through Kivalina Lagoon near the village of Kivalina and is estimated to have over 100,000 overwintering Dolly Varden annually.

Since the start of the ADF&G Trophy Fish Program in 1967, 140 of 219 qualifying Dolly Varden have come from Kotzebue Sound and Chukchi Sea drainages. Additionally, the current Alaska sport fish angling record for Dolly Varden was 12.4 kg (27 lb 4 oz), taken from the Wulik River in 2002 and surpassing the previous record (also taken from the Wulik River in 2000). In spite of this, sport fishing effort has been consistently low, which is probably due to the remote location and difficult access of fishing sites (Scanlon 2009). Dolly Varden sport fish harvests in the last 10 years in Norton Sound averaged almost 3,200 annually but averaged less than 1,200 in the Kotzebue/Chukchi Sea areas (Appendix F3).

## **Historical Escapement**

Since 1990, aerial survey counts of overwintering Dolly Varden on the Wulik River has ranged from 135,135 fish in 1992 to 1,500 fish in 2003 (Appendix F7). Weather and water conditions have precluded flying aerial surveys during many years. Weather permitting, Division of Sport Fish conducts aerial surveys of Noatak River spawning grounds in summer, and Kivalina and Wulik rivers overwintering areas in fall. Since 2000, however, only Wulik River has been surveyed.

## **WHITEFISH**

Although inconnu belong to the whitefish family, this section deals with several smaller species of genera *Coregonus* and *Prosopium*. Genus *Coregonus* contains “broad” and “humpback” whitefish or *C. nasus* and *C. pidschian*, respectively. In addition, 3 whitefish species known as “ciscoes” belong to these genera: least cisco *C. sardinella*, Arctic cisco *C. autumnalis*, and Bering cisco *C. laurettae*. “Round” whitefish *Prosopium cylindraceus* are the sole representatives of genus *Prosopium* in this area.

## **Spawning Areas and Timing**

Whitefish occur throughout most bodies of fresh water in Norton Sound, Port Clarence, Kotzebue, and Arctic districts and can also be found at various times of year in inshore marine

waters. Several whitefish species spawn in freshwater in late August to October when lakes and streams are close to freezing.

### **Commercial Fishery**

Limited commercial whitefish harvests have been allowed since statehood, normally under auspices of a permit that delineates harvest levels, open areas, legal gear, etc. Commercial whitefish fisheries were generally limited to large open-water areas (e.g., Grantley Harbor in Port Clarence District) or ocean waters. Beach seines were stipulated as legal gear in some instances in order to reduce the number of incidental species taken. Little comparative commercial catch and effort data were recorded, but harvest levels were historically low. Most commercial catches were made in Golovnin Bay in Norton Sound District, in Kuzitrin River in Port Clarence District, and in Hotham Inlet and Selawik River in Kotzebue District. Fish were sold to local markets for human consumption, dog food, or, more recently, crab bait. During the 2006–2007 season, one local Nome fisherman, who waived confidentiality, sold 3,723 lb of whitefish. No further whitefish harvests occurred until the 2010–2011 season, and since then only one or two fishermen have harvested whitefish commercially for no more than 2,200 lb total (Appendix F9).

In the Arctic District, a commercial fishery for freshwater finfish has existed in the Colville River delta (located approximately 60 miles west of Prudhoe Bay) since 1964 (Menard et al. 2013). Historically, commercial fishing generally took place during late June and July for broad and humpback whitefish and October through early December for Arctic and least cisco. However, since 1990 commercial fishing effort has predominantly occurred in October and November for Arctic and least cisco. Set gillnets are used as capture gear, and fishing during fall months occurs under the ice. All fish were harvested with the intent to sell commercially and are reported daily on a catch form. However, not all fish reported on permits for this area were sold. Those fish not commercially sold were retained and used for subsistence purposes. No commercial harvest has been reported since 2007 from the Coville River (Appendix H1).

### **Subsistence Fishery**

Whitefish are important for subsistence use and taken mainly by beach seine or set gillnets. Catches are usually dried and used for human consumption or dog food. In some areas, fish are “gutted” and dried early in summer, but later in summer, fish are filleted and dried with eggs and viscera intact.

Subsistence catch enumeration is difficult because fishermen do not count fish individually, but by “tubs,” “bags,” “strings,” or other estimators of gross abundance. Additionally, many fish are dried and consumed or stored in caches before the survey period. Reported subsistence harvests were generally the result of a limited and sporadic survey effort and should be regarded as minimum values and not comparable from year to year. In 1997, subsistence harvests of whitefish were included for the first time in Division of Subsistence household salmon harvest surveys in Kotzebue Sound villages (Appendix F8).

The relative importance of whitefish is higher in Kotzebue Sound District than in many areas of the state (Georgette and Shiedt 2005). Average subsistence harvests of whitefish estimated for the village of Noatak and the 5 Kobuk River villages combined from 1997 to 2004 was almost 54,000 fish (Appendix F8). Harvest numbers are considered minimal and are not comparable year to year. Since 2004, subsistence harvest surveys have not been conducted in the Kotzebue Sound District until 2012, when the Division of Subsistence conducted a comprehensive subsistence fish harvest survey in the Kobuk River villages and Noatak.



## **Historical Escapement**

Whitefish escapements have not been monitored in the past, but limited ADF&G observations and fishermen interviews do not indicate declining populations.

## **SAFFRON COD**

Saffron cod, or tomcod as they are called locally, are extensively utilized as a subsistence resource in Norton Sound-Port Clarence and Arctic-Kotzebue areas. Tomcod are taken through the ice by jigging, and with gillnets in open water and under the ice.

No extensive commercial fishery on tomcod in Norton Sound-Port Clarence and Arctic-Kotzebue areas has ever occurred, but during the 1980s, a limited commercial fishery occurred in Norton Sound (Menard et al. 2013). According to local fishermen, these fish were used for dog food, crab bait, and human consumption. In the mid-1990s, NSEDC established markets for several fish species not commercially utilized in the past. The need for crab bait was the primary factor in initiating the saffron cod fishery near Unalakleet. A total of 1,402 lb of saffron cod were sold during the 1993–1994 season. The NSEDC market was not available the following winter and was probably a factor in the reduced harvest of 52 lb (Appendix F10).

No commercial harvest was reported from 1995 through 2008. Since the fall of 2009, total annual tomcod harvest has ranged from 1,748 lb to almost 34,000 lb (Appendix F10), all sold to Norton Sound Seafood Products (NSSP) in Nome for use as crab bait. NSSP would only buy tomcod that were caught through the ice by jigging gear.

## **Miscellaneous Finfish Species**

Other finfish species taken for subsistence in Norton Sound, Port Clarence, Kotzebue, and Arctic areas include capelin, rainbow smelt (boreal smelt), northern pike, starry flounder, yellow fin sole, Arctic flounder, Alaska plaice, Arctic grayling, burbot, blackfish and halibut (Appendix G1).

## **Subsistence Fishery**

Subsistence utilization of these species has been documented, although effort and catch vary widely in scale and importance with locality. Some species are important to the subsistence community in certain localities during specific seasons of the year. In Nome Subdistrict, both Nome and Solomon rivers were closed to subsistence fishing for Arctic grayling in 2001 when abundance was determined to be low.

## **Commercial Fishery**

Burbot, or freshwater cod, have been commercially sold sporadically in the past in Kotzebue, Port Clarence, and Norton Sound districts under commercial permits.

## **Sport Fishery**

Sport fisheries for Arctic grayling exist in Norton Sound-Port Clarence and Arctic-Kotzebue areas, but they are relatively small. Average annual sport fish harvests for Arctic grayling in the last 5 years were under 1,000 fish in both Norton Sound and Kotzebue Districts. In Norton Sound, average Arctic grayling sport fish harvests for the last 10 years are roughly a fourth of that of Dolly Varden, but in Kotzebue District, average Arctic grayling sport fish harvests for the last 10 years is over half that of Dolly Varden (Appendix F3).

## **SECTION 2: SALMON FISHERIES**

---

### **2013 NORTON SOUND SALMON FISHERY**

#### **2013 Norton Sound Fisheries Outlook**

The 2013 outlook was for a commercial harvest level of 40,000 to 70,000 chum salmon, 50,000 to 100,000 pink salmon, and 30,000 to 60,000 coho salmon. Salmon outlooks and harvest projections for the 2013 season were based on qualitative assessments of parent-year escapements and age composition, subjective determinations of freshwater overwintering and ocean survival conditions, and, in the case of the commercial fishery, anticipated market interest and processing capacity.

For the first time in over 20 years, commercial fishing for chum salmon was expected to occur in Nome Subdistrict. Commercial periods for chum salmon were not expected to exceed 24 hours in length.

As in previous years, the bulk of commercial salmon harvests were expected to come from southern Norton Sound (Subdistricts 4–6). The relatively large southern Norton Sound watersheds (e.g., Inglutalik, Ungalik, Shaktoolik, and Unalakleet rivers) generally support larger runs of salmon. This fact, coupled with stable, healthy salmon runs (except Chinook salmon) and more liberal fisheries management plans, allows for more commercial harvest opportunity in the southern Norton Sound subdistricts. In contrast, salmon runs, particularly chum salmon runs, have been more unstable in the smaller drainages to the north in Subdistricts 2 (Golovin) and 3 (Elim) since the early 2000s. Subdistricts 2 and 3 chum salmon runs have either been very strong, providing large surpluses available for commercial use (e.g., 2006, 2007, 2010, 2011); or very weak, with runs often below levels needed to achieve escapement goals, such as in 2004, 2005, 2008, and 2009. The extent and frequency of commercial chum and pink salmon periods in Subdistricts 2 and 3 is also largely predicated on the Subdistricts 2 and 3 management plan, which directs ADF&G to ensure that chum salmon escapement goals and subsistence needs are achieved.

#### **Commercial Fishery Season Summary**

Weak Chinook salmon runs occurred throughout Norton Sound in 2013, requiring inseason restrictions and early closures to southern Norton Sound subsistence fisheries. As expected in odd-numbered years, pink salmon runs were only sufficient to provide for subsistence needs and limited directed commercial fishing openings. However, substantial increases in commercial chum and coho salmon harvests occurred in 2013, and sockeye salmon abundance was sufficient to reach escapement goals and avoid an early closure to the Pilgrim River sockeye salmon fishery for the first time since 2008.

The 2013 Norton Sound District commercial salmon fishery came in well above the forecast range of 40,000 to 70,000 chum salmon and within the forecast range of 30,000 to 60,000 coho salmon. Norton Sound commercial salmon harvest was 118,672 chum, 193 sockeye, 53,754 coho, and 8,251 pink salmon (Table 1). Also, there were 151 Chinook, 54 sockeye, 48 coho, 87 pink, and 37

chum salmon kept for personal use. The buyer was not able to buy Chinook salmon in Subdistricts 5 and 6, per department emergency order, and chose not to buy Chinook salmon in other subdistricts.

Large chum salmon harvests in conjunction with high prices paid for coho salmon accounted for nearly all \$1,183,236 paid to 124 permit holders in 2013 (Appendices A2 and A3). The 2013 exvessel value ranks third highest and represents the third year since 2010 in which exvessel value has exceeded 1 million dollars (Appendix A3).

Commercial chum salmon harvest in 2013 was the highest in over 25 years. Southern Norton Sound (Subdistricts 4–6) accounted for the bulk of the commercial harvest. Subdistrict 4 (Norton Bay) had a record chum and coho salmon harvest. Commercial chum salmon harvest was also robust in Subdistrict 6 (Unalakleet), which ranked third highest. The chum salmon run was weaker in northern Norton Sound with the exception of Subdistrict 1 (Nome) and Port Clarence District. Subdistrict 2 (Golovin) and Subdistrict 3 (Elim) had the second-highest and fourth-highest coho salmon harvests on record, respectively.

The pink salmon harvest of 8,251 fish was well below the forecast of 50,000 to 100,000 fish. The low harvest was probably due to the existence of only one pink salmon directed fishing period. Both the buyer and fishermen expressed interest in having chum salmon directed fishing periods because of the higher price per pound of chum salmon.

The coho salmon harvest of 53,802 fish was below the recent 5- and 10-year average harvests of 73,080 fish and 76,695 fish, respectively (Appendix A13). Although there were near record to record harvests in Subdistricts 2 and 4, they make up a smaller portion of the overall catch in Norton Sound. The larger harvests usually came from Subdistricts 5 and 6, but in 2013, catches in those subdistricts ranged from only 36% to 66% compared to the recent 5- and 10-year averages (Appendix A10 and A11). Some of this can be attributed to weather during the month of August, particularly in Subdistrict 5. High surf conditions only diminished for brief periods during many scheduled openings in August, which caused several permit holders to lose interest and pursue other endeavors.

The number of permit holders (124) participating in the commercial fishery this year was above average and was the highest number of participants since 1993 (Appendix A2). The previous 5-year average in Norton Sound was 108 permits fished, and the previous 10-year average was 78 permits fished (Appendix A2). The increased fishing effort in the salmon fishery since 2010 is largely the result of stronger chum salmon runs, improved market interest, and high dock prices for salmon, particularly coho salmon. Dock prices per pound for Norton Sound salmon in 2013 were \$0.22, \$0.55, \$1.49, and \$1.77 for pink, chum, sockeye, and coho salmon, respectively (Appendix A4). Chinook salmon were not purchased by the buyer in 2013. Average commercial weights by species were 7.6 lb for coho salmon, 2.9 lb for pink salmon, and 6.9 lb for chum salmon (Appendix A5).

Only one salmon buyer operated in Norton Sound during the 2013 season. The Unalakleet fish plant operated by Norton Sound Seafood Products was the base of commercial fisheries operations. Salmon were both delivered to the Unalakleet dock and tendered from Subdistricts 2–5. Subdistrict 1 catch was delivered by the single permit holder to the Nome plant.

## **Subsistence Fishery Season Summary**

Subsistence salmon fishermen in Port Clarence District, Cape Woolley Subdistrict, and Subdistricts 1–3 were required to possess a subsistence salmon fishing permit for each household that fished in these locations. Households may obtain and fish permits for multiple areas. Return rates for these permits have been close to 100% in most years, and in 2013 the return rate was 100% for the third year in a row (Table 2).

In southern Norton Sound, in 2013, postseason household surveys were conducted in Koyuk, Shaktoolik, and Unalakleet, and attempts were made to contact 100% of the households. Catch information for Subdistricts 4–6 are in Appendices A9–A11. Unlike in 2012, the villages of Stebbins and St. Michael were not surveyed in 2013.

In Norton Sound District, there are limits on subsistence salmon harvests only in Subdistrict 1 (Nome), where salmon limits have been in place since 1985. Also, hook and line subsistence fishermen must follow sport fish bag limits except in the Subdistrict 1 subsistence zones, where they can catch the subsistence limit. In 2013, an average chum salmon run was forecasted for Subdistrict 1 and the subdistrict was not closed to salmon fishing in mid-June for the eighth year in a row. From 1991 through 2005, Subdistrict 1 was closed to subsistence salmon fishing in mid-June in order for ADF&G to determine the run strength of chum salmon before allowing fishing. Furthermore, Tier II regulations were not in effect in 2013 because the chum salmon run was projected to exceed the amount necessary for subsistence (ANS).

In Port Clarence District, subsistence permits are required and a separate permit is required for Pilgrim River and for Salmon Lake. There are no salmon harvest limits in Port Clarence District, except for Kuzitrin River, Pilgrim River, and Salmon Lake.

Beginning in 2007, regulations allowed for cash sales of up to \$200 worth of subsistence-taken finfish per household, per year, in the Norton Sound–Port Clarence Area only, and starting in 2013 the amount allowed was raised to \$500. In 2013, ADF&G increased efforts to remind residents about the permit requirement when selling subsistence-caught finfish and 18 permits were issued, which resulted in cash sales of almost \$2,000 (Appendix A34).

## **Season Summary by Subdistrict**

### ***Nome–Norton Sound Subdistrict 1***

In Subdistrict 1, 2013 chum salmon run abundance was projected to achieve the subdistrict-wide biological escapement goal (BEG) range of 23,000–35,000 chum salmon and amounts necessary for subsistence (ANS) range of 3,430–5,716 chum salmon. As such, a Tier II fishery was not implemented in 2013. There has not been a Tier II fishery implemented since 2005, and Tier II subsistence fishing restrictions were rescinded early during the 2004 and 2005 seasons.

Regulation changes made at 2013 Alaska Board of Fisheries meeting allowed for subsistence gillnet fishing 7 days a week in marine waters in the eastern half of Subdistrict 1, and beach seining was allowed in all subsistence locations during the chum salmon run when gillnet fishing was open. Excellent marine subsistence catches of chum salmon were reported in late June and early July in eastern Subdistrict 1. Aerial surveys were conducted in mid-July of the eastern Nome Subdistrict drainages (Flambeau, Eldorado, and Bonanza rivers) and Sinuk River in the western Nome Subdistrict. Several thousand chum salmon were observed on these surveys in the lower reaches of these drainages. The Eldorado River, Nome River, and Snake River weir counts

exceeded the chum salmon escapement goal ranges in 2013. Consequently, chum salmon subsistence gillnet fishing proceeded on the standard freshwater schedule, and the marine schedule for western Subdistrict 1 was extended from 3 days a week to 5 days a week. Several beach seining opportunities were also issued via emergency order to increase the efficiency of subsistence chum and pink salmon harvests during optimal drying weather periods.

The Subdistrict 1 BEG of 23,000–35,000 chum salmon has been achieved 4 of the last 5 years. However, achievement of the goal is often a result of better and more productive chum salmon runs east of Cape Nome disproportionately contributing to the BEG. The chum salmon escapement goal range for the Eldorado River, which is east of Cape Nome, is double the combined escapement goal range of the Nome and Snake rivers, both of which are west of Cape Nome, highlighting the disparity in river productivity within the subdistrict. In the last 5 years, the Eldorado River has exceeded the chum salmon escapement goal range in 4 years, and the Nome and Snake rivers have met or exceeded their escapement goal ranges in 3 years (Appendix A22–A23 and A26). Although chum salmon runs are greater east of Cape Nome (Appendix A32), for pink salmon the run strength is much greater west of Cape Nome (Appendix A33). Both the Nome and Sinuk rivers have much larger runs of pink salmon, particularly in even-numbered years, compared to rivers east of Cape Nome. Nome River has the only pink salmon escapement goal in Subdistrict 1, and the odd-year goal of 3,200 pink salmon was easily exceeded (Appendix A26).

No coho salmon escapement goals have been established in Subdistrict 1, but the escapement in Nome and Snake rivers was about in the mid-range compared to 10 previous years of sufficient escapement estimates with no large-scale flooding events.

In 2013 there were 477 subsistence salmon permits issued for the Nome Subdistrict, slightly below the 483 permits issued last year and below the record 494 permits issued during the 2010 season. All 477 permits issued were returned (Table 2).

Reported subsistence harvest was 48 Chinook, 3,065 chum, 845 pink, 1,804 coho, and 211 sockeye salmon (Appendix A6). The chum salmon harvest was the fourth highest since 1990, but it was somewhat less than expected based on the large abundance of chum salmon available for harvest. However, increased fishing opportunity in 2013 for sockeye salmon at Pilgrim River may have shifted some effort from Subdistrict 1. The pink salmon harvest was comparable to most odd-numbered years in the last 20 years, and the coho salmon harvest was average compared to the recent 5- and 10-year harvest averages.

### ***Golovin–Norton Sound Subdistrict 2***

The Subdistrict 2 regulatory salmon management plan limits commercial harvest to a maximum of 15,000 chum salmon before mid-July in an attempt to protect chum salmon stocks and allow for some harvest while flesh quality is at its best. By mid-July, the chum salmon run can be assessed and fishing time adjusted accordingly. The counting tower project on the Niukluk River had been used to evaluate escapement in the Golovin Subdistrict from 1995–2012, but the project was eliminated in 2013. The Niukluk River is a tributary of Fish River, a major salmon-producing river in the Golovin Subdistrict. Telemetry studies in the early 2000s showed an average of 33% of the chum salmon in the Fish River drainage pass the Niukluk River tower (Todd et al. 2005).

There was no commercial chum salmon fishing in Golovin Subdistrict from 2002 to 2007, largely because escapements, in most of those years, had fallen short of the lower bound SEG of greater than 30,000 fish for the Niukluk River (Appendix A25). Consequently, ADF&G has implemented a conservative approach with respect to determining when commercial fishing may occur. Early indicators of 2013 chum salmon abundance to Golovin Subdistrict were limited to scant subsistence catch reports of fair catches. ADF&G had concerns about the chum run because the adjacent Subdistrict 3 counting tower at Kwiniuk River was having near record low counts. In 16 of the previous 18 years, if Kwiniuk River counting tower reached or did not reach its chum salmon escapement goal, then Niukluk River counting tower was in agreement for those years in reaching or not reaching the chum salmon escapement goal. Aerial surveys did show larger numbers of chum salmon in the Fish River drainage compared to Kwiniuk River escapement counts and 2 directed chum salmon fishing periods were allowed beginning the third week of July with mediocre catches.

As a result of chum salmon conservation concerns, the pink salmon directed commercial fishery (4.5 in or smaller gillnet mesh size) could not commence until July 14 per the management plan for Subdistricts 2 and 3. On July 17, commercial pink salmon fishing commenced in Golovin Subdistrict with a 48 hour opening. The harvest of 1,028 pink salmon was below average.

Commercial coho salmon fishing periods commenced on August 1 with 24-hour fishing periods. Coho salmon catches were above average, and there were 3 more 36-hour fishing periods the first 2 weeks of August followed by three 48-hour fishing periods the last 2 weeks of August. The coho salmon harvest was the second highest on record, and aerial surveys of Niukluk River and Ophir Creek confirmed that the combined aerial survey goal of 950–1900 coho salmon used in the early 2000s had been exceeded.

The commercial catch in Golovin Subdistrict for 2013 including personal use was 5,362 coho, 1,180 pink, and 3,113 chum salmon caught by 14 permit holders (Table 4). The number of permit holders participating in the fishery was tied with last year and both were the highest since 1998.

This was the tenth year that subsistence salmon permits were required, and 153 permits were issued for Golovin Subdistrict in 2013. Reported harvest was 47 Chinook, 3,256 chum, 3,655 pink, 964 coho, and 15 sockeye salmon (Appendix A7). The number of salmon reported harvested (7,937) ranked fourth lowest in the 2000s.

### ***Elim–Norton Sound Subdistrict 3***

The Subdistrict 3 management plan directs ADF&G to project that chum salmon escapement goals will be reached and ensure that harvestable surpluses will be in excess of subsistence needs before directed chum or pink salmon commercial fishing is allowed. Further, in times of low chum salmon abundance, directed pink salmon commercial fishing may not occur before July 7 in the subdistrict. By this date, historical data indicate that the bulk of the chum salmon run is in river, and commercial pink salmon fishing would be expected to have little impact on chum salmon escapement or subsistence needs.

Early indicators of chum salmon abundance to Elim Subdistrict were limited to scant subsistence catch reports of fair catches in marine waters. However, early projections of chum salmon escapement as indexed by the Kwiniuk River tower counts indicated a very weak run, with the tower-based OEG range of 11,500–23,000 chum salmon unlikely to be achieved. Although there

could have been pink salmon directed commercial fishing periods after July 6, the permit holders expressed no interest because of the lower run size during an odd-numbered year. Late in the season there were 2 directed chum salmon commercial fishing periods, restricted to an area west of the Kwiniuk River mouth. Chum catches were sparse with a harvest of 850 chum salmon (Table 5).

The first directed coho salmon fishing period began on August 1 with a 24-hour opening. Good catches combined with sufficient escapement counts at Kwiniuk River counting tower allowed coho salmon directed fishing periods ranging from 36 to 48 hours for the remainder of the month.

The commercial catch in Elim Subdistrict including personal use was 6 Chinook, 27 sockeye, 6,675 coho, 601 pink, and 1,434 chum salmon caught by 21 permit holders (Table 5). The 2013 Elim coho salmon harvest ranks fourth best historically but is slightly below the 5-year average harvest because the 3 highest harvests occurred in the last 5 years (Appendix A8).

There were 64 subsistence salmon permits issued for Elim Subdistrict in 2013. The number of salmon reported harvested (3,921) was the second lowest since harvest estimation methods were standardized in 1994. Estimated subsistence harvests by species were 39 Chinook, 15 sockeye, 1,515 coho salmon, 1,134 pink salmon, and 1,218 chum salmon. Chinook salmon harvest was a record low since estimation methods were standardized (Appendix A8). Also, the Kwiniuk River Chinook salmon escapement count (15) was the lowest since the 1970s.

#### ***Norton Bay–Norton Sound Subdistrict 4***

Historically, Norton Bay Subdistrict has had difficulty attracting a buyer due to its remoteness and its reputation for watermarked fish. Until recently, Norton Bay Subdistrict has typically been managed based on Shaktoolik and Unalakleet Subdistricts salmon run assessments due to a lack of ground-based escapement projects in Norton Bay. However, in 2011, an enumeration tower project was initiated by NSEDC on the Inglutalik River to provide an index of salmon escapement to Norton Bay. Currently, the Inglutalik River escapement counts are considered ancillary to comparative catch statistics for inseason management until a longer time series of escapement data becomes established.

In 2008, a small-scale commercial salmon fishery occurred in Norton Bay Subdistrict for the first time since 1997, and 4 permit holders participated. ADF&G again opened the commercial salmon fishery in 2009 and 7 permit holders participated. In 2010, there were 5 permit holders participating in the fishery, which was limited due to a combination of inadequate tendering capacity in early July, mechanical breakdowns on tender vessels in August, and reduced fishery participation probably due to concurrent fisheries prosecuted in the Elim and Shaktoolik Subdistricts (permit data on file with ADF&G, Division of Commercial Fisheries; Nome).

In 2011 effort nearly doubled to 12 permit holders, and in 2012 there were 18 permit holders fishing in Norton Bay Subdistrict and a record 49,970 pink salmon were harvested. In 2013 there was a record catch of 36,021 chum and 5,485 coho salmon by 18 permit holders (Table 6).

The first chum salmon fishing period began on June 25 and periods were 48 hours twice a week until mid-July, when fishing periods were increased to 72 hours twice a week. Longer periods were allowed because chum salmon escapement at Inglutalik counting tower was tracking double the commercial catch. Coho salmon directed fishing periods of 48 hours twice a week began on August 2, and the last fishing period ended on September 5.

Cumulative commercial catch by species for Norton Bay Subdistrict including personal use was 8 Chinook, 4 sockeye, 5,485 coho, 487 pink, and 36,021 chum salmon. The final escapement estimate at Inglutalik River tower was 860 Chinook, 5,904 coho, 268,537 pink, and 61,259 chum salmon (Appendix A29). The coho salmon estimate was a minimum estimate because high water precluded counting the entire run.

This was the sixth consecutive year that household subsistence salmon surveys were conducted in the village of Koyuk. Surveys were conducted from 1994 to 2003, but funding limitations precluded surveys of Koyuk during the 2004–2007 seasons. There were 76 households that were successfully contacted out of a possible 82 in 2013. Results from these households were expanded to estimate harvests by species, gear type, and location (e.g., Inglutalik River, Ungalik River, Koyuk River, Mukluktulik River, and marine waters) for those households not surveyed (Appendix A9).

An estimated 123 Chinook, 2 sockeye, 826 coho, 1,341 pink, and 3,853 chum salmon were reported as subsistence harvest in Norton Bay Subdistrict in 2013 and ranked fourth highest out of the last 6 years (Appendix A9).

#### ***Shaktoolik and Unalakleet–Norton Sound Subdistricts 5 and 6***

Both Subdistricts 5 and 6, which share a common boundary, consistently attract commercial markets due to larger volumes of fish and better transportation services. Management actions typically encompass both subdistricts because salmon tend to intermingle, and harvest in one subdistrict affects the movement of fish in the adjacent subdistrict. Results from ADF&G's test net in Unalakleet River (Kent 2010), North River tower counts, and subsistence fishermen interviews in Unalakleet had been used to set early fishing periods in both subdistricts. However, the test net project was discontinued in 2013. This year ADF&G used the North River tower counts to assess run strength along with commercial and subsistence catches and, later in the run, counts from the Unalakleet River weir, which is much farther upstream. Radiotelemetry projects in the Unalakleet River drainage have shown that a large percentage of the Chinook salmon run spawns in the North River compared to chum and coho salmon (Estensen et al. 2005; Estensen and Hamazaki 2007; Joy et al. 2005; Joy and Reed 2006, 2007; Wuttig 1998 and 1999). Aerial surveys are only useful for late-season escapement assessment because of the long travel time between the fishing and spawning grounds.

In Subdistricts 5 and 6, directed commercial Chinook salmon fishing has only occurred in 2 of the previous 11 years, and in only 1 year since 2001. Restrictive action was taken in the subsistence and sport fisheries from 2003 to 2004 and from 2006 to 2013. As forecasted, a weak run of Chinook salmon to Shaktoolik and Unalakleet Subdistricts in 2013 precluded commercial fisheries directed on Chinook salmon but also led to a significant amount of foregone chum salmon harvest surplus. As a consequence of the poor Chinook salmon run, directed chum salmon fishing was delayed until July 1 per the Shaktoolik and Unalakleet Subdistricts management plan.

Estimated 2013 Chinook salmon escapements from the Unalakleet River mainstem and its major tributary, North River, were 767 and 564 fish, respectively, and were the lowest ever recorded (data on file with ADF&G, Division of Commercial Fisheries; Nome). Subsistence Chinook salmon harvests in Subdistrict 5 and 6 were the lowest recorded since survey methods were standardized in 1994, with 136 and 468 fish, respectively (Appendices A10 and A11).



Despite the late start to commercial fishing in Subdistricts 5 and 6, the chum salmon commercial harvest was well above average. The Subdistrict 5 harvest (23,268) ranked third highest and the Subdistrict 6 harvest (54,873) ranked the highest out of the last 20 years of commercial harvests, respectively (Appendices A10 and A11). Initially ADF&G started with 24-hour chum salmon periods the first week of July to protect Chinook salmon. Good catches of chum salmon in both subdistricts combined with low Chinook salmon incidental harvest resulted in the department increasing to 48-hour fishing periods the second week of July and then increasing to two 72-hour fishing periods per week in both Subdistricts 5 and 6 beginning the third week of July until the end of the month.

The commercial fishing schedule for coho salmon for Subdistricts 5 and 6 went into effect in early August with two 48-hour fishing periods per week. During the last decade commercial coho salmon harvests were the highest on record in Norton Sound District (Appendix A13), but they have recently returned to average levels prior to the record harvests from 2006–2008. In Subdistrict 5 coho salmon harvest declined for the third consecutive year with a harvest of 6,890 coho (Appendix A10). Subdistrict 5 fishermen had to contend with relatively severe local surf conditions in August, which may in part explain why Shaktoolik coho salmon harvests declined this season despite increased commercial harvests elsewhere in Norton Sound compared to last year (Appendices A6–A11).

The Subdistrict 6 commercial harvest of 29,390 coho salmon ranked twelfth highest in the last 20 years and represented a 32% increase from the 2012 harvest of 22,274 (Appendix A11).

## **Escapement**

Table 3 and Appendix A17 summarize escapement assessments for the major index river systems of Norton Sound and Port Clarence Districts in 2013. Appendices A22–A31 present passage numbers for Chinook, chum, coho, pink, and sockeye salmon at various enumeration projects in Norton Sound. Aerial survey assessments are indices and relative to historical escapement sizes.

Escapement projects in Norton Sound include counting towers on North, Inglutalik, and Kwiniuk rivers; sonar on Shaktoolik River; and weirs on Unalakleet, Snake, Nome, Solomon, Eldorado, and Pilgrim rivers, and in Glacial Creek, which flows from Glacial Lake into Sinuk River.

Escapement project operations were a result of multiple collaborators, including ADF&G, NSEDC, BLM, and Unalakleet IRA. All projects supplied important daily information to ADF&G that was very useful for management of local salmon resources and will become more important the longer they operate. Funding sources for projects come from USFWS Office of Subsistence Management, NSEDC, and ADF&G.

Aerial survey assessment conditions were fair to poor during July and August of 2013; as a result, there were very few aerial surveys flown.

### ***Chinook Salmon***

Chinook salmon escapement was estimated to be very weak in many locations in 2013. A record low 15 Chinook salmon were counted at the Kwiniuk River tower, which was well below the lower end of the SEG range of 300–550 fish (Appendix A24). Also, a record low escapement of 564 Chinook salmon at the North River tower was less than half the lower end of the escapement goal range of 1,200–2,600 Chinook salmon (Appendix A30). Final escapement at the Unalakleet

River weir was 767 Chinook salmon, which was the lowest count in the 4-year project history (Appendix A31). Surprisingly, preemptive measures, additional marine mesh-size restrictions, and eventual early closures were not sufficient to achieve Chinook salmon escapement goals in Subdistrict 6.

### *Chum Salmon*

Chum salmon escapement goals were achieved in 5 of 8 established Norton Sound chum salmon runs. The Kwiniuk River tower-based goal was not achieved. The former Niukluk River tower-based goal could not be determined because the project is no longer operational. Tubutulik River's escapement goal was not evaluated because the aerial survey was not conducted during the peak spawning stage.

Subdistrict 1 ended up having its largest chum salmon escapement in over 20 years. Estimated subdistrict-wide escapement of chum salmon was 108,120 fish, 209% above the upper bound of the subdistrict-wide biological escapement goal (BEG) range of 23,000–35,000 chum salmon (Table 3; Appendix A21). Subdistrict 1 escapements of chum salmon have exceeded the upper bound of the escapement goal range in 8 of the last 13 years of the established goal. As in previous years, the majority (62%) of the chum salmon escapement occurred in rivers east of Cape Nome. However, Sinuk River, west of Cape Nome, had the largest estimated escapement for an individual river system, contributing 31,691 chum salmon or 29% of the subdistrict-wide escapement (Appendix A32).

Escapement at Kwiniuk River tower was 5,631 chum salmon, only slightly above the record low count of 5,577 chum salmon observed in 2012 (Appendix A24). This was the second consecutive season that escapement fell well short of the optimal escapement goal (OEG) range of 11,500–23,000 chum salmon. However, escapement of chum salmon to the neighboring Tubutulik River may have narrowly met the OEG range of 9,200–18,400 fish, based on a July 9 aerial survey of 4,532 chum salmon in the lower reaches of the watershed (Table 3). To the west in Subdistrict 2, the peak spawning ground aerial survey of the Niukluk River tributary of the Fish River was 17,203 chum salmon (Table 3). This year's survey count suggests that actual ground-based escapement of chum salmon to the Niukluk River tower was near the former tower-based sustainable escapement goal (SEG) threshold of  $\geq 23,000$  fish.

As with harvest patterns, southern Norton Sound drainages showed comparably strong chum salmon escapements in 2013. For example, Norton Sound's largest chum salmon producer, the Unalakleet River drainage, had its second largest escapement on record. The estimated escapement was 124,471 chum salmon (based on aggregate tower and weir counts) in 2013 (Table 3).

In Port Clarence District, chum salmon runs were also strong in 2013. Escapement of chum salmon to the Pilgrim River was 47,557 fish, which established a new record high count for the Pilgrim River floating weir project (Appendix B2). The previous record escapement was 45,361 chum salmon set in 2006.

### *Coho Salmon*

Coho salmon are found in nearly all of the chum salmon producing streams throughout Norton Sound, with the primary commercial contributors being the Unalakleet and Shaktoolik rivers. Escapement data are not available over a long time series for several streams because few projects counted the coho salmon run prior to the early 2000s due to funding limitations. More

recent Norton Sound escapement assessment projects have been funded to monitor coho salmon as well as chum salmon and are becoming increasingly important to fisheries management.

There are only 2 coho salmon escapement goals in Norton Sound, and both are aerial survey goals. The North River goal of 550–1,100 was achieved with an aerial survey estimate of 867 fish in 2013 (Table 3). The Kwiniuk River goal of 650–1,300 was probably achieved because, although no survey was flown, the final tower count was 3,940 fish (Table 3).

The previous aerial survey goal for Niukluk River and Ophir Creek was 950–1,900 coho salmon, but it was eliminated with a Niukluk River tower goal of 2,400–7,200 coho salmon. A combined aerial survey count of 2,353 coho salmon of Niukluk River and Ophir Creek indicates that both former escapement goals would have been reached (Table 3).

Both the Snake (1,203) and Nome (2,624) rivers' weir projects had counts near the median for coho salmon (Appendix A23 and A26).

### ***Pink Salmon***

For over 25 years, pink salmon runs to Norton Sound have followed an odd- and even-numbered year cycle, with even-numbered year runs typically much higher in abundance than odd-numbered years. Pink salmon escapement estimates were successfully obtained from all ground-based escapement projects in 2013. There are 3 pink salmon escapement goals in Norton Sound: Nome River (3,200), Kwiniuk River (8,400), and North River (25,000). In almost all years the goals are reached, and the goals were likewise reached in 2013 (Table 3).

### ***Sockeye Salmon***

River spawning sockeye salmon are typically found in small numbers throughout Norton Sound District. Glacial Lake (Nome Subdistrict) and Salmon Lake (Port Clarence District) support populations of lake-spawning sockeye salmon and constitute the northernmost populations of any significance of sockeye salmon in North America. Salmon Lake spawning populations seldom exceeded 10,000 fish in years previous to 2003, whereas from 2003 to 2007 there were near-record to record runs of sockeye salmon. Likewise, Glacial Lake saw an upswing in sockeye salmon returns beginning in 2004, and a record count of 11,135 sockeye salmon occurred in 2005 (Appendix A28).

In 2008, sockeye salmon escapement dropped off at both Glacial Lake and Salmon Lake, and in 2009 sockeye salmon counts further declined at both Glacial Lake weir and Pilgrim River weir. The Glacial Lake weir is operated at Glacial Creek near the outlet of the lake and about 1 mile upstream from the confluence with the Sinuk River, and 826 sockeye salmon were counted in 2009, the lowest count since the weir project started in 2000 (Appendix A28). The 2009 Salmon Lake sockeye salmon run was also the lowest since Pilgrim River weir began operations in 2003, with 953 sockeye salmon counted through the weir (Appendix B2).

Sockeye salmon escapements in these 2 systems increased in 2010, although not by much. Sockeye salmon escapement in 2010 at Glacial Lake was 1,047 fish, tying 2002 for the third lowest count since the project began in 2000 (Appendix A28). Pilgrim River weir sockeye salmon escapement in 2010 was 1,654 fish, which was the second lowest on record (Appendix B2).

The escapement at Glacial Lake weir in 2012 of 1,636 sockeye salmon was just slightly less than the 2011 escapement; the same goes for the Pilgrim River weir escapement of 7,085 sockeye salmon.

Improving sockeye salmon runs occurred at both Glacial and Salmon lakes in 2013. An estimated 2,544 sockeye salmon were enumerated at Glacial Lake weir, and 12,428 sockeye salmon were enumerated at the Pilgrim River weir in 2013 (Appendices A28 and B2). The 2013 Glacial Lake weir count was a 55% increase from the 1,636 sockeye salmon counted in 2012. Similarly, the 2013 Pilgrim weir count represents a 75% increase from the 2012 weir count of 7,085 sockeye salmon (Appendix B2). An August 13 aerial survey count of 6,971 sockeye salmon was greater than the 5,830 and 5,144 sockeye salmon observed during the 2011 and 2012 aerial surveys, respectively. The 2013 season signifies the third consecutive season in which the Grand Central River/Salmon Lake aerial survey SEG range of 4,000–8,000 has been achieved, but unlike the 2 previous years, the Pilgrim River was not closed to subsistence salmon fishing. A total of 1,366 sockeye salmon was also observed at Glacial Lake, making 2013 the first season since 2010 that the Glacial Lake aerial survey SEG range (800–1600 sockeye salmon) has been evaluated and achieved.

## **Enforcement**

Fishing regulations are primarily enforced by the Department of Public Safety, Alaska Wildlife Troopers (AWT). Two AWT officers patrolled the Norton Sound District 2013 commercial salmon fisheries in Unalakleet, and one AWT officer patrolled the Nome area. In addition, Nome ADF&G Division of Commercial Fisheries has 7 deputized staff with the ability to issue citations, of which 2 worked the commercial salmon fishery in Shaktoolik and Unalakleet Subdistricts. The subsistence fishery had no official patrol, but random checks were conducted by 2 ADF&G personnel.

## **2014 NORTON SOUND SALMON OUTLOOK**

Salmon outlooks and harvest projections for the 2014 salmon season are based on qualitative assessments of parent-year escapements, subjective determinations of freshwater overwintering and ocean survival, and, in the case of the commercial fishery, the projections of local market conditions. The Chinook salmon run is expected to be very weak and similar to the 2013 run, with no commercial fishing targeting Chinook salmon expected. Additional preemptive subsistence restrictions are also likely for southern Norton Sound in order to conserve Chinook salmon to reach escapement goals. These restrictions include preemptive closures or reductions in fishing time in marine waters, inriver closures to gillnets with a mesh size greater than 4.5 in, and 6 in or less mesh size restrictions in marine waters. However, beach seining subsistence opportunity will be provided early in the run to allow the take of other, more plentiful species like pink and chum salmon.

Chum salmon runs are expected to be average in southern Norton Sound Subdistricts (Norton Bay, Shaktoolik, and Unalakleet) based on the recent 5-year trend of average to above-average chum salmon abundance in southern Norton Sound and sibling relationship analyses. As a result, directed chum salmon fishing is anticipated to commence as early as the third week of June in Norton Bay Subdistrict but no earlier than July 1 in Shaktoolik and Unalakleet Subdistricts because of Chinook salmon conservation concerns. In 2014, northern Norton Sound chum salmon runs are expected to be below average to average. Chum salmon abundance is anticipated to be sufficient to reach escapement goals and perhaps provide for a limited chum salmon

commercial harvest in Subdistricts 2 (Golovin) and 3 (Elim). A limited commercial fishery for chum salmon is possible in Nome Subdistrict dependent on a sufficient chum salmon run to obtain escapement goals throughout the subdistrict. Overall projected commercial harvest of chum salmon in Norton Sound is expected to range between 80,000–110,000 fish with an increased contribution to this harvest expected for Norton Bay Subdistrict due to improvements in tendering capacity, a good forecast, and a flexible management plan.

ADF&G expects the pink salmon run to be average for an even-numbered year, and dependent on buyer interest the harvest could be 250,000–500,000 fish. No subsistence fishing restrictions for pink salmon are expected.

The coho salmon run in 2014 is expected to be average based on recent 5-year trends in abundance and ocean conditions, as well as parent-year escapements and freshwater rearing conditions for the 2010 brood year. Northern Norton Sound contributions to the coho salmon run are expected to be below average to average, based on less-than-favorable freshwater rearing conditions. Conversely, southern Norton Sound runs of coho salmon in 2014 are expected to range from average (Subdistricts 5 and 6) to above average (Subdistrict 4) based on favorable freshwater rearing conditions. Considering these factors collectively, the commercial harvest is expected to range from 60,000 to 90,000 coho salmon. Coho salmon subsistence fishing restrictions are not expected.

## **2013 PORT CLARENCE SALMON FISHERY**

### **Commercial Fishery Season Summary**

No commercial salmon fishing was allowed in 2013. ADF&G had projected, and later confirmed by counts from the Pilgrim River weir, that the sockeye salmon run for Pilgrim River in 2013 would not reach the inriver goal of 30,000 sockeye salmon that is necessary for a commercial fishery to occur.

### **Subsistence Fishery Season Summary**

Subsistence fishing permits have been required for Pilgrim River since 1964, and beginning in 2003 the number of permits issued has skyrocketed with the record sockeye salmon runs in the mid-2000s. In 2013 a record 265 permits were issued, surpassing the previous record in 2008 when 255 permits were issued (Menard et al. 2012). Pilgrim River estimated subsistence harvests by species were 8 coho salmon, 48 pink salmon, 2,761 sockeye salmon, and 282 chum salmon (Table 2). This was the first year since 2008 that subsistence salmon gillnet fishing was not closed during part of the season on Pilgrim River.

The size of the Pilgrim River sockeye salmon run greatly affects the number of issued subsistence permits. The first year of the great runs of sockeye salmon (2003), there were 100 permits issued. In 2004, there were 223 permits issued (permit data on file with ADF&G, Division of Commercial Fisheries; Nome). For comparison, in 2002 only 25 permits were issued, and a counting tower in operation that year at the same location as the present-day weir estimated less than 4,000 sockeye salmon passing (Appendix B2).

Although permits have been required in the Pilgrim River drainage for 50 years, 2013 was only the tenth year that permits were required throughout Port Clarence District. The number of subsistence salmon permits issued for all waters of Port Clarence District, excluding Pilgrim

River and Salmon Lake, was 162 permits, compared to 147 permits issued last year. Salmon Lake was opened in September in the northeast section and 4 permits were issued (Table 2).

## **Escapement**

Aerial surveys are not typically flown in Port Clarence District except for Salmon Lake because higher priority is assigned to Nome Subdistrict and surrounding areas where commercial fishing occurs. Aerial surveys had shown an increasing trend of sockeye salmon returns to Salmon Lake since 1990 (Appendix B1). However, the sockeye salmon run crashed beginning in 2009, and ADF&G has had to have subsistence fishing restrictions on Pilgrim River in 4 of the last 5 years. An aerial survey in 2012 of Salmon Lake and Grand Central River estimated 4,730 sockeye salmon in Salmon Lake and 1,100 sockeye salmon in Grand Central River, a tributary to Salmon Lake. The combined aerial survey count of 6,971 sockeye salmon in 2013 was higher than both the 2012 (5,830) and 2011 (5,144) counts, and was the third time in a row the escapement goal has been reached since 2008 (survey data on file with ADF&G, Division of Commercial Fisheries; Nome). The combined aerial survey escapement goal for Salmon Lake and Grand Central River is 4,000–8,000 sockeye salmon (Table 3).

Salmon Lake had an average sockeye salmon spawning population of roughly 12,500 fish in the 5 years previous to 2003. But from 2003 to 2007, sockeye salmon escapements skyrocketed, and average weir count for the 5-year period was almost 56,000 sockeye salmon (Appendix B2). Average count for the next 5 years decreased greatly, to less than 8,000 sockeye salmon. From 2012 to 2013, the count improved by 75%, from 7,085 to 12,428 sockeye salmon (Appendix B2).

## **Enforcement**

In 2013, one AWT officer patrolled Pilgrim River in Port Clarence District.

## **2014 PORT CLARENCE SALMON OUTLOOK**

The guideline harvest range (GHR) set by BOF for the Port Clarence commercial sockeye salmon fishery allows for a harvest of up to 10,000 sockeye salmon. Based on recent history, ADF&G expects that the inriver goal of 30,000 sockeye salmon for Pilgrim River will not be met; therefore, no commercial fishing is expected in 2014. In addition, based on escapement and smolt data, the sockeye salmon run is expected to drop, and subsistence fishing restrictions may occur during the second week of July or later, if necessary. Chum and pink salmon are expected to have sufficient runs allowing for subsistence fishing.

ADF&G will compare the 2014 run with sockeye salmon escapement counts at the weir from the last few years and determine whether any subsistence fishing restrictions are needed.

## **2013 KOTZEBUE SOUND SALMON FISHERY**

### **Commercial Fishery Season Summary**

The Kotzebue Sound District commercial salmon fishery opened on July 10 and closed after the August 31 fishing period. Similar to the previous year, there was a very strong run of chum salmon and again commercial fishing time was limited, mostly in August, because of insufficient cargo space on airplanes to get the catch out to processing facilities.

During most of July, there was sufficient airplane capacity for the fleet to fish 6 days a week and fishing periods were 4 to 8 hours in length. Starting in late July until the third week of August, periods were 4 hours or less because of the high catches during the traditional peak weeks of harvest. During the last week of August, fishing periods were increased to 12 hours a day as a result of decreasing catches, reduced fishing effort, and sufficient airplane capacity.

There were 66 permit holders who sold chum salmon in 2013 (Appendix C1). Of these, 65 permit holders sold fish to the major buyer, Great Pacific Seafoods. One catcher–seller sold to a second buyer, Maniilaq Services, Incorporated, and also to Kotzebue area residents. In late August, with the decreasing catches, Maniilaq began to buy from additional permit holders who had been previously selling to Great Pacific. Maniilaq purchased from 21 permit holders during the season. There was a 20% drop in the number of permit holders selling fish this year compared to last year (Appendix C1). The price per pound for chum salmon dropped 16% from last year (Appendix C3) and may have been a factor in the number of permit holders fishing this season.

The overall chum salmon run to Kotzebue Sound in 2013 was estimated to be well above average based on commercial harvest rates, subsistence fishermen reporting good catches, and the Kobuk test fish index being the highest in the 21-year project history (Table 10).

The commercial harvest of 319,062 chum salmon was the highest since the 1980s (Appendix C1). There were 67 chum salmon kept for personal use included in the commercial harvest total. Also kept for personal use were 16 Chinook salmon, 13 sockeye salmon, 42 pink salmon, 43 coho salmon, 302 Dolly Varden, 705 sheefish, and 50 whitefish (Table 9). However, there were likely some additional fish kept for personal use that did not get reported on fish tickets.

A total of 2,555,304 pounds of chum salmon (average weight 8.0 lbs) were sold at an average of \$0.27 per pound (Appendices C2 and C3). The total exvessel value was \$689,163. The average value for each participating permit holder was \$10,442. The total exvessel value was almost two and a half times that of the average from the last 20 years (Appendix C4).

In the Kotzebue fishery, gear is limited to set nets with an aggregate of no more than 150 fathoms per permit holder. Fishermen generally operate with an end on or near shore and with all 3 shackles connected. Fishermen also set in deeper channels in the mud flats farther out from shore. Most gear used in the district is 5.875 in (14.9 cm) or 6 in (15.2 cm) stretch mesh gillnet.

Age, sex, and length (ASL) composition was taken from commercial catch samples but was not used to manage the fishery. The majority of the chum salmon each year are usually 4- and 5-year-old fish. In 2013, commercial catch samples were 4% age-0.2 fish, 54% age-0.3 fish, 40% age-0.4 fish and 2% age-0.5 fish. Historical comparisons showed the catch samples falling within the range of previous years (<http://www.adfg.alaska.gov/CommFishR3/WebSite/AYKDBMSWebsite/Default.aspx>).

### **Subsistence Fishery Season Summary**

Subsistence household salmon surveys were regularly conducted in Kotzebue District from 1990 to 2004 by the Division of Subsistence (DOS), and again starting in 2012, when comprehensive subsistence fish harvest data were collected from 6 Kotzebue area villages by DOS. In 2012, total subsistence chum salmon reported caught was 26,693 salmon, more than in 2003 and 2004, the last 2 years that the same 6 villages were surveyed (Appendices C5 and C6), and subsistence chum harvest per household averaged 70 salmon for Kobuk River villages (Appendix C7).

Subsistence survey information from 2013 is not yet available, but funding is available for DOS to conduct surveys through 2014 (Nikki Braem, Subsistence Resource Specialist, ADF&G, Fairbanks; personal communication).

### **Escapement**

This year's test fishing chum salmon CPUE cumulative index at ADF&G test fish project on Kobuk River near Kiana was 2,698 points and was a record index at the Kobuk River test fish project (Table 10). The Kobuk River test net catch samples were 6% age-0.2 fish, 33% age-0.3 fish, 60% age-0.4 fish and 1% age-0.5 fish. Historical comparisons show the catch samples falling within the range of previous years. High and turbid water prevented aerial surveys of the Kobuk River and Noatak River drainages in 2013.

### **Enforcement**

One AWT officer patrolled the Kotzebue Sound District 2013 commercial salmon fishery.

### **2014 Kotzebue Salmon Outlook**

The outlook for the 2014 season is based on the parent-year returns and returning age classes observed in the commercial catch samples and in the test fishing catch samples from the Kobuk River in 2013. During the 2014 season, the 4-year-old component of the run is expected to be above average based on the 3-year-old return. The 5-year-old component of the run is expected to be average based on the 4-year-old return this past season. The 3-year-old and 6-year-old age classes are much smaller components of the run and are expected to be average (age data on file with ADF&G, Division of Commercial Fisheries; Nome). The commercial harvest is expected to fall within the range of 250,000 to 275,000 chum salmon, if market conditions can accept that level of harvest.



## SECTION 3: PACIFIC HERRING FISHERIES

---

### 2013 NORTON SOUND PACIFIC HERRING FISHERY

#### Sac Roe

In contrast to last year when there was no herring sac roe fishery, there was a fishery in 2013 and 490 tons of sac roe herring were harvested by 40 permit holders (Table 11). The majority of the sac roe herring harvest was from the Cape Denbigh area (Subdistrict 3). Historical information for the Norton Sound commercial sac roe fishery can be found in Appendix D2 and Menard et al. 2013. Other historical fisheries information is presented in Appendices D1 and D3.

#### Spawn-on-Kelp

There was no market interest expressed in the commercial spawn-on-wild-kelp (*Fucus* sp.) or *Macrocystis* spawn-on-kelp fisheries in 2013.

#### Bait Fishery

Two tons of herring were bought as bait during the sac roe fishery.

#### Commercial Fishery Management

ADF&G projection for the 2013 herring spawning biomass for Norton Sound was 58,594 tons. At 20% exploitation rate, the guideline harvest level (GHL) for the Norton Sound District fishery was 11,719 tons with 11,399 tons allocated to the sac roe fishery. NSEDC was successful at developing a market for 800–1,200 tons of sac roe herring in 2013. However, only 41–61% of the desired harvest was obtained before the floating processor had to depart for Bristol Bay salmon fishery on June 21 (Appendix D2). As in 2010, extensive pack ice in the northern Bering Sea delayed the arrival of tender and processing vessels in Norton Sound. Additionally, extensive shorefast ice in southern Norton Sound also severely hampered fishing effort during the second week of June when herring spawning biomass was building in the St. Michael (Subdistrict 1) and Cape Denbigh (Subdistrict 3) Subdistricts. With the exception of Shaktoolik, permit holders had difficulty departing from southern Norton Sound villages until after June 10 due to shorefast ice. Even though the harvest was small, roe recovery averaged 13.2%, the third highest on record and the third consecutive season that roe percentages have exceeded 13% (Appendix D2).

A temporary lead in shorefast ice allowed the ADF&G field crew to deploy for Cape Denbigh from Unalakleet on June 10 to conduct test fishing operations. The 2013 season was the first season since 2008 in which test fishing operations were conducted throughout the entire herring spawning season. Test fishing operations commenced on morning of June 11. The test fishery crew's first successful variable-mesh gillnet (VMG) haul occurred on June 12 when 51 herring were sampled from a catch of 306 herring on the western end of Cape Denbigh in front of Elim Beach. Approximately 85% of the catch occurred in the 2.0-inch mesh panel of the VGM. Test fish operations were conducted daily morning and evening from June 12–18 and a total of 452 VMG samples were collected, the majority originating from the Elim Beach area. During this

time, the crew also collected 615 samples from tender vessels operating in the Cape Denbigh Subdistrict throughout the duration of the fishery (data on file with ADF&G, Division of Commercial Fisheries; Nome). The crew departed the Cape Denbigh area on June 18 because the fishery was winding down and they wanted to avoid forecasted hazardous surf conditions that would have significantly delayed their return to Unalakleet.

### **Catch Reporting and Enforcement**

No AWT officers were on Norton Sound herring grounds during the 2013 fishery because of the ice conditions preventing a sac roe fishery. Catch reporting for the bait fishery was sufficient for the limited harvest, and a final report was submitted to ADF&G.

### **Biomass Determination**

There were no Norton Sound herring aerial surveys conducted this season by NSEDC or ADF&G biologists because of ice conditions that made surveying and accurately estimating herring biomass impossible.

### **2014 NORTON SOUND PACIFIC HERRING OUTLOOK**

The 2014 projected biomass for Norton Sound District is 52,138 tons. A 20% exploitation rate would result in a GHL of 10,428 tons. A maximum of 320 tons of herring are reserved to allow for the pound fishery to harvest a maximum of 90 tons of product (combined weight of herring roe and kelp). This leaves 10,108 tons for sac roe harvest. The beach seine harvest is allocated 10% of the sac roe projected harvest, or 1,011 tons. The 2014 herring fishery will be opened by emergency order and the fishery will close by emergency order when up to 20% of the available herring biomass has been harvested. Varied harvest rates may be applied to individual subdistricts based on biomass distribution, roe quality, weather, and sea ice conditions. Herring ages 6 and 7 are expected to make up 30% of the returning biomass (Appendix D13). Herring age 8 and older are expected to make up 64% of the biomass. If there are more favorable ice conditions in 2014, ADF&G expects to conduct a more comprehensive test fishery, and a commercial sampling program is anticipated for the 2014 season to obtain more representative age class data from the spawning biomass and harvest.

## SECTION 4: KING CRAB FISHERIES

---

### NORTON SOUND CRAB FISHERY

#### Abundance

The ADF&G length-based population model estimated harvestable legal male crab biomass for the 2013 summer commercial crab fishery at 4.13 million lb (1.56 million crab), a decrease of 2% from the revised model biomass estimate of 4.22 million lb (1.56 million crab) for legal male crab for 2012. An exploitation rate of 12% on the harvestable population equates to a guideline harvest level (GHL) of 495,600 lb of crab. This follows the revised harvest strategy set in regulation by the BOF in March of 2012. By regulation, the CDQ fishery is allocated 7.5% of the GHL; therefore, the CDQ harvest quota was set at 37,170 lb preseason, with the open access fishery allocation set at 458,430 lb.

#### Summer Open Access Commercial Fishery

The 2013 summer open access commercial crab fishery was opened by emergency order at noon on July 3 in the Norton Sound Section, with a GHL of 458,430 pounds of crab. Two companies, Norton Sound Seafood Products (NSSP) and Aquatech, were registered to buy crab, and 5 fishermen registered to sell crab dockside as catcher–sellers or catcher–processors. NSSP operated a seafood processing plant in Nome and 2 tenders in eastern Norton Sound. A fisherman based in Unalakleet shipped live crab via airplane to Aquatech in Anchorage. The majority of crab was delivered to NSSP, whereas the catcher–sellers sold crab directly to local residents as well as to NSSP. The 2 catcher–processors sold to a live market in Korea and to NSSP.

The first open access delivery was made on July 5 and final delivery was made September 15, the day after the open access portion of the fishery was closed by emergency order at 6:00 PM, for a total season length of 74 days, compared to 44 days in 2012 (Table 12). This year, the season start was delayed for a few days because NSSP found poor meat fill in some of their test fishery crab; however, once the open access season was under way, both land-based buyers purchased crab continuously with no further reports of poor crab meat fill until the very end of the season.

In 2013, catch rate was poor for the entire fleet and by the end of July the projected trend line showed that if the catch rate did not change, the quota would not be reached until the second half of September. On August 1, to allow the fleet to efficiently obtain the allowable harvest of red king crab, the closure line south of Nome from 166°15'W (Sledge Island) and 163°30'W (near Square Rock) was moved by emergency order north by 3 miles, to 64°18'N from 64°15'N (Appendix E11). On August 7, to allow more of the fleet additional fishing opportunity, the closure line between 162°38'W (Carson Creek) and 162°W (the eastern border of statistical area 626401) was moved by emergency order north by 5 miles to 64°15'N, and the closure line between 162°W and 161°30'W (Cape Denbigh) was also moved north, by 3 miles to 64°18'N (Appendix E11). Catch rates increased slightly and temporarily for the portion of the fleet that fished in the western half of Norton Sound but did not improve for the crabbers fishing in the east side of Norton Sound. By the second half of August, the trend line projected the quota would not be

reached. Because most of the registered buyers were still interested in purchasing crab, the fishing time was extended from September 3 to September 10, by which time a few newly molted crab had shown up in the fishery. A shorter extension was granted, to September 13, but safety concerns due to adverse marine conditions resulted in the final closure date of September 14.

The open access harvest from fish ticket reports was 124,485 red king crab or 373,278 lb (81% of the open access quota; Table 12). Of this total, 882 lb were reported as deadloss and 6,248 lb as personal use. A total of 33 vessels and 33 permit holders made 435 landings, and average weight for commercially caught crab was 3.0 lb. Including CDQ, the number of pots registered was 1,420, and there were 15,058 pot pulls, for a season CPUE of 9 crab. In 2013, the total catch rate tracked very closely to 2009; these 2 years had the lowest catch rates out of the last 6 years (Appendix E9). Average price paid (including CDQ catch) was \$5.63 per lb, the highest amount ever paid (Appendix E3). Exvessel value of the fishery (including CDQ) was \$2.165 million, second only to the record-setting amount paid out in 2012 of \$2.556 million, the highest since 1994 when Norton Sound was designated a superexclusive area, which effectively changed the character of the fishery from a large vessel to a small vessel fishery (Appendix E10).

### **CDQ Fishery**

For the fourth time in the last 5 years, the CDQ fishery opened concurrently with the open access fishery in 2013. The first CDQ delivery was made on July 16 and the last delivery was made September 15, when 18,585 lb (50% of the allocated 37,170 lb) had been harvested, for a total length of 74 days (Table 12 and Appendix E1). In 2013, as in the previous 6 years, YDFDA transferred their CDQ quota to NSEDC; however, due to the poor catch rate all season, NSEDC only harvested their original half of the CDQ quota, resulting in the first time in 10 years that the fishery did not harvest all, or nearly all, of the allocation. Seven permit holders were registered to fish the CDQ fishery, but only 4 fished, making a total of 25 landings and 862 pots lifts. Average price paid to fishermen was \$5.50 per lb, for an exvessel value of \$101,250 for the CDQ fishery. This was the thirteenth year a CDQ harvest occurred since the CDQ fishery was implemented in 1998.

### **Harvest Areas and Commercial Catch Sampling**

Fish ticket reports document fourteen statistical areas were fished in the open access and CDQ fisheries (Table 13), compared to 9 areas in 2012. Unlike last year, the top harvest came from statistical area 656401 (38%). As the crabbers moved around more this year trying to find the crab, their harvests came from a larger area instead of being concentrated as in years past. Several statistical areas yielded harvests of 9–15%: 646401 (15%), 656402 (10%), 626401 (9%), 636401 (9%), and 666401 (9%). Except for statistical area 656402, which was temporarily opened this year, all the other listed statistical areas are directly south of the original closed boundary line (Appendix E11). The catch from statistical areas east of 164°W longitude made up 23% of the harvest, less than half of the 56% last year (Appendices E1 and E12).

Carapace length (CL) measurements and shell age were collected from 6,072 commercially caught crab during the open access and CDQ fisheries. Carapace age was classified as new (2–12 months old) or old (over 13 months old). Male new-shell crab made up 91% of the total legal crab sampled, and old-shell crab made up 9%. Recruit crab are new-shell legal crab < 116 mm CL. Postrecruit crab are legal new-shell male crab  $\geq$  116 mm CL and all legal old-shell males.

Recruit crab made up 32% of the legal crab sampled and postrecruit crab made up 68%, the highest percentage of postrecruit crab since 2006 (Appendix E4). Overall mean carapace length of legal male crab was 120 mm. For comparison of historical length composition of Norton Sound red king crab summer commercial harvests from 1990 to 2013, see Appendices E15–E20.

## **Enforcement**

No AWT trooper made dockside checks during the 2013 summer crab fishery; however, an ADF&G staff member who worked the king crab fishery was deputized to cite violations if necessary.

## **Winter Commercial Fishery**

The winter commercial season opened November 15, 2012, and 34 fishermen registered. One land-based processor (NSSP) and 2 land-based catcher-processors registered to buy crab, and 7 fishermen registered to sell crab dockside as catcher-sellers. Based on fish tickets submitted, the first landing was made December 22, 2012. From then until the last landing on May 15, when the season closed, the 26 fishermen that fished made a total of 495 landings, with an overall CPUE of 4 and an average weight of 2.75 lb/crab. Price of crab averaged \$6.73/lb, slightly higher than the \$6.47/lb price in 2012, and total exvessel value of the fishery was \$402,256, almost 3 times the amount from 2012 (\$150,569). A total of 22,639 crab (62,179 lb) were harvested, with percentages of crab sold (and CPUE) each month as follows: December 1% (2), January 11% (3), February 25% (4), March 26% (5), April 24% (3), and May 13% (3). Total number of crab harvested was over twice that of 2012 and over six times the average harvest from 1990 to 2012 (Appendix E5). As an indicator of nearshore ice stability, commercial fishermen reported losing 23 out of 389 pots (6%) during the 2012–2013 winter season. Pots were fished from 10 miles east to 18 miles west of Nome, excluding the area closed to commercial fishing from 3.5 miles east to 2.0 miles west of Nome. Similar to years previous to 2012, the majority of crabbers (16) and harvest (93%) came from the Nome area, with the remaining crabbers and harvest coming from Elim, Golovin, Shaktoolik, and St. Michael/Stebbins areas. Unlike last year, ice was unstable in most of eastern Norton Sound; therefore, no crabbers from Unalakleet even registered to fish the winter commercial crab fishery.

The harvest is generally divided between local residents who buy crab directly from the crabbers, the seafood plant (NSSP) in Nome, and other nonlocal markets such as Anchorage. Starting in 2013, the 2 catcher-processors shipped live crab to Korea. Most crabbers consider the winter commercial crabbing a sideline and hold other jobs. Usually, a few of the winter crab fishermen sell the majority of the crab.

## **Subsistence Fishery**

Both a summer and a winter subsistence red king crab fishery occur in Norton Sound, although the majority of the effort and harvest is from the winter fishery (Appendices E5 and E6). During the 2012–2013 winter crab season, all but one of the 149 permits issued were returned, and the 104 permit holders that actually fished reported retaining 7,662 crab, similar to the 7,371 crab retained in 2012. The number caught, which included crab thrown back to the ocean, was 21,752 crab, the greatest amount caught since 1984, when records started to be kept of total crab caught; however, the amount kept was only the eighth highest since 1984. Incidental reports from subsistence crabbers in Nome indicated an unusually high number of smaller male crab and female crab caught in their pots, which resulted in many crab thrown back in the sea. Residents

of Brevig Mission, Diomedes, Elim, Golovin, St. Michael, Shaktoolik, Stebbins, Unalakleet, and White Mountain had a combined harvest of 983 crab, which was 13% of the total harvest. Out of 197 pots reported fishing, 4 (2%) were reportedly lost during the season due to moving ice (Appendix E7). Percentages of subsistence crab harvested each month are as follows: December 2%, January 6%, February 31%, March 35%, April 21%, and May 5%. Like 2012, more than 99% of the crab were caught with pots in 2013 (permit data on file with ADF&G, Division of Commercial Fisheries; Nome).

During the 2013 Norton Sound summer subsistence crab season, 47 permits were issued and 46 were returned. The 26 fishermen that actually fished reported harvesting a total of 1,865 crab, over half of which came from Nome area and over a third of which came from Unalakleet area, with the remainder harvested by villagers from Brevig Mission, Elim, or White Mountain. Crab kept per fisherman averaged 72 crab for summer 2013 (Appendix E6).

### **Sport Fishery**

Sport fishermen can fish for crab, and a harvest log issued by the Nome office similar to a subsistence permit is required. Sport fishermen are only allowed to keep 6 male crab daily, and they must be of legal size (4.75 in or greater). The only recent harvest by sport fishermen was in 2005. That year, 9 harvest logs were issued and 8 were returned, showing that 6 nonresident sport fishermen caught 918 crab and kept 106, for an average harvest of 18 crab per fisherman (permit data on file with ADF&G, Division of Commercial Fisheries; Nome).

### **Future Resource Investigations**

The triennial Norton Sound Trawl Survey will take place in the summer of 2014. Red king crab biomass estimates from the trawl survey are an integral part of the data used in the length-based population model to project the summer king crab legal biomass (during years when no trawl survey occurs) and appropriate GHL for the summer commercial king crab fishery.

A winter pot study that had been conducted annually in nearshore waters of Nome since the early 1980s was replaced with a much larger tagging project in the summer and fall of 2012 and 2013. Results of the winter project have been used in the length-based model to project the summer legal biomass and appropriate GHL for the upcoming summer commercial crab fishery. Size composition by year from the winter king crab project is shown in Appendix E8. This summer tagging project is planned for June 2014, and the results will be compared with previous winter tagging projects.

## **ST. LAWRENCE ISLAND CRAB FISHERY**

### **Commercial Fishery**

In 2006, the BOF split the St. Lawrence Island section between north and south of 66° N latitude. In the northern section, now known as the Kotzebue section, the commercial season was from noon June 15 through August 1. The southern section was merged with Norton Sound section. This change was initiated by Norton Sound area fishermen to expand fishing opportunity to an area with little commercial utilization since 1995. No harvest was reported from this new area in 2013. No permit holders fished in the Kotzebue section in 2013.

## SECTION 5: MISCELLANEOUS SPECIES

---

### INCONNU (SHEEFISH)

#### Commercial Fishery

Although inconnu, commonly known as sheefish, may have been harvested and sold in the winter of 2012–2013, no fish tickets were submitted to ADF&G. In Kotzebue Sound District, no fishermen reported selling inconnu (Appendix F1). Sheefish are not commonly found in either Norton Sound or Port Clarence Districts.

#### Subsistence and Sport Fishery

In 2012 and 2013, there were comprehensive subsistence surveys for fish and wildlife harvests of 6 Kotzebue area villages conducted by the Division of Subsistence. In 2012, surveyed households in 5 Kobuk River villages and Noatak reported harvesting 11,693 sheefish, more than any other year since 1990 (Appendix F2). However, because survey effort was limited during many years, harvest numbers should be considered minimal and are not comparable year to year. Information is not yet available for 2013.

Sport fish harvest reports indicate a harvest of 218 sheefish in Kotzebue Sound District in 2013 (Brendon Scanlon, Sport Fish Biologist, ADF&G, Fairbanks; personal communication). Sheefish sport harvests in the last 10 years have averaged approximately 850 annually (Appendix F3).

#### Escapement

No aerial surveys are flown to determine sheefish escapement. An ADF&G test fishing project on the Kobuk River helps to give an index of abundance, but the test fishery is operated to determine the index of chum salmon abundance and begins operation well after sheefish have begun to pass the site. In 2013, test fishing on Kobuk River resulted in 310 sheefish caught in 208 drifts, for a cumulative CPUE of 330. The CPUE ranked eleventh out of 16 years sheefish catches were recorded (data on file with ADF&G, Division of Commercial Fisheries; Nome).

### DOLLY VARDEN

#### Commercial Fishery

Dolly Varden *Salvelinus malma* are occasionally incidentally caught in commercial salmon fisheries in Norton Sound and Kotzebue Districts. During the 2013 commercial salmon fishery, Kotzebue District reported 302 Dolly Varden caught but not sold, similar to last year when 300 were caught and not sold (Appendix F4).

#### Subsistence and Sport Fishery

Subsistence harvest data for Dolly Varden were not recorded for Norton Sound or Port Clarence, and household surveys for Dolly Varden subsistence catches were not conducted in Arctic communities. A comprehensive survey of fish harvests was done in Kobuk River villages and Noatak by the Division of Subsistence in 2012 and 2013. In 2012, surveyed Noatak households

reported harvesting 6,437 Dolly Varden (Appendix F5). Information is not yet available for 2013.

Sport fish harvest was 1,184 Dolly Varden in Norton Sound and 1,074 Dolly Varden in Kotzebue/Chukchi Sea areas in 2013 (Appendix F3). Overall, Dolly Varden sport fish harvests in the last 10 years in Norton Sound averaged over 3,000 annually, with most fish harvested out of the Unalakleet River (Appendix F6).

## **Escapement**

Dolly Varden escapement is determined from aerial surveys conducted by ADF&G Sport Fish Division in the Kotzebue area, and weir or tower counts in Norton Sound. In 2013, a survey on the Wulik River counted a total of 23,312 Dolly Varden, although the count should be considered a rough estimate because counting conditions were poor due to the presence of river ice (Appendix F7).

## **WHITEFISH**

### **Commercial Fishery**

Commercial whitefish harvest information for the 2012–2013 season was 105 pounds in Norton Sound District (Appendix F9).

### **Subsistence and Sport Fishery**

Subsistence harvest data for whitefish were not recorded for Norton Sound, Port Clarence or Arctic Districts, but a comprehensive survey of fish subsistence harvests by the Division of Subsistence was conducted in Kobuk River villages and Noatak in 2012 and 2013. In 2012, survey data showed that 41,229 whitefish were harvested for 6 villages in Kotzebue District (Appendix F8). Information is not yet available for 2013. Harvest numbers are considered minimal and are not comparable year to year. For the sport fishery, no harvest data are collected in Norton Sound, Port Clarence, or Kotzebue Sound Districts for whitefish.

## **SAFFRON COD**

### **Commercial Fishery**

During the 2012–2013 season, 25 permit holders harvested 33,939 lb of saffron cod *Eleginus gracilis*, commonly known as tomcod, in Norton Sound and sold them to a commercial buyer at \$0.50/lb for use as bait (Appendix F10).

### **Subsistence and Sport Fishery**

In Norton Sound areas tomcod are primarily fished by “jigging” through the ice. Because no subsistence permit is required and a sport fish license is not needed for Alaska residents in northern Norton Sound from Cape Prince of Wales to Bald Head, harvests of tomcod are not reported or documented. In 2013, Norton Sound household subsistence surveys were conducted; however, subsistence harvest information of tomcod was not collected.



## **CAPELIN**

### **Commercial Fishery**

No reported commercial fishery has occurred for capelin *Mallotus villosus*, although there are substantial stocks in northern Norton Sound (Pahlke 1985).

### **Subsistence**

Because no subsistence permit for capelin is required, accurate harvests of capelin are not reported or documented. In 2013, one of the latest capelin spawning events observed on Nome beaches occurred on July 19, compared to mid-June in most years when capelin are observed spawning on Nome beaches. Many residents harvested capelin with various gear, such as nets, buckets, plastic bags, and shovels. No other information on capelin harvest is available.

## **ACKNOWLEDGEMENTS**

Employees of the Alaska Department of Fish and Game, U.S. Fish and Wildlife Service including the Office of Subsistence Management, Bureau of Land Management, National Park Service, Norton Sound Economic Development Corporation, Unalakleet IRA, LGL, and other agencies and organizations worked long and irregular hours at various locations throughout the Norton Sound, Kotzebue, and Port Clarence Areas collecting data presented in this report. We gratefully acknowledge their hard work and funding support.

## REFERENCES CITED

- Barton, L. H. 1978. Finfish resource surveys in Norton Sound and Kotzebue; final report, Alaska Marine Environment Assessment Project, Research Unit 19. Alaska Department of Fish and Game, Division of Commercial Fisheries, AYK Region OCS Report No. 13, Anchorage.
- Bockstoce, J. 1979. The archeology of Cape Nome, Alaska. The University Museum, University of Pennsylvania, Philadelphia.
- Bue, F. J., T. L. Lingnau, C. F. Lean, and E. L. Brennan. 1997. Annual management report 1996, Norton Sound–Port Clarence–Kotzebue. Alaska Department of Fish and Game, Division of Commercial Fisheries Management and Development, Regional Information Report 3A97-30, Anchorage.
- Clark, J. H. 2001. Biological escapement goal for chum salmon in District 1 of Norton Sound. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 3A01-09, Anchorage.
- DeCicco, F. 2001. Fishery management report for sport fisheries in the Northwest Alaska regulatory areas, 1999-2000. Alaska Department of Fish and Game, Fishery Management Report No. 01-01, Anchorage.
- Estensen, J. L., G. L. Todd, and C. S. Monsivais. 2005. Estimation of abundance and distribution of chum salmon in the Unalakleet River drainage, 2004. Alaska Department of Fish and Game, Fishery Data Series No. 05-52, Anchorage.
- Estensen, J. L., and T. Hamazaki. 2007. Estimation of abundance and distribution of chum salmon (*Oncorhynchus keta*) in the Unalakleet River drainage, 2005. Alaska Department of Fish and Game, Fishery Data Series No. 07-03, Anchorage.
- Gaudet, D. M., and G. Schaefer. 1982. Migrations of salmon in Norton Sound, Alaska determined by tagging in 1978-1979. Alaska Department of Fish and Game, Division of Commercial Fisheries, Informational Leaflet No. 198, Juneau.
- Georgette, S., D. Caylor, and S. Tahbone. 2003. Subsistence salmon harvest summary, northwest Alaska 2002. Alaska Department of Fish and Game, Division of Subsistence and Kawerak, Inc.
- Georgette, S., and A. Shiedt. 2005. Whitefish: traditional ecological knowledge and subsistence fishing in the Kotzebue Sound Region, Alaska. Alaska Department of Fish and Game and Maniilaq Association, Technical Paper No. 290, Juneau.
- Joy, P., A. L. J. Brase, and D. J. Reed. 2005. Estimation of coho salmon abundance and spawning distribution in the Unalakleet River 2004. Alaska Department of Fish and Game, Fishery Data Series No. 05-38, Anchorage.
- Joy, P., and D. J. Reed. 2006. Estimation of coho salmon abundance and spawning distribution in the Unalakleet River 2005. Alaska Department of Fish and Game, Fishery Data Series No. 06-38, Anchorage.
- Joy, P., and D. J. Reed. 2007. Estimation of coho salmon abundance and spawning distribution in the Unalakleet River 2004-2006, final report for study 05-101 USFWS, Office of Subsistence Management Fishery Information Service Division. Alaska Department of Fish and Game, Fishery Data Series No. 07-48, Anchorage.
- Kent, S. 2010. Unalakleet River salmon studies, 2002-2008. Alaska Department of Fish and Game, Fishery Data Series No. 10-83, Anchorage.
- Magdanz, J. S., and D. E. Punguk. 1981. Nome River fishery II. Alaska Department of Fish and Game, Division of Subsistence, Technical Paper No. 5, Nome.
- Menard, J., J. Soong, and S. Kent. 2010. 2008 Annual management report Norton Sound, Port Clarence, and Kotzebue. Alaska Department of Fish and Game, Fishery Management Report No. 10-49, Anchorage.
- Menard, J., J. Soong, and S. Kent. 2012. 2011 Annual management report Norton Sound, Port Clarence, and Kotzebue. Alaska Department of Fish and Game, Fishery Management Report No. 12-39, Anchorage.
- Menard, J., J. Soong, S. Kent, and A. Brown. 2013. 2012 Annual management report Norton Sound, Port Clarence, and Kotzebue. Alaska Department of Fish and Game, Fishery Management Report No. 13-28, Anchorage.

## REFERENCES CITED (Continued)

- NPFMC (North Pacific Fisheries Management Council). 2013. Stock assessment and fishery evaluation report for the king and Tanner crab fisheries of the Bering Sea and Aleutian Islands Regions. Stock Assessment and Fishery Evaluation Reports. North Pacific Fishery Management Council. 605 W. 4th Ave., #306, Anchorage.
- Pahlke, K. A. 1985. Preliminary studies of capelin (*Mallotus villosus*) in Alaska waters. Alaska Department of Fish and Game, Division of Commercial Fisheries, Informational Leaflet 250, Juneau.
- Pedersen, S., and A. Linn Jr. 2005. North Slope (Kaktovik) subsistence fish harvest assessment. USFWS Office of Subsistence Management, Fisheries Resource Monitoring Program, Annual Report No. FIS 01-101, Anchorage.
- Pedersen, S., and S. C. Hugo. 2005. North Slope (Anaktuvuk Pass) subsistence fish harvest assessment. USFWS Office of Subsistence Management, Fisheries Resource Monitoring Program, Annual Report No. FIS 02-050-3, Anchorage.
- Ray, D. J. 1975. The Eskimos of Bering Strait, 1650-1898. University of Washington Press, Seattle.
- Scanlon, B. 2009. Fishery management report for sport fisheries in the Northwest/North Slope Management Area, 2008. Alaska Department of Fish and Game, Fishery Management Report No. 09-48, Anchorage.
- Thomas, D. C. 1982. The role of local fish and wildlife resources in the community of Shaktoolik, Alaska. Alaska Department of Fish and Game, Division of Subsistence, Technical Paper No. 13, Nome.
- Todd, G. L., C.S. Monsivais, and D. F. Kaplan. 2005. Estimation of chum salmon abundance, migration timing, and spawning distribution in the Fish River complex, Norton Sound Alaska, 2002-2004. Alaska Department of Fish and Game, Fishery Data Series No. 05-67, Anchorage.
- Wuttig, K. G. 1998. Escapement of Chinook salmon in the Unalakleet River in 1997. Alaska Department of Fish and Game, Fishery Data Series No. 98-08, Anchorage.
- Wuttig, K. G. 1999. Escapement of Chinook salmon in the Unalakleet River in 1998. Alaska Department of Fish and Game, Fishery Data Series No. 99-10, Anchorage.
- Zheng, J., G. H. Kruse, and L. Fair. 1998. Using multiple data sets to assess red king crab *Paralithodes camtschaticus* in Norton Sound, Alaska: a length-based stock synthesis approach. Pages 591-612 [In] F. Funk, T. J. Quinn II, J. Heifetz, J. N. Ianelli, J. E. Powers, J. F. Schweigert, P. J. Sullivan, and C. I. Zhang, editors. Fishery Stock Assessment Models. Alaska Sea Grant College Program Report No. AK-SG-98-01, University of Alaska Fairbanks.



## **TABLES**

Table 1.–Norton Sound commercial salmon harvest summary by subdistrict, 2013.

		Subdistricts						Total
		1	2	3	4	5	6	
Number of Fishermen <sup>a</sup>		1	14	21	18	24	57	124
Chinook <sup>b</sup>	Number	0	0	0	0	0	0	0
	Weight (lbs)	0	0	0	0	0	0	0
Sockeye	Number	0	0	21	4	40	128	193
	Weight (lbs)	0	0	168	30	286	932	1,416
Coho	Number	0	5,362	6,651	5,485	6,890	29,366	53,754
	Weight (lbs)	0	41,189	51,185	40,290	54,052	223,295	410,011
Pink	Number	0	1,180	598	454	14	6,005	8,251
	Weight (lbs)	0	3,039	1,708	1,613	45	17,719	24,124
Chum	Number	0	3,113	1,412	36,021	23,268	54,858	118,672
	Weight (lbs)	0	22,144	10,164	252,242	162,299	374,082	820,931
Total	Number	0	9,655	8,682	41,964	30,212	90,357	180,870
	Weight (lbs)	0	66,372	63,225	294,175	216,682	616,028	1,256,482

*Note:* An additional 151 Chinook, 54 sockeye, 48 coho, 87 pink, and 37 chum salmon were retained for personal use. Average commercial weights by species were 7.34 lb for sockeye salmon, 7.63 lb for coho salmon, 2.92 lb for pink salmon, and 6.92 lb for chum salmon.

<sup>a</sup> Number of fishermen is unique number of permit holders that fished in each subdistrict. Some permit holders fished in more than 1 subdistrict. Subdistrict 1 catch is confidential and is not included in total.

<sup>b</sup> The buyer did not buy Chinook salmon in 2013; all were retained for personal use.

Table 2.–Subsistence salmon harvest for northern Norton Sound, 2013.

	Permits Fished <sup>a</sup>	Number of salmon harvested					Total
		Chinook	Sockeye	Coho	Pink	Chum	
Marine Waters	50	36	161	609	372	2,194	3,372
Bonanza River	14	0	0	76	41	140	257
Cripple Creek	6	0	0	19	0	0	19
Eldorado River – above weir	1	0	0	1	0	0	1
Eldorado River – below weir	17	1	0	108	56	344	509
Flambeau River	4	0	0	19	1	25	45
Safety Sound	1	0	0	0	0	0	0
Nome River – above weir	11	0	1	9	22	4	36
Nome River – below weir	162	10	18	480	211	177	896
Penny River	9	0	0	27	43	0	70
Sinuk River	34	0	26	80	19	43	168
Snake River – above weir	0	0	0	0	0	0	0
Snake River – below weir	74	1	4	301	80	134	520
Solomon River	14	0	1	75	0	4	80
<b>Nome Subdistrict Total <sup>b</sup></b>	<b>302</b>	<b>48</b>	<b>211</b>	<b>1,804</b>	<b>845</b>	<b>3,065</b>	<b>5,973</b>
<b>Cape Woolley <sup>c</sup></b>	<b>3</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>0</b>	<b>1</b>	<b>1</b>
Marine Waters	16	40	12	79	93	929	1,153
Kachavik River	8	2	1	7	1,627	834	2,471
McKinley River	5	0	0	24	2	0	26
Chinik Creek	5	1	0	88	11	0	100
Fish River	56	4	0	601	1,690	1400	3,695
Niukluk River	21	0	2	141	232	92	467
Other Creeks/Rivers	3	0	0	24	0	1	25
<b>Golovin Subdistrict Total <sup>d</sup></b>	<b>97</b>	<b>47</b>	<b>15</b>	<b>964</b>	<b>3,655</b>	<b>3,256</b>	<b>7,937</b>
Marine Waters	15	5	4	523	342	592	1,466
Kwiniuk River – above tower	3	0	0	11	70	0	81
Kwiniuk River – below tower	35	18	7	649	572	515	1,761
Next Creek	1	0	0	0	0	0	0
Tubutulik River	10	12	0	265	134	110	521
Iron Creek	5	0	0	67	16	1	84
Other Creeks/Rivers	2	4	4	0	0	0	8
<b>Elim Subdistrict Total <sup>e</sup></b>	<b>45</b>	<b>39</b>	<b>15</b>	<b>1,515</b>	<b>1,134</b>	<b>1,218</b>	<b>3,921</b>
Port Clarence – Marine Waters	93	36	2,041	609	1,587	4,402	8,675
Tuksuk Channel	13	2	428	34	133	1,446	2,043
Imuruk Basin	0	0	0	0	0	0	0
Agiapuk River	3	0	0	0	20	448	468
Kuzitrin River	6	0	9	0	0	10	19
Salmon Lake	2	0	4	0	0	0	4
Pilgrim River – above weir	62	0	807	8	6	172	993
Pilgrim River – below weir	89	0	1,954	0	42	110	2,106
<b>Port Clarence District Total <sup>f</sup></b>	<b>265</b>	<b>38</b>	<b>5,243</b>	<b>651</b>	<b>1,788</b>	<b>6,588</b>	<b>14,308</b>
<b>Total</b>	<b>712</b>	<b>172</b>	<b>5,484</b>	<b>4,934</b>	<b>7,422</b>	<b>14,128</b>	<b>32,140</b>

<sup>a</sup> There were 7 locations where subsistence permits were issued in 2013 for northern Norton Sound: 1-Nome Subdistrict; 2-Cape Woolley; 3-Golovin Subdistrict; 4-Elim Subdistrict; 5-Pilgrim River; 6-Salmon Lake; and 7-Port Clarence District. Except for Pilgrim River and Salmon Lake, each permit is valid for both marine and fresh waters. Permits fished include those permit holders who fished but reported no harvest.

<sup>b</sup> All 477 Nome Subdistrict permits issued were returned.

<sup>c</sup> All 19 Cape Woolley permits issued were returned.

<sup>d</sup> All 153 Golovin Subdistrict permits issued were returned.

<sup>e</sup> All 64 Elim Subdistrict permits issued were returned.

<sup>f</sup> All 265 Pilgrim River and 4 Salmon Lake permits issued were returned. All 162 Port Clarence District permits issued were returned.

Table 3.—Salmon counts of rivers and associated salmon escapement goal ranges (SEG, BEG or OEG), Norton Sound and Port Clarence, 2013.

Stream	Chinook Salmon				Chum Salmon				
	Weir/ Tower Count	Escapement Goal Range	Aerial Survey Count <sup>a</sup>	Escapement Goal Range	Weir/ Tower Count	Escapement Goal Range	Aerial Survey Count <sup>a</sup>	Aerial Survey Expansion	Escapement Goal Range
Salmon L.			0				0		
Grand Central R.			0				0		
Agiapuk R.									
American R.									
Pilgrim R.	47				47,557				
Glacial L.	0				35				
Sinuk R.			1				19,500	31,691	
Cripple R.									
Penny R.									
Anvil Creek									
Dry Creek									
Snake R.	8				2,755	1,600–2,500 <sup>b</sup>			
Nome R.	9				4,811	2,900–4,300 <sup>b</sup>			
Flambeau R.							16,088	27,928	
Eldorado R.	9				27,928	6,000–9,200 <sup>b</sup>	16,859		
Bonanza R.							5,284	13,437	
Solomon R.	0				1,377		156		
Nome Subdistrict						23,000–35,000 <sup>c</sup>		108,120	
<u>Fish R.</u>			15	Combined			2,550		
<u>Boston Cr.</u>			19	100–250			16,100		
Niukluk R.			68			23,000	17,203		
Ophir Cr.									
Kwiniuk R.	15	300–550			5,631	11,500–23,000 <sup>d</sup>			
Tubutulik R.			2			9,200–18,400 <sup>e</sup>	4,532		
Ungalik R.			17				28,283		
Inglutalik R.	860				61,259		24,562		
Pikmiktalik R.									
Shaktoolik R.			151	400–800			11,878		
<u>Unalakleet R.</u>	767			Combined	113,953				Combined
<u>Old Woman R.</u>			168	550–1,100			367		2,400–4,800
North R.	564		339	1,200–2,600	10,518		2,425		

-continued-



Table 3.—Page 2 of 2.

Stream	Coho Salmon			Sockeye Salmon			Pink Salmon		
	Weir/ Tower Count	Aerial Survey Count <sup>a</sup>	Escapement Goal Range	Weir/ Tower Count	Aerial Survey Count <sup>a</sup>	Escapement Goal Range	Weir/ Tower Count	Escapement Goal Range	Aerial Survey Count <sup>a</sup>
Salmon L.		0			5,820	Combined			0
Grand Central R.		0			1,151	4,000–8,000			0
Agiapuk R.									
American R.									
Pilgrim R.	890			12,428			1,060		
Glacial L.				2,544	1,366	800–1,600	2		
Sinuk R.		1,054			39				23,000
Cripple R.									
Penny R.									
Anvil Creek									
Dry Creek									
Snake R.	1,203			163			1,333		
Nome R.	2,624			38			10,257	3,200	
Flambeau R.									
Eldorado R. <sup>b</sup>	15			0			1,025		
Bonanza R.		945							800
Solomon R.	178			3			2,733		
Fish R.									
Boston Cr.			Tower Goal						
Niukluk R.		2,279	2,400–7,200		0				9,700
Ophir Cr.		74							
Kwiniuk R.	3,940		650–1,300	0			13,212	8,400	
Tubutulik R.							700		700
Ungalik R.		1,063							49,890
Inglutalik R.	5,904			0			268,537		109,980
Pikmiktalik R.									
Shaktoolik R.		271					10,400		
Unalakleet R.	25,566			243			144,225		
Old Woman R.									
North R.	8,834	867	550–1,100	12			46,668	25,000	5,025

Note: Data not available for all streams. Sustainable escapement goal (SEG), biological escapement goal (BEG), and optimal escapement goal (OEG) are listed.

<sup>a</sup> All aerial surveys are rated fair to good, unless otherwise noted.

<sup>b</sup> The Alaska Board of Fisheries (BOF) also established an OEG with the same range as the BEG.

<sup>c</sup> BOF-established OEG is the same range as the BEG and is based on a combination of weir counts and expanded aerial survey counts.

<sup>d</sup> This represents the OEG in regulation. The BEG is 10,000–20,000 for the Kwiniuk River and 8,000–16,000 for the Tubutulik River.

<sup>e</sup> The goal listed is actual fish and not aerial counts. However, at this time there is no counting project on the river.

Table 4.–Commercial salmon set gillnet catches from Golovin, Subdistrict 2, Norton Sound, 2013.

Period	Target species	Dates fished	Length (hours)	Permits fished	Chinook harvest	Chum harvest	Pink harvest	Sockeye harvest	Coho harvest
1	Pink	7/17–7/19	48	8	0	506	1,028	0	10
2	Chum	7/20–7/22	48	10	0	1,771	146	0	188
3	Chum	7/23–7/24	24	9	0	301	6	0	130
4	Coho	8/01–8/02	24	7	0	174	0	0	527
5	Coho	8/05–8/07	36	9	0	204	0	0	861
6	Coho	8/08–8/10	36	7	0	104	0	0	1,476
7	Coho	8/11–8/13	36	11	0	23	0	0	940
8	Coho	8/14–8/16	48	10	0	26	0	0	609
9	Coho	8/19–8/21	48	7	0	2	0	0	406
10	Coho	8/23–8/25	48	8	0	2	0	0	215
Totals			396	14	0	3,113	1,180	0	5,362

Table 5.–Commercial salmon set gillnet catches from Elim, Subdistrict 3, Norton Sound, 2013.

Period	Target species	Dates fished	Length (hours)	Permits fished	Chinook harvest	Chum harvest	Pink harvest	Sockeye harvest	Coho harvest
1	Chum	7/20–7/22	48	6	0	618	184	2	23
2	Chum	7/24–7/25	24	11	0	232	300	1	141
3	Coho	8/01–8/02	24	12	0	54	36	0	457
4	Coho	8/05–8/07	36	12	0	86	43	1	537
5	Coho	8/08–8/10	36	14	0	46	20	0	944
6	Coho	8/11–8/13	36	15	0	45	9	6	750
7	Coho	8/14–8/16	48	18	0	40	1	0	880
8	Coho	8/19–8/21	48	15	0	70	0	1	870
9	Coho	8/23–8/25	48	16	0	97	2	2	884
10	Coho	8/27–8/29	48	16	0	105	2	8	745
11	Coho	8/30–8/31	40	9	0	19	1	0	420
Totals			436	21	0	1,412	598	21	6,651

*Note:* An additional 6 Chinook, 22 chum, 3 pink, 6 sockeye, and 24 coho salmon were retained for personal use in 2013.

Table 6.—Commercial salmon set gillnet catches from Norton Bay, Subdistrict 4, Norton Sound, 2013.

Period	Target species	Dates fished	Length (hours)	Permits fished	Chinook harvest	Chum harvest	Pink harvest	Sockeye harvest	Coho harvest
1	Chum	6/25–6/27	48	5	0	1,388	0	0	0
2	Chum	6/28–6/30	48	8	0	4,323	0	0	0
3	Chum	7/02–7/04	48	9	0	5,137	0	0	0
4	Chum	7/05–7/07	48	12	0	5,755	0	0	0
5	Chum	7/09–7/11	48	9	0	3,660	0	0	2
6	Chum	7/12–7/15	48	11	0	4,350	0	0	5
7	Chum	7/16–7/19	72	9	0	3,280	0	2	91
8	Chum	7/20–7/23	72	3	0	1,935	131	0	228
9	Chum	7/25–7/27	72	12	0	1,948	308	0	821
10	Chum	7/28–7/31	64	10	0	1,513	8	0	1,220
11	Chum	8/02–8/04	72	13	0	689	4	0	556
12	Coho	8/05–8/07	48	6	0	467	0	0	569
13	Coho	8/09–8/11	48	11	0	528	3	0	758
14	Coho	8/12–8/14	48	7	0	349	0	0	337
15	Coho	8/15–8/17	48	6	0	247	0	0	391
16	Coho	8/19–8/21	48	6	0	221	0	0	333
17	Coho	8/23–8/25	48	4	0	179	0	0	148
18	Coho	8/27–8/29	48	3	0	52	0	2	26
19	Coho	8/30–9/01	48			No One Fished			
20	Coho	9/03–9/05	48			No One Fished			
Totals			1,072	18	0	36,021	454	4	5,485

Note: An additional 8 Chinook and 33 pink salmon were retained for personal use in 2013.

Table 7.—Commercial salmon set gillnet catches from Shaktoolik, Subdistrict 5, Norton Sound, 2013.

Period	Target species	Dates fished	Length (hours)	Permits fished	Chinook harvest	Chum harvest	Pink harvest	Sockeye harvest	Coho harvest
1	Chum	7/02–7/03	24			No One Fished			
2	Chum	7/05–7/06	24	6	0	1,866	0	0	0
3	Chum	7/08–7/10	48	14	0	4,769	0	2	1
4	Chum	7/12–7/14	48	18	0	8,308	0	12	3
5	Chum	7/15–7/18	72	15	0	2,177	0	6	81
6	Chum	7/19–7/22	72	1	0	320	0	0	16
7	Chum	7/23–7/26	72	14	0	1,393	0	6	420
8	Chum	7/27–7/30	72	16	0	1,480	14	7	794
9	Coho	7/31–8/02	48	15	0	989	0	3	522
10	Coho	8/04–8/06	48	6	0	342	0	1	286
11	Coho	8/07–8/09	48	11	0	365	0	0	485
12	Coho	8/11–8/13	48	16	0	313	0	2	480
13	Coho	8/14–8/16	48	14	0	308	0	0	514
14	Coho	8/18–8/20	48	17	0	432	0	1	1,354
15	Coho	8/21–8/23	48	11	0	78	0	0	517
16	Coho	8/25–8/27	48	11	0	67	0	0	851
17	Coho	8/28–8/30	48	18	0	55	0	0	554
18	Coho	9/01–9/03	48	17	0	6	0	0	12
19	Coho	9/04–9/06	48			No One Fished			
Totals			960	21	0	23,268	14	40	6,890

*Note:* An additional 6 Chinook and 5 sockeye salmon were retained for personal use in 2013.

Table 8.—Commercial salmon set gillnet catches from Unalakleet, Subdistrict 6, Norton Sound, 2013.

Period	Target species	Dates fished	Length (hours)	Permits fished	Chinook harvest	Chum harvest	Pink harvest	Sockeye harvest	Coho harvest
1	Chum	7/01–7/02	24	19	0	5,804	0	0	0
2	Chum	7/05–7/06	24	17	0	5,602	287	0	0
3	Chum	7/08–7/09	24	21	0	1,103	121	2	1
4	Chum	7/11–7/13	48	16	0	8,831	1,514	32	31
5	Chum	7/15–7/18	72	0	0	8,279	1,648	29	276
6	Chum	7/19–7/22	72	19	0	8,123	1,377	9	939
7	Chum	7/23–7/26	72	27	0	4,670	434	19	1,909
8	Chum	7/27–7/30	72	27	0	3,687	309	10	3,257
9	Coho	7/31–8/02	48	30	0	1,318	161	3	1,215
10	Coho	8/04–8/06	48	27	0	1,928	39	7	1,692
11	Coho	8/07–8/09	48	28	0	880	22	2	1,964
12	Coho	8/11–8/13	48	36	0	1,161	49	2	3,715
13	Coho	8/14–8/16	48	34	0	567	13	0	1,852
14	Coho	8/18–8/20	48	36	0	983	19	3	3,914
15	Coho	8/21–8/23	48	33	0	686	0	5	2,551
16	Coho	8/25–8/27	48	18	0	685	4	2	4,088
17	Coho	8/28–8/30	48	25	0	391	6	1	1,328
18	Coho	9/01–9/03	48	33	0	91	1	1	403
19	Coho	9/04–9/06	48	32	0	69	1	1	231
Totals			936	55	0	54,858	6,005	128	29,366

*Note:* There were an additional 131 Chinook, 15 chum, 51 pink, 43 sockeye, and 24 coho salmon retained for personal use in 2013.

Table 9.–Kotzebue District commercial chum salmon catch and average weight by date, 2013.

Date	Number of fishermen	Catch	Pounds	Average weight
7/10	10	1,360	10,870	8.0
7/11	12	1,535	12,321	8.0
7/12	9	2,368	19,386	8.2
7/14	10	2,077	16,942	8.2
7/15	17	2,038	16,191	7.9
7/16	18	5,994	48,926	8.2
7/17	23	4,908	39,714	8.1
7/18	29	10,476	86,911	8.3
7/19	26	6,013	49,559	8.2
7/21	33	5,585	46,119	8.3
7/22	34	6,216	51,847	8.3
7/23	38	17,712	145,926	8.2
7/24	9	2,683	21,022	7.8
7/25	41	11,894	98,133	8.3
7/26	40	9,082	75,349	8.3
7/28	41	17,283	141,096	8.2
7/30	47	10,927	89,513	8.2
7/31	45	15,773	130,429	8.3
8/01	38	10,062	83,466	8.3
8/04	38	9,453	75,974	8.0
8/05	40	9,672	78,190	8.1
8/06	34	7,300	60,105	8.2
8/07	41	12,427	99,541	8.0
8/08	43	10,875	85,858	7.9
8/11	35	11,830	92,091	7.8
8/12	40	11,067	87,602	7.9
8/13	40	10,917	86,257	7.9
8/14	44	12,057	93,852	7.8
8/15	44	15,805	125,682	8.0
8/16	39	8,385	65,160	7.8
8/18	35	9,706	76,075	7.8
8/19	35	4,229	32,865	7.8
8/20	31	3,210	24,603	7.7
8/21	18	3,160	24,213	7.7
8/22	29	6,134	47,977	7.8
8/23	14	2,779	20,764	7.5
8/25	11	2,180	16,085	7.4
8/26	6	113	768	6.8
8/27	22	439	3,033	6.9
8/28	17	792	5,535	7.0
8/29	14	1,142	8,168	7.2
8/30	8	0	0	
8/31	8	0	0	
Total	65	297,658	2,394,118	8.0

*Note:* Catch by date is for Great Pacific Seafoods only. Maniilaq had a catch of 21,312 chum salmon (160,952 pounds) by 21 permit holders. There was one catcher–seller who reported a catch of 25 chum salmon (230 pounds). Both buyers and the catcher–seller signed waivers allowing ADF&G to report their harvest figures. Also harvested during the 2013 commercial fishery and kept for personal use were 16 Chinook, 67 chum, 13 sockeye, 42 pink, and 43 coho salmon, and 302 Dolly Varden, 705 sheefish, and 50 whitefish.

Table 10.—Historical chum salmon catch for Kobuk River drift test fishery, 1993–2013.

Year	Dates of operation	Number of drifts	Cumulative CPUE <sup>a</sup>	Midpoint date
1993	7/12–8/12	164	494	8/03
1994	7/13–8/30	248	1,207	8/04
1995	7/12–8/16	196	1,188	8/02
1996	7/09–8/14	208	2,581	7/31
1997	7/09–8/14	202	797	8/03
1998	7/10–8/15	182	538	7/29
1999	7/11–8/13	176	1,357	8/02
2000	7/07–8/14	228	1,481	8/01
2001	7/05–8/13	232	1,575	7/26
2002	7/05–8/12	218	875	7/23
2003	7/09–8/13	214	749	8/02
2004	7/02–8/12	242	855	8/05
2005	7/07–8/15	207	1,207	8/06
2006	7/07–8/19	217	743	8/16
2007	7/11–8/20	207	1,342	8/09
2008	7/09–8/14	200	2,269	7/30
2009	7/10–8/20	242	971	8/06
2010	7/15–8/24	234	1,401	8/05
2011	7/13–8/21	220	2,499	8/10
2012	7/17–8/16	151	2,398	8/08
2013	7/17–8/25	208	2,698	8/06

<sup>a</sup> Cumulative catch per unit of effort (CPUE) is calculated as the sum of daily CPUE during the period of data collection, and daily CPUE (I) is calculated as the number of fish that would have been caught if 100 fathoms of gillnet had been fished for 60 minutes.  $I = (6,000 * C) / (L * T)$ , where  $C$  = number of chum salmon caught,  $L$  = length of gillnet in fathoms, and  $T$  = mean fishing time in minutes.

Table 11.—Commercial herring sac roe harvest summary by subdistrict, Norton Sound District, 2013.

	St. Michael Subdistrict (333-70)			Unalakleet Subdistrict (333-72)			Cape Denbigh Subdistrict (333-74)			Norton Sound District Total		
	Number of permit holders	Sac roe short tons	Percent roe	Number of permit holders	Sac Roe short tons	Percent roe	Number of permit holders	Sac Roe short tons	Percent roe	Number of permit holders	Sac Roe short tons	Percent roe
Total	27	107.1	13.6	17	82	13.4	17	301.5	13.0	40	490.6	13.2



Table 12.—Daily catch for the open access and CDQ summer commercial king crab harvests, Norton Sound Section, Eastern Bering Sea, July 3–September 14, 2013.

Date <sup>a</sup>	Landings	Number of crab	Crab harvested (lbs)	Cumulative total (lbs)	Number pots Pulled	Average weight (lbs)	CPUE
Open Access							
7/05	1	144	453	453	20	3.1	7
7/06	4	755	2,187	2,640	113	2.9	7
7/07	9	3,036	9,124	11,764	359	3.0	8
7/08	3	880	2,591	14,355	89	2.9	10
7/09	3	374	1,097	15,452	57	2.9	7
7/10	2	666	2,016	17,468	79	3.0	8
7/11	9	3,189	9,438	26,906	305	3.0	10
7/12	11	4,151	12,604	39,510	391	3.0	11
7/13	5	1,144	3,417	42,927	159	3.0	7
7/14	3	1,446	4,537	47,464	120	3.1	12
7/15	1	269	823	48,287	40	3.1	7
7/16	10	3,999	12,319	60,606	328	3.1	12
7/17	2	195	562	61,168	41	2.9	5
7/18	17	6,871	20,496	81,664	596	3.0	12
7/19	11	2,976	8,777	90,441	296	2.9	10
7/20	4	1,884	5,329	95,770	160	2.8	12
7/21	3	651	1,912	97,682	118	2.9	6
7/23	3	776	2,360	100,042	81	3.0	10
7/24	15	4,314	12,853	112,895	560	3.0	8
7/25	12	3,423	10,356	123,251	399	3.0	9
7/26	4	1,313	3,971	127,222	131	3.0	10
7/27	3	502	1,657	128,879	32	3.3	16
7/28	12	2,896	8,578	137,457	394	3.0	7
7/29	2	304	903	138,360	60	3.0	5
7/30	15	3,071	9,096	147,456	468	3.0	7
7/31	9	1,905	5,690	153,146	327	3.0	6
8/01	6	876	2,661	155,807	160	3.0	5
8/02	8	2,091	6,199	162,006	356	3.0	6
8/03	12	2,427	7,117	169,123	351	2.9	7
8/04	11	2,971	8,748	177,871	336	2.9	9
8/05	1	311	929	178,800	39	3.0	8
8/06	6	1,846	5,411	184,211	210	2.9	9
8/07	5	1,529	4,452	188,663	134	2.9	11
8/08	4	696	2,096	190,759	158	3.0	4
8/09	7	2,815	8,209	198,968	280	2.9	10
8/10	9	1,969	5,955	204,923	320	3.0	6
8/11	10	3,727	11,093	216,016	369	3.0	10
8/12	4	1,168	3,581	219,597	117	3.1	10

-continued-

Table 12.–Page 2 of 3.

Date <sup>a</sup>	Landings	Number of crab	Crab harvested (lbs)	Cumulative total (lbs)	Number pots pulled	Average weight (lbs)	CPUE
Open Access							
8/13	6	1,917	5,714	225,311	194	3.0	10
8/14	8	2,153	6,368	231,679	216	3.0	10
8/15	10	2,468	7,439	239,118	371	3.0	7
8/16	9	1,811	5,326	244,444	285	2.9	6
8/17	11	1,369	4,002	248,446	326	2.9	4
8/18	3	1,146	3,422	251,868	160	3.0	7
8/19	3	451	1,317	253,185	80	2.9	6
8/20	7	1,130	3,365	256,550	252	3.0	4
8/21	11	1,317	3,896	260,446	372	3.0	4
8/22	3	331	997	261,443	120	3.0	3
8/23	1	4	13	261,456	40	3.3	0
8/24	3	616	2,076	263,532	88	3.4	7
8/25	9	1,069	3,244	266,776	243	3.0	4
8/27	3	396	1,170	267,946	69	3.0	6
8/28	10	2,359	7,181	275,127	351	3.0	7
8/29	5	1,403	4,223	279,350	135	3.0	10
8/30	2	915	2,784	282,134	78	3.0	12
8/31	5	2,798	8,568	290,702	189	3.1	15
9/04	8	3,899	12,228	302,930	270	3.1	14
9/05	3	1,777	5,237	308,167	71	2.9	25
9/06	6	2,617	7,694	315,861	200	2.9	13
9/07	8	2,851	8,430	324,291	237	3.0	12
9/08	4	1,485	4,440	328,731	131	3.0	11
9/09	6	2,195	6,817	335,548	204	3.1	11
9/10	12	2,920	8,826	344,374	314	3.0	9
9/11	5	1,368	4,065	348,439	155	3.0	9
9/12	1	123	367	348,806	12	3.0	10
9/13	6	2,854	8,660	357,466	177	3.0	16
9/14	3	1,747	5,386	362,852	91	3.1	19
9/15	8	3,436	10,426	373,278	212	3.0	16
Total	435	124,485	373,278	373,278	14,196	3.0	9

-continued-

Table 12.–Page 3 of 3.

Date <sup>a</sup>	Landings	Number of crab	Crab harvested (lbs)	Cumulative total (lbs)	Number pots pulled	Average weight (lbs)	CPUE
CDQ							
7/16	1	734	2,266	2,266	40	3.1	18
7/18	1	272	799	3,065	40	2.9	7
7/20	1	126	373	3,438	40	3.0	3
7/23	1	318	929	4,367	40	2.9	8
7/25	1	123	362	4,729	40	2.9	3
7/26	1	289	774	5,503	40	2.7	7
7/28	1	326	973	6,476	40	3.0	8
7/31	1	323	957	7,433	40	3.0	8
8/07	1	465	1,336	8,769	40	2.9	12
8/08	1	186	550	9,319	40	3.0	5
8/09	1	284	852	10,171	40	3.0	7
8/12	1	204	639	10,810	40	3.1	5
8/14	1	100	289	11,099	40	2.9	3
8/19	1	138	423	11,522	40	3.1	3
8/21	1	15	45	11,567	20	3.0	1
8/25	2	226	647	12,214	63	2.9	4
8/29	1	28	75	12,289	17	2.7	2
9/01	1	83	259	12,548	26	3.1	3
9/04	1	512	1,744	14,292	39	3.4	13
9/05	1	47	158	14,450	15	3.4	3
9/07	1	258	795	15,245	39	3.1	7
9/09	1	323	1,049	16,294	42	3.2	8
9/13	1	606	1,874	18,168	31	3.1	20
9/15	1	132	417	18,585	10	3.2	13
Total	25	6,118	18,585	18,585	862	3.0	7

Source: Fish ticket data.

<sup>a</sup> Both the open access and CDQ (community development quota) fisheries closed by emergency order 9/14, and the last deliveries were made 9/15.

Table 13.—Commercial harvest of red king crab from Norton Sound Section by statistical area, Norton Sound District, 2013.

Statistical Area	Number of crab <sup>a</sup>	Crab harvested (lbs.)	Number pots pulled	CPUE	Average weight (lbs)
616401	2,654	7,729	174	15	2.91
626331	230	686	58	4	2.98
626401	12,260	36,802	2,117	6	3.00
636330	4,129	12,035	492	8	2.91
636401	11,486	34,027	1,912	6	2.96
646330	1,446	4,195	173	8	2.90
646401	19,767	59,737	1,469	13	3.02
646402	1,803	5,271	148	12	2.92
656330	2,770	8,515	324	9	3.07
656401	49,031	147,569	5,696	9	3.01
656402	12,623	37,743	1,404	9	2.99
666401	10,938	33,469	918	12	3.06
666402	470	1,419	93	5	3.02
666431	996	2,669	80	12	2.68
Total	130,603	391,863	15,058	9	3.00

*Note:* Data for summer fishery only. CPUE is catch per unit of effort.

<sup>a</sup> Includes 6,118 crab (18,585 lb) from the CDQ (community development quota) fishery.

## **APPENDIX A: NORTON SOUND FISHERIES**

Appendix A1.—Commercial salmon catch by species, Norton Sound District, 1990–2013.

Year	Chinook	Sockeye	Coho	Pink	Chum	Total
1990	8,895	434	56,712	501	65,123	131,665
1991	6,068	203	63,647	0	86,871	156,789
1992	4,541	296	105,418	6,284	83,394	199,933
1993	8,972	279	43,283	157,574	53,562	263,670
1994	5,285	80	102,140	982,389	18,290	1,108,184
1995	8,860	128	47,862	81,644	42,898	181,392
1996	4,984	1	68,206	487,441	10,609	571,241
1997	12,573	161	32,284	20	34,103	79,141
1998	7,429	7	29,623	588,013	16,324	641,396
1999	2,508	0	12,662	0	7,881	23,051
2000	752	14	44,409	166,548	6,150	217,873
2001	213	44	19,492	0	11,100	30,849
2002	5	1	1,759	0	600	2,365
2003	12	21	17,060	0	3,560	20,653
2004 <sup>a</sup>	22	47	42,016	0	6,296	48,381
2005	151	12	85,523	0	3,983	89,669
2006	20	3	130,808	0	10,042	140,873
2007	19	2	126,136	3,769	22,431	152,357
2008	83	60	120,309	75,525	25,124	221,101
2009 <sup>a</sup>	84	126	87,041	17,364	34,122	138,737
2010	140	103	62,079	31,557	117,743	211,622
2011	185	369	58,917	7,141	110,555	177,167
2012 <sup>a</sup>	197	134	37,056	205,498	62,772	305,657
2013	151	247	53,802	8,338	118,709	181,247
Avg 2008–2012	138	158	73,080	67,417	70,063	210,857
Avg 2003–2012	91	88	76,695	34,085	39,663	150,622

*Note:* Harvest numbers may include a small number of salmon reported on fish tickets that were retained for personal use and not commercially sold.

<sup>a</sup> All Chinook salmon caught were retained for personal use and not sold.

Appendix A2.–Number of commercial salmon permits fished, Norton Sound, 1990–2013.

Year	Subdistrict						District Total <sup>a</sup>
	1	2	3	4	5	6	
1990	0	15	23	0	28	73	128
1991	0	16	24	0	25	75	126
1992	2	1	21	9	25	71	110
1993	1	8	26	15	37	66	153
1994	1	5	21	0	39	71	119
1995	2	7	12	0	26	58	105
1996	1	4	12	0	20	54	86
1997	0	11	21	9	19	57	102
1998	0	16	23	0	28	52	82
1999	0	0	0	0	15	45	60
2000	0	12	13	0	26	49	79
2001	0	5	5	0	13	29	51
2002	0	0	0	0	7	5	12
2003	0	0	0	0	10	20	30
2004	0	0	0	0	11	25	36
2005	0	0	0	0	12	28	40
2006	0	0	0	0	22	40	61
2007	0	0	11	0	15	47	71
2008	0	4	12	4	23	58	91
2009	0	5	17	7	21	49	88
2010	0	10	19	5	35	59	115
2011	0	13	32	12	30	65	123
2012	0	14	24	18	21	55	123
2013	1	14	21	18	24	57	124
Avg 2008–2012	0	9	21	9	26	57	108
Avg 2003–2012	0	5	12	5	20	45	78

<sup>a</sup> District total is the number of fishermen that actually fished in Norton Sound; some fishermen may have fished more than one subdistrict.

Appendix A3.—Round weight and value of commercially caught salmon by species, Norton Sound District, 1990–2013.

Year	Pounds caught (Round wt. in lbs)				Salmon roe (lbs)	Value of catch (\$)
	Chinook	Coho	Pink	Chum		
1990	168,745	426,902	<sup>a</sup>	482,060	75	474,064
1991	107,541	469,495	<sup>a</sup>	597,272	221	413,479
1992	57,571	820,406	18,230	595,345	2,641	448,395
1993	151,504	287,702	406,820	347,072	2,608	368,723
1994	98,492	766,050	2,185,066	122,540	0	863,060
1995	174,771	356,190	198,121	290,445	0	356,164
1996	95,794	573,372	1,196,115	84,349	0	340,347
1997	225,136	235,517	50	253,006	880	363,908
1998	127,831	232,705	1,330,624	106,687	0	358,982
1999	48,421	88,037	0	57,656	0	76,860
2000	11,240	307,565	369,800	40,298	0	149,907
2001	3,803	152,293	0	79,558	0	56,921
2002	50	12,972	0	4,555	0	2,941
2003	136	139,775	0	23,687	0	64,473
2004	0	302,379	0	42,385	0	122,506
2005	2,511	659,278	0	28,071	0	296,154
2006	167	869,427	0	68,500	0	389,707
2007	206	1,002,078	10,537	151,386	0	572,195
2008	970	855,980	187,979	171,151	0	759,451
2009	0	679,416	46,698	240,502	0	722,167
2010	1,697	472,939	87,954	799,550	0	1,220,487
2011	1,659	438,481	19,768	774,906	0	1,269,730
2012	0	245,078	492,372	425,233	0	758,908
2013	0	410,791	24,201	823,453	0	1,183,236

<sup>a</sup> Information not available.



Appendix A4.—Estimated mean prices paid to commercial salmon fishermen in dollars, Norton Sound District, 1990–2013.

Year	Chinook	Coho	Pink	Chum	Sockeye
1990	1.01	0.50	(0.75 for roe)	0.23	
1991	0.87	0.36 (3.00 for roe)		0.27 (3.00 for roe)	
1992	0.66	0.33 (1.50 for roe)	0.16	0.22	
1993	0.72	0.22 (1.76 for roe)	0.15	0.24	0.40
1994	1.02	0.52	0.15	0.29	
1995	0.66	0.43	0.18	0.18	
1996	0.54	0.28	0.10	0.08	
1997	1.00	0.47	0.06	0.11	
1998	0.74	0.29	0.14	0.09	
1999	0.82	0.35		0.11	
2000	1.30	0.30	0.10	0.15	
2001	1.00	0.25		0.19	0.37
2002	0.39	0.20		0.07	
2003	0.64	0.44		0.14	0.45
2004		0.39		0.14	
2005	1.22	0.44		0.15	0.45
2006	1.49	0.44		0.14	
2007	0.55	0.53	0.14	0.24	0.55
2008	0.73	0.77	0.23	0.34	0.56
2009		0.93	0.18	0.33	0.34
2010	2.25	1.47	0.32	0.62	0.63
2011	3.01	1.70	0.25	0.68	1.04
2012		1.47	0.36	0.52	1.45
2013		1.77	0.22	0.55	1.49
Avg 2008–2012	1.50	1.59	0.34	0.62	1.01

*Note:* Blank cells indicate no known purchases were made.

Appendix A5.—Mean commercial salmon harvest weights, Norton Sound District, 1990–2013.

Year	Mean round weight in pounds <sup>a</sup>			
	Chinook	Coho	Pink	Chum
1990	19.0	7.5	NA	7.4
1991	17.7	7.4	<sup>c</sup>	6.9
1992 <sup>b</sup>	12.7	7.8	2.9	7.1
1993	16.9	6.6	2.6	6.5
1994	18.6	7.5	2.2	6.7
1995	19.7	7.4	2.4	6.7
1996	19.2	8.4	2.4	7.9
1997	17.9	7.3	2.5	7.4
1998	17.2	7.9	2.3	6.5
1999	19.3	6.9	<sup>c</sup>	7.3
2000	14.9	6.9	2.2	6.5
2001	17.8	7.8	<sup>c</sup>	7.2
2002 <sup>b</sup>	10.0	7.4	<sup>c</sup>	7.6
2003 <sup>b</sup>	11.3	8.2	<sup>c</sup>	6.7
2004	<sup>c</sup>	7.2	<sup>c</sup>	6.7
2005	16.6	7.7	<sup>c</sup>	7.0
2006 <sup>b</sup>	14.4	6.6	<sup>c</sup>	6.8
2007 <sup>b</sup>	10.8	7.9	2.8	6.7
2008 <sup>b</sup>	14.7	7.1	2.5	6.8
2009	<sup>c</sup>	7.8	2.7	7.0
2010	14.4	7.6	2.8	6.8
2011	11.4	7.3	2.8	7.0
2012	<sup>c</sup>	6.6	2.4	6.8
2013	<sup>c</sup>	7.6	2.9	6.9

<sup>a</sup> Based on age-weight-length samples or fish tickets.

<sup>b</sup> Low Chinook salmon weight due to utilization of restricted mesh size.

<sup>c</sup> None sold.

Appendix A6.—Commercial and subsistence salmon catch by species, by year in Nome Subdistrict, Norton Sound District, 1990–2013.

	NOME (SUBDISTRICT 1)																	
	Commercial						Subsistence						Combined					
Year	Chinook	Sockeye	Coho	Pink	Chum	Total	Chinook	Sockeye	Coho	Pink	Chum	Total	Chinook	Sockeye	Coho	Pink	Chum	Total
1990	0	0	0	0	0	0	58	234	510	2,233	4,246	7,281	58	234	510	2,233	4,246	7,281
1991	0	0	0	0	0	0	83	166	1,279	194	3,715	5,437	83	166	1,279	194	3,715	5,437
1992	1	2	693	185	881	1,762	152	163	1,481	7,351	1,684	10,831	153	165	2,174	7,536	2,565	12,593
1993	0	2	611	0	132	745	52	80	2,070	873	1,766	4,841	52	82	2,681	873	1,898	5,586
1994	0	1	287	0	66	354	23	69	983	6,556	1,673	9,304	23	70	1,270	6,556	1,739	9,658
1995	0	1	369	0	122	492	26	148	1,365	336	3,794	5,669	26	149	1,734	336	3,916	6,161
1996	0	0	9	13	3	25	9	185	828	3,510	2,287	6,819	9	185	837	3,523	2,290	6,844
1997	0	0	0	0	0	0	10	50	325	175	2,696	3,256	10	50	325	175	2,696	3,256
1998	0	0	0	0	0	0	15	14	1,057	4,797	964	6,847	15	14	1,057	4,797	964	6,847
1999 <sup>a</sup>	0	0	0	0	0	0	11	85	161	58	337	652	11	85	161	58	337	652
2000	0	0	0	0	0	0	7	26	747	2,657	535	3,972	7	26	747	2,657	535	3,972
2001	0	0	0	0	0	0	2	92	425	113	858	1,490	2	92	425	113	858	1,490
2002	0	0	0	0	0	0	4	79	666	3,161	1,114	5,024	4	79	666	3,161	1,114	5,024
2003	0	0	0	0	0	0	63	76	351	507	565	1,562	63	76	351	507	565	1,562
2004	0	0	0	0	0	0	100	106	1,574	15,047	685	17,512	100	106	1,574	15,047	685	17,512
2005	0	0	0	0	0	0	62	177	1,287	5,075	803	7,404	62	177	1,287	5,075	803	7,404
2006 <sup>b</sup>	0	0	0	0	0	0	24	159	3,865	9,329	890	14,267	24	159	3,865	9,329	890	14,267
2007	0	0	0	0	0	0	18	297	1,103	850	2,938	5,206	18	297	1,103	850	2,938	5,206
2008	0	0	0	0	0	0	39	127	3,423	12,592	739	16,920	39	127	3,423	12,592	739	16,920
2009	0	0	0	0	0	0	32	64	1,132	487	387	2,102	32	64	1,132	487	387	2,102
2010	0	0	0	0	0	0	39	77	1,983	6,281	3,124	11,504	39	77	1,983	6,281	3,124	11,504
2011	0	0	0	0	0	0	19	47	1,229	1,389	1,428	4,112	19	47	1,229	1,389	1,428	4,112
2012	0	0	0	0	0	0	11	171	1,150	8,376	2,521	12,229	11	171	1,150	8,376	2,521	12,229
2013	<sup>c</sup>	<sup>c</sup>	<sup>c</sup>	<sup>c</sup>	<sup>c</sup>	<sup>c</sup>	48	211	1,804	805	3,065	5,973	<sup>c</sup>	<sup>c</sup>	<sup>c</sup>	<sup>c</sup>	<sup>c</sup>	<sup>c</sup>
5-year avg <sup>d</sup>	0	0	0	0	0	0	28	97	1,783	5,825	1,640	9,373	28	97	1,783	5,825	1,640	9,373
10-year avg <sup>e</sup>	0	0	0	0	0	0	41	130	1,710	5,993	1,408	9,282	41	130	1,710	5,993	1,408	9,282

Note: Commercial harvest numbers may include a small number of salmon reported on fish tickets that were retained for personal use and not commercially sold.

<sup>a</sup> Beginning in 1999, Tier II chum salmon fishing restrictions limited the number of permit holders that could fish for chum salmon.

<sup>b</sup> Beginning in 2006, Tier II chum salmon fishing restrictions were suspended.

<sup>c</sup> Confidential.

<sup>d</sup> 2008–2012.

<sup>e</sup> 2003–2012.

Appendix A7.—Commercial and subsistence salmon catch by species, by year in Golovin Subdistrict, Norton Sound District, 1990–2013.

Year	Commercial						GOLOVIN (SUBDISTRICT 2)						Combined					
							Subsistence											
	Chinook	Sockeye	Coho	Pink	Chum	Total	Chinook	Sockeye	Coho	Pink	Chum	Total	Chinook	Sockeye	Coho	Pink	Chum	Total
1990	52	21	0	0	15,993	16,066	a	a	a	a	a	a	a	a	a	a	a	a
1991	49	1	0	0	14,839	14,889	a	a	a	a	a	a	a	a	a	a	a	a
1992	6	9	2,085	0	1,002	3,102	a	a	a	a	a	a	a	a	a	a	a	a
1993	1	4	2	8,480	2,803	11,290	a	a	a	a	a	a	a	a	a	a	a	a
1994 <sup>b</sup>	0	0	3,424	0	111	3,535	253	168	733	8,410	1,337	10,901	253	168	4,157	8,410	1,448	14,436
1995 <sup>b</sup>	0	0	1,616	4,296	1,987	7,899	165	34	1,649	7,818	10,373	20,039	165	34	3,265	12,114	12,360	27,938
1996 <sup>b</sup>	0	0	638	0	0	638	86	134	3,014	17,399	2,867	23,500	86	134	3,652	17,399	2,867	24,138
1997 <sup>b</sup>	19	2	102	20	8,003	8,146	138	427	555	4,570	4,891	10,581	157	429	657	4,590	12,894	18,727
1998 <sup>b</sup>	1	0	3	106,761	723	107,488	184	37	1,292	13,340	1,893	16,746	185	37	1,295	120,101	2,616	124,234
1999 <sup>b</sup>	0	0	0	0	0	0	60	48	1,234	469	3,656	5,467	60	48	1,234	469	3,656	5,467
2000 <sup>b</sup>	0	0	1,645	17,408	164	19,217	169	18	2,335	10,906	1,155	14,583	169	18	3,980	28,314	1,319	33,800
2001 <sup>b</sup>	0	43	30	0	7,094	7,167	89	72	880	1,665	3,291	5,997	89	115	910	1,665	10,385	13,164
2002 <sup>b</sup>	0	0	0	0	0	0	69	66	1,640	14,430	1,882	18,087	69	66	1,640	14,430	1,882	18,087
2003 <sup>b</sup>	0	0	0	0	0	0	166	28	309	5,012	1,477	6,992	166	28	309	5,012	1,477	6,992
2004 <sup>c</sup>	0	0	0	0	0	0	164	6	654	19,936	880	21,640	164	6	654	19,936	880	21,640
2005 <sup>c</sup>	0	0	0	0	0	0	96	15	686	11,467	1,852	14,116	96	15	686	11,467	1,852	14,116
2006 <sup>c</sup>	0	0	0	0	0	0	136	38	1,760	14,670	722	17,326	136	38	1,760	14,670	722	17,326
2007 <sup>c</sup>	0	0	0	0	0	0	188	321	1,179	3,980	4,217	9,885	188	321	1,179	3,980	4,217	9,885
2008 <sup>c</sup>	0	0	256	2,699	623	3,578	146	95	2,337	10,155	350	13,083	146	95	2,593	12,854	973	16,661
2009 <sup>c</sup>	0	0	2,452	0	87	2,539	237	33	1,377	3,787	1,694	7,128	237	33	3,829	3,787	1,781	9,667
2010 <sup>c</sup>	3	2	5,586	2,039	17,212	24,842	59	32	2,020	9,620	1,133	12,864	62	34	7,606	11,659	18,345	37,706
2011 <sup>c</sup>	7	0	859	3	20,075	20,944	99	74	1,345	5,652	2,122	9,292	106	74	2,204	5,655	22,197	30,236
2012 <sup>c</sup>	2	14	573	31,055	3,791	35,435	57	52	1,143	7,635	1,056	9,943	59	66	1,716	38,690	4,847	45,378
2013 <sup>c</sup>	0	0	5,362	1,180	3,113	9,655	47	15	964	3,655	3,256	7,937	47	15	6,326	4,835	6,369	17,592
5-year avg <sup>d</sup>	2	3	1,945	7,159	8,358	17,468	120	57	1,644	7,370	1,271	10,462	122	60	3,590	14,529	9,629	27,930
10-year avg <sup>e</sup>	1	2	973	3,580	4,179	8,734	135	69	1,281	9,191	1,550	12,227	136	71	2,254	12,771	5,729	20,961

Note: Commercial harvest numbers may include a small number of salmon reported on fish tickets that were retained for personal use and not commercially sold.

<sup>a</sup> Subsistence surveys were not conducted.

<sup>b</sup> Subsistence harvests were estimated from Division of Subsistence household surveys. Previous surveys often were partial surveys and did not capture late-season harvests like coho salmon.

<sup>c</sup> Beginning in 2004 a permit was required for the subdistrict, replacing household surveys. The permit system helped to record harvest by residents living outside the subdistrict.

<sup>d</sup> 2008–2012.

<sup>e</sup> 2003–2012.

Appendix A8.—Commercial and subsistence salmon catch by species, by year in Elim Subdistrict, Norton Sound District, 1990–2013.

Year	Commercial						ELIM (SUBDISTRICT 3)						Combined					
							Subsistence											
	Chinook	Sockeye	Coho	Pink	Chum	Total	Chinook	Sockeye	Coho	Pink	Chum	Total	Chinook	Sockeye	Coho	Pink	Chum	Total
	a	a	a	a	a	a	a	a	a	a	a	a	a	a	a	a	a	a
1990	202	0	0	501	3,723	4,426												
1991 <sup>b</sup>	161	0	0	0	804	965	312	0	2,153	3,555	2,660	8,680	473	0	2,153	3,555	3,464	9,645
1992 <sup>b</sup>	0	0	3,531	0	6	3,537	100	0	1,281	6,152	1,260	8,793	100	0	4,812	6,152	1,266	12,330
1993 <sup>b</sup>	3	0	4,065	0	167	4,235	368	0	1,217	1,726	1,635	4,946	371	0	5,282	1,726	1,802	9,181
1994 <sup>b</sup>	0	0	5,345	0	414	5,759	322	104	1,180	9,345	3,476	14,427	322	104	6,525	9,345	3,890	20,186
1995 <sup>b</sup>	4	44	3,742	2,962	1,171	7,923	284	17	1,353	2,046	3,774	7,474	288	61	5,095	5,008	4,945	15,397
1996 <sup>b</sup>	0	0	1,915	68,609	0	70,524	417	52	1,720	9,442	2,319	13,950	417	52	3,635	78,051	2,319	84,474
1997 <sup>b</sup>	844	0	1,409	0	2,683	4,936	619	50	1,213	1,314	2,064	5,260	1,463	50	2,622	1,314	4,747	10,196
1998 <sup>b</sup>	105	0	1,462	145,669	2,311	149,547	414	49	1,831	6,891	1,376	10,561	519	49	3,293	152,560	3,687	160,108
1999 <sup>b</sup>	0	0	0	0	0	0	424	13	975	1,564	744	3,720	424	13	975	1,564	744	3,720
2000 <sup>b</sup>	10	0	5,182	46,369	535	52,096	248	46	1,429	5,983	1,173	8,879	258	46	6,611	52,352	1,708	60,975
2001 <sup>b</sup>	7	0	1,696	0	681	2,384	427	70	1,352	1,390	898	4,137	434	70	3,048	1,390	1,579	6,521
2002 <sup>b</sup>	0	0	0	0	0	0	565	14	1,801	8,345	1,451	12,176	565	14	1,801	8,345	1,451	12,176
2003 <sup>b</sup>	0	0	0	0	0	0	660	39	1,143	2,524	1,687	6,053	660	39	1,143	2,524	1,687	6,053
2004 <sup>c</sup>	0	0	0	0	0	0	412	0	704	7,858	683	9,657	412	0	704	7,858	683	9,657
2005 <sup>c</sup>	0	0	0	0	0	0	225	9	1,011	3,721	598	5,564	225	9	1,011	3,721	598	5,564
2006 <sup>c</sup>	0	0	0	0	0	0	179	13	1,769	5,216	1,267	8,444	179	13	1,769	5,216	1,267	8,444
2007 <sup>c</sup>	1	0	5,908	1,648	4,567	12,124	260	0	2,295	1,742	2,334	6,631	261	0	8,203	3,390	6,901	18,755
2008 <sup>c</sup>	5	0	4,602	14,536	304	19,447	269	0	1,804	7,655	1,284	11,012	274	0	6,406	22,191	1,588	30,459
2009 <sup>c</sup>	0	1	9,582	35	597	10,215	545	13	2,434	1,522	600	5,114	545	14	12,016	1,557	1,197	15,329
2010 <sup>c</sup>	9	5	10,180	11,658	23,453	45,305	97	7	1,679	7,830	3,925	13,538	106	12	11,859	19,488	27,378	58,843
2011 <sup>c</sup>	4	12	8,336	165	23,531	32,048	160	3	1,688	704	3,671	6,226	164	15	10,024	869	27,202	38,274
2012 <sup>c</sup>	3	1	2,003	52,775	2,262	57,044	42	0	1,302	10,848	1,494	13,686	45	1	3,305	63,623	3,756	70,730
2013 <sup>c</sup>	6	27	6,675	601	1,434	8,743	39	15	1,515	1,134	1,218	3,921	45	42	8,190	1,735	2,652	12,664
5-year avg. <sup>d</sup>	4	4	6,941	15,834	10,029	32,812	223	5	1,781	5,712	2,195	9,915	227	8	8,722	21,546	12,224	42,727
10-year avg. <sup>e</sup>	2	2	4,061	8,082	5,471	17,618	285	8	1,583	4,962	1,754	8,593	287	10	5,644	13,044	7,226	26,211

Note: Commercial harvest numbers may include a small number of salmon reported on fish tickets that were retained for personal use and not commercially sold.

<sup>a</sup> Subsistence surveys were not conducted.

<sup>b</sup> Subsistence harvests were estimated from Division of Subsistence household surveys. Previous surveys often were partial surveys and did not capture later season harvest like coho salmon.

<sup>c</sup> Beginning in 2004 a permit was required for the subdistrict, replacing household surveys. The permit system helped to record harvest by residents living outside the subdistrict.

<sup>d</sup> 2008–2012.

<sup>e</sup> 2003–2012.

Appendix A9.—Commercial and subsistence salmon catch by species, by year in Norton Bay Subdistrict, Norton Sound District, 1990–2013.

Year	NORTON BAY (SUBDISTRICT 4)																	
	Commercial						Subsistence						Combined					
	Chinook	Sockeye	Coho	Pink	Chum	Total	Chinook	Sockeye	Coho	Pink	Chum	Total	Chinook	Sockeye	Coho	Pink	Chum	Total
1990	0	0	0	0	0	0	a	a	a	a	a	a	a	a	a	a	a	a
1991	0	0	0	0	0	0	a	a	a	a	a	a	a	a	a	a	a	a
1992	27	0	0	0	1,787	1,814	a	a	a	a	a	a	a	a	a	a	a	a
1993	267	0	0	290	1,378	1,935	a	a	a	a	a	a	a	a	a	a	a	a
1994 <sup>b</sup>	0	0	0	0	0	0	308	1	370	6,049	4,581	11,309	308	1	370	6,049	4,581	11,309
1995 <sup>b</sup>	0	0	0	0	0	0	475	46	985	3,514	5,828	10,848	475	46	985	3,514	5,828	10,848
1996 <sup>b</sup>	0	0	0	0	0	0	295	3	676	3,929	4,161	9,064	295	3	676	3,929	4,161	9,064
1997 <sup>b</sup>	194	0	0	0	531	725	656	54	322	1,795	4,040	6,867	850	54	322	1,795	4,571	7,592
1998 <sup>b</sup>	0	0	0	0	0	0	684	0	388	2,009	6,192	9,273	684	0	388	2,009	6,192	9,273
1999 <sup>b</sup>	0	0	0	0	0	0	327	0	167	1,943	4,153	6,590	327	0	167	1,943	4,153	6,590
2000 <sup>b</sup>	0	0	0	0	0	0	397	2	267	2,255	4,714	7,635	397	2	267	2,255	4,714	7,635
2001 <sup>b</sup>	0	0	0	0	0	0	460	14	276	5,203	4,445	10,398	460	14	276	5,203	4,445	10,398
2002 <sup>b</sup>	0	0	0	0	0	0	557	0	509	6,049	3,971	11,086	557	0	509	6,049	3,971	11,086
2003 <sup>b</sup>	0	0	0	0	0	0	373	46	510	4,184	3,397	8,510	373	46	510	4,184	3,397	8,510
2004	0	0	0	0	0	0	a	a	a	a	a	a	a	a	a	a	a	a
2005	0	0	0	0	0	0	a	a	a	a	a	a	a	a	a	a	a	a
2006	0	0	0	0	0	0	a	a	a	a	a	a	a	a	a	a	a	a
2007	0	0	0	0	0	0	a	a	a	a	a	a	a	a	a	a	a	a
2008	7	0	600	1,232	507	2,346	187	2	1,084	4,489	3,330	9,092	194	2	1,684	5,721	3,837	11,438
2009	0	0	1,714	558	1,850	4,122	259	2	891	2,508	3,183	6,843	259	2	2,605	3,066	5,033	10,965
2010	0	7	1,606	2,597	6,007	10,217	341	21	461	3,115	3,180	7,118	341	28	2,067	5,712	9,187	17,335
2011	5	9	4,836	652	7,558	13,060	239	1	549	1,132	3,529	5,450	6	558	5,968	4,181	13,008	13,066
2012	10	16	4,378	49,970	8,417	62,791	103	0	310	2,623	2,721	5,757	113	16	4,688	52,593	11,138	68,548
2013	8	4	5,485	487	36,021	42,005	123	2	826	1,341	3,853	6,145	131	6	6,311	1,828	39,874	48,150
5-year avg. <sup>c</sup>	6	8	3,284	13,752	6,085	23,134	282	7	824	3,467	3,986	8,565	228	152	4,253	17,818	10,551	30,338

Note: Commercial harvest numbers may include a small number of salmon reported on fish tickets that were retained for personal use and not commercially sold.

<sup>a</sup> Subsistence surveys were not conducted.

<sup>b</sup> Subsistence harvests were estimated from Division of Subsistence household surveys. Previous surveys often were partial surveys that did not capture later season harvests like coho salmon.

<sup>c</sup> 2008–2012.

Appendix A10.—Commercial and subsistence salmon catch by species, by year in Shaktoolik Subdistrict, Norton Sound District, 1990–2013.

SHAKTOOLIK (SUBDISTRICT 5)																		
	Commercial						Subsistence						Combined					
Year	Chinook	Sockeye	Coho	Pink	Chum	Total	Chinook	Sockeye	Coho	Pink	Chum	Total	Chinook	Sockeye	Coho	Pink	Chum	Total
1990	2,644	49	4,695	0	21,748	29,136	a	a	a	a	a	a	a	a	a	a	a	a
1991	1,324	55	11,614	0	31,619	44,612	a	a	a	a	a	a	a	a	a	a	a	a
1992	1,098	56	14,660	0	27,867	43,681	a	a	a	a	a	a	a	a	a	a	a	a
1993	2,756	20	11,130	106,743	20,864	141,513	a	a	a	a	a	a	a	a	a	a	a	a
1994 <sup>b</sup>	885	8	22,065	502,231	5,411	530,600	1,175	1	2,777	9,133	1,221	14,307	2,060	9	24,842	511,364	6,632	544,907
1995 <sup>b</sup>	1,239	5	10,856	37,377	14,775	64,252	1,303	72	2,682	7,176	2,534	15,885	2,542	77	13,538	44,553	17,309	80,137
1996 <sup>b</sup>	1,340	1	13,444	304,982	3,237	323,004	1,114	31	3,615	8,370	4,425	17,555	2,454	32	17,059	313,352	7,662	340,559
1997 <sup>b</sup>	2,449	0	4,694	0	5,747	12,890	1,146	62	2,761	5,779	1,612	11,360	3,595	62	7,455	5,779	7,359	24,250
1998 <sup>b</sup>	910	0	3,624	236,171	7,080	247,785	982	92	1,872	6,270	1,034	10,250	1,892	92	5,496	242,441	8,114	258,035
1999 <sup>b</sup>	581	0	2,398	0	2,181	5,160	818	183	1,556	5,092	467	8,116	1,399	183	3,954	5,092	2,648	13,276
2000 <sup>b</sup>	160	3	7,779	85,493	2,751	96,186	440	20	2,799	5,432	2,412	11,103	600	23	10,578	90,925	5,163	107,289
2001 <sup>b</sup>	90	0	2,664	0	1,813	4,567	936	143	2,090	10,172	1,553	14,894	1,026	143	4,754	10,172	3,366	19,461
2002 <sup>b</sup>	1	0	680	0	261	942	1,230	4	2,169	8,769	800	12,972	1,231	4	2,849	8,769	1,061	13,914
2003 <sup>b</sup>	2	0	4,031	0	485	4,518	881	50	2,941	12,332	587	16,791	883	50	6,972	12,332	1,072	21,309
2004	0	0	12,734	0	1,372	14,106	943	12	1,994	7,291	139	10,379	943	12	14,728	7,291	1,511	24,485
2005	50	0	21,818	0	791	22,659	807	0	1,913	12,075	202	14,997	857	0	23,731	12,075	993	37,656
2006	8	0	32,472	0	3,321	35,801	382	36	1,968	4,817	351	7,554	390	36	34,440	4,817	3,672	43,355
2007	5	0	31,810	0	6,076	37,891	515	28	1,443	2,708	465	5,159	520	28	33,253	2,708	6,541	43,050
2008	6	24	37,624	8,219	6,042	51,915	422	2	1,504	4,920	201	7,049	428	26	39,128	13,139	6,243	58,964
2009	4	36	13,063	5,146	10,941	29,190	417	57	2,141	6,101	374	9,090	421	93	15,204	11,247	11,315	38,280
2010	4	18	11,868	4,622	40,483	56,995	327	115	1,940	6,406	1,680	10,468	331	133	13,808	11,028	42,163	67,463
2011	45	69	15,368	29	25,388	40,899	235	100	1,241	2,681	490	4,747	280	169	16,609	2,710	25,878	45,646
2012	25	29	7,828	19,253	20,141	47,276	214	9	1,110	4,609	634	6,576	239	38	8,938	23,862	20,775	53,852
2013	6	45	6,890	14	23,268	30,223	136	108	2,146	3,346	983	6,719	142	153	9,036	3,360	24,251	36,942
5-year avg. <sup>c</sup>	17	35	17,150	7,454	20,599	45,255	323	57	1,587	4,943	676	7,586	340	92	18,737	12,397	21,275	52,841
10-year avg. <sup>d</sup>	15	18	18,862	3,727	11,504	34,125	514	41	1,820	6,394	512	9,281	529	59	20,681	10,121	12,016	43,406

Note: Commercial harvest numbers may include a small number of salmon reported on fish tickets that were retained for personal use and not commercially sold.

<sup>a</sup> Subsistence surveys were not conducted.

<sup>b</sup> Subsistence harvests were estimated from Division of Subsistence household surveys. Previous surveys often were partial surveys that did not capture later season harvests by fishermen.

<sup>c</sup> 2008–2012.

<sup>d</sup> 2003–2012.

Appendix A11.—Commercial and subsistence salmon catch by species, by year in Unalakleet Subdistrict, Norton Sound District, 1990–2013.

Year	UNALAKLEET (SUBDISTRICT 6)																	
	Commercial						Subsistence						Combined					
	Chinook	Sockeye	Coho	Pink	Chum	Total	Chinook	Sockeye	Coho	Pink	Chum	Total	Chinook	Sockeye	Coho	Pink	Chum	Total
1990	5,998	358	52,015	0	23,659	82,030	2,476	a	a	a	a	a	8,474	a	a	a	a	a
1991	4,534	147	52,033	0	39,609	96,323	a	a	a	a	a	a	a	a	a	a	a	a
1992	3,409	229	84,449	6,284	52,547	146,918	a	a	a	a	a	a	a	a	a	a	a	a
1993	5,944	251	26,290	42,061	28,156	102,702	a	a	a	a	a	a	a	a	a	a	a	a
1994 <sup>b</sup>	4,400	71	71,019	480,158	12,288	567,936	3,035	404	11,386	27,163	3,325	45,313	7,435	475	82,405	507,321	15,613	613,249
1995 <sup>b</sup>	7,617	78	31,280	37,009	24,843	100,827	3,114	591	9,833	16,625	5,458	35,621	10,731	669	41,113	53,634	30,301	136,448
1996 <sup>b</sup>	3,644	0	52,200	113,837	7,369	177,050	3,023	181	11,187	18,026	4,227	36,644	6,667	181	63,387	131,863	11,596	213,694
1997 <sup>b</sup>	9,067	159	26,079	0	17,139	52,444	4,191	196	6,746	10,600	1,603	23,336	13,258	355	32,825	10,600	18,742	75,780
1998 <sup>b</sup>	6,413	7	24,534	99,412	6,210	136,576	4,066	201	7,489	13,654	3,038	28,448	10,479	208	32,023	113,066	9,248	165,024
1999 <sup>b</sup>	1,927	0	10,264	0	5,700	17,891	2,691	537	8,140	10,060	3,692	25,120	4,618	537	18,404	10,060	9,392	43,011
2000 <sup>b</sup>	582	11	29,803	17,278	2,700	50,374	2,429	212	5,878	10,540	3,000	22,059	3,011	223	35,681	27,818	5,700	72,433
2001 <sup>b</sup>	116	1	15,102	0	1,512	16,731	2,810	359	6,270	11,269	2,918	23,626	2,926	360	21,372	11,269	4,430	40,357
2002 <sup>b</sup>	4	1	1,079	0	339	1,423	2,367	280	4,988	15,915	3,877	27,427	2,371	281	6,067	15,915	4,216	28,850
2003 <sup>b</sup>	10	21	13,029	0	3,075	16,135	2,585	297	6,192	21,779	1,785	32,638	2,595	318	19,221	21,779	4,860	48,773
2004	22	47	29,282	0	4,924	34,275	2,829	417	6,653	22,755	2,154	34,808	2,851	464	35,935	22,755	7,078	69,083
2005	101	12	63,705	0	3,192	67,010	2,193	656	7,886	25,447	2,660	38,842	2,294	668	71,591	25,447	5,852	105,852
2006	12	3	98,336	0	6,721	105,072	2,537	326	9,905	22,547	2,712	38,027	2,549	329	108,241	22,547	9,433	143,099
2007	13	2	88,418	2,121	11,788	102,342	1,666	292	5,859	11,674	2,057	21,547	1,678	294	94,277	13,795	13,845	123,889
2008	65	36	77,227	48,839	17,648	143,815	1,402	137	7,452	15,116	2,805	26,912	1,467	173	84,679	63,955	20,453	170,727
2009	80	89	60,230	11,625	20,647	92,671	1,892	200	6,923	11,707	2,708	23,430	1,972	289	67,153	23,332	23,355	116,101
2010	124	71	32,839	10,641	30,588	74,263	1,257	297	3,780	9,002	3,159	17,495	1,381	368	36,619	19,643	33,747	91,758
2011	124	279	29,518	6,292	34,003	70,216	607	189	2,486	5,608	3,316	12,206	731	468	32,004	11,900	37,319	82,422
2012	157	74	22,274	52,445	28,161	103,111	808	192	4,558	9,460	3,973	18,991	965	266	26,832	61,905	32,134	122,102
2013	131	171	29,390	6,056	54,873	90,621	468	221	6,117	7,724	3,129	17,659	599	392	35,507	13,780	58,002	108,280
5-year avg. <sup>c</sup>	110	110	44,418	25,968	26,209	96,815	1,193	203	5,040	10,179	3,192	19,807	1,303	313	49,457	36,147	29,402	116,622
10-year avg. <sup>d</sup>	71	63	51,486	13,196	16,075	80,891	1,778	300	6,169	15,510	2,733	26,490	1,848	364	57,655	28,706	18,808	107,381

Note: Commercial harvest numbers may include a small number of salmon reported on fish tickets that were retained for personal use and not commercially sold.

<sup>a</sup> Subsistence surveys were not conducted.

<sup>b</sup> Subsistence harvests were estimated from Division of Subsistence household surveys. Previous surveys often were partial surveys that did not capture later season harvests by fishermen.

<sup>c</sup> 2008–2012.

<sup>d</sup> 2003–2012.



Appendix A12.–Subsistence salmon catch by species and year for St. Michael and Stebbins in Norton Sound District, 1994–2013.

Year	Chinook	Chum	Pink	Sockeye	Coho	Total
St Michael						
1994	769	4,309	2,673	127	1,022	8,900
1995	1,267	5,778	391	45	2,235	9,716
1996	1,400	6,352	1,503	3	1,641	10,899
1997	970	2,816	84	41	547	4,458
1998	542	1,502	961	143	1,406	4,554
1999	1,053	3,036	365	111	798	5,363
2000	160	1,381	80	16	1,180	2,817
2001	282	2,246	229	17	490	3,264
2002	227	1,136	583	20	989	2,955
2003	295	1,994	577	89	1,438	4,393
2004		Subsistence surveys were not conducted.				
2005	998	3,614	1,742	61	1,497	7,912
2006	271	2,628	480	347	1,256	4,982
2007	452	2,119	265	9	622	3,467
2008		Subsistence surveys were not conducted.				
2009	825	921	169	24	1,088	3,027
2010		Subsistence surveys were not conducted.				
2011		Subsistence surveys were not conducted.				
2012	80	2,172	457	20	911	3,640
2013		Subsistence surveys were not conducted.				
Stebbins						
1994	1,525	5,989	5,552	288	3,948	17,302
1995	1,211	5,042	758	207	2,570	9,788
1996	1,030	7,401	2,375	424	3,746	14,976
1997	1,164	3,230	243	116	1,826	6,579
1998	1,410	3,909	3,125	295	3,116	11,855
1999	760	3,312	459	200	1,312	6,043
2000	298	2,913	364	341	2,429	6,345
2001	570	3,999	202	0	2,759	7,530
2002	450	3,586	7,459	300	2,324	14,119
2003	265	2,399	2,685	171	1,215	6,735
2004		Subsistence surveys were not conducted.				
2005	485	5,164	4,353	59	2,702	12,763
2006	355	4,236	4,321	140	4,856	13,908
2007	763	4,980	1,881	0	2,006	9,630
2008		Subsistence surveys were not conducted.				
2009	713	1,461	328	0	1,114	3,616
2010		Subsistence surveys were not conducted.				
2011		Subsistence surveys were not conducted.				
2012	109	3,456	3,659	0	1,256	8,480
2013		Subsistence surveys were not conducted.				

Note: Harvest numbers shown have been expanded to include households not contacted.

Appendix A13.—Commercial, subsistence, and sport salmon catch by species, by year for Subdistricts 1–6 in Norton Sound District, 1990–2013.

Year	SUBDISTRICTS 1–6																	
	Commercial						Subsistence						Sport fish					
	Chinook	Sockeye	Coho	Pink	Chum	Total	Chinook	Sockeye	Coho	Pink	Chum	Total	Chinook	Sockeye	Coho	Pink	Chum	Total
1990 <sup>a</sup>	8,895	434	56,712	501	65,123	131,665	2,534	234	510	2,233	4,246	7,281	364	198	3,305	7,647	925	12,439
1991 <sup>a</sup>	6,068	203	63,647	0	86,871	156,789	395	166	3,432	3,749	6,375	14,117	404	237	5,800	1,738	1,415	9,594
1992 <sup>a</sup>	4,541	296	105,418	6,284	83,394	199,933	252	163	2,762	13,503	2,944	19,624	204	131	4,671	6,403	523	11,932
1993 <sup>a</sup>	8,972	279	43,283	157,574	53,562	263,670	420	80	3,287	2,599	3,401	9,787	595	10	3,783	2,250	691	7,329
1994	5,285	80	102,140	982,389	18,290	1,108,184	5,116	747	17,429	66,656	15,613	105,561	600	18	5,547	7,051	536	13,752
1995	8,860	128	47,863	81,644	42,898	181,393	5,367	908	17,867	37,515	31,761	95,536	438	104	3,705	928	394	5,569
1996	4,984	1	68,206	487,441	10,609	571,241	4,944	586	21,040	60,676	20,286	107,532	662	100	7,289	5,972	662	14,685
1997	12,573	161	32,284	20	34,103	79,141	6,760	839	11,922	24,233	16,906	60,660	1,106	30	4,393	1,458	278	7,265
1998	7,429	7	29,623	588,013	16,324	641,396	6,345	393	13,929	46,961	14,497	82,125	590	16	4,441	6,939	682	12,668
1999	2,508	0	12,662	0	7,881	23,051	4,331	866	12,233	19,186	13,049	49,665	630	0	5,582	3,039	211	9,462
2000	752	14	44,409	166,548	6,150	217,873	3,690	324	13,455	37,773	12,989	68,231	889	45	7,441	2,886	1,097	12,358
2001	213	44	19,492	0	11,100	30,849	4,724	750	11,293	29,812	13,963	60,542	271	39	4,802	360	1,709	7,181
2002	5	1	1,759	0	600	2,365	4,792	443	11,773	56,669	13,095	86,772	802	0	4,211	4,303	818	10,134
2003	12	21	17,060	0	3,560	20,653	4,728	536	11,446	46,338	9,498	72,546	239	572	3,039	2,222	292	6,364
2004 <sup>a</sup>	22	47	42,016	0	6,296	48,381	4,448	541	11,579	72,887	4,541	93,996	535	404	5,806	8,309	498	15,552
2005 <sup>a</sup>	151	12	85,523	0	3,983	89,669	3,383	857	12,783	57,785	6,115	80,923	216	0	3,959	473	36	4,684
2006 <sup>a</sup>	20	3	130,808	0	10,042	140,873	3,258	572	19,267	56,579	5,942	85,618	427	22	11,427	5,317	344	17,537
2007 <sup>a</sup>	19	2	126,136	3,769	22,431	152,357	2,647	938	11,879	20,954	12,011	48,428	147	15	6,179	1,331	96	7,768
2008	83	60	120,309	75,525	25,124	221,101	2,465	363	17,604	54,927	8,709	84,068	580	63	10,756	6,855	341	18,595
2009	84	126	87,041	17,364	34,122	138,737	3,382	369	14,898	26,112	8,946	53,707	277	0	6,664	1,321	417	8,679
2010	140	103	62,079	31,557	117,743	211,622	2,120	549	11,863	42,254	16,201	72,987	61	0	5,876	2,717	118	8,772
2011	185	369	58,917	7,141	110,555	177,167	1,359	414	8,538	17,166	14,556	42,033	61	58	3,582	566	139	4,406
2012	197	134	37,056	205,498	62,772	305,657	1,235	424	9,573	43,551	12,399	67,182	0	28	5,099	3,220	209	8,556
2013	151	247	53,802	8,338	118,709	181,247	861	572	13,372	18,045	15,504	48,354	0	23	7,567	1,806	2,267	11,663
5-year avg. <sup>b</sup>	138	158	73,080	67,417	70,063	210,857	2,112	424	12,495	36,802	12,162	63,995	196	30	6,395	2,936	245	9,802
10-year avg. <sup>c</sup>	91	88	76,695	34,085	39,663	150,622	2,903	556	12,943	43,855	9,892	70,149	254	116	6,239	3,233	249	10,091

Note: Commercial harvest may include a small number of salmon reported on fish tickets that were retained for personal use and not commercially sold. ND is no data or insufficient data.

<sup>a</sup> Not all subdistricts were surveyed.

<sup>b</sup> 2008–2012.

<sup>c</sup> 2003–2012.

Appendix A14.—Sport salmon harvest by species, by year for the Unalakleet River, 1990–2013.

Year	Chinook	Coho	Chum	Pink	Total
1990	276	1,826	298	1,180	3,580
1991	296	2,180	497	437	3,410
1992	117	1,555	379	779	2,830
1993	382	643	116	89	1,230
1994	379	2,425	220	402	3,426
1995	259	2,033	207	222	2,721
1996	384	3,411	463	59	4,317
1997	842	2,784	228	1,055	4,909
1998	513	2,742	447	434	4,136
1999	415	2,691	211	2,946	6,263
2000	345	4,150	403	961	5,859
2001	250	2,766	714	188	3,918
2002	544	2,937	607	1,378	5,466
2003	97	1,604	191	29	1,921
2004	356	3,524	47	2,003	5,930
2005	216	3,959	36	473	4,684
2006	394	4,985	224	891	6,494
2007	147	4,117	85	618	4,967
2008	580	6,029	175	2,077	8,861
2009	236	5,095	260	586	6,177
2010	61	3,006	59	535	3,661
2011	54	2,493	77	391	3,015
2012	0	3,283	118	20	3,421
2013	0	4,068	354	886	5,308
Avg 2008–2012	186	3,981	138	722	5,027
Avg 2003–2012	214	3,810	127	762	4,913

Appendix A15.–Sport salmon harvest by species, by year for the Fish/Niukluk rivers, 1990–2013.

Year	Chinook	Coho	Chum	Pink	Total
1990	0	267	216	638	1,121
1991	14	977	272	356	1,619
1992	0	753	15	357	1,125
1993	9	1,185	514	278	1,986
1994	10	1,122	119	231	1,482
1995	18	818	27	136	999
1996	11	1,652	166	404	2,233
1997	71	462	0	58	591
1998	0	316	0	0	316
1999	44	1,365	0	80	1,489
2000	174	1,165	0	51	1,390
2001	0	969	439	161	1,569
2002	75	298	45	254	672
2003	39	216	101	196	552
2004	22	291	435	353	1,101
2005	37	400	0	58	495
2006	0	948	0	134	1,082
2007	0	786	11	30	827
2008	0	1,986	166	969	3,121
2009	30	939	72	25	1,066
2010	0	1,069	0	99	1,168
2011	0	700	29	10	739
2012	0	1,163	74	636	1,873
2013	0	1,227	0	0	1,227
Avg 2008–2012	6	1,171	68	348	1,593
Avg 2003–2012	13	850	89	251	1,202

Appendix A16.—Sport salmon harvest by species, by year for the Nome River, 1990–2013.

Year	Chinook	Coho	Chum	Pink	Total
1990	39	407	122	2,651	3,219
1991	22	417	241	356	1,036
1992	16	713	0	4,397	5,126
1993	93	602	0	723	1,418
1994	0	326	0	4,103	4,429
1995	0	143	0	230	373
1996	0	598	0	3,280	3,878
1997	10	295	0	83	388
1998	0	189	0	1,985	2,174
1999	0	219	0	0	219
2000	0	342	0	578	920
2001	0	297	0	0	297
2002	0	217	0	312	529
2003	0	68	0	12	80
2004	0	270	0	3,369	3,639
2005	0	1,001	0	1,193	2,194
2006	0	2,768	0	2,422	5,190
2007	0	797	0	402	1,199
2008	0	1,793	0	2,954	4,747
2009	0	229	0	178	407
2010	13	602	0	1,716	2,331
2011	0	68	0	85	153
2012	0	259	0	1,264	1,523
2013	0	279	139	302	720
Avg 2008–2012	3	590	0	1,239	1,832
Avg 2003–2012	1	786	0	1,360	2,146

Appendix A17.—Comparative salmon aerial survey escapement indices of Norton Sound streams unless noted otherwise, 1990–2013.

Year <sup>a</sup>	Sinuk River				Nome River			
	Chinook	Chum	Pink	Coho	Chinook	Chum	Pink	Coho
1990	ND	95	29,040	161	ND	541	13,085	377
1991	3	5,420	14,680	701	11	3,520	4,690	611
1992	1	470	292,400	422	3	813	255,700	691
1993	7	1,570	5,120	104	8	1,520	8,941	276
1994	10	1,140	492,000	307	2	350	265,450	631
1995	ND	3,110	1,250	290	ND	1,865	182	517
1996	5	1,815	74,100	367	1	799	34,520	723
1997	ND	2,975	1,200	57	4	956	65	544
1998	ND	630	372,850	322	3	335	179,680	515
1999	ND	1,697	180	217	ND	375	345	620
2000	ND	10	12,608	912	ND	658	6,380	1,032
2001	ND	3,746	115 <sup>b</sup>	750	ND	946 <sup>b</sup>	790 <sup>b</sup>	1,307 <sup>b</sup>
2002	ND	1,682	28,487	1,290 <sup>b</sup>	ND	127 <sup>b</sup>	295 <sup>b</sup>	1,796
2003	ND	677	9,885	190	8	337	2,841	604
2004	ND	100 <sup>b</sup>	1,267,100 <sup>b</sup>	2,085	ND	3 <sup>b</sup>	707,350 <sup>b</sup>	1,687
2005	ND	1,072 <sup>b</sup>	211,000 <sup>b</sup>	2,045	2 <sup>b</sup>	2,082 <sup>b</sup>	212,000 <sup>b</sup>	3,541
2006	0 <sup>b</sup>	1115 <sup>b</sup>	515,000 <sup>b</sup>	2,147	0 <sup>b</sup>	394 <sup>b</sup>	441,550 <sup>b</sup>	3,650
2007	3 <sup>b</sup>	7,210 <sup>b</sup>	6,810 <sup>b</sup>	668	4 <sup>b</sup>	1,449 <sup>b</sup>	3,378 <sup>b</sup>	1,442
2008	ND	ND	1,496,000 <sup>b</sup>	1,633	ND	106 <sup>b</sup>	528,000 <sup>b</sup>	2,051
2009	0 <sup>b</sup>	344 <sup>b</sup>	6,730 <sup>b</sup>	508 <sup>b</sup>	ND	ND	ND	877 <sup>b</sup>
2010	0 <sup>b</sup>	3,955 <sup>b</sup>	168,600 <sup>b</sup>	5,507 <sup>b</sup>	0 <sup>b</sup>	2,998 <sup>b</sup>	98,272 <sup>b</sup>	0 <sup>b</sup>
2011	0 <sup>b</sup>	6,265 <sup>b</sup>	21,100 <sup>b</sup>	479 <sup>b</sup>	0 <sup>b</sup>	1,317 <sup>b</sup>	9,575 <sup>b</sup>	870 <sup>b</sup>
2012	0 <sup>b</sup>	3,650 <sup>b</sup>	506,500 <sup>b</sup>	ND	No survey occurred.			
2013	0 <sup>b</sup>	19,500 <sup>b</sup>	23,000 <sup>b</sup>	1,054 <sup>b</sup>	No survey occurred.			

-continued-

Appendix A17.–Page 2 of 4.

Year <sup>a</sup>	Flambeau River				Eldorado River			
	Chinook	Chum	Pink	Coho	Chinook	Chum	Pink	Coho
1990	ND	905	ND	96	17	884	2,050	44
1991	ND	2,828	7,180	ND	76	5,755	1,590	98
1992	ND	55	ND	42	2	4,887	6,615	113
1993	ND	819	640	11	38	2,895	120	111
1994	ND	3,612	4	213	ND	5,140	53,890	242
1995	ND	1,876	1,102	186	4	9,025	50	247
1996	ND	647	355	71	21	20,710	40,100	254
1997	ND	2,250 <sup>b</sup>	200 <sup>b</sup>	751	40	5,967	10	37
1998	ND	2,828	7,180	ND	ND	3,000	123,950	71
1999	ND	55	ND	42	2	1,741	6	45
2000	ND	819	640	11	2	3,383	16,080	24
2001	ND	3,612	4	213	2	4,450	8	232
2002	ND	1,876	1,102	186	8	139	58,700	463
2003	ND	647	355	71	12	1,257	821	71
2004	ND	2,250 <sup>b</sup>	200 <sup>b</sup>	751	ND	109 <sup>b</sup>	52,000 <sup>b</sup>	755
2005	ND	2,261 <sup>b</sup>	100 <sup>b</sup>	154	2 <sup>b</sup>	5,445 <sup>b</sup>	2,050 <sup>b</sup>	376
2006	0 <sup>b</sup>	16,000 <sup>b</sup>	8,800 <sup>b</sup>	ND	0 <sup>b</sup>	2,355 <sup>b</sup>	156,500 <sup>b</sup>	523
2007	1 <sup>b</sup>	4,452 <sup>b</sup>	0 <sup>b</sup>	38	2 <sup>b</sup>	6,315 <sup>b</sup>	318 <sup>b</sup>	34
2008	0 <sup>b</sup>	4,235 <sup>b</sup>	106,200 <sup>b</sup>	918	No survey occurred.			
2009	0 <sup>b</sup>	860 <sup>b</sup>	1,598 <sup>b</sup>	627 <sup>b</sup>	14 <sup>b</sup>	1,069 <sup>b</sup>	210 <sup>b</sup>	301 <sup>b</sup>
2010	0 <sup>b</sup>	13,600 <sup>b</sup>	36,000 <sup>b</sup>	ND	0 <sup>b</sup>	30,600 <sup>b</sup>	84,582 <sup>b</sup>	ND
2011	0 <sup>b</sup>	5,283 <sup>b</sup>	1,810 <sup>b</sup>	292 <sup>b</sup>	0 <sup>b</sup>	9,225 <sup>b</sup>	260 <sup>b</sup>	120 <sup>b</sup>
2012	0 <sup>b</sup>	7,911 <sup>b</sup>	ND	ND	No survey occurred.			
2013	0 <sup>b</sup>	16,088 <sup>b</sup>	ND	ND	4 <sup>b</sup>	16,859 <sup>b</sup>	52 <sup>b</sup>	ND

-continued-

Appendix A17.–Page 3 of 4.

Year <sup>a</sup>	Fish River				Boston Creek			
	Chinook	Chum	Pink	Coho	Chinook	Chum	Pink	Coho
1990		No survey occurred.			112	1,455	8,440	ND
1991	58	10,470	51,190	ND	152	2,560	3,210	ND
1992	4	390	1,387,000	ND	68	1,540	50,850	ND
1993	48	12,695	13,440	ND	227	4,563	1,930	ND
1994	55	16,500	910,000	ND	95	4,270	355,600	ND
1995	40	13,433	780	1,829	78	4,221	ND	230
1996	189	5,840 <sup>c</sup>	684,780	ND	ND	3,505 <sup>c</sup>	35,980	ND
1997	110	19,515	800	465	452	4,545	ND	ND
1998	96	28,010	663,050	ND	255	1,570	175,330	ND
1999	ND	50	20	821	ND	ND	ND	319
2000	ND	ND	ND	805	ND	ND	ND	414
2001	8	3,220	1,744	1,055	33	3,533	1,038	155
2003	95	3,200	1,014	ND	145	750	701	ND
2004	19	621	404,930	90	93	55	135,000	140
2005	0	6,875	319,170	ND	46	1,675	5,850	ND
2010		No survey occurred.			29 <sup>b</sup>	3,010 <sup>b</sup>	5,110 <sup>b</sup>	73 <sup>b</sup>
2013	15 <sup>b</sup>	2,550 <sup>b</sup>	ND	ND	19 <sup>b</sup>	16,100 <sup>b</sup>	ND	ND

Year <sup>a</sup>	Niukluk River				Kwiniuk River			
	Chinook	Chum	Pink	Coho	Chinook <sup>d</sup>	Chum <sup>d</sup>	Pink <sup>d</sup>	Coho <sup>d</sup>
1990	15	6,200	115,250	170	744	13,735	416,511	746 <sup>e</sup>
1991	42	10,700	37,410	1,783 <sup>f</sup>	587	18,802	53,499	809 <sup>e</sup>
1992	ND	7,770	803,200	812	479	12,077	1,464,717	532 <sup>e</sup>
1993	15	19,910	2,840	2,104	565	15,823	43,065	1,238 <sup>e</sup>
1994	7	16,470	1,294,100	274	627	33,010	2,304,099	2,547
1995	48	25,358	200	2,136	468	42,161	17,509	1,625 <sup>e</sup>
1996	25	9,732 <sup>c</sup>	153,150	2,047	567	27,256	907,894	1,410 <sup>e</sup>
1997	131	16,550	ND	983	972	20,118	9,536	610 <sup>e</sup>
1998	51	2,556	205,110	593	296	24,248	655,933	610 <sup>e</sup>
1999	ND	640	ND	619	115	8,763	608	223 <sup>e</sup>
2000	ND	ND	ND	3,812	144	12,878	750,173	541 <sup>e</sup>
2001	6	2,448	2,856	809	258	16,598	8,423	9,532
2002	ND	ND	ND	1,122	778	37,995	111,410	6,459
2003	55	2,315	272	146	744	12,123	22,329	5,490
2004	15	173	277,900	828	663	10,362	3,054,684	11,240
2005	6	3,225	154,000	ND	342	12,083	341,048	12,950
2006	ND	ND	ND	737 <sup>g</sup>	195	39,519	1,347,090	22,341
2007	ND	ND	ND	ND	258	27,756	54,225	9,429
2008	ND	ND	ND	1,715	237	9,483	1,444,213	10,461
2009		No survey occurred.			444	8,739	42,960	9,036
2010		No survey occurred.			135	71,388	634,220	8,049
2011	4 <sup>b</sup>	9,735 <sup>b</sup>	375 <sup>b</sup>	838 <sup>b</sup>	57	31,604	30,023	3,288
2012	ND	ND	ND	928 <sup>b</sup>	54	5,577	393,302	777
2013	68 <sup>b</sup>	17,203 <sup>b</sup>	9,700 <sup>b</sup>	2,279 <sup>b</sup>	15	5,631	13,212	3,940

-continued-



Appendix A17.–Page 4 of 4.

Year <sup>a</sup>	Tubutulik River				North River			
	Chinook	Chum	Pink	Coho	Chinook	Chum	Pink	Coho
1990	397	4,350	186,400	ND	255	1,345	25,685	ND
1991	661	7,085	26,870	ND	656	2,435	119,140	2,510
1992	260	2,595	138,600	ND	329	ND	631,140	398
1993	1,061	8,740	18,650	1,395	900	445	13,570	1,397
1995	377	16,158	4,020	930	622	1,370	18,300	690 <sup>h</sup>
1996	439	10,790	226,750	ND	106	270 <sup>c</sup>	125,500	917
1997	1,946	3,105	16,890	ND	1,605	9,045	17,870	ND
1998	894	10,180	1,124,800	ND	591	50	153,150	233
1999	No survey occurred.				18	1,480	3,790	533
2001	77	863	ND	ND	367	330	ND	ND
2002	42	180	182,000	ND	122	217	4,590	800
2003	50	1,352	60	292	131	222	11,010	ND
2004	321	1,117	391,000	779	189	283	264,000	1,386
2005	78	1,336	48,203	ND	156	310	381,150	1,963
2007	823	7,045	32,250	4,552	554	295	50,100	2,349
2008	ND	ND	ND	4,197	ND	ND	ND	2,774
2009	627	3,161	12,695	ND	438	3,263	189,939	2,830
2010	122	16,097	16,520	50	124	1,627	1,480	200
2011	141 <sup>b</sup>	14,127 <sup>b</sup>	3,875 <sup>b</sup>	1,606	433	9,785	20,920	898
2012	ND	ND	ND	2,889 <sup>b</sup>	No survey occurred.			
2013	2	4,532	700	ND	339	2,425	5,025	867

*Note:* Years for which there are no survey or weir count data are excluded.

<sup>a</sup> Represents “high count” for season.

<sup>b</sup> Helicopter survey.

<sup>c</sup> Numerous pink salmon made enumerating of chum salmon difficult; pink count may include some chum.

<sup>d</sup> Total counts obtained from counting tower.

<sup>e</sup> Aerial survey, not tower count.

<sup>f</sup> Includes counts from Casadepaga and Ophir Creeks.

<sup>g</sup> Includes counts from Ophir Creek.

<sup>h</sup> Poor survey conditions or partial survey, poor counting tower conditions.

Appendix A18.—Combined aerial survey numbers of chum, pink, coho, and Chinook salmon for Norton Sound, 1990–2013.

Year <sup>a</sup>	Chum	Pink	Coho	Chinook
1990	29,510	796,461	1,594	1,540
1991	69,575	319,459	6,512	2,246
1992	30,597	5,030,222	3,010	1,146
1993	68,980	108,316	6,636	2,869
1994	80,492	5,675,143	4,214	796
1995	118,577	43,393	8,680	1,637
1996	81,364	2,283,129	5,789	1,353
1997	85,026	46,571	3,447	5,260
1998	73,407	3,661,033	2,344	2,186
1999	14,801	4,949	3,439	135
2000	17,748	785,881	7,551	146
2001	39,746	14,978	14,053	751
2002	42,216	386,584	12,116	950
2003	22,880	49,288	6,864	1,240
2004	15,073	6,554,164	19,741	1,300
2005	36,364	1,674,571	21,029	632
2006	59,383	2,468,940	29,398	195
2007	54,522	147,081	18,512	1,645
2008	13,824	3,574,413	23,749	237
2009	17,436	254,132	14,179	1,523
2010	143,275	1,044,784	13,879	410
2011	87,341	87,563	8,391	633
2012	17,138	899,802	4,594	54
2013	100,888	51,689	8,140	462

<sup>a</sup> Rivers surveyed were the Sinuk, Nome, Flambeau, Eldorado, Fish, Niukluk, Kwiniuk, Tubutulik, North, and Boston Creek. Not all rivers were surveyed for all the years. Kwiniuk numbers are from tower counts.

Appendix A19.—Total escapement for chum, pink, coho, and Chinook salmon from weir and tower projects at Kwiniuk, Niukluk, Nome, and Snake rivers (starting 1995), North River (starting 1996), and Eldorado River (starting 1997).

Year	Chum	Pink	Coho <sup>a</sup>	Chinook
1995	138,318	49,409	7,333	626
1996 <sup>b</sup>	124,571	2,535,593	16,175	2,027
1997	109,961	163,728	11,434	5,550
1998	98,166	3,070,848	4,496	2,741
1999	55,352	73,077	10,069	1,846
2000	65,007	1,883,867	19,678	1,324
2001	70,451	79,706	30,645	1,718
2002	93,931	2,239,565	21,625	2,925
2003	49,749	392,827	13,761	2,466
2004	40,494	6,432,486	28,399	2,022
2005	68,585	2,594,334	44,351	1,530
2006	126,045	5,763,830	56,484	1,256
2007	123,394	708,669	37,112	2,324
2008	41,660	3,930,689	49,737	1,250
2009	41,800	275,835	39,236	3,050
2010	191,571	1,490,227	31,058	1,481
2011	99,261	191,243	11,494	955
2012 <sup>c</sup>	50,916	994,745	6,003	1,078
2013 <sup>d</sup>	49,836	72,495	16,616	605

<sup>a</sup> Most projects did not operate during the coho season until 2001.

<sup>b</sup> In 1996 the majority of pink salmon for Nome River escaped through the pickets and were not counted

<sup>c</sup> Most projects were only operational for a short duration during coho season because of high water.

<sup>d</sup> Starting in 2013, there was no longer a counting tower at Niukluk.

Appendix A20.—Total escapement (6 rivers) and catch (commercial, subsistence, and sport fish) for chum, pink, coho, and Chinook salmon for Norton Sound District, 1995–2013.

Year <sup>a, b</sup>	Chum	Pink	Coho	Chinook
1995	213,371	169,496	76,768	15,291
1996 <sup>c</sup>	156,128	3,089,682	112,710	12,617
1997 <sup>d</sup>	161,248	189,439	60,033	25,989
1998 <sup>d</sup>	129,669	3,712,761	52,489	17,105
1999	76,493	95,302	40,546	9,315
2000	85,243	2,091,074	84,983	6,655
2001	97,223	109,878	66,232	6,926
2002	108,444	2,300,537	39,368	8,524
2003	63,099	441,387	45,306	7,445
2004	51,829	6,513,682	87,800	7,027
2005	78,719	2,652,592	146,616	5,280
2006	142,373	5,825,726	217,986	4,961
2007	157,932	734,723	181,306	5,137
2008	75,834	4,067,996	198,406	4,378
2009	85,285	320,632	147,839	6,793
2010	325,633	1,566,755	110,876	3,802
2011	224,511	216,116	82,531	2,560
2012	126,296	1,247,014	57,731	2,510
2013	186,316	100,684	91,357	1,617

<sup>a</sup> Kwiniuk, Niukluk, Nome, and Snake rivers (starting 1995), North River (starting 1996), Eldorado River (starting 1997).

<sup>b</sup> Not all subdistricts from 2004 to 2007 were surveyed for subsistence use.

<sup>c</sup> In 1996, the majority of pink salmon for Nome River escaped through the pickets and were not counted.

<sup>d</sup> Subsistence totals for 1997 and 1998 include data from Savoonga and Gambell.

Appendix A21.–Nome Subdistrict chum salmon estimated escapement, 1999–2013.

Year	Rivers	Aerial Survey Counts	Estimated Escapement <sup>a</sup>	Year	Rivers	Aerial Survey Counts	Estimated Escapement <sup>a</sup>
1999	Nome		1,048	2000	Nome	658	4,056
	Snake <sup>b</sup>		484		Snake <sup>b</sup>		1,911
	Eldorado <sup>b</sup>		4,218		Eldorado <sup>b</sup>	3,383	11,617
	Flambeau	51	637		Flambeau	819	3,947
	Solomon	51	637		Solomon	150	1,294
	Sinuk	1,697	6,370		Sinuk <sup>c</sup>		7,198
	Bonanza	361	2,304		Bonanza	1,130	4,876
			<u>15,698</u>				<u>34,898</u>
2001	Nome	946	2,859	2002	Nome		1,720
	Snake <sup>b</sup>	752	2,182		Snake <sup>b</sup>	402	2,776
	Eldorado <sup>b</sup>	4,450	11,635		Eldorado <sup>b</sup>		10,215
	Flambeau	3,612	10,465		Flambeau	1,876	6,804
	Solomon	280	1,949		Solomon	325	2,150
	Sinuk	3,746	10,718		Sinuk	1,682	6,333
	Bonanza	1,084	4,745		Bonanza	595	3,199
			<u>44,553</u>				<u>33,197</u>
2003	Nome	888	1,957	2004	Nome		3,903
	Snake	440	2,201		Snake		2,146
	Eldorado	1,257	3,591		Eldorado		3,277
	Flambeau	647	3,380		Flambeau	2,250	7,667
	Solomon	73	806		Solomon <sup>c</sup>		1,436
	Sinuk	677	3,482		Sinuk <sup>c</sup>		3,197
	Bonanza	220	1,664		Bonanza <sup>c</sup>		2,166
			<u>17,081</u>				<u>23,792</u>
2005	Nome	2,082	5,584	2006	Nome	394	5,677
	Snake	1,842	2,967		Snake	840	4,160
	Eldorado	5,445	10,369		Eldorado	2,355	42,105
	Flambeau	2,261	7,692		Flambeau	16,000	27,828
	Solomon	775	3,806		Solomon	305	2,062
	Sinuk	1,072	4,710		Sinuk	1,115	4,834
	Bonanza	1,370	5,534		Bonanza	60	708
			<u>40,662</u>				<u>87,374</u>
2007	Nome	1,449	7,034	2008	Nome	106	2,607
	Snake	1,702	8,147		Snake		1,244
	Eldorado	6,315	21,312		Eldorado		6,746
	Flambeau	4,452	12,006		Flambeau	4,235	11,618
	Solomon	673	3,469		Solomon <sup>c</sup>		959
	Sinuk	7,210	16,481		Sinuk <sup>c</sup>		5,367
	Bonanza	2,628	8,491		Bonanza <sup>c</sup>		3,636
			<u>76,940</u>				<u>32,177</u>

-continued-

Appendix A21.–Page 2 of 2.

Year	Rivers	Aerial Survey Counts	Estimated Escapement <sup>a</sup>	Year	Rivers	Aerial Survey Counts	Estimated Escapement <sup>a</sup>
2009	Nome		1,565	2010	Nome	2,998	5,906
	Snake		891		Snake	2,625	6,973
	Eldorado	1,069	4,943		Eldorado <sup>d</sup>	30,600	42,612
	Flambeau	860	4,075		Flambeau	13,600	25,009
	Solomon	89	918		Solomon	454	2,678
	Sinuk	344	2,232		Sinuk	3,955	11,107
	Bonanza	1,851	6,744		Bonanza	686	3,513
			<u>21,368</u>				<u>97,798</u>
2011	Nome		3,582	2012	Nome		2,015
	Snake		4,343		Snake		1,235
	Eldorado		16,227		Eldorado		13,393
	Flambeau	6,283	15,056		Flambeau	7,911	17,517
	Solomon	1,010	4,529		Solomon	165	1,377
	Sinuk	6,265	15,028		Sinuk	3,650	10,537
	Bonanza	2,113	7,357		Bonanza	1,550	6,002
			<u>66,122</u>				<u>52,076</u>
2013	Nome		4,811				
	Snake		2,755				
	Eldorado		26,121				
	Flambeau	16,088	27,928				
	Solomon		1,377				
	Sinuk	19,500	31,691				
	Bonanza	5,284	13,437				
			<u>108,120</u>				

<sup>a</sup> Escapement is estimated by adding Nome, Snake, and Eldorado weir counts and the aerial survey expansion estimates of the other 4 rivers. Aerial survey expansion is calculated as aerial survey count to 0.657142 power multiplied by 48.059 (Clark, 2001), unless otherwise footnoted.

<sup>b</sup> Escapement was estimated by counting tower.

<sup>c</sup> Because of the lack of aerial survey estimates, method used (from Clark, 2001) was Solomon = 0.368 multiplied by Nome escapement, Sinuk = 1.476 multiplied by Bonanza escapement, and Bonanza = 0.198 multiplied by Eldorado and Flambeau escapements combined.

<sup>d</sup> Weir was breached and aerial survey expansion count was used.

Appendix A22.–Historical escapement of salmon and Dolly Varden at Eldorado River counting tower, 1997–2002 and weir, 2003–2013.

Year	Operating Period	Chinook	Chum	Pink	Coho	Sockeye	Dolly Varden
1997	June 29–Aug 19	98	14,302	1,022	194	n/a	n/a
1998	June 29–Aug 12	8	13,808	137,283	21	n/a	n/a
1999	July 10–Sept 01	28	4,218	977	510	n/a	n/a
2000	June 29–Aug 25	33	11,617	55,992	192	n/a	n/a
2001	July 08–Sept 13	50	11,635	488	1,509	n/a	n/a
2002	June 24–Sept 10	26	10,215	119,098	540	10	377
2003	June 21–Sept 08	29	3,591	173	115	0	60
2004	June 22–Sept 09	25	3,277	60,866	1,151	57	0
2005	June 23–Sept 02	32	10,369	12,356	689	10	23
2006	June 26–Aug 03	41	42,105	222,348	55	1	65
2007	June 26–Aug 06	14	21,312	833	2	22	60
2008	June 27–July 31	36	6,746	244,641	38	3	14
2009	July 02–Aug 03	31	4,943	1,119	2	0	72
2010 <sup>a</sup>	June 30–July 24	23	42,612	48,136	2	8	72
2011	June 30–Aug 03	3	16,227	489	1	0	2
2012	July 04–Aug 15	0	13,393	59,952	1	0	30
2013	July 01–Aug 06	9	26,121	1,025	15	0	2

<sup>a</sup> Numerous breaches in weir during the season resulted in minimal counts, except for chum salmon count that was determined by aerial survey expansion from the aerial survey count.

Appendix A23.—Historical escapement of salmon and Dolly Varden at Snake River counting tower 1995–2002 and weir 2003–2013.

Year	Operating Period	Chinook	Chum	Pink	Coho	Sockeye	Dolly Varden
1995	July 01–Aug 18	0	4,393	917	856	0	NA
1996	July 03–Aug 22	5	2,772	44,558	1,638	0	NA
1997	July 07 –Aug 18	12	6,184	6,742	1,157	0	NA
1998	July 01–Aug 11	0	11,067	219,679	178	0	NA
1999	July 01–Aug 14	20	484	116	90	0	NA
2000	June 29–Aug 25	28	1,911	4,723	406	0	NA
2001	July 08–Sept 05	33	2,182	1,295	1,335	0	NA
2002	June 28–Sept 16	9	2,776	4,103	851 <sup>a</sup>	8	149
2003	June 26–Sept 11	50	2,201	2,856	489	84	111
2004	June 23–Sept 03	17	2,146	126,917	474	22	290
2005	June 27–Sept 11	31	2,967	13,813	2,948	275	28
2006	July 01–Sept 11	32	4,160	74,028	4,776	302	614
2007	July 01–Sept 14	61	8,147	4,634	1,781	1,354	121
2008	July 06–Sept 06	13	1,244	145,761	5,206	143	452
2009	July 08–Aug 30 <sup>b</sup>	6	891	769	50	2	14
2010	July 03–Sept 11	43	6,973	51,099	2,243	124	198
2011	July 08–Sept 11	1	4,343	7,011	343	14	5
2012	July 06–Aug 15 <sup>c</sup>	1	1,235	5,954	14	3	3
2013	July 19–Sept 10	8	2,755	1,333	1,203	163	1

<sup>a</sup> Includes 442 coho salmon estimated by aerial survey to be holding below the weir site after the weir was removed.

<sup>b</sup> Weir was not fish tight last week of August and hundreds of coho salmon passed through the weir without being counted.

<sup>c</sup> Weir was knocked out for 13 days in late July and early August. An interpolation was made for chum salmon.



Appendix A24.–Historical salmon escapement at Kwiniuk River counting tower, 1990–2013.

Year	Operating Period	Chum	Pink	Chinook	Coho
1990	June 21–July 25	13,957	416,512	900	0
1991	June 18–July 27	19,801	53,499	708	0
1992	June 27–July 28	12,077	1,464,716	479	0
1993	June 27–July 27	15,824	43,063	600	0
1994	June 23–Aug 09	33,012	2,303,114	625	2,547
1995	June 21–July 26	42,500	17,511	498	114
1996	June 20–July 25	28,493	907,893	577	461
1997	June 18–July 27	20,119	9,535	974	0
1998	June 18–July 27	24,247	655,934	303	0
1999	June 25–July 28	8,763	607	116	0
2000	June 22–July 27	12,879	750,173	144	41
2001	June 27–Sept 15	16,598	8,423	261	9,532
2002	June 17–Sept 11	37,995	1,114,410	778	6,459
2003	June 15–Sept 15	12,123	22,329	744	5,490
2004	June 16–Sept 14	10,362	3,054,684	663	11,240
2005	June 17–Sept 13	12,083	341,048	342	12,950
2006	June 22–Sept 12	39,519	1,347,090	195	22,341
2007	June 21–Sept 10	27,756	54,255	258	9,429
2008	June 23–Sept 07	9,483	1,444,213	237	10,461
2009	June 24–Sept 13	8,739	42,962	444	8,677
2010	June 25–Sept 7	71,388	634,220	135	8,049
2011	June 20–Sept 11	31,604	30,023	57	3,288
2012	June 23–Aug 16	5,577	393,302	54	777
2013	June 24–Sept 16	5,631	13,212	15	3,940

Appendix A25.–Historical salmon escapement at Niukluk River counting tower, 1995–2012.

Year	Operating Period	Chum	Pink	Chinook	Coho
1995	June 29–Sept 12	86,332	17,088	123	4,713
1996	June 23–Sept 12	80,178	1,154,922	243	12,781
1997	June 28–Sept 09	57,305	10,468	259	3,994
1998	July 04–Aug 13	45,588	1,624,438	260	840
1999	July 04–Sept 04	35,239	20,351	40	4,260
2000	July 04–Aug 27	29,573	961,603	48	11,382
2001	July 10–Sept 08	30,662	41,625	30	3,468
2002	June 25–Sept 10	35,307	645,141	621	7,391
2003	June 25–Sept 10	20,018	75,855	179	1,282
2004	June 25–Sept 08	10,770	975,895	141	2,064
2005	June 28–Sept 09	25,598	270,424	41	2,727
2006	June 26–Sept 08	29,199	1,371,919	39	11,169
2007	July 01–Sept 04	50,994	43,617	30	3,498
2008	July 01–Sept 06	12,078	669,234	33	13,779
2009	July 03–Sept 02	15,879	24,204	204	6,861
2010	July 01–Sept 01	48,561	434,205	15	9,042
2011	June 28–Sept 06	23,607	15,425	18	2,405
2012	July 04–Aug 17	19,576	249,212	21	1,729

*Note:* The Niukluk River counting tower project was discontinued after 2012.

Appendix A26.–Historical salmon escapement at Nome River counting tower, 1993–1995, and weir, 1996–2013.

Year	Operating Period	Chum	Pink	Chinook	Coho	Sockeye
1993	July 25–Aug 28	1,859	13,036	63	4,349	
1994	June 24–Aug 15	2,893	142,604	54	726	
1995	June 22–Sept 06	5,093	13,893	5	1,650	
1996	June 26–July 23	3,339	95,681 <sup>a</sup>	5	66	
1997	June 27–Aug 27	5,147	8,035	22	321	
1998	July 01–Aug 11	1,930	359,469	70	96	
1999	July 02–Aug 25	1,048	2,033	3	417	6
2000	June 29–Aug 25	4,056	41,673	25	698	19
2001	July 08–Sept 11	2,859	3,138	7	2,418	55
2002	June 29–Sept 11	1,720	35,057	7	3,418	29
2003	July 05–Sept 10	1,957	11,402	12	548	47
2004	June 25–Sept 12	3,903	1,051,146	51	2,283	114
2005	June 27–Sept 11	5,584	285,759	69	5,848	381
2006	July 02–Sept 07	5,677	578,555	43	8,308	188
2007	July 03–Sept 16	7,034	24,395	13	2,437	534
2008	July 02–Sept 17	2,607	1,186,554	28	4,605	90
2009	July 01–Sept 20	1,565	16,490	10	1,370	103
2010	June 30–Sept 16	5,906	171,760	9	4,114	43
2011	July 01–Sept 12	3,582	14,403	12	1,833	22
2012	July 04–Aug 15	2,015	149,119	6	224	48
2013	July 05–Sept 16	4,811	10,257	9	2,624	38

<sup>a</sup> In 1996 the majority of pink salmon escaped through the pickets and was not counted.

Appendix A27.–Salmon escapement at Solomon River weir, 2013.

Year	Operating Period	Chum	Pink	Chinook	Coho	Sockeye
2013	July 05–Aug 26	1,377	2,733	0	178	3

*Note:* The Solomon River weir was initiated in 2013.

Appendix A28.–Historical sockeye salmon escapement at Glacial Lake weir, 2000–2013.

Year	Operating Period	Chum <sup>a</sup>	Pink <sup>b</sup>	Sockeye
2000	July 11–July 30			884
2001	July 02–July 28	1		2,487
2002	June 25–July 26			1,047
2003	June 24–July 28			2,004
2004	June 18–July 25	1		8,115
2005	June 20–July 25			11,135
2006	July 04–July 18			6,849
2007	July 05–July 20			4,533
2008	June 27–July 28	10	614	1,794
2009	June 20–July 27			826
2010	June 26–July 28			1,047
2011	June 28–July 26	4		1,697
2012 <sup>c</sup>	July 01–Aug 09	25	165	1,636
2013 <sup>d</sup>	June 20–Aug 12	35	2	2,544

<sup>a</sup> Chum salmon will pass upstream through the Glacial Lake weir and often exit the lake back downstream through the weir.

<sup>b</sup> Pink salmon have been observed often in even-numbered years, but 2008 was the first year the crew was instructed to enumerate pink salmon passage.

<sup>c</sup> A video project was tested during 2012 and was in operation 11 days (July 31 to August 9) after human occupation of the weir site. Included in totals are 34 sockeye, 12 pink, and 10 chum salmon that were counted by camera during that time.

<sup>d</sup> A video project was in operation from July 14 to August 12.

Appendix A29.–Historical salmon escapement at Inglutalik River counting tower, 2011–2013.

Year	Operating Period	Chum	Pink	Chinook	Coho
2011	June 24–Aug 14	64,892	494,099	1,467	870
2012	June 23–Aug 23	32,832	90,349	1,134	1,431
2013	June 21–Aug 11	61,259	268,537	860	5,904

*Note:* Due to speciation problems in 2013, the Chinook and coho counts are probably inaccurate.

Appendix A30.–Historical salmon escapement at North River counting tower, 1996–2013.

Year	Operating Period	Chum	Pink	Chinook	Coho
1996	June 16–July 25	9,789	332,539	1,197	1,229
1997	June 16–Aug 21	6,904	127,926	4,185	5,768
1998	June 15–Aug 12	1,526	74,045	2,100	3,361
1999	June 30–Aug 31	5,600	48,993	1,639	4,792
2000	June 17–Aug 12	4,971	69,703	1,046	6,959
2001	July 05–Sept 15	6,515	24,737	1,337	12,383
2002	June 19–Aug 29	5,918	321,756	1,484	2,966
2003	June 15–Sept 13	9,859	280,212	1,452	5,837
2004	June 15–Sept 14	10,036	1,162,978	1,125	11,187
2005	June 15–Sept 15	11,984	1,670,934	1,015	19,189
2006	June 18–Sept 11	5,385	2,169,890	906	9,835
2007	June 16–Sept 05	8,151	580,935	1,948	19,965
2008	June 19–Sept 13	9,502	240,286	903	15,648
2009	June 19–Sept 11	9,783	190,291	2,355	22,276
2010	June 19–Sept 07	16,131	150,807	1,256	7,608
2011	June 17–Sept 08	19,898	123,892	864	3,624
2012	June 21–Aug 19	9,120	137,006	996	3,258
2013	July 01–Aug 05	10,518	46,668	564	8,834

Appendix A31.–Historical salmon escapement at Unalakleet River weir, 2010–2013.

Year	Operating Period	Chum	Pink	Chinook	Coho	Sockeye
2010	June 22–July 31	70,811	832,904	1,021	5,382	130
2011	June 17–Aug 07	108,770	363,906	1,111	10,418	190
2012	June 24–Aug 15	70,669	672,083	807	17,766	245
2013	June 20–Aug 22	113,953	144,225	767	25,566	243

Appendix A32.—Chum salmon escapement by river, Nome Subdistrict, 1993–2013.

Year	Rivers West of Cape Nome			Rivers East of Cape Nome				Total <sup>e</sup>
	Sinuk <sup>a</sup>	Snake <sup>b</sup>	Nome <sup>c</sup>	Flambeau <sup>a</sup>	Eldorado <sup>d</sup>	Bonanza <sup>a</sup>	Solomon <sup>a</sup>	
1993	6,052	2,115	5,925	6,103	9,048	3,007	2,525	34,775
1994	4,905	3,519	2,893	12,889	13,202	5,178	1,066	43,652
1995	9,464	4,395	5,093	16,474	18,955	11,182	2,106	67,669
1996	6,658	2,772	3,339	13,613	32,970	7,049	2,141	68,542
1997	9,212	6,184	5,147	9,455	14,302	4,140	2,111	50,551
1998	6,720	11,067	1,930	9,129	13,808	4,552	925	48,131
1999	6,370	484	1,048	637	4,218	2,304	637	15,698
2000	7,198	1,911	4,056	3,947	11,617	4,876	1,294	34,899
2001	10,718	2,182	2,859	10,465	11,635	4,745	1,949	44,553
2002	6,333	2,776	1,720	6,804	10,243	3,199	2,150	33,225
2003	3,482	2,201	1,957	3,380	3,591	1,664	806	17,081
2004	3,197	2,145	3,903	7,667	3,273	2,166	1,436	23,787
2005	4,710	2,948	5,584	7,692	10,426	5,534	1,914	38,808
2006	4,834	4,128	5,677	27,828	41,985	708	2,062	87,222
2007	16,481	8,147	7,034	12,006	21,312	8,491	3,469	76,940
2008	5,367	1,244	2,607	11,618	6,746	3,636	959	32,177
2009	2,232	891	1,565	4,075	4,943	6,744	918	21,368
2010	11,107	6,973	5,906	25,009	42,612	3,513	2,678	97,798
2011	15,028	4,343	3,582	15,056	16,227	7,357	4,529	66,122
2012	10,537	1,235	2,015	17,517	13,393	6,002	1,377	52,076
2013	31,691	2,755	4,811	27,928	26,121	13,437	1,377	108,120
Total	182,296	74,415	78,651	249,292	330,627	109,484	38,429	1,063,194

<sup>a</sup> Sinuk, Flambeau, Bonanza, and Solomon rivers' escapements are estimated by aerial survey, but in 2013, Solomon River escapement was a weir count.

<sup>b</sup> Snake River escapements are estimated by aerial survey (1993–1994), tower counts (1995–2002), and weir counts (2003–2012). Escapement goal range is 1,600–2,500 chum salmon.

<sup>c</sup> Nome River escapements are estimated by aerial survey expansion (1993), tower counts (1994–1995), and weir counts (1996–2012). Escapement goal range is 2,900–4,300 chum salmon.

<sup>d</sup> Eldorado River escapements are estimated by aerial survey (1993–1996), tower counts (1997–2002), and weir counts (2003–2012). Escapement goal range is 6,000–9,200 chum salmon.

<sup>e</sup> Subdistrict 1 BEG is 23,000–35,000 chum salmon.

Appendix A33.–Pink salmon escapement by year and river, Nome Subdistrict, 1993–2013.

Year	Rivers West of Cape Nome			Rivers East of Cape Nome				Total
	Sinuk <sup>a</sup>	Snake <sup>b</sup>	Nome <sup>c</sup>	Flambeau <sup>a</sup>	Eldorado <sup>d</sup>	Bonanza <sup>a</sup>	Solomon <sup>a</sup>	
1993	5,120		13,036	5,584	120	ND	ND	23,860
1994	492,100	63,860	142,604	19,202	53,890	20	ND	771,676
1995	1,250	917	13,893	8,086	4,243	619	350	29,358
1996	74,400	44,558	95,681	17,182	46,100	40,510	15,230	333,661
1997	1,200	6,742	8,035	2,117	1,022	ND	80	19,196
1998	342,100	219,679	359,469	8,720	137,283	167,130	45,175	1,279,556
1999	180	116	2,033	1,251	977	245	90	4,892
2000	12,175	4,723	41,673	2,159	55,992	12,410	2,899	132,031
2001	115	1,295	3,138	924	488	221	ND	6,181
2002	28,487	4,103	35,057	2,233	119,098	17,095	9,170	215,243
2003	9,907	2,856	11,402	194	173	1,540	157	26,229
2004	1,267,100	126,917	1,051,146	7,351	60,866	185,000	109,000	2,807,380
2005	211,285	13,813	285,759	873	12,356	55,000	11,100	590,186
2006	515,000	74,028	578,555	6,556	222,348	268,500	165,215	1,830,202
2007	6,810	4,634	24,395	336	833	1,360	2,400	40,768
2008	1,496,000	145,761	1,186,554	3,510	244,641	212,000	81,000	3,369,466
2009	6,740	769	16,490	175	1,119	3,276	1,565	30,134
2010	168,600	51,099	171,760	4,797	48,136	106,000	21,804	572,196
2011	21,100	7,011	14,403	58	489	11,050	5,580	59,691
2012	506,500	5,954	149,119	2,657	59,318	54,700	15,000	793,248
2013	143,921	1,333	10,257	ND	1,025	800	2,733	160,069
Total	5,310,090	780,168	4,214,459	93,965	1,070,517	1,137,476	488,548	13,095,223

<sup>a</sup> Sinuk, Flambeau, Bonanza, and Solomon rivers' escapements are estimated by aerial survey, but in 2013, Solomon River escapement was a weir count.

<sup>b</sup> Snake River escapements are estimated by aerial survey (1993–1994), tower counts (1995–2002), and weir counts (2003–2013).

<sup>c</sup> Nome River escapements are estimated by tower counts (1993–1995) and weir counts (1996–2013). Escapement goal range is 13,000 pink salmon in even-numbered years and 3,200 pink salmon in odd-numbered years.

<sup>d</sup> Eldorado River escapements are estimated by aerial survey (1993–1996), tower counts (1997–2002), and weir counts (2003–2013).

Appendix A34.—Number of customary trade permits issued, Norton Sound District and Port Clarence District, 2007–2013.

Year	Norton Sound District								Port Clarence District			Value
	Nome	Golovin	Elim	Koyuk	Shaktoolik	Unalakleet	St. Michael	Stebbins	Teller	Brevig Mission	Wales	
2007	3	0	2	0	0	0	0	0	0	0	0	\$200.00
2008	3	0	0	0	0	0	0	0	0	0	0	\$0.00
2009	0	0	0	0	0	0	0	0	0	0	0	Confidential
2010	1	0	0	0	0	0	0	0	0	0	0	Confidential
2011	0	0	0	0	0	0	1	0	0	0	0	Confidential
2012	2	0	0	0	0	0	0	0	0	0	0	Confidential
2013	4	4	1	0	0	0	0	0	3	6	0	\$1,790.00





## **APPENDIX B: PORT CLARENCE FISHERIES**

Appendix B1.—Comparative sockeye salmon aerial survey indices, Port Clarence District, 1990–2013.

Year	Salmon Lake	Grand Central River	Total
1990	2,834	926	3,760
1991	3,790	1,570	5,360
1992	1,500	<sup>a</sup>	1,500
1993	2,885	216	3,092
1994	3,740	1,230	4,970
1995	5,433	628 <sup>b</sup>	6,061
1996	6,610	770	7,380
1997	8,760	1,520	10,280
1998	5,210	1,977	7,187
1999	31,720	1,780	33,500
2000	12,772	<sup>a</sup>	12,772
2001	9,400	155	9,555
2002	3,520	71	3,591
2003	19,275	1,015	20,290
2004	23,005	2,855	25,860
2005	41,500	740	42,240
2006	39,400	2,380	41,780
2007	14,920	5,692	20,612
2008	9,420	2,252	11,672
2009	136	50	186
2010	73	711	784
2011	4,604	540	5,144
2012	4,730	1,100	5,830
2013	5,820	1,151	6,971

<sup>a</sup> No survey occurred.

<sup>b</sup> Early count.

Appendix B2.—Historical escapement of salmon and Dolly Varden at Pilgrim River counting tower (1997–2002) and weir (2003–2013).

Year	Operating Period	Chinook	Chum	Pink	Coho	Sockeye	Dolly Varden
1997	July 12–Aug 21	356	15,619 <sup>a</sup>	5,557	452	15,619 <sup>a</sup>	NA
1998	Did not operate						
1999	July 13–Aug 06	6	2,617	35,577	104	4,650	NA
2000	July 05–Aug 18	72	861	374	21	12,141	NA
2001	Did not operate						
2002	July 04–Aug 04	150	5,590	3,882	246	3,888	NA
2003	June 21–Sept 14	1,016	15,200	14,100	677	42,729	550
2004	June 21–Sept 14	925	10,239	50,760	1,573 <sup>b</sup>	85,417	264
2005	June 24–Sept 05	216	9,685	13,218	304	55,951	112
2006	June 30–Sept 09	275	45,361	17,701	973	52,323	505
2007	June 29–Sept 10	501	35,334	3,616	605	43,432	339
2008	June 25–Sept 01	137	24,550	92,471	260	20,452	409
2009	June 26–Aug 31	52	5,427	483	18	953	130
2010	June 24–Sept 01	44	25,379	29,239	272	1,654	285
2011	June 28–Sept 01	44	41,740	3,364	269	8,449	229
2012	June 26–Aug 18	64	25,521	46,135	95	7,085	65
2013	June 27–Sept 08	47	47,557	1,060	890	12,428	27

<sup>a</sup> Chum and sockeye salmon escapements were combined due to species identification problems during 1997.

<sup>b</sup> Coho salmon were misidentified. Nearly 30% of scale samples in 2004 were actually sockeye salmon.

Appendix B3.—Estimated number of subsistence fishing families and harvest in Port Clarence District, 1994–2013.

Year		Number of fishing families interviewed	Chinook	Sockeye	Coho	Pink	Chum	Total
1994	<sup>a</sup>	127	203	2,220	1,892	4,309	2,294	10,918
1995	<sup>a</sup>	122	76	4,481	1,739	3,293	6,011	15,600
1996	<sup>a</sup>	117	194	2,634	1,258	2,236	4,707	11,029
1997	<sup>a</sup>	126	158	3,177	829	755	2,099	7,018
1998	<sup>a</sup>	138	289	1,696	1,759	7,815	2,621	14,180
1999	<sup>a</sup>	155	89	2,392	1,030	786	1,936	6,233
2000	<sup>a</sup>	134	72	2,851	935	1,387	1,275	6,520
2001	<sup>a</sup>	160	84	3,692	1,299	1,183	1,910	8,168
2002	<sup>a</sup>	159	133	3,732	2,194	3,394	2,699	12,152
2003	<sup>a,b</sup>	204	177	4,495	1,434	4,113	2,430	12,649
2004	<sup>c</sup>	376 <sup>d</sup>	278	8,688	1,131	5,918	2,505	18,520
2005	<sup>c</sup>	335 <sup>d</sup>	152	8,492	726	6,615	2,479	18,464
2006	<sup>c</sup>	345 <sup>d</sup>	102	9,940	1,061	4,939	4,353	20,395
2007	<sup>c</sup>	363 <sup>d</sup>	85	9,484	705	1,468	4,454	16,196
2008	<sup>c</sup>	408 <sup>d</sup>	125	5,069	512	7,527	2,449	15,682
2009	<sup>c</sup>	326 <sup>d</sup>	40	1,643	804	1,882	3,060	7,429
2010	<sup>c</sup>	290 <sup>d</sup>	63	824	596	5,202	5,232	11,917
2011	<sup>c</sup>	267 <sup>d</sup>	57	1,611	393	2,610	4,338	9,008
2012	<sup>c</sup>	335 <sup>d</sup>	44	1,422	703	5,200	7,802	15,171
2013	<sup>c</sup>	431 <sup>d</sup>	38	5,243	651	1,788	6,588	14,308

<sup>a</sup> Harvest estimate from ADF&G Division of Subsistence survey.

<sup>b</sup> Includes harvest reported from 59 Pilgrim River permits. In total, 101 permits were issued and 79 were returned.

<sup>c</sup> Beginning in 2004 a permit was required for Port Clarence District (including Pilgrim River and Salmon Lake) that replaced household surveys.

<sup>d</sup> The number is all permits issued for the Port Clarence District (including Pilgrim River and Salmon Lake permits).

Appendix B4.–Application of 20-05-00 liquid blend of phosphorous and nitrogen fertilizer to Salmon Lake, 1997–2013.

Year	Fertilizer (tons)	Organization
1997	40	NSEDC/ADF&G/BLM
1998	40	NSEDC/ADF&G/BLM
1999	40	NSEDC/ADF&G/BLM
2000	40	NSEDC/ADF&G/BLM
2001	40	NSEDC/ADF&G/BLM
2002	0	
2003	0	
2004	27	NSEDC/ADF&G
2005	0	
2006	0	
2007	16	NSEDC
2008	8	NSEDC
2009	28	NSEDC
2010	19	NSEDC
2011	11	NSEDC
2012	10	NSEDC
2013	10	NSEDC



## **APPENDIX C: KOTZEBUE FISHERIES**

Appendix C1.–Kotzebue District chum salmon catch statistics, 1990–2013.

Year	Total Catch	Number of fishermen	Season catch per fisherman
1990	163,263	153	1,067
1991	239,923	142	1,690
1992	289,184	149	1,941
1993 <sup>a</sup>	73,071	114	641
1994	153,452	109	1,408
1995	290,730	92	3,160
1996	82,110	55	1,493
1997	142,720	68	2,099
1998	55,907	45	1,242
1999	138,605	60	2,310
2000	159,802	64	2,497
2001	211,672	66	3,207
2002	8,390	3	2,797
2003	25,423	4	6,356
2004	51,038	43	1,187
2005	75,971	41	1,853
2006	137,961	42	3,285
2007	147,087	46	3,198
2008	190,550	48	3,970
2009	187,562	62	3,025
2010	270,343	67	4,035
2011	264,225	89	2,970
2012	227,965	83	2,747
2013	319,062	66	4,834
Avg 1990–2012	155,955	72	2,529

<sup>a</sup> Includes 2,000 chum salmon from the Sikusuilag Springs Hatchery terminal fishery.



Appendix C2.–Kotzebue District chum salmon type of processing and weights, 1990–2013.

Year	Chum salmon (pounds)	Other <sup>a</sup>	Fresh frozen salmon roe (pounds)
1990	1,453,040	538	
1991	1,951,041	714	
1992	2,397,302	2,714	
1993 <sup>b</sup>	613,968	1,507	1,000
1994 <sup>c</sup>	1,166,494	73	
1995	2,329,898	93	
1996 <sup>d</sup>	97,510	51	
1997	1,141,741	649	
1998	447,256	2,971	
1999	1,108,898	87	
2000	1,370,637	106	
2001	1,847,361	64	
2002	74,341	0	
2003	218,091	0	
2004	419,059	1,450	
2005	621,573	1,258	
2006	1,040,023	0	
2007	1,209,842	0	
2008	1,541,922	0	
2009	1,505,734	0	
2010	2,160,264	0	
2011	2,158,365	0	
2012	1,751,473	0	
2013	2,555,304	0	

*Note:* Data not available for all years.

<sup>a</sup> Chinook and pink salmon, and Dolly Varden.

<sup>b</sup> Includes 11,160 pounds from the Sikusuilaq Springs Hatchery terminal fishery. Pounds of roe stripped are from a verbal report.

<sup>c</sup> Includes 31,500 pounds commercially caught but not reported on fish tickets.

<sup>d</sup> Includes 17,600 pounds commercially caught but not sold on fish tickets.

Appendix C3.—Kotzebue District mean prices paid per pound in dollars to salmon fishermen by species, 1990–2013.

Year	Chum salmon		Chinook salmon	Inconnu	Dolly Varden
	Average weight	Average price			
1990	8.9	0.31	2.00		0.25
1991	8.1	0.22	1.64	0.50	0.18
1992	8.3	0.22	1.89	0.58	0.10
1993	8.5	0.38	2.37	0.50	0.10
1994	7.8	0.20	1.14		0.17
1995	8.0	0.13	1.00	0.50	0.20
1996	8.0	0.09	1.00	0.44	0.25
1997	8.0	0.16	1.02		0.20
1998 <sup>a</sup>	8.0	0.15	1.00		0.20
1999 <sup>a</sup>	8.0	0.16	1.00		0.20
2000	8.6	0.18	1.00		0.20
2001	8.7	0.17	1.00		
2002	8.9	0.10	0.00		
2003	8.6	0.12	0.00		0.50
2004	8.2	0.15	0.72		0.26
2005	8.2	0.20	0.50		0.30
2006	7.5	0.22			
2007	8.2	0.20			
2008	8.1	0.25			
2009	8.0	0.25			
2010	8.0	0.40			
2011	8.2	0.40			
2012	7.7	0.32			
2013	8.0	0.27			

*Note:* Information is not available for some species in some years.

<sup>a</sup> Each chum salmon was assumed to weigh 8 pounds, but no fish were weighed individually.

Appendix C4.–Kotzebue District commercial fishery dollar value estimates, 1990–2013.

Year	Gross value of catch to fishermen <sup>a</sup>	Number of fishermen	Average value per fisherman
1990	\$438,044	153	\$2,863
1991	\$437,948	142	\$3,084
1992	\$533,731	149	\$3,582
1993 <sup>b</sup>	\$235,061	114	\$2,062
1994	\$233,512	109	\$2,142
1995	\$316,031	92	\$3,435
1996	\$56,310	55	\$1,024
1997	\$187,978	68	\$2,764
1998	\$70,587	45	\$1,569
1999	\$179,781	60	\$2,996
2000	\$246,786	64	\$3,856
2001	\$322,650	66	\$4,889
2002	\$7,572	3	\$2,524
2003	\$26,377	4	\$6,594
2004	\$64,420	43	\$1,498
2005	\$124,820	41	\$3,044
2006	\$229,086	42	\$5,454
2007	\$243,149	46	\$5,286
2008	\$385,270	48	\$8,026
2009	\$376,554	62	\$6,073
2010	\$860,125	67	\$12,838
2011	\$867,085	89	\$9,743
2012	\$567,664	83	\$6,839
2013	\$689,163	66	\$10,442
Avg 1990–2012	\$304,806	72	\$4,443

<sup>a</sup> Values represent chum salmon value and incidental species such as char, whitefish, and other salmon.

<sup>b</sup> Includes \$3,648 from Sikusuilaq Springs Hatchery terminal fishery.

Appendix C5.–Kotzebue District commercial (1990–2013) and subsistence salmon catches (1990–2004 and 2012–2013).

Year	Commercial catch			Subsistence catch			Total documented catch	
	Chum	Other <sup>a</sup>	Total	Chum	Number of fishermen interviewed	Average catch per fisherman		
1990	163,263	32	163,295	8,268	<sup>b</sup>	<sup>b</sup>	163,295	
1991	239,923	44	239,967	14,740	<sup>b</sup>	<sup>b</sup>	239,967	
1992	289,184	204	289,388	14,303	<sup>b</sup>	<sup>b</sup>	289,388	
1993	73,071 <sup>c</sup>	131	131	15,430	<sup>b</sup>	<sup>b</sup>	131	
1994	153,452 <sup>d</sup>	3	3	36,226 <sup>e</sup>	375	97	36,229	
1995	290,730	5	290,735	102,881	593	173	393,616	
1996	82,110 <sup>f</sup>	3	3	99,740	596	167	99,743	
1997	142,720	45	142,765	57,906	530	109	200,671	
1998	55,907	210	56,117	48,980	592	83	105,097	
1999	139,120	5	139,125	94,342	353	267	233,467	
2000	159,802	10	159,812	65,975	422	156	225,787	
2001	211,672	6	211,678	49,232	408	121	260,910	
2002	8,390	0	8,390	16,880 <sup>e,g</sup>	191	88	25,270	
2003	25,423	0	25,423	19,201 <sup>e</sup>	446	43	44,624	
2004	51,038	116	51,154	24,637 <sup>e</sup>	440	63	75,791	
2005	75,971	7	75,978	Subsistence surveys were not conducted.				
2006	137,961	17	137,978	Subsistence surveys were not conducted.				
2007	147,087	20	147,107	Subsistence surveys were not conducted.				
2008	190,550	742	191,292	Subsistence surveys were not conducted.				
2009	187,562	106	187,668	Subsistence surveys were not conducted.				
2010	270,343	583	270,926	Subsistence surveys were not conducted.				
2011	264,321	166	264,487	Subsistence surveys were not conducted.				
2012	227,965	476	228,441	26,693 <sup>e</sup>	360	74	255,134	
2013	319,062	114	319,176	Information is not yet available.				
Average 2003–'12	157,822	223	158,045	Average 1995–'04	57,977	457	127	166,498

<sup>a</sup> Includes Chinook, pink, and sockeye salmon that were not sold but retained for personal use.<sup>b</sup> Information not available.<sup>c</sup> Includes 2,000 chum salmon from the Sikusuilag Springs Hatchery terminal fishery.<sup>d</sup> Includes 4,000 chum salmon commercially harvested on August 5 but not sold.<sup>e</sup> Does not include the town of Kotzebue.<sup>f</sup> Includes 2,200 chum salmon commercially harvested on July 29 but not sold.<sup>g</sup> Only 2 of 6 villages surveyed.

Appendix C6.–Kotzebue District subsistence chum salmon catches by village, 1990–2004 and 2012–2013.

Year	Village					Kobuk River Villages	Noatak Village	Village					District Total
	Noorvik	Kiana	Ambler	Shungnak	Kobuk			Kotzebue	Deering	Kivalina	Buckland	Shishmaref	
1990	4,353	a	a	a	a	4,353	3,915	a	a	a	a	a	8,268
1991	6,855	a	a	4,248	a	11,103	3,637	a	a	a	a	a	14,740
1992	8,370	a	a	3,890	a	12,260	2,043	a	a	a	a	a	14,303
1993	8,430	a	a	3,730	a	12,160	3,270	a	a	a	a	a	15,430
1994	8,157	1,891	2,860	7,982	5,722	26,612	6,126	a	3,488	a	a	a	36,226
1995	15,485	5,985	8,558	5,880	2,959	38,867	6,359	50,708	a	a	a	6,947	102,881
1996	13,611	5,935	9,062	8,649	1,819	39,076	10,091	50,573	a	a	a	a	99,740
1997	14,323	3,064	2,713	5,513	629	26,242	5,309	26,355	a	a	a	a	57,906
1998	9,845	3,414	2,432	4,676	1,031	21,398	2,614	24,968	a	a	a	a	48,980
1999	17,843	3,788	590	3,868	1,869	27,958	1,616	64,768	a	a	a	a	94,342
2000	10,391	2,876	5,009	2,944	318	21,538	7,293	37,144	a	a	a	a	65,975
2001	16,540	5,500	a	4,310	2,843	29,193	2,326	17,713	a	a	a	a	49,232
2002	13,943	b	b	b	b	b	2,937	b	a	a	a	a	16,880
2003	7,982	3,010	1,719	2,860	1,453	17,024	2,177	a	a	a	a	a	19,201
2004	6,025	3,896	3,446	4,186	3,087	20,640	3,997	a	a	a	a	a	24,637
2012	9,584	2,442	1,621	2,595	2,637	18,879	7,814	a	a	a	a	a	
2013	Information is not yet available.												

*Note:* No subsistence surveys were conducted from 2005 to 2011. Kotzebue area villages were surveyed by the Division of Subsistence in 2012 and 2013, but data are not yet available for 2013.

<sup>a</sup> Not surveyed.

<sup>b</sup> The Kotzebue Sound communities of Ambler, Kiana, Kobuk, Kotzebue, and Shungnak, although normally included, were not surveyed in 2002 (Georgette et al. 2003).

Appendix C7.–Kotzebue District average subsistence chum salmon harvest per household by village, 1990–2004 and 2012–2013.

Year	Kotzebue	Noatak	Noorvik	Kiana	Ambler	Shungnak	Kobuk	Deering
1990	<sup>a</sup>	135	198	<sup>a</sup>	<sup>a</sup>	<sup>a</sup>	<sup>a</sup>	<sup>a</sup>
1991	<sup>a</sup>	145	311	<sup>a</sup>	<sup>a</sup>	283	<sup>a</sup>	<sup>a</sup>
1992	<sup>a</sup>	89	310	<sup>a</sup>	<sup>a</sup>	243	<sup>a</sup>	<sup>a</sup>
1993	<sup>a</sup>	136	312	<sup>a</sup>	<sup>a</sup>	196	<sup>a</sup>	<sup>a</sup>
1994	<sup>a</sup>	90	133	32	99	154	260	92
1995	71	69	123	59	110	111	110	<sup>a</sup>
1996	73	115	117	58	111	154	76	<sup>a</sup>
1997	41	71	125	35	39	117	28	<sup>a</sup>
1998	35	27	79	34	30	84	41	<sup>a</sup>
1999	78	18	151	42	8	76	81	<sup>a</sup>
2000	48	72	93	33	72	64	11	<sup>a</sup>
2001	23	24	152	62	<sup>a</sup>	94	109	<sup>a</sup>
2002	<sup>a</sup>	29	121	<sup>a</sup>	<sup>a</sup>	<sup>a</sup>	<sup>a</sup>	<sup>a</sup>
2003	<sup>a</sup>	21	58	32	26	57	43	<sup>a</sup>
2004	<sup>a</sup>	50	56	46	56	75	111	<sup>a</sup>
-	-	-	-	-	-	-	-	-
2012	<sup>a</sup>	94	115	38	31	56	88	<sup>a</sup>
2013	<sup>a</sup>	Information is not yet available.						<sup>a</sup>

*Note:* No subsistence surveys were conducted from 2005 to 2011.

<sup>a</sup> Not surveyed.

Appendix C8.–Kotzebue District chum salmon aerial survey counts, 1990–2009.

Stream <sup>a</sup>	1990 <sup>b</sup>	1991 <sup>b</sup>	1992 <sup>b</sup>	1993	1994 <sup>c</sup>	1995	1996	1997	1998	1999
Noatak Drainage										
Noatak River below Kelly River	23,345 <sup>b</sup>	82,750	34,335	25,415		147,260	306,900 <sup>c</sup>	<sup>c</sup>	<sup>b</sup>	
Eli River	3,000	2,940	701	4,795		7,860	30,040 <sup>c</sup>	<sup>c</sup>	<sup>b</sup>	
Kelly River and Lake	325 <sup>d</sup>	654	726	9		8,384	1,427	2,792	2,631	
Noatak River System Total	26,670	86,344	35,762	30,219		163,504	338,367		<sup>b</sup>	84,085
Kobuk Drainage										
Kobuk to Pah River	4,610	9,840	1,030	3,896		12,190	20,700	2,248 <sup>b</sup>	<sup>b</sup>	
Pah River to just below Selby River	305	2,780	3,820	1,535		4,537	4,600	404 <sup>b</sup>	<sup>b</sup>	
Selby River mouth & slough	420	1,040	1,500	1,800		1,250	4,100	662 <sup>b</sup>	<sup>b</sup>	
Selby River	7,505	1,460	868	824		3,364	14,950	853 <sup>b</sup>	730	
Selby River mouth to Beaver Creek		5,250	3,845	929		10,898	15,480	2,582 <sup>b</sup>		
Beaver Creek mouth	2,515							914 <sup>b</sup>	<sup>b</sup>	
Above Beaver Creek		4,155	740	3,174		3,486	14,940	850 <sup>b</sup>	<sup>b</sup>	
Upper Kobuk River Total	15,355	24,525	11,803	12,158		35,725	74,770	8,513 <sup>b</sup>	<sup>b</sup>	27,340
Squirrel River	5,500	4,606	2,765	4,463		10,605	10,740	4,779 <sup>b</sup>		13,513
Salmon River	6,335	5,845	1,345	13,880		13,988	23,790	1,181 <sup>b</sup>	<sup>b</sup>	4,989
Tutuksuk River	2,275	744	1,162	1,196		3,901	21,805	163 <sup>b</sup>	<sup>b</sup>	2,906
Kobuk River System Total	29,465	35,720	17,075	31,697		64,219	131,105	14,636	<sup>b</sup>	48,748

-continued-

Appendix C8.–Page 2 of 2.

Stream <sup>a</sup>	2001	2002	2003	2004	2006	2008	2009	Goals <sup>e</sup>
Noatak Drainage								
Noatak River below Kelly River		700	34,575	49,541	36,125 <sup>b</sup>	257,695	67,265	
Eli River				2,917	1,285 <sup>b</sup>	13,052	2,607	
Kelly River and Lake		1,116	1,566	2,987	2,375 <sup>b</sup>	1,865	3,986	
Noatak River System Total			36,141	55,445	39,785 <sup>b</sup>	272,612	73,858	42,000–91,000
Kobuk Drainage								
Kobuk to Pah River	2,790		5,501	7,493	8,525 <sup>b</sup>	19,421	7,468	
Pah River to just below Selby River	1,380	857	828	1,885		5,795	10,852	
Selby River mouth & slough	1,780	2,100	1,110	3,846				
Selby River			427	3,760	500 <sup>b</sup>	1,750	208	
Selby River mouth to Beaver Creek	7,470		1,274	6,215		13,201	26,627	
Beaver Creek mouth								
Above Beaver Creek		490	2,462			3,180		
					39,725 <sup>f</sup>			
Upper Kobuk River Total	13,420	3,447	11,602	23,199	48,750 <sup>b</sup>	43,347	45,155	9,700–21,000
Squirrel River			<sup>b</sup>					4,900–10,500
Salmon River			<sup>b</sup>					3,300–7,200
Tutuksuk River			<sup>b</sup>					1,400–3,000
Kobuk River System Total	13,420	3,447	11,602	23,199	48,750 <sup>b</sup>	43,347	45,155	19,600–39,200

Note: No surveys were flown in 2000, 2005, 2007, or since 2009.

<sup>a</sup> Three aerial surveys are attempted yearly at different intervals for each tributary to assess escapements prior to the peak, at the peak, and after the peak of the run. Indices listed in this table are the largest survey observed for each tributary during the given year.

<sup>b</sup> Poor survey conditions or incomplete, early, or late survey.


<sup>c</sup> Unacceptable survey conditions.

<sup>d</sup> Surveyed well before peak of migration.

<sup>e</sup> Aerial survey goals were revised in 2007.

<sup>f</sup> Unclear where these fish were observed.





Appendix C9.—Kobuk River chum salmon drift test fish cumulative catch per unit effort (CPUE), 2007–2013.



## **APPENDIX D: HERRING FISHERIES**

Appendix D1.—Norton Sound herring and spawn-on-kelp harvests (in tons) by U.S. commercial fishermen, 1990–2013.

Year <sup>a</sup>	Sac Roe Herring	Food or bait herring	Total herring	Spawn on kelp
1990	5,253	1,026	6,279	0
1991	5,465	207	5,672	0
1992 <sup>b</sup>	0	0	0	0
1993	4,713	321	5,034	0
1994	958	2	960	0
1995	6,647	116	6,763	0
1996 <sup>c</sup>	6,061	109	6,220	0
1997 <sup>d</sup>	3,709	262	3,976	0
1998	2,623	8	2,631	9.04 <sup>e</sup>
1999	2,693 <sup>f</sup>	53	2,751	3.74
2000	4,487 <sup>g</sup>	0	4,487	2.25
2001	2,245	0	2,245	2.20
2002	1,059	64	1,123	0
2003	1,587	21	1,608	0.88
2004 <sup>b</sup>	0	11	11	0
2005	1,951	0	1,951	0
2006	646	25	671	0.57
2007 <sup>b</sup>	0	33	33	0.14
2008 <sup>b</sup>	0	91	91	0.18
2009 <sup>b</sup>	0	28	28	0
2010	623	65	688	0
2011	739	67	806	0
2012 <sup>b</sup>	0	7	7	0
2013	490	2	492	0

<sup>a</sup> From 1990 to present, the fishery has occurred in southeastern Norton Sound.

<sup>b</sup> No commercial fishery took place in 1992, and no sac roe fishery took place in 2004, 2007–2009, and 2012.

<sup>c</sup> Total includes an estimated 50 short tons (st) of wastage.

<sup>d</sup> Total includes an estimated 5 st of wastage and approximately 1,000 lb taken as bait.

<sup>e</sup> Includes 2,100 lb of wild kelp and 16,083 lb of *Macrocystis* kelp.

<sup>f</sup> Includes an estimated 5 st of wastage.

<sup>g</sup> Includes an estimated 15 st of wastage.

Appendix D2.—Commercial herring fishery summary information, Norton Sound District, 1990–2013.

Year	Estimated biomass (tons)	Catch gillnet (tons)	Beach seine (tons)	Wild kelp (tons)	<i>Macrocystis</i> kelp (lbs.)	Number of fishermen	Dollar value (millions)	Number of buyers	Average roe %	Peak catch day	Fishery duration
1990	39,384	6,032	347			365	3.60	8	8.8	5/29	5/28–05/30
1991	42,854	5,150	522			279	2.40	8	9.3	5/25	5/23–05/25
1992	57,974	0 <sup>a</sup>	0 <sup>a</sup>			<sup>a</sup>	0.00	<sup>a</sup>	<sup>a</sup>	6/20 <sup>b</sup>	<sup>a</sup>
1993	46,549	4,291	742			264	1.50	5	9.9	5/25	5/24–06/05
1994	31,088	921	40			215	0.30	6	10.3	6/8	6/05–06/09
1995	37,779	6,033	614			215	4.20	6	10.4	5/24	5/23–05/30
1996	26,596	5,581	589			287	4.50	10	10.6	5/25	5/24–05/25
1997	47,748	3,459	513			220	0.61	9	9.9	5/22	5/20–05/24
1998	52,033	2,632	0	1.00	16,083	47	0.20	2	9.2	5/25	5/22–06/09
1999	34,314	2,755	0		7,482	122	0.61	4	10.5	6/17	6/13–06/22
2000	32,680	4,390	81		4,500	97	0.89	4	9.5	6/11	6/07–06/15
2001	26,305	2,245	0		4,400	76	0.35	3	12.3	6/12	6/12–06/16
2002	27,068	1,123	0		0	46	0.16	2	10.6	5/24	5/22–06/03
2003	32,918	1,608	0		1,750	32	0.22	2	10.5	5/18	5/16–05/25
2004 <sup>a</sup>	34,180	11 <sup>c</sup>	0		0	4	0.00	0	<sup>a</sup>	5/24 <sup>b</sup>	<sup>c</sup>
2005	43,013	1,951	0		0	56	0.32	1	11.4	6/04	6/03–06/10
2006	38,833 <sup>d</sup>	671 <sup>e</sup>	0	0.57	0	41	0.14	1	10.2	6/09	6/08–06/11
2007 <sup>a</sup>	38,415 <sup>d</sup>	33	0	0.14	0	7	0.02	1	<sup>a</sup>	6/09	6/09–06/15
2008 <sup>a</sup>	37,401 <sup>d</sup>	91	0		0	14	0.18	1	<sup>a</sup>	6/11	6/10–06/24
2009 <sup>a</sup>	36,917 <sup>d</sup>	28	0		0	6	0.02	1	<sup>a</sup>	6/12	6/12–06/15
2010	42,889 <sup>d</sup>	688	0		0	30	0.19	1	13.5	6/17	6/11–06/19
2011	53,786	807	0		0	35	0.27	1	14.8	6/04	6/01–06/10
2012 <sup>a</sup>	52,949 <sup>d</sup>	7	0		0	8	0.01	1	<sup>a</sup>	6/25	6/16–06/25
2013	58,594 <sup>d</sup>	492	0		0	40	0.15	1	13.2	6/15	6/14–06/20

<sup>a</sup> No fishery in 1992 and very limited fishery in 2012 due to late sea ice breakup, and no sac roe fishery in 2004 and 2007–2009 due to lack of a buyer.

<sup>b</sup> Date of peak aerial survey biomass estimate, typically 1 or 2 days prior to peak catch. The 2004 catch was by king crab permit holders for bait.

<sup>c</sup> All fish caught were kept as bait; none were sold.

<sup>d</sup> Conditions did not allow for a peak survey; therefore, biomass was estimated by extrapolation.

<sup>e</sup> Twenty-five tons out of total sac roe herring catch was sold off as bait to NSEDC.

Appendix D3.–Norton Sound commercial herring harvest (tons) by subdistrict, by year, 1990–2013.

Year <sup>a</sup>	Subdistricts							Totals
	1	2	3	4	5	6	7	
1990	4,498	950	931	0	0	0	0	6,379 <sup>b</sup>
1991	0	880	4,792	0	0	0	0	5,672 <sup>c</sup>
1992 <sup>d</sup>	0	0	0	0	0	0	0	0
1993	2,288	587	1,881	0	278	0	0	5,034 <sup>e</sup>
1994	250	36	634	0	40	0	0	960
1995	2,359	604	1,524	0	2,108	167	0	6,762
1996	3,074	111	2,831	0	153	0	0	6,170 <sup>f</sup>
1997	2,046	62	1,864	0	0	0	1 <sup>g</sup>	3,976 <sup>h</sup>
1998	1,543	0	1,081	0	0	0	0	2,624
1999	285	323	2,050	0	0	0	8	2,746 <sup>i</sup>
2000 <sup>j</sup>	2,623	81	1,767	0	0	0	0	4,471
2001 <sup>j</sup>	898	0	1,347	0	0	0	0	2,245
2002 <sup>j</sup>	373	0	750	0	0	0	0	1,123
2003 <sup>j</sup>	283	0	1,325	0	0	0	0	1,608
2004	0	0	0	0	0	0	11	11
2005 <sup>j</sup>	783	9	1,149	0	10	0	0	1,951
2006	191	0	480	0	0	0	0	671
2007	0	33	0	0	0	0	0	33
2008	0	91	0	0	0	0	0	91
2009	0	28	0	0	0	0	0	28
2010	314	300	74	0	0	0	0	688
2011	600	84	123	0	0	0	0	807
2012	6	0	0	0	0	0	1	7
2013	107	84	302	0	0	0	0	492

<sup>a</sup> Includes herring taken for sac roe and bait.

<sup>b</sup> Does not include an estimated wastage of 60 short tons (st) in abandoned gillnets.

<sup>c</sup> Does not include an estimated wastage of 125 st in abandoned gillnets.

<sup>d</sup> No commercial fishery in 1992.

<sup>e</sup> Does not include an estimated wastage of 45 st in abandoned beach seine sets.

<sup>f</sup> Does not include an estimated 50 st of wastage.

<sup>g</sup> Approximately 1,000 lb of herring bait was taken under 5 AAC 27.971 in June (not during sac roe fishery).


<sup>h</sup> Does not include an estimated 5 st of wastage.

<sup>i</sup> There were 75.8 tons added to the sac roe total due to dewatering by buyers. Three tons were added to the bait total due to dewatering by the buyer. Does not include an estimated 5 st of wastage.

<sup>j</sup> There was 10% added to sac roe total due to dewatering by buyers.

Appendix D4.–Port Clarence District commercial herring fishery, 1986–1996.


Year	Fishery	Gillnet Permits	Purse Seine Permits	Harvest (pounds)
1986	Fall Bait	1		130
1987	Sac Roe	3	3	291,000
1987	Fall Bait	Unknown		1,100
1988	Sac Roe	3	3	160,000
1994	Fall Bait	4		8,706
1995	Spring Bait	8		19,193
1995	Fall Bait	2		9,119
1996	Spring Bait	4		5,546



Appendix D5.—Norton Sound herring age class composition by percentage of commercial catch, commercial gear combined (beach seine and gillnet), 1987–1993.


*Note:* No commercial fishing occurred in 1992.






Appendix D6.—Norton Sound herring age class composition by percentage of commercial catch, commercial gear combined (beach seine and gillnet), 1994–1999.

*Note:* No commercial catch from beach seine gear in 1998 and 1999.




Appendix D7.—Norton Sound herring age class composition by percentage of commercial catch, commercial gear combined (beach seine and gillnet), 2000–2006.

*Note:* No commercial catch from beach seine gear after 2000. No fishery in 2004.




Appendix D8.—Norton Sound herring age class composition by percentage of commercial catch, commercial gear (gillnet only), 2013.


*Note:* No commercial samples were available 2007–2012.




Appendix D9.—Norton Sound herring age class composition by percentage of total catch, variable mesh gillnets, 1987–1992.



Appendix D10.—Norton Sound herring age class composition by percentage of total catch, variable mesh gillnets, 1993–1998.




Appendix D11.—Norton Sound herring age class composition by percentage of total catch, variable mesh gillnets, 1999–2004.



Appendix D12.—Norton Sound herring age class composition by percentage of total catch, variable mesh gillnets, 2005–2012.

*Note:* Herring age class composition by percentage of total catch for 2010 and 2012 was not available.



Appendix D13.—Norton Sound Pacific herring age composition comparison of the 2013 variable-mesh gear and the projected age composition of the 2014 return.



## **APPENDIX E: KING CRAB FISHERIES**

Appendix E1.—Historical commercial summer harvest of red king crab from Norton Sound Section, Eastern Bering Sea, by statistical areas, 1990–2013 (catch in pounds).

150

Statistical Area	1990	1992	1993	1994	1995	1996 <sup>a</sup>	1997	1998	1999
616331				48					633
616401					35				
626331						61			
626401					18,971	45,045	18,066	8,065	508
626402									
636330						4,560	3,838	2,449	
636401		1,159	1,373	3,340	24,329	70,677	59,206	10,771	14,201
636402				1,754	3,466				
646301					4,628	13,888			
646330					1,493	2,894	314		3,021
646401			1,963	37,510	105,045	22,834	1,052	3,194	221
646402			730	139,661	66,821				
656300									
656330		4,814	265		19,745	15,446	4,661	4,078	1,300
656401	171	53,119	105,341	34,686	32,289	9,985	4,035	1,127	2,739
656402			193,079	110,289	44,000				
666230									
666300						25,519			
666330	27,185	4,305	31,758		730				
666401	162,263	10,632	746	396		3,001	1,816		930
666402			535	1,221					
666431					1,124				
676300						546			
676330									
676400	3,212					9,775			
676430									
676501									
686330									
686431									
Total (tons)	96	37	168	164	161	112	46	15	12

-continued-

Appendix E1.–Page 2 of 3.

151

Statistical Area	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
616331	4,557		3,506	646			2357		5,658	888
616401								231	416	6,170
626331			2,455				1415	27,018	3,235	3,047
626401	4,689	61,620	53,722	15,899	23,113	94,130	118202	61,704	96,327	103,043
626402				1,352						
636330		2,253				126	26680	10,253	2,350	5,026
636401	130,463	91,343	50,906	83,949	166,489	227,204	224531	123,092	197,948	96,279
636402										
646301										
646330		1,868	1,955		2,226	4,097	2629	5,290	1,505	933
646401		4,287		3,952	1,964	149	1660		18,728	46,264
646402										
656300				14	932		284	1,909		
656330	1,990	20,869	12,374	21,176	46,288	47,411	17752	4,911		10,617
656401	95,979	55,158	63,038	40,566	21,579	9,405	28434	70,065	68,968	107,557
656402				1,441		380	807	2,254		
666230							1721			
666300							18245			
666330	5,839	7,030	1,332	1,296	12,359	142	5041	511		1,514
666401	69,007	43,771	35,970	83,998	42,452	727	600	2,498		10,021
666402			30,070	12,873	23,344	16,025	1050	2,959		6,228
666431			4,274	45						
676300										
676330										
676400								180		
676430										
676501						1,008				
686330										
686431							340			
Total (tons)	156	144	130	134	170	200	226	156	198	199

-continued-

Appendix E1.—Page 3 of 3.

152

Statistical Area	2010	2011	2012	2013	Total
616331					18,293
616401				7,729	14,581
626331		2,489		686	40,406
626401	52,054	85,271	115,524	36,802	1,012,755
626402					1,352
636330	2,584		1,454	12,035	73,608
636401	182,040	146,973	148,183	34,027	2,088,482
636402					5,220
646301					18,516
646330	1,205		1,204	4,195	34,829
646401	77,437	83,099	98,811	59,737	567,907
646402				5,271	212,483
656300					3,139
656330	17,660	1,546	8,168	8,515	269,586
656401	82,747	77,149	85,920	147,569	1,197,625
656402				37,743	389,993
666230					1,721
666300					43,764
666330		2,042	1,000		102,084
666401			15,726	33,469	518,023
666402	1,577	2,271		1,419	99,572
666431				2,669	8,112
676300					546
676330					0
676400					13,167
676430					0
676501					1,008
686330					0
686431					340
Total					6,737,111
(tons)	209	200	238	196	3,369

Note: Not all statistical areas had recorded harvest. No commercial fishery occurred in 1991.

<sup>a</sup> Does not include approximately 2,490 lb not reported on fish tickets.

Appendix E2.—The results of the population assessment surveys conducted for red king crab in Norton Sound since 1976.

Year	Date	Research Agency	Gear	Population Abundance Estimates			Legal Male Biomass (millions of pounds)
				Number of crab <sup>a</sup>			
				Pre-2 males <sup>b</sup>	Pre-1 Males <sup>b</sup>	Legal Males <sup>c</sup>	
1976	9/02–09/05	NMFS	Trawl	331,555	808,091	1,742,755	5,228,265
	9/16–10/07						
1979 <sup>d</sup>	7/26–08/05	NMFS	Trawl			809,799	2,429,397
1980 <sup>e</sup>	7/04–07/14	ADF&G	Pots			1,900,000	5,700,000
1981	6/28–07/14	ADF&G	Pots			1,285,195	3,855,585
1982	7/06–07/20	ADF&G	Pots			353,273	1,059,819
1982	9/05–09/11	NMFS	Trawl	356,724	832,581	877,722	2,633,166
1985	7/01–07/14	ADF&G	Pots			907,579	2,722,737
1985	9/16–10/01	NMFS	Trawl	466,858	707,140	1,051,857	3,155,571
1988	8/16–08/30	NMFS	Trawl	565,255	493,030	978,748	2,936,244
1991	8/22–08/30	NMFS	Trawl	294,801	303,682	1,287,486	3,862,458
1996	9/07–09/18	ADF&G	Trawl	452,580	325,699	536,235	1,608,705
1999	7/28–08/07	ADF&G	Trawl	103,832	940,198	1,594,341	4,783,023
2002	7/27–08/06	ADF&G	Trawl	427,703	518,638	771,569	2,314,707
2006	7/25–08/08	ADF&G	Trawl	775,076	569,833	726,251	2,178,753
2008	7/24–08/11	ADF&G	Trawl	795,777	697,442	811,727	2,435,182
2011	7/18–08/15	ADF&G	Trawl	431,153	311,550	1,310,634	3,931,902

*Note:* Data not available for all years.

<sup>a</sup> Population estimates are valid for the date of the survey (i.e., either before or after the summer commercial fishery).

<sup>b</sup> Pre-2 male crab were defined as 76–89 mm in carapace length (CL), and pre-1 male crab were defined as 90–104 mm in CL.

<sup>c</sup> Legal male red king crab were defined as  $\geq 121$  mm (4.75 in) in carapace width for the pot surveys and all ADF&G trawl surveys (except for 1996, when legal male crab were defined as at least 105 mm CL), and  $\geq 104$  mm CL for all of the NMFS trawl surveys (except the 1979 survey, which defined legal male crab as at least 100 mm CL).

<sup>d</sup> Pre-2 male and pre-1 male crab data are unavailable for the 1979 NMFS trawl survey.

<sup>e</sup> The 1980 pot survey estimate has been revised from the original estimate of 13.4 million pounds, which was thought inaccurate due to under-reporting of recovered tagged crab.

Appendix E3.—Historical summer commercial red king crab fishery economic performance, Norton Sound Section, Eastern Bering Sea, 1990–2013.

Year	Guideline	Legal male		Commercial							Total	Total	Season length	
	harvest	population est.		harvest (lbs) <sup>a,b</sup>							exvessel	fishery value		
	level	No. crab		Open		Total number of			Total number of pots		price/lb	(millions \$)	Days	Dates
	(lbs) <sup>b</sup>	(millions)	lbs <sup>b</sup>	access	CDQ	Vessels	Permits	Landings	Registered	Pulls				
1990	0.20			0.19		4	4	<sup>c</sup>	1,388	3,172	<sup>c</sup>	<sup>c</sup>	4	8/01–8/05
1991	0.34	1.3	3.9			No Summer Fishery								
1992	0.34			0.07		27	27	<sup>c</sup>	2,635	5,746	1.75	0.130	2	8/01–8/03
1993	0.34			0.33		14	20	208	560	7,063	1.28	0.430	52	7/01–8/28 <sup>d</sup>
1994	0.34			0.32		34	52	407	1,360	11,729	2.02	0.646	31	7/01–7/31
1995	0.34			0.32		48	81	665	1,900	18,782	2.87	0.926	67	7/01–9/05
1996	0.34	0.5	1.5	0.22		41	50	264	1,640	10,453	2.29	0.519	57	7/01–9/03 <sup>e</sup>
1997	0.08			0.09		13	15	100	520	2,982	1.98	0.184	44	7/01–8/13 <sup>f</sup>
1998	0.08			0.03	0.00	8	11	50	360	1,639	1.47	0.041	65	7/01–9/03 <sup>g</sup>
1999	0.08	1.6	4.8	0.02	0.00	10	9	53	360	1,630	3.08	0.073	66	7/01–9/04 <sup>h</sup>
2000	0.33	1.4	4.2	0.29	0.01	15	22	201	560	6,345	2.32	0.715	91	7/01– 9/29 <sup>i</sup>
2001	0.30	1.3	3.8	0.28	0.00	30	37	319	1,200	11,918	2.34	0.674	97	7/01– 9/09 <sup>j</sup>
2002	0.24	1.0	3.1	0.24	0.01	32	49	201	1,120	6,491	2.81	0.729	77	6/15–9/03 <sup>k</sup>
2003	0.25	1.0	3.1	0.25	0.01	25	43	236	960	8,494	3.09	0.823	68	6/15–8/24 <sup>l</sup>
2004	0.35	1.6	4.4	0.31	0.03	26	39	227	1,120	8,066	3.12	1.063	51	6/15–8/08 <sup>m</sup>
2005	0.37	1.7	4.8	0.37	0.03	31	42	255	1,320	8,867	3.14	1.264	73	6/15–8/27 <sup>n</sup>
2006	0.45	1.6	4.5	0.42	0.03	28	40	249	1,120	8,867	2.26	1.021	68	6/15–8/22 <sup>m</sup>
2007	0.32	1.1	3.1	0.29	0.02	38	30	251	1,200	9,118	2.49	0.750	52	6/15–8/17 <sup>m</sup>
2008	0.41	1.5	4.1	0.36	0.03	23	30	248	920	8,721	3.20	1.231	73	6/23–9/03 <sup>o</sup>
2009	0.38	1.3	3.8	0.37	0.03	22	27	359	920	11,934	3.17	1.225	98	6/15–9/20 <sup>p</sup>
2010	0.40	1.7	4.5	0.39	0.03	23	32	286	1,040	9,698	3.73	1.528	58	6/28–8/24 <sup>q</sup>

-continued-

Appendix E3.–Page 2 of 2.

Year	Guideline	Legal Male		Commercial							Total	Total		
	Harvest	Population Est.		Harvest (lbs) <sup>a,b</sup>							Exvessel	Fishery Value	Season Length	
	Level	No. crab		Open		Total Number of			Total Number of Pots		Price/lb	(millions \$)	Days	Dates
	(lbs) <sup>b</sup>	(millions)	lbs <sup>b</sup>	Access	CDQ	Vessels	Permits	Landings	Registered	Pulls				
2011	0.36	1.5	4.0	0.37	0.03	24	25	173	1,040	6,808	5.23	2.016	33	6/28–7/30 <sup>r</sup>
2012	0.47	1.4	3.7	0.44	0.03	40	29	312	1,200	10,041	5.41	2.556	72	6/29–9/08 <sup>s</sup>
2013	0.50	1.6	4.1	0.37	0.02	37	33	460	1,420	15,058	5.63	2.165	74	7/03–9/14 <sup>t</sup>

<sup>a</sup> Deadloss included in total. Data not available for all years.

<sup>b</sup> Millions of pounds.

<sup>c</sup> Information not available.

<sup>d</sup> Fishing actually began 7/8.

<sup>e</sup> Fishing began 7/9 due to fishermen strike.

<sup>f</sup> First delivery was made 7/10.

<sup>g</sup> First delivery was made 7/16.

<sup>h</sup> The season was extended 24 hours due to bad weather.

<sup>i</sup> Open access fishery (OA) closed 8/29. CDQ fishery opened 9/1–9/29.

<sup>j</sup> OA closed 9/1. CDQ fishery opened 9/1–9/9.

<sup>k</sup> OA was 7/1–8/6. CDQ fishery opened 6/15–6/28 and 8/9–9/3.

<sup>l</sup> OA was 7/1–8/13. CDQ fishery opened 6/15–6/28 and 8/15–8/24.

<sup>m</sup> CDQ fishery opened 6/15–6/28. OA opened 7/1 to the end date.

<sup>o</sup> OA opened 6/23–8/18. CDQ opened 8/17–9/3.

<sup>n</sup> OA was 7/1–8/15. CDQ fishery opened 6/15–6/28 and 8/17–8/27.

<sup>p</sup> CDQ opened 6/15 – 7/28. OA opened 6/15 to the end date.

<sup>q</sup> CDQ opened 6/28 – 7/16. OA opened 7/1 to the end date.

<sup>r</sup> CDQ opened 6/28 – 7/8. OA opened 6/28 to the end date.

<sup>s</sup> CDQ opened 6/29 to the end date. OA opened 6/29–8/11.

<sup>t</sup> CDQ and OA opened and closed at the same time.

Appendix E4.—Average length and percentage of recruit and postrecruit male red king crab from summer commercial fishery catch samples in Norton Sound Section, Eastern Bering Sea, 1990–2013.

Year	Average Length (mm)	Recruits <sup>a</sup>	Postrecruits <sup>b</sup>
1990	121	21	79
1991 <sup>c</sup>			
1992	120	28	72
1993	119	31	69
1994	119	20	80
1995	118	36	64
1996	117	30	70
1997	116	49	51
1998	117	32	68
1999	118	42	58
2000	116	41	60
2001	119	33	67
2002	120	33	67
2003	117	48	52
2004	117	49	51
2005	118	36	64
2006	119	25	75
2007	117	45	55
2008	115	45	55
2009	116	43	57
2010	115	49	51
2011	116	43	57
2012	118	33	67
2013	120	32	68

<sup>a</sup> Recruits = all new-shell, legal size, male king crab of carapace length < 116mm.

<sup>b</sup> Postrecruits = all other male king crab of legal size.

<sup>c</sup> No summer commercial fishery.



Appendix E5.–Winter commercial and subsistence red king crab harvests, Norton Sound, Eastern Bering Sea, 1990–2013.

Commercial				Subsistence						
Year <sup>a</sup>	Number of fishermen	Number of crab harvested	Pounds of crab harvested	Winter <sup>b</sup>	Number of permits issued	Number of permits returned	Number of permits fished	Total crab caught <sup>c</sup>	Total crab harvested <sup>d</sup>	Average number kept/ permits fished
1990	13	3,626	9,792	1989–90	136	118	107	16,635	12,152	114
1991	11	3,800	10,064	1990–91	119	104	79	9,295	7,366	93
1992	13	7,478	21,177	1991–92	158	105	105	15,051	11,736	112
1993	8	1,788	4,926	1992–93	88	79	37	1,193	1,097	30
1994	25	5,753	17,214	1993–94	118	95	71	4,894	4,113	58
1995	42	7,538	18,845	1994–95	166	131	97	7,777	5,426	56
1996	9	1,999	5,064	1995–96	84	44	35	2,936	1,679	48
1997	2	<sup>e</sup>	<sup>e</sup>	1996–97	38	22	13	1,617	745	57
1998	5	984	2,349	1997–98	94	73	64	20,327	8,622	135
1999	5	2,714	7,041	1998–99	95	80	71	10,651	7,533	106
2000	10	3,045	7,894	1999–00	98	64	52	9,816	5,723	107
2001	3	1,098	2,943	2000–01	50	27	12	366	256	21
2002	11	2,591	6,860	2001–02	114	101	67	8,805	3,669	55
2003	13	6,853	16,827	2002–03	107	73	64	9,052	4,140	65
2004 <sup>f</sup>	2	522	1,293	2003–04	96	77	41	1,775	1,181	29
2005	4	2,121	5,619	2004–05 <sup>g</sup>	170	102	60	6,496	3,973	66
2006	1	<sup>e</sup>	<sup>e</sup>	2005–06	98	97	67	2,083	1,239	18
2007	8	3,313	8,023	2006–07	129	127	116	21,444	10,690	92
2008	9	5,796	14,676	2007–08	139	137	108	18,621	9,485	88
2009	7	4,951	12,348	2008–09	105	105	70	6,971	4,752	68
2010	10	4,834	12,028	2009–10	125	123	85	9,004	7,044	83
2011	9	3,365	8,669	2010–11	148	148	95	9,183	6,640	70
2012	35	9,157	24,142	2011–12	204	204	138	11,341	7,371	53
2013	26	22,639	62,179	2012–13	149	148	104	21,752	7,662	74
Avg 1990–2012	11	3,620	9,486	Avg 1990–2012	116	97	72	8,928	5,506	71

<sup>a</sup> Fishing may occur from November 15 to May 15.

<sup>b</sup> The winter subsistence fishery is open December through May.

<sup>c</sup> The number of crab actually caught; some may have been released.

<sup>d</sup> The number of crab harvested is the number of crab retained.

<sup>e</sup> Confidential under AS 16.05.815.

<sup>f</sup> Confidentiality was waived by the fishermen.

<sup>g</sup> Permits were only given out of the Nome ADF&G office, except during the 2004–2005 season, when permits were also given out in Elim, Golovin, Shaktoolik, and White Mountain.

Appendix E6.—Summer subsistence red king crab harvests, Norton Sound, Eastern Bering Sea, 2004–2013.

Year	Number Permits Issued	Number Permits Returned	Number Permits Fished	Total Crab Caught	Total Crab Harvested	Average Number Kept/ Permits Fished
2004	38	18	5	996	350	70
2005	14	12	4	753	304	76
2006	6	4	3	67	62	21
2007	19	19	5	1,425	1,008	202
2008	30	30	14	1,816	1,176	84
2009	20	20	13	1,874	653	50
2010	27	27	15	1,086	660	44
2011	43	42	27	4,026	2,658	98
2012	45	44	13	1,346	912	70
2013	47	46	26	3,102	1,865	72
Avg. 2008–2012	33	33	16	2,030	1,212	69

Appendix E7.—Number of crab pots lost during the subsistence and commercial winter crab fisheries and ADF&G winter studies, 2006–2013.

Year	Subsistence	Commercial	ADF&G Winter Study	Total
2005–06	50	ND	6	56
2006–07	132	ND	7	139
2007–08	6	ND	4	10
2008–09	8	ND	2	10
2009–10	23	30	2	55
2010–11	8	3	0	11
2011–12	19	64	0	83
2012–13	4	23	No winter study in 2013.	27

Appendix E8.—Size composition by percent of red king crab from winter research pots near Nome, Norton Sound, Eastern Bering Sea, 1990–2012.


Year	Undersized <sup>a</sup>			Legal <sup>a</sup>		
	Prerecruit 2	Prerecruit 1	Total	Recruits	Post recruits	Total
1990	16	33	49	25	26	51
1991	5	30	36	34	31	65
1992	<sup>b</sup>	<sup>b</sup>	<sup>b</sup>	<sup>b</sup>	<sup>b</sup>	<sup>b</sup>
1993	3	9	12	17	71	88
1994	<sup>b</sup>	<sup>b</sup>	<sup>b</sup>	<sup>b</sup>	<sup>b</sup>	<sup>b</sup>
1995	10	11	23 <sup>c</sup>	32	45	77
1996	22	33	64 <sup>c</sup>	10	26	36
1997	32	21	64 <sup>c</sup>	14	22	36
1998	36	44	82 <sup>c</sup>	9	9	18
1999	7	42	50 <sup>c</sup>	39	11	50
2000	16	20	37 <sup>c</sup>	39	25	64
2001	23	16	39 <sup>c</sup>	14	48	61
2002	43	26	79 <sup>c</sup>	9	12	21
2003	20	42	66 <sup>c</sup>	20	14	34
2004	9	40	50 <sup>c</sup>	37	13	50
2005	16	24	41 <sup>c</sup>	25	34	59
2006	29	33	63 <sup>c</sup>	16	22	38
2007	16	53	78 <sup>c</sup>	11	11	22
2008	36	31	71 <sup>c</sup>	18	12	30
2009	11	42	54 <sup>c</sup>	24	22	46
2010	10	32	43 <sup>c</sup>	30	27	57
2011	15	26	44 <sup>c</sup>	23	33	56
2012	25	29	57 <sup>c</sup>	14	29	43

*Note:* No winter study occurred in 2013.

<sup>a</sup> Undersized crab are male crab less than 4-¾ inch carapace width (CW). Legal crab are male king crab greater than or equal to 4-¾ inch CW.


<sup>b</sup> No winter crab research study occurred in 1992 or 1994.

<sup>c</sup> Includes prerecruit 3.




Appendix E9.—Current and historical catch performance for the Norton Sound summer commercial crab fishery, 2008–2013.

*Note:* CDQ catch is not included in years prior to 2009 because the open access and CDQ portions of the crab fishery did not occur concurrently in those years.




Appendix E10.—Norton Sound crab exvessel value and fishery price per pound, 1994–2013.




Appendix E11.—Closed water regulations in effect for the Norton Sound summer commercial crab fishery, with dotted line showing temporary boundaries for 2013 fishery only.

*Note:* Line drawn around the coastline delineates the 3-mile state waters zone.




Appendix E12.—The percent of crab harvested during the Norton Sound summer commercial red king crab fishery east of 164° W longitude, 1990–2013.






Appendix E13.—Norton Sound male red king crab size distribution from trawl assessment surveys conducted by the National Marine Fisheries Service in 1991, and by ADF&G in 1996, 1999, and 2002.


*Note:* Mean length information is not available for 1991.




Appendix E14.—Norton Sound male red king crab size distribution from trawl assessment surveys conducted by ADF&G in 2006, 2008, and 2011.




Appendix E15.—Length composition of Norton Sound red king crab summer commercial harvests, 1990–1994.  
*Note:* No fishery in 1991.




Appendix E16.—Length composition of Norton Sound red king crab summer commercial harvests, 1995–1998.




Appendix E17.—Length composition of Norton Sound red king crab summer commercial harvests, 1999–2002.



Appendix E18.—Length composition of Norton Sound red king crab summer commercial harvests, 2003–2006.



Appendix E19.—Length composition of Norton Sound red king crab summer commercial harvests, 2007–2010.



Appendix E20.—Length composition of Norton Sound red king crab summer commercial harvests, 2011–2013.



## **APPENDIX F: MISCELLANEOUS FISHERIES**

Appendix F1.–Kotzebue District winter commercial sheefish harvest statistics, 1990–2013.

Year <sup>b</sup>	Number of Fishermen	Number of Fish	Pounds <sup>a</sup>		Price per Pound (\$)	Estimated Value (\$)
			Total	Average		
1990 <sup>c</sup>	6	687	5,617	8.2		
1991	5	852	8,224	9.7	0.50	4,112
1992	3	289	2,850	9.9	0.65	1,853
1993	1	210 <sup>d</sup>	1,700	8.1	0.50	850
1994 <sup>e</sup>						
1995	1	226	2,240	9.9	0.50	1,120
1996	2	308	3,002	9.7	0.44	1,321
1997 <sup>e</sup>						
1998	1	254	2,400	9.4	0.43	1,032
1999 <sup>e</sup>						
2000 <sup>e</sup>						
2001	1	19	200	10.5	1.00	200
2002	4	30	300	10.0	1.00	300
2003	1	122	1,250	10.2	0.56	700
2004	1	37	474	12.8	1.91	905
2005 <sup>f</sup>			All Information Confidential			
2006–2011 <sup>e</sup>						
2012 <sup>f</sup>			All Information Confidential			
2013 <sup>e</sup>						

<sup>a</sup> Data are not exact; in some instances total catch poundage was determined from average weight and catch data. Similarly, various price-per-pound figures were determined from price-per-fish and average weight data.

<sup>b</sup> Season was from October 1 to September 30. Year indicated would be the year the commercial season ended. For example, the year 1980 would represent October 1, 1979, to September 30, 1980.

<sup>c</sup> Data unavailable or incomplete.

<sup>d</sup> Number of fish is not always reported. Estimates were based on average weight from reported sales that documented the number of fish.

<sup>e</sup> No reported commercial catches.

<sup>f</sup> Less than 4 deliveries; data confidential under Alaska Statute 16.05.815. Prior to 2005, confidentiality was waived by permit holders.

Appendix F2.–Kotzebue District reported subsistence harvests of sheefish, 1991–2004 and 2012–2013.

Year <sup>a</sup>	Number of fishermen interviewed	Reported harvest	Average catch per fisherman
1991	40	2,180	55
1992	43	2,821	66
1993	46	2,441	53
1994	171	3,181	19
1995 <sup>b</sup>	314	9,465	30
1996 <sup>b</sup>	389	6,953	18
1997 <sup>b</sup>	338	9,805	25
1998 <sup>b</sup>	435	5,350	14
1999 <sup>b</sup>	191	8,256	19
2000 <sup>b</sup>	237	7,446	17
2001 <sup>b</sup>	363	3,838	9
2002	101	3,882	38
2003	488	7,823 <sup>c</sup>	0
2004 <sup>d</sup>	440	10,163	23
2012 <sup>d</sup>	360	11,693	32
2013	Information is not yet available		

*Note:* Subsistence surveys were not conducted from 2005 to 2011.

<sup>a</sup> Due to limited survey effort during many years, total catch and effort should be regarded as minimum numbers only and are not comparable year to year.

<sup>b</sup> Subsistence sheefish harvests are from villages on Kobuk River.

<sup>c</sup> Includes 10 fish reported from commercial salmon fishery and used for subsistence.

<sup>d</sup> Subsistence surveys were not conducted in the town of Kotzebue.

Appendix F3.—Non-salmon sport fish harvests in Norton Sound and Kotzebue/Chukchi Sea, 1990–2013.

Year	Norton Sound		Kotzebue / Chukchi Sea		
	Dolly Varden	Arctic Grayling	Dolly Varden	Arctic Grayling	Inconnu/ Sheefish
1990	3,765	1,378	806	622	151
1991	10,365	5,121	1,149	1,981	603
1992	2,382	492	582	968	1,904
1993	5,907	1,584	914	916	1,029
1994	3,071	1,331	2,365	814	564
1995	2,908	1,037	939	910	1,142
1996	4,285	1,485	913	2,136	485
1997	4,467	1,262	598	1,903	906
1998	2,240	298	440	1,788	414
1999	6,708	1,600	796	1,247	635
2000	7,952	1,203	1,599	1,233	1,201
2001	3,174	994	1,693	1,244	1,305
2002	2,252	1,565	1,884	1,994	500
2003	5,531	1,778	533	1,473	2,509
2004	4,318	824	1,285	1,983	1,634
2005	2,617	595	239	269	393
2006	3,180	419	2,328	760	810
2007	2,808	314	2,924	836	1,066
2008	3,319	965	852	293	61
2009	3,373	1,185	1,406	445	957
2010	1,835	232	493	366	595
2011	4,041	1,398	865	486	385
2012	252	520	781	626	104
2013	1,184	500	1,074	563	218
Average					
2008–2012	2,564	860	879	443	420
2003–2012	3,127	823	1,171	754	851

Appendix F4.–Kotzebue District incidentally caught and sold Dolly Varden during the commercial salmon fishery, 1990–2013.

Year	Number of fish sold	Estimated Total catch <sup>a</sup>	Pounds sold	Average weight <sup>b</sup>	Average price
1990	604	<sup>c</sup>	4,219	7.0	0.25
1991	6,136	<sup>c</sup>	40,747	6.6	0.18
1992	1,977	<sup>c</sup>	11,951	6.0	0.10
1993	76	<sup>c</sup>	540	7.1	0.10
1994	149	<sup>c</sup>	767	5.1	0.17
1995	2,090	<sup>c</sup>	13,195	6.3	0.20
1996	188	<sup>c</sup>	1,153	6.1	0.25
1997	3,320	<sup>c</sup>	23,203	7.0	0.20
1998	349	<sup>c</sup>	2,640	7.6	0.20
1999	1,502	<sup>c</sup>	11,352	7.6	0.20
2000	7	<sup>c</sup>	44	6.3	0.20
2001	0	<sup>c</sup>	0	<sup>d</sup>	0.00
2002	0	30	0	<sup>d</sup>	0.00
2003	20	176	160	8.0	0.50
2004	124	<sup>c</sup>	846	6.8	0.26
2005	181	<sup>c</sup>	1,158	6.4	0.30
2006	0	278	0	<sup>d</sup>	0.00
2007	0	960	0	<sup>d</sup>	0.00
2008	0	1,629	0	<sup>d</sup>	0.00
2009	0	960	0	<sup>d</sup>	0.00
2010	0	1,323	0	<sup>d</sup>	0.00
2011	0	400	0	<sup>d</sup>	0.00
2012	0	300	0	<sup>d</sup>	0.00
2013	0	302	0	<sup>d</sup>	0.00

<sup>a</sup> Estimate includes fish caught but not sold based on interviews of fishermen or fish tickets.

<sup>b</sup> Some data extrapolated from average reported weight.

<sup>c</sup> No estimates were made of Dolly Varden caught but not sold.

<sup>d</sup> Dolly Varden caught but not sold were not weighed.

Appendix F5.—Subsistence harvests of Dolly Varden from the villages of Kivalina and Noatak, 1991–2013.

Year <sup>a</sup>	Kivalina		Noatak <sup>b,c</sup>
	Number	Pounds	Number
1991			4,814
1992			4,395
1993			4,275
1995			5,762
1996			5,031
1997			4,763
1998			3,872
2000			3,315
2001			2,702
2002			3,242
2003			6,386
2004			11,697
2007	20,527	67,739	10,234
2012			6,437
2013	Information is not yet available		

*Note:* Data are not available for all years.

<sup>a</sup> Subsistence surveys were not conducted in 1994, 1999, 2005–2006, and 2008–2011. The Division of Subsistence did a comprehensive survey of Noatak fish harvests in 2012 and of both villages in 2013, but data are not yet available for 2013.

<sup>b</sup> No data are available on poundage.

<sup>c</sup> Based on ADF&G, Division of Subsistence, household surveys in Noatak.

Appendix F6.—Dolly Varden sport fish harvests in Norton Sound, by river, 1990–2013.

Year	Location									Total
	Marine Water	Nome	Pilgrim	Unalakleet	Fish- Niukluk	Sinuk	Snake	Solomon	Other Streams	
1990	183	1,078	166	614	348				1,227	3,616
1991	0	1,220	856	1,474	1,474	729	1,252	2,219	1,141	10,365
1992	204	557	131	746	270	139	115	131	89	2,382
1993	205	917	448	427	1,003	536	331	893	1,147	5,907
1994	90	431	63	410	699	305	117	197	759	3,071
1995	0	462	74	976	346	158	131	366	395	2,908
1996	12	873	388	1,506	402	485	97	49	473	4,285
1997	189	328	65	936	2,071	346	81	186	265	4,467
1998	0	302	14	588	160	311	0	383	482	2,240
1999	330	791	45	2,384	1,952	88	44	154	920	6,708
2000	1,069	340	0	4,462	1,687	59	199	0	136	7,952
2001	166	43	270	1,002	1,197	86	108	162	140	3,174
2002	67	511	72	789	259	47	18	18	471	2,252
2003	0	1,223	482	134	110	712	13	0	2,857	5,531
2004	72	226	0	3,593	120	42	0	53	212	4,318
2005	95	553	12	500	1,148	141	27	0	141	2,617
2006	0	959	0	1,307	0	531	51	153	179	3,180
2007	14	625	0	731	193	144	461	481	159	2,808
2008	0	46	0	1,062	1,061	107	46	0	997	3,319
2009	0	253	0	2,794	108	50	50	0	118	3,373
2010	0	165	0	1,411	12	117	0	24	106	1,835
2011	0	0	11	2,219	1,631	0	10	0	170	4,041
2012	0	111	0	88	0	9	33	0	11	252
2013	0	17	0	483	0	0	0	0	684	1,184
Average										
2008–2012	0	115	2	1,515	562	57	28	5	280	2,564
2003–2012	18	416	51	1,384	438	185	69	71	495	3,127

*Note:* Data are not available for all years.

Appendix F7.—Aerial survey counts of overwintering and spawning Dolly Varden in the Kotzebue District, 1990–2013.

Year <sup>a</sup>	Noatak River spawner survey <sup>b</sup>	Overwintering	
		Wulik River <sup>c</sup>	Kivalina River <sup>c</sup>
1990	7,261	d	d
1991	9,605	126,985	35,275
1992	d	135,135	e
1993	9,560	144,138	16,534
1994	d	66,752	d
1995	6,500	128,705	28,870
1996	12,184	61,005	d
1997	d	95,412	d
1998	d	104,043	d
1999	9,059 <sup>f</sup>	70,704	d
2000	d	d	d
2001	d	92,614	d
2002	d	44,257	d
2003	d	1,500 <sup>g</sup>	d
2004	d	101,806	d
2005	d	120,848	d
2006	d	108,352	d
2007	d	99,311	d
2008	d	71,493	d
2009	d	63,977	d
2010	d	36,866	d
2011	d	64,499	d
2012	d	21,084	d
2013	d	23,312 <sup>h</sup>	d

<sup>a</sup> Counts are considered minimal because data listed include both poor and good surveys.

<sup>b</sup> Includes spawner counts on the Kelly, Kugurorok, and Nimiuktuk rivers, and tributaries of the Noatak River.

<sup>c</sup> Surveys conducted by Division of Sport Fish.

<sup>d</sup> Not surveyed.

<sup>e</sup> Poor weather hampered or prevented survey.

<sup>f</sup> Poor conditions on the Nimiuktuk did not allow a count.

<sup>g</sup> Spawning survey conducted very early (8/20/03).

<sup>h</sup> Counting conditions were poor due to presence of river ice.



Appendix F8.—Subsistence whitefish catch and effort in the Kotzebue District, 1991–1993, 1997–2004, and 2012–2013.

Year <sup>a</sup>	Number of fishermen interviewed	Number of whitefish harvested	Average catch per fisherman
1991 <sup>b</sup>	63	16,015	254
1992 <sup>b</sup>	66	17,485	265
1993 <sup>b</sup>	70	19,060	272
1997	413 <sup>c</sup>	84,851	205
1998	435 <sup>c</sup>	39,754	91
1999	191 <sup>c</sup>	56,326	295
2000	237 <sup>c</sup>	70,097	296
2001	363 <sup>c</sup>	30,976	85
2002	101 <sup>d</sup>	25,607	254
2003	446	73,242	164
2004	440 <sup>c</sup>	50,501	115
2012	360 <sup>c</sup>	41,229	115
2013	Information is not yet available		

*Note:* Subsistence surveys were not conducted from 1994 to 1996 and from 2005 to 2011.

<sup>a</sup> Whitefish harvest information was collected during chum salmon subsistence surveys and is considered a fraction of the annual catch. Whitefish numbers include all species of whitefish, except sheefish.

<sup>b</sup> Subsistence interviews from Noatak, Noorvik, and Shungnak villages only.

<sup>c</sup> Subsistence harvest information is from Ambler, Kiana, Kobuk, Noatak, Noorvik, and Shungnak.

<sup>d</sup> Subsistence harvest information is from Noatak and Noorvik.

Appendix F9.–Norton Sound District winter commercial whitefish harvest statistics, 2007–2013.

Year <sup>a</sup>	Number of fishermen	Total pounds	Price per pound (\$)	Estimated value (\$)
2006–2007	1	3,723	0.44	2,635
2007–2008 <sup>b</sup>				
2008–2009 <sup>b</sup>				
2009–2010 <sup>b</sup>				
2010–2011	1	2,009	0.50	1,005
2011–2012	1	2,148	0.50	859
2012–2013	2	105	0.50	53

<sup>a</sup> Season was from September 15 to June 15. Confidentiality was waived by fishermen.

<sup>b</sup> No reported sales.

Appendix F10.–Norton Sound District winter commercial saffron cod harvest statistics, 1994–1995 and 2010–2013.

Year <sup>a</sup>	Number of fishermen	Total pounds	Price per pound (\$)	Estimated value (\$)
1993–1994	<sup>b</sup>	1,402	<sup>b</sup>	<sup>b</sup>
1994–1995	<sup>b</sup>	52	0.50	26
2009–2010 <sup>c</sup>	1	1,748	0.30	524
2010–2011	5	8,031	0.50	4,016
2011–2012	9	3,780	0.47	1,772
2012–2013	25	33,939	0.50	16,970

<sup>a</sup> Season was from September 15 to June 15.

<sup>b</sup> Information is not available.

<sup>c</sup> Confidentiality was waived by the fisherman.

## **APPENDIX G: OVERVIEW OF 2013**

Appendix G1.—List of common and scientific names of finfish species of the Norton Sound, Port Clarence, Kotzebue, and Arctic Districts.

Common Name	Scientific Name
Arctic lamprey	<i>Lampetra camtschatica</i>
Arctic char	<i>Salvelinus alpinus</i>
Arctic cod	<i>Boreogadus saida</i>
Arctic flounder	<i>Liopsetta glacialis</i>
Arctic grayling	<i>Thymallus arcticus</i>
Alaska plaice	<i>Pleuronectes quadrituberculatus</i>
Burbot	<i>Lota lota</i>
Bering cisco	<i>Coregonus laurettae</i>
Bering poacher	<i>Ocella dodecaedria</i>
Bering wolfish	<i>Anarjicas orientalis</i>
Blackfish	<i>Dallia pectoralis</i>
Boreal smelt (rainbow-toothed)	<i>Osmerus mordax</i>
Broad whitefish	<i>Coregonus nasus</i>
Capelin	<i>Mallotus villosus</i>
Dolly Varden	<i>Salvinus malma</i>
Pond smelt	<i>Hypomesus olidus</i>
Humpback whitefish	<i>Coregonus pidschian</i>
Inconnu (sheefish)	<i>Stenodus leucichthys</i>
Lake trout	<i>Salvelinus namaycush</i>
Least cisco	<i>Coregonus sardinella</i>
Longhead dab	<i>Liranda proboscidea</i>
Ringtail snailfish	<i>Liparis rutteri</i>
Northern Pike	<i>Esox lucius</i>
Longnose sucker	<i>Casostomus catostomus</i>
Pricklebacks	<i>Stichaeidae</i>
Pacific herring	<i>Clupea harengus pallasii</i>
Rock flounder	<i>Lepidosetta bilineata</i>
Rock greenling (terpug)	<i>Hexagrammus lagocephalus</i>
Round whitefish	<i>Prosopium cylindraceum</i>
Sculpins	<i>Cottidae</i>
Pink salmon	<i>Oncorhynchus gorbuscha</i>
Chum salmon	<i>Oncorhynchus keta</i>
Coho salmon	<i>Oncorhynchus kisutch</i>
Sockeye salmon	<i>Oncorhynchus nerka</i>
Chinook salmon	<i>Oncorhynchus tshawytscha</i>
Saffron cod	<i>Eleginus gracilis</i>
Starry flounder	<i>Platichthys stellatus</i>
Sandlance	<i>Amrodytes hexapterus</i>
Sturgeon poacher	<i>Angonus acipenserinus</i>
Threespine stickleback	<i>Gasterosteus aculeatus</i>
Ninespine stickleback	<i>Pungitius pungitius</i>
Tubenose poacher	<i>Pallasina barbata aix</i>
Whitespotted greenling	<i>Hexagrammus stelleri</i>
Yellowfin sole	<i>Limanda aspera</i>

Appendix G2.—Alaska Department of Fish and Game and associated cooperative studies conducted within the Norton Sound, Port Clarence, Kotzebue and Arctic Districts, 2013.

---

HERRING

Herring Test Fishing

- a) Location: Norton Sound ocean waters, with field camp at Cape Denbigh, and base camp in Unalakleet.
- b) Description: To determine age class composition through test fishing with variable mesh-gillnets and collection of commercial catch samples. Alaska Department of Fish and Game (ADF&G) project.

SALMON

Eldorado River Weir

- a) Location: Eldorado River, approximately 15 miles upstream from the Safety Sound highway bridge, and approximately 3 miles above the furthest upstream connecting channel to the Flambeau River.
- b) Description: Determine daily and seasonal timing and magnitude of chum and pink salmon escapements. Collect age, sex, and length data from chum salmon from weir trap. Cooperative project operated by NSEDC with assistance from ADF&G.

Glacial Lake Weir and Video Enumeration Project

- a) Location: At outlet of Glacial Lake.
- b) Description: Determine daily and seasonal timing and magnitude of sockeye salmon escapement. Compare aerial survey totals with weir counts in order to improve survey accuracy. Weir is cooperative project operated by ADF&G with assistance from NSEDC. Video project is operated by ADF&G.

Inglutalik River Tower

- a) Location: Inglutalik River, approximately 18 miles upstream from the mouth at Norton Bay.
- b) Description: Determine daily and seasonal timing and magnitude of Chinook, chum, pink, and coho salmon escapements. Collect age, sex, and length data from Chinook, chum, and coho salmon from beach seine. Cooperative project operated by NSEDC with assistance from ADF&G.

Kwiniuk River Tower

- a) Location: Kwiniuk River, approximately five miles upstream from mouth.
- b) Description: Determine daily and seasonal timing and magnitude of salmon escapements. Determine age, sex, and length of Chinook and chum salmon in the Kwiniuk River escapement from beach seining. ADF&G project with additional funding from NSEDC.

Nome River Weir

- a) Location: Nome River, approximately one mile upstream of the VOR site.
- b) Description: To determine daily and seasonal timing and magnitude of salmon escapement. Compare aerial survey totals with weir counts in order to improve survey accuracy. Collect age and sex data through escapement sampling of weir trap. ADF&G project with additional funding from NSEDC.

North River Tower

- a) Location: North River, approximately two miles below bridge.
- b) Description: Determine daily and seasonal timing and magnitude of salmon escapements. Cooperative project operated by NSEDC with assistance from ADF&G.

---

-continued-

**Pilgrim River Weir**

- a) Location: Pilgrim River, approximately six miles downstream of Pilgrim River bridge at mile 65 of the Kougarok Road / Nome–Taylor Highway.
- b) Description: Determine daily and seasonal timing and magnitude of the salmon escapements. Collect age, sex, and length data from weir trap. Cooperative project operated by NSEDC with assistance from ADF&G.

**Snake River Weir**

- a) Location: Snake River, approximately five miles upstream of boat harbor, where river turns north.
- b) Description: Determine daily and seasonal timing and magnitude of salmon escapements. Cooperative project operated by ADF&G and NSEDC.

**Solomon River Weir**

- a) Location: Solomon River, at approximately mile 35.5 on the Nome–Council road.
- b) Description: Determine daily and seasonal timing and magnitude of salmon escapements. ADF&G project.

**Unalakleet River Weir**

- a) Location: Unalakleet River, approximately 15 miles upstream from village of Unalakleet.
- b) Description: Determine daily and seasonal timing and magnitude of Chinook, chum, and pink escapements. Collect age, sex, and length data from Chinook and chum salmon from weir trap. Cooperative ADF&G, BLM, NSEDC, and Unalakleet IRA project.

**Kobuk River Test Fish**

- a) Location: Lower Kobuk River, approximately two miles downriver of Kiana.
- b) Description: Evaluate chum salmon abundance migrating into the Kobuk River drainage using systematic drift gillnet catches and qualitatively assess the impact of the Kotzebue District commercial salmon fishery on chum abundance into the Kobuk River drainage for fisheries management purposes. Describe migratory timing in the lower Kobuk River. Sample for age, sex, and length. ADF&G project.

**Salmon Lake Limnology Project / Sockeye Salmon Restoration**

- a) Location: Salmon Lake, throughout; and smolt trap two miles downstream from lake, on Pilgrim River.
- b) Description: Restore sockeye salmon population to higher historical levels. Biological (age, weight, and length) samples taken from emigrating smolt and enumerated by mark–recapture. Hydroacoustic-tow net studies conducted to estimate rearing fry population and gather growth data. Fertilization of Salmon Lake. NSEDC project.

**Subsistence Salmon Fishing Surveys**

- a) Location: Norton Sound District.
- b) Description: Determine subsistence utilization of salmon for formulating management procedures and goals. Subsistence salmon permits were issued in northern Norton Sound and Port Clarence District by Commercial Fisheries Division. Koyuk, Shaktoolik, and Unalakleet were also surveyed by Commercial Fisheries Division. ADF&G project.

---

-continued-

**CRAB**

**Offshore Summer King Crab Study**

- a) Location: Tagging occurred along transects 5 and 10 miles from shore from Cape Nome to Elim; observers were placed on commercial fishing vessels throughout the open fishing area of Norton Sound.
- b) Description: Investigate movement, size composition, potential essential habitat, and handling of red king crab in eastern Norton Sound. Cooperative project between ADF&G and NSEDC with funding provided by North Pacific Research Board.

**Norton Sound Red King Crab Trawl Survey (conducted in 2011)**

- a) Location: Ocean waters of Norton Sound, 10-mile grid.
  - b) Description: Triennial trawl survey to establish abundance of red king crab. Biological (sex and size) samples and species presence–absence data taken. Cooperative ADF&G and NSEDC project with financial assistance from the National Oceanic and Atmospheric Administration.
-

Appendix G3.—Commercial processors and buyers operating in Norton Sound and Kotzebue Sound, 2013.

Company	Address	Type of Processing	District
Aqua Tech	P.O. Box 10119 Anchorage, AK 99510	Fresh Crab	Norton Sound
Norton Sound Seafood Products	Nome, AK 99762 and Unalakleet, AK 99684	Frozen/Fresh Salmon Herring Roe King Crab	Norton Sound
Great Pacific Seafoods	Anchorage, AK	Buy and Fly Frozen/Fresh Salmon	Kotzebue Sound
Maniilaq Services, Inc. dba Arctic Coast Wild Salmon	Seattle, WA 98101	Buy and Fly Frozen/Fresh Salmon	Kotzebue Sound



<b>NORTON SOUND 2013 SUBSISTENCE SALMON HARVEST SURVEY</b>		Community ID# 357
Alaska Department of Fish and Game		Household ID# _____
<b>Community:</b> UNALAKLEET <b>Survey Date:</b> _____ <b>Household Size:</b> _____ <b>Interviewer:</b> _____ <b>(If new household) PO Box:</b> _____		
Household participation is voluntary. Individual household data will not be released without permission of household head.		
1. Did your household fish for salmon for subsistence use this year? (Include fishing with a rod and reel) <span style="float: right;"><input type="checkbox"/> YES <input type="checkbox"/> NO</span>		
2. Does your household <u>usually</u> subsistence fish for salmon? <span style="float: right;"><input type="checkbox"/> YES <input type="checkbox"/> NO</span>		
<b>FOR SALMON FISHING HOUSEHOLDS ONLY ("Yes" to #1)</b>		
3. Please estimate how many salmon your household caught for subsistence use this year, including with a rod and reel. It is important not to double count fish harvests. Report only your share of the catch if fishing with others. Include salmon you gave away, ate fresh, fed to dogs, lost to spoilage, or obtained from helping others process fish.		
	<b>NUMBER OF SALMON YOUR HOUSEHOLD HARVESTED (BY GEAR TYPE)</b>	<b>NUMBER OF SALMON YOUR HOUSEHOLD HARVESTED (BY LOCATION)</b>
<b>SPECIES</b>	<b>SUBSISTENCE GILL NET or SEINE</b> <small>(Number of fish)</small>	<b>ROD &amp; REEL</b> <small>(Number of fish)</small>
CHUM SALMON Dog		
CHINOOK SALMON King		
PINK SALMON Humpy		
SCKEYE SALMON Red		
COHO SALMON Silver		
<b>4. Comments or Suggestions?</b>      		

<b>NORTON SOUND 2013 SUBSISTENCE SALMON HARVEST SURVEY</b>		Community ID# 307				
Alaska Department of Fish and Game-		Household ID# _____				
<b>Community:</b> SHAKTOOLIK _____						
<b>Survey Date:</b> _____ <b>Interviewer:</b> _____		<b>Household Size:</b> _____ <b>(If new household) PO Box:</b> _____				
Household participation is voluntary. Individual household data will not be released without permission of household head.						
1. Did your household fish for salmon for subsistence use this year? (Include fishing with a rod and reel)						
		<input type="checkbox"/> YES <input type="checkbox"/> NO				
2. Does your household <u>usually</u> subsistence fish for salmon?						
		<input type="checkbox"/> YES <input type="checkbox"/> NO				
<b>FOR SALMON FISHING HOUSEHOLDS ONLY ("Yes" to #1)</b>						
3. Please estimate how many salmon your household caught for subsistence use this year, including with a rod and reel. It is important not to double count fish harvests. Report only your share of the catch if fishing with others. Include salmon you gave away, ate fresh, fed to dogs, lost to spoilage, or obtained from helping others process fish.						
	<b>NUMBER OF SALMON YOUR HOUSEHOLD HARVESTED (BY GEAR TYPE)</b>	<b>NUMBER OF SALMON YOUR HOUSEHOLD HARVESTED (BY LOCATION)</b>				
<b>SPECIES</b>	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; text-align: center;"> <b>SUBSISTENCE GILL NET or SEINE</b>  <small>(Number of fish)</small> </td> <td style="width: 50%; text-align: center;"> <b>ROD &amp; REEL</b>  <small>(Number of fish)</small> </td> </tr> </table>	<b>SUBSISTENCE GILL NET or SEINE</b> <small>(Number of fish)</small>	<b>ROD &amp; REEL</b> <small>(Number of fish)</small>	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; text-align: center;"> <b>MARINE WATERS</b> </td> <td style="width: 50%; text-align: center;"> <b>SHAKTOOLIK RIVER</b> </td> </tr> </table>	<b>MARINE WATERS</b>	<b>SHAKTOOLIK RIVER</b>
<b>SUBSISTENCE GILL NET or SEINE</b> <small>(Number of fish)</small>	<b>ROD &amp; REEL</b> <small>(Number of fish)</small>					
<b>MARINE WATERS</b>	<b>SHAKTOOLIK RIVER</b>					
CHUM SALMON Dog						
CHINOOK SALMON King						
PINK SALMON Humpy						
SOCKEYE SALMON Red						
COHO SALMON Silver						
<b>4. Comments or Suggestions?</b>						

<b>NORTON SOUND 2013 SUBSISTENCE SALMON HARVEST SURVEY</b>		Community ID# 204				
Alaska Department of Fish and Game-		Household ID# _____				
<b>Community:</b> KOYUK <b>Survey Date:</b> _____ <b>Interviewer:</b> _____						
		<b>Household Size:</b> _____ <b>(If new household) PO Box:</b> _____				
Household participation is voluntary. Individual household data will not be released without permission of household head.						
1. Did your household fish for salmon for subsistence use this year? (Include fishing with a rod and reel)						
		<input type="checkbox"/> YES	<input type="checkbox"/> NO			
2. Does your household <u>usually</u> subsistence fish for salmon?						
		<input type="checkbox"/> YES	<input type="checkbox"/> NO			
<b>FOR SALMON FISHING HOUSEHOLDS ONLY ("Yes" to #1)</b>						
3. Please estimate how many salmon your household caught for subsistence use this year, including with a rod and reel. It is important not to double count fish harvests. Report only your share of the catch if fishing with others. Include salmon you gave away, ate fresh, fed to dogs, lost to spoilage, or obtained from helping others process fish.						
<b>SPECIES</b>	<b>NUMBER OF SALMON YOUR HOUSEHOLD HARVESTED (BY GEAR TYPE)</b>		<b>NUMBER OF SALMON YOUR HOUSEHOLD HARVESTED (BY LOCATION)</b>			
	<b>SUBSISTENCE GILL NET or SEINE (Number of fish)</b>	<b>ROD &amp; REEL (Number of fish)</b>	<b>MARINE WATERS</b>	<b>KOYUK RIVER</b>	<b>INGLUTALIK RIVER</b>	<b>UNGALIK RIVER</b>
<b>CHUM</b> Dog						
<b>CHINOOK</b> King						
<b>PINK</b> Humpy						
<b>SOCKEYE</b> Red						
<b>COHO</b> Silver						
<b>4. Comments or Suggestions?</b>						

**RED KING CRAB**

Emergency Order: 3-C-Z-01-13    Effective Date: July 3, 2013

EXPLANATION: This emergency order opens both the CDQ fishery and the commercial open access crab fishery in Norton Sound from 12:00 noon Wednesday, July 3 until 12:00 noon Monday, September 2, or when the CDQ and the open access quota is reached.

JUSTIFICATION: By regulation the open access king crab fishery can open anytime on or after June 15 by Emergency Order. Currently two land-based processor-buyers are registered and both buyers are ready to purchase open access crab. The guideline harvest level for the 2013 Norton Sound open access fishery is 458,430 pounds. By regulation the CDQ crab fishery can open anytime the CDQ group is ready to harvest the crab. The CDQ crab can only be harvested by permit holders approved by Norton Sound Economic Development Corporation and the quota is 37,170 pounds. The CDQ group has notified the department they are ready to harvest crab.

Emergency Order: 3-C-Z-02-13    Effective Date: August 1, 2013

EXPLANATION: This emergency order moves the closed boundary line between 166°15'W (Sledge Island) and 163°30'W (near Square Rock) north by 3 miles, to 64°18'N from the current 64°15'N.

JUSTIFICATION: As of July 30, approximately 145,000 pounds have been harvested out of the total GHIL of 495,600 pounds. Based on the current catch rate, the GHIL is projected to be reached in mid- to late-September. The department is reducing the area closed to commercial crab fishermen by an incremental amount in order to give fishermen access to additional fishing grounds so they can more efficiently harvest the allowable quota of red king crab. Analysis of results from the spring tagging project and known distribution patterns from trawl surveys suggests this expanded area will allow additional access to legal king crab while minimizing handling of non-target king crab such as females and juvenile crab. Due to the particularly large biomass of crab available for all fisheries, this small increase in area available to commercial harvest is not expected to adversely impact the subsistence king crab fisheries or the future health of the king crab population.

Emergency Order: 3-C-Z-03-13    Effective Date: August 7, 2013

EXPLANATION: This emergency order moves the closed boundary line between 162°38'W (Carson Creek) and 162°W (the eastern border of statistical area 626401) north by 5 miles to 64°15'N from the current 64°10'N. Additionally, the closure line between 162°W and 161°30'W (Cape Denbigh) will be moved north, by 3 miles to 64°18'N from the current 64°15'N.

JUSTIFICATION: As of August 6, approximately 191,000 pounds have been harvested out of the total GHIL of 495,600 pounds. Based on the current catch rate, the GHIL is projected to be reached in mid- to late-September. The department is reducing the area on the east side of Norton Sound closed to commercial crab fishermen by an incremental amount in order to give east side fishermen access to additional fishing grounds so they can more efficiently harvest the allowable quota of red king crab. Results from the spring tagging project indicate this newly opened area is likely to have an appreciable amount of large male red king crab.

Emergency Order: 3-C-Z-04-13    Effective Date: September 3, 2013

EXPLANATION: This emergency order extends both the CDQ fishery and the commercial open access crab fishery in Norton Sound from 12:00 noon Tuesday, September 3 until 12:00 noon Tuesday, September 10, unless there are conservation concerns or there is no longer any buyer interest. In addition, the closed boundary line between 166°15'W (Sledge Island) and 163°30'W (near Square Rock) that was moved north by 3 miles, to 64°18'N from 64°15'N on August 1, will remain at 64°18'N. The closed boundary line between 162°38'W (Carson Creek) and 162°W (the eastern border of statistical area 626401) that was moved north by 5 miles to 64°15'N from 64°10'N, and the closure line between 162°W and 161°30'W (Cape Denbigh) that was moved north by 3 miles to 64°18'N

---

-continued-

from 64°15'N on August 7, will both remain at the new location until September 10.

**JUSTIFICATION:** Currently four processor-buyers are registered and most buyers continue to be interested in purchasing open access crab. The guideline harvest level for the 2013 Norton Sound open access fishery is 458,430 pounds and for the CDQ fishery, the quota is 37,170 pounds. Through the morning of September 3, total reported open access harvest is ~289,800 pounds, and total CDQ harvest is 13,480 pounds. Roughly 168,600 pounds remain of the open access quota and 23,690 pounds remain of the CDQ quota. Based on the current catch rate, the GHF is not expected to be reached this year. However, the season is being extended to provide fishermen with additional harvest opportunity on the remaining quotas unless there are indications that crab are molting. In 2009, during a similar temperature regime to this current year, molting crab began appearing in the harvest during mid-September. Therefore, the fishery may be able to extend for an additional week without harvesting appreciable quantities of molting crab.

**Emergency Order:** 3-C-Z-05-13    **Effective Date:** September 10, 2013

**EXPLANATION:** This emergency order extends both the CDQ fishery and the commercial open access crab fishery in Norton Sound from 12:00 noon Tuesday, September 10 until 6:00 PM Friday, September 13, unless there are conservation concerns or there is no longer any buyer interest. In addition, the closed boundary line between 166°15'W (Sledge Island) and 163°30'W (near Square Rock) that was moved north by 3 miles, to 64°18'N from 64°15'N on August 1, will remain at 64°18'N. The closed boundary line between 162°38'W (Carson Creek) and 162°W (the eastern border of statistical area 626401) that was moved north by 5 miles to 64°15'N from 64°10'N, and the closure line between 162°W and 161°30'W (Cape Denbigh) that was moved north by 3 miles to 64°18'N from 64°15'N on August 7, will both remain at the new location until September 13.

**JUSTIFICATION:** Currently four processor-buyers are registered and all buyers continue to be interested in purchasing crab. The guideline harvest level for the 2013 Norton Sound open access fishery is 458,430 pounds and for the CDQ fishery, the quota is 37,170 pounds. Through the morning of September 10, total reported open access harvest is ~336,100 pounds, and total CDQ harvest is 16,252 pounds. Roughly 122,300 pounds remain of the open access quota and 20,918 pounds remain of the CDQ quota. The GHF will not be reached this year. However, the season is being extended again to provide fishermen with additional harvest opportunity on the remaining quotas. A handful of newly molted crab has shown up in the commercial fishery, but meat fill samples from the major buyer have remained consistent in the 60-70% range by weight since early August. Therefore, the fishery may be able to extend for an additional 3 days without harvesting appreciable quantities of molting crab.

**Emergency Order:** 3-C-Z-06-13    **Effective Date:** September 13, 2013

**EXPLANATION:** This emergency order extends by one day the Norton Sound CDQ and commercial open access king crab fishery, which will close at 6:00PM on Saturday, September 14. Permit holders must have all pots unbaited and secured open by 6:00PM, Saturday, September 14 and all pots must be removed from the water by Saturday, September 21, 2013.

**JUSTIFICATION:** The guideline harvest level for the 2013 Norton Sound king crab fishery of 495,600 pounds will not be reached. However, because of sudden adverse marine conditions, the commercial open access and CDQ king crab fishery in Norton Sound originally scheduled to close at 6:00 PM Friday, September 13 will now have fishing time extended to 6:00 PM Saturday, September 14. All pot gear must have doors open and bait containers removed by 6:00 PM Saturday, September 14. All crab must be delivered by 6:00 PM Sunday, September 15 and all crab gear used in the commercial fishery must be out of the water by Saturday, September 21. Weather is expected to lay down by Saturday morning.

## **HERRING**

**Emergency Order:** 3-H-Z-1-13    **Effective Date:** June 7, 2013

**EXPLANATION:** This emergency order opens the Norton Sound District to commercial gillnet fishing for sac roe herring beginning 6:00 p.m. Friday, June 7, 2013 until Monday, July 1, 2013, unless superseded by another

---

-continued-

emergency order.

**JUSTIFICATION:** As of this morning, water temperatures between 5–6 degrees Celsius have been recorded amidst the floating ice and farther away from melt water plumes, water temperatures as high as 7 degrees have been observed. Approximately 50 tons of herring were also observed by NSEDC biologists on June 5 near Cape Denbigh. Taken collectively, all available assessment information suggests that large schools of ripe herring are accumulating offshore in deeper, relatively warm water, but will soon approach the southeastern Norton Sound coastline to spawn in abundance, possibly as early as this weekend. Once major spawning begins, excellent catch rates with high roe recovery are expected in the fishery for 2–4 days, which will be followed by a sharp decline in catch rates and percent roe as the larger, older age-class herring vacate the spawning grounds. Any herring not purchased by the buyer must be retained for personal or subsistence uses.

#### **KOTZEBUE SALMON**

Emergency Order: 3-S-X-01-13   Effective Date: July 10, 2013

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District for 8 hours from the hours of 8 p.m. Wednesday, July 10 until 4 a.m. Thursday, July 11.

**JUSTIFICATION:** One major commercial salmon buyer has registered to purchase Kotzebue chum salmon this season. The buyer has limited quantities of ice and airline schedules will affect the buyer's ability to ship fish out. Regulation allows the season to be open from July 10 through August 31. The buyer has notified the department that they would like to begin purchasing fish on the evening of July 10. This 8 hour opening will serve as test of earlier run strength.

Emergency Order: 3-S-X-02-13   Effective Date: July 11, 2013

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District for 6 hours from the hours of 8 p.m. Thursday, July 11 until 2 a.m. Friday, July 12.

**JUSTIFICATION:** The first commercial salmon fishing period of the season was 8 hours last night and 10 permit holders caught 1,360 chum salmon. The catch was average for this date. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-03-13   Effective Date: July 12, 2013

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District for 8 hours from the hours of 6 p.m. Friday, July 12 until 2 a.m. Saturday, July 13.

**JUSTIFICATION:** The second commercial salmon fishing period of the season was 6 hours last night and 12 permit holders caught 1,535 chum salmon. The catch was average for this date. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-04-13   Effective Date: July 13, 2013

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District from the hours of 8 a.m. to 1 p.m. daily beginning on Saturday, July 13 until 1 p.m. Saturday, July 20.

**JUSTIFICATION:** There is a small market buyer interested in buying salmon from one permit holder. Compared to the normal fishing effort of over 30 permit holders during some periods during this time last year the small effort should not jeopardize making escapement goals or subsistence opportunity.

Emergency Order: 3-S-X-05-13   Effective Date: July 14, 2013

---

-continued-

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District for 8 hours from the hours of 6 p.m. Sunday, July 14 until 2 a.m. Monday, July 15.

**JUSTIFICATION:** The third commercial salmon fishing period of the season was 8 hours Friday night and 9 permit holders caught 2,368 chum salmon. The catch was average for this date. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

**Emergency Order:** 3-S-X-06-13   **Effective Date:** July 15, 2013

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District for 8 hours from the hours of 6 p.m. Monday, July 15 until 2 a.m. Tuesday, July 16.

**JUSTIFICATION:** The fourth commercial salmon fishing period of the season was 8 hours Sunday night and 10 permit holders caught 2,077 chum salmon. The catch was average for this date. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

**Emergency Order:** 3-S-X-07-13   **Effective Date:** July 16, 2013

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District for 8 hours from the hours of 6 p.m. Tuesday, July 16 until 2 a.m. Wednesday, July 17.

**JUSTIFICATION:** The fifth commercial salmon fishing period of the season was 8 hours Monday night and 17 permit holders caught 2,038 chum salmon. The catch was average for this date. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

**Emergency Order:** 3-S-X-08-13   **Effective Date:** July 17, 2013

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District for 8 hours from the hours of 6 p.m. Wednesday, July 17 until 2 a.m. Thursday, July 18.

**JUSTIFICATION:** The sixth commercial salmon fishing period of the season was 8 hours Tuesday night and 18 permit holders caught 5,994 chum salmon. The catch was more than double the previous 8-hour opening, but was average for this date. Usually commercial catch jumps up the in mid-July and will continue to build until early August during normal run timing years. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

**Emergency Order:** 3-S-X-09-13   **Effective Date:** July 18, 2013

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District for 8 hours from the hours of 6 p.m. Thursday, July 18 until 2 a.m. Friday, July 19.

**JUSTIFICATION:** The seventh commercial salmon fishing period of the season was 8 hours Wednesday night and 23 permit holders caught 4,908 chum salmon. The catch was average for this date. The cumulative catch is now over 20,000 chum salmon. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

**Emergency Order:** 3-S-X-10-13   **Effective Date:** July 19, 2013

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District for 4 hours from the hours of 8 p.m. Friday, July 19 until 12 a.m. Saturday, July 20.

**JUSTIFICATION:** The eighth commercial salmon fishing period of the season was 8 hours Thursday night and 29 permit holders caught 10,476 chum salmon. Usually during the second week of July the catch often does jump to 10,000 plus fish. The cumulative catch is now over 30,000 chum salmon. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

---

-continued-

Emergency Order: 3-S-X-11-13    Effective Date: July 21, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District from the hours of 7 a.m. to 1 p.m. daily beginning on Sunday, July 21 until 1 p.m. Saturday, July 27.

JUSTIFICATION: There is a small market buyer interested in buying salmon from one permit holder. Compared to the normal fishing effort of over 30 permit holders during some periods during this time last year the small effort should not jeopardize making escapement goals or subsistence opportunity.

Emergency Order: 3-S-X-12-13    Effective Date: July 21, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 6 hours from the hours of 8 p.m. Sunday, July 21 until 2 a.m. Monday, July 22.

JUSTIFICATION: The ninth commercial salmon fishing period of the season was 4 hours Friday night and 26 permit holders caught 6,013 chum salmon. Increasing run strength necessitated the buyer to reduce fishing time from 8 hours to 4 hours. Because the run is building normally the buyer has capacity concerns and will likely continue with openings less than 8 hours. The Kobuk River test fishing project has been operational since Wednesday with catches better than last year over the same time period. Last year's season catch index ranked third out of 20 years. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-13-13    Effective Date: July 22, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 6 hours from the hours of 8 p.m. Monday, July 22 until 2 a.m. Tuesday, July 23.

JUSTIFICATION: The tenth commercial salmon fishing period of the season was 6 hours Sunday night and 33 permit holders caught 5,586 chum salmon. The Kobuk River test fishing project has been operational since Wednesday with catches better than last year over the same time period. Last year's season catch index ranked third out of 20 years. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-14-13    Effective Date: July 23, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 6 hours from the hours of 6 p.m. Tuesday, July 23 until 12 a.m. Wednesday, July 24.

JUSTIFICATION: The eleventh commercial salmon fishing period of the season was 6 hours Monday night and 34 permit holders caught 6,216 chum salmon. Catch and fishing effort were average for this time period. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-15-13    Effective Date: July 24, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 4 hours from the hours of 8 p.m. Wednesday, July 24 until 12 a.m. Thursday, July 25.

JUSTIFICATION: The twelfth commercial salmon fishing period of the season was 4 hours Tuesday night and 38 permit holders caught 17,712 chum salmon. The catch was the second highest catch for one fishing period since a buyer returned to the Kotzebue commercial fishery in 2004. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-16-13    Effective Date: July 25, 2013

---

-continued-



**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District for 6 hours from the hours of 6 p.m. Thursday, July 25 until 12 a.m. Friday, July 26.

**JUSTIFICATION:** Catch during last night's 4-hour opening was 2,683 chum salmon by 9 permit holders. Windy conditions likely resulted in a reduced fishing effort last night. The commercial harvest this season is expected to be within the range of 225,000 to 250,000 chum salmon if market conditions can accept that level of harvest. The total commercial harvest to date is nearly 70,000 chum salmon. Last year by this date the harvest was 60,000 chum salmon and total harvest for the season was nearly 228,000 chum salmon. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

**Emergency Order:** 3-S-X-17-13    **Effective Date:** July 28, 2013

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District from the hours of 7 a.m. to 1 p.m. daily beginning on Sunday, July 28 until 1 p.m. Saturday, August 10.

**JUSTIFICATION:** There is a small market buyer interested in buying salmon from one permit holder. The major buyer is restricting fishing time with the peak weeks of salmon passage now and plans to request fishing openings for limited hours in the evening. Compared to the normal fishing effort of over 30 permit holders during some periods this season the small effort should not jeopardize making escapement goals or subsistence opportunity.

**Emergency Order:** 3-S-X-18-13    **Effective Date:** July 26, 2013

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District for 4 hours from the hours of 6 p.m. until 10 p.m. Friday, July 26.

**JUSTIFICATION:** Catch during last night's 6-hour opening was 11,894 chum salmon by 41 permit holders. The commercial harvest this season is expected to be within the range of 225,000 to 250,000 chum salmon if market conditions can accept that level of harvest. The commercial harvest to date is over 80,000 chum salmon. Last year by this date the harvest was 63,000 chum salmon and total harvest for the season was nearly 228,000 chum salmon. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

**Emergency Order:** 3-S-X-19-13    **Effective Date:** July 28, 2013

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District for 6 hours from the hours of 6 p.m. Sunday, July 28 until 12 a.m. Monday, July 29.

**JUSTIFICATION:** Catch during Friday night's 4-hour opening was 9,083 chum salmon by 40 permit holders. Chum salmon catch index at the department test fish project on Kobuk River near Kiana ranks sixth highest in 21-year project history indicating a good chum salmon run to the Kobuk River system. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

**Emergency Order:** 3-S-X-20-13    **Effective Date:** July 30, 2013

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District for 5 hours from the hours of 6 p.m. until 11 p.m. Tuesday, July 30.

**JUSTIFICATION:** Catch during Sunday night's 6-hour opening was 17,283 chum salmon by 41 permit holders. Chum salmon catch index at the department test fish project on Kobuk River near Kiana ranks sixth highest in 21-year project history indicating a good chum salmon run to the Kobuk River system. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

**Emergency Order:** 3-S-X-21-13    **Effective Date:** July 31, 2013

---

-continued-

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District for 6 hours daily from the hours of 6 p.m. to 12 a.m. beginning on Wednesday, July 31 until 12 a.m. Tuesday, August 6.

**JUSTIFICATION:** Cumulative test fish catch ranks sixth highest out of 21 years at the Kobuk River test fish project. The cumulative test fish index is expected to reach 600 today and that is the minimum amount needed to ensure escapement and allow continued commercial fishing. Continued short duration openings the coming week should not jeopardize subsistence opportunity.

**Emergency Order:** 3-S-X-22-13    **Effective Date:** August 1, 2013

**EXPLANATION:** This emergency order amends emergency order 3-S-X-21-13 by reducing commercial fishing time from 6 hours to 4 hours for period on Thursday, August 1 in the Kotzebue District. Effective 6:00 p.m. Thursday, August 1, this emergency order reopens the Kotzebue District for 4 hours from 6:00 p.m. to 10:00 p.m. Effective Friday, August 2, this emergency order permits commercial salmon fishing to resume on a fishing schedule of 6 hours a day from 6:00 p.m. to 12:00 midnight, until 12:00 a.m. Tuesday, August 6.

**JUSTIFICATION:** Cumulative test fish catch ranks sixth highest out of 21 years at the Kobuk River test fish project. The cumulative test fish index is expected to reach 600 today and that is the minimum amount needed to ensure escapement and allow continued commercial fishing. Continued short duration openings the coming week should not jeopardize subsistence opportunity.

**Emergency Order:** 3-S-X-23-13    **Effective Date:** August 4, 2013

**EXPLANATION:** This emergency order amends emergency order 3-S-X-22-13 by reducing commercial fishing time from 6 hours to 4 hours for period on Sunday, August 4 in the Kotzebue District. Effective 6:00 p.m. Sunday, August 4, this emergency order reopens the Kotzebue District for 4 hours from 6:00 p.m. to 10:00 p.m. Effective Monday, August 5, this emergency order permits commercial salmon fishing to resume on a fishing schedule of 6 hours a day from 6:00 p.m. to 12:00 midnight, until 12:00 a.m. Tuesday, August 6.

**JUSTIFICATION:** Cumulative test fish catch ranks fourth highest out of 21 years at the Kobuk River test fish project. The cumulative test fish index is at 997 index points which is well above the minimum amount (600 points) needed to ensure escapement and allow continued commercial fishing. Continued short duration openings the coming week should not jeopardize subsistence opportunity.

**Emergency Order:** 3-S-X-24-13    **Effective Date:** August 5, 2013

**EXPLANATION:** This emergency order supersedes emergency order 3-S-X-23-13 by reducing commercial fishing time from 6 hours to 4 hours for period on Sunday, August 4 in the Kotzebue District. Effective 6:00 p.m. Sunday, August 4, this emergency order reopens the Kotzebue District for 4 hours from 6:00 p.m. to 10:00 p.m.

**JUSTIFICATION:** Cumulative test fish catch ranks fourth highest out of 21 years at the Kobuk River test fish project. The cumulative test fish catch has reached 1,100 index points yesterday and that is well above the minimum index of 600 fish needed to ensure escapement and allow continued commercial fishing. Continued short duration openings the coming week should not jeopardize subsistence opportunity.

**Emergency Order:** 3-S-X-25-13    **Effective Date:** August 6, 2013

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District for 4 hours from the hours of 6 p.m. until 10 p.m. Tuesday, August 6.

**JUSTIFICATION:** Catch during last night's 4-hour opening was 9,672 chum salmon by 40 permit holders. The commercial harvest this season is tracking ahead the department forecast range of 225,000 to 250,000 chum salmon. The commercial harvest to date is over 165,000 chum salmon. Last year by this date the harvest was 151,000 chum salmon and total harvest for the season was nearly 228,000 chum salmon. Continuing with short duration

---

-continued-

commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-26-13    Effective Date: August 7, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 4 hours from the hours of 7 p.m. until 11 p.m. Wednesday, August 7.

JUSTIFICATION: Catch during last night's 4-hour opening was 7,300 chum salmon by 34 permit holders. The commercial harvest this season is tracking ahead the department forecast range of 225,000 to 250,000 chum salmon. The commercial harvest to date is over 172,000 chum salmon. Last year by this date the harvest was 158,000 chum salmon and total harvest for the season was nearly 228,000 chum salmon. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-27-13    Effective Date: August 8, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 4 hours from the hours of 5 p.m. until 9 p.m. Thursday, August 8.

JUSTIFICATION: Catch during last night's 4-hour opening was 12,427 chum salmon by 41 permit holders. The commercial harvest this season is tracking ahead the department forecast range of 225,000 to 250,000 chum salmon. The commercial harvest to date is over 185,000 chum salmon. Last year by this date the harvest was 175,000 chum salmon and total harvest for the season was nearly 228,000 chum salmon. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-28-13    Effective Date: August 11, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District from the hours of 7 a.m. to 1 p.m. daily beginning on Sunday, August 11 until 1 p.m. Saturday, August 17.

JUSTIFICATION: There is a small market buyer interested in buying salmon from a few permit holders. The major buyer is restricting fishing time with the peak weeks of salmon passage now and plans to request fishing openings for limited hours in the evening. Compared to the normal fishing effort of over 40 permit holders during some periods this season the small effort should not jeopardize making escapement goals or subsistence opportunity.

Emergency Order: 3-S-X-29-13    Effective Date: August 11, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 4 hours from the hours of 5 p.m. until 9 p.m. Sunday, August 11.

JUSTIFICATION: Catch during Thursday night's 4-hour opening was 10,875 chum salmon by 43 permit holders. The commercial harvest to date is over 195,000 chum salmon. Last year by this date the harvest was 188,000 chum salmon and total harvest for the season was nearly 228,000 chum salmon. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-30-13    Effective Date: August 12, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 4 hours from the hours of 5 p.m. until 9 p.m. Monday, August 12.

JUSTIFICATION: Catch during Sunday night's 4-hour opening was 11,830 chum salmon by 35 permit holders. The catch was average for this date. The commercial harvest to date is over 208,000 chum salmon. The department forecast was for a commercial harvest of 225,000 to 250,000 chum salmon in 2013. Last year by this date the harvest was 188,000 chum salmon and total harvest for the season was nearly 228,000 chum salmon. Catch at the Kobuk River test fish project continues to be well above average and ranks second best in the 21-year project

---

-continued-

history. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-31-13   Effective Date: August 13, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 4 hours from the hours of 5 p.m. until 9 p.m. Tuesday, August 13.

JUSTIFICATION: Catch during Monday night's 4-hour opening was 11,067 chum salmon by 40 permit holders. The catch was average for this date. The commercial harvest to date is over 219,000 chum salmon. The department forecast was for a commercial harvest of 225,000 to 250,000 chum salmon in 2013. Last year by this date the harvest was 195,000 chum salmon and total harvest for the season was nearly 228,000 chum salmon. Catch at the Kobuk River test fish project continues to be well above average and ranks second best in the 21-year project history. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-32-13   Effective Date: August 14, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 4 hours from the hours of 5 p.m. until 9 p.m. Wednesday, August 14.

JUSTIFICATION: Catch during Tuesday night's 4-hour opening was 10,917 chum salmon by 40 permit holders. The catch was average for this date. The commercial harvest to date is over 230,000 chum salmon. The department forecast was for a commercial harvest of 225,000 to 250,000 chum salmon in 2013 and this year's catch has exceeded last year's catch of nearly 228,000 chum salmon. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-33-13   Effective Date: August 15, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 4 hours from the hours of 5 p.m. until 9 p.m. Thursday, August 15.

JUSTIFICATION: Catch during last night's 4-hour opening was 12,057 chum salmon by 44 permit holders. The catch was above average for this date. The commercial harvest to date is over 242,000 chum salmon. The department forecast was for a commercial harvest of 225,000 to 250,000 chum salmon in 2013 and the forecast will easily be exceeded this year. Kobuk River test catch index ranks second best and projections indicate the catch index will likely be a record at project conclusion. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-34-13   Effective Date: August 16, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 3 hours from the hours of 6 p.m. until 9 p.m. Friday, August 16.

JUSTIFICATION: Catch during last night's 4-hour opening was 15,805 chum salmon by 44 permit holders. The catch was above average for this date. During the past week the catch was the best this season with over 60,000 chum salmon harvested. The commercial harvest to date is over 258,000 chum salmon. The department forecast of a 225,000 to 250,000 chum salmon harvest in 2013 has been exceeded. Kobuk River test catch index ranks second best and projections indicate the catch index will likely be a record at project conclusion. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-35-32   Effective Date: August 18, 2013

---

-continued-

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District from the hours of 7 a.m. to 1 p.m. daily beginning on Sunday, August 18 until 1 p.m. Saturday, August 24.

**JUSTIFICATION:** There is a small market buyer interested in buying salmon from a few permit holders. The major buyer is restricting fishing time with the peak week of salmon catch this week and plans to request fishing openings for limited hours in the evening. Compared to the normal fishing effort of over 40 permit holders during some periods this season the small effort should not jeopardize making escapement goals or subsistence opportunity.

**Emergency Order:** 3-S-X-36-13    **Effective Date:** August 18, 2013

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District for 3 hours from the hours of 6 p.m. until 9 p.m. Sunday, August 18.

**JUSTIFICATION:** Catch during Friday night's 3-hour opening was 8,385 chum salmon by 38 permit holders. The catch was above average for the time fished. During the past week the catch was the best this season with over 70,000 chum salmon harvested. The commercial harvest to date is over 266,000 chum salmon. The department forecast of a 225,000 to 250,000 chum salmon harvest in 2013 has been exceeded. Kobuk River test catch index ranks second best and projections indicate the catch index will likely be a record at project conclusion. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

**Emergency Order:** 3-S-X-37-13    **Effective Date:** August 19, 2013

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District for 2 hours from the hours of 7 p.m. until 9 p.m. Monday, August 19.

**JUSTIFICATION:** Catch during Sunday night's 3-hour opening was 9706 chum salmon by 36 permit holders. The catch was above average for this time of the year. The commercial harvest to date is over 276,000 chum salmon and is a record for the 2000s. Kobuk River test catch index ranks second best and projections indicate the catch index be a record by the conclusion test fishing today or tomorrow. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

**Emergency Order:** 3-S-X-38-13    **Effective Date:** August 20, 2013

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District for 3 hours from the hours of 6 p.m. until 9 p.m. Tuesday, August 20.

**JUSTIFICATION:** Catch during Monday night's 2-hour opening was 4,229 chum salmon by 35 permit holders. The catch was above average for the time fished. The commercial harvest to date is over 280,000 chum salmon. The department forecast of a 225,000 to 250,000 chum salmon harvest in 2013 has been exceeded. Kobuk River test catch index had the best catch index this season in the 21-year project history. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

**Emergency Order:** 3-S-X-39-13    **Effective Date:** August 21, 2013

**EXPLANATION:** This emergency order opens commercial salmon fishing in the Kotzebue District for 4 hours from the hours of 5 p.m. until 9 p.m. Wednesday, August 21.

**JUSTIFICATION:** Catch during Tuesday night's 3-hour opening was 3,467 chum salmon. The catch was average for the time fished. The commercial harvest to date is over 286,000 chum salmon and has exceeded the department forecast of a 225,000 to 250,000 chum salmon harvest. Kobuk River test catch index had the best catch index this season in the 21-year project history. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

---

-continued-

Emergency Order: 3-S-X-40-13    Effective Date: August 22, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 5 hours from the hours of 4 p.m. until 9 p.m. Thursday, August 22.

JUSTIFICATION: Catch during Wednesday night's 4-hour opening was nearly 4,000 chum salmon by 23 permit holders. The catch was average for this date. The commercial harvest for the season is nearly 290,000 chum salmon and with the upcoming period will likely push past 291,000 chum salmon harvested making it the best commercial harvest in 25 years. This season's Kobuk River test catch index has been the best in the 21-year project history. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-41-13    Effective Date: August 23, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 5 hours from the hours of 4 p.m. until 9 p.m. Friday, August 23.

JUSTIFICATION: Catch during Thursday night's 5-hour opening was 7,052 chum salmon by 35 permit holders. The catch was well above average for this date. The commercial harvest for the season is over 296,000 chum salmon and with the upcoming period will likely push past 300,000 chum salmon harvested for only the tenth time in the 52-year history of the fishery. This season's Kobuk River test catch index has been the best in the 21-year project history. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-42-13    Effective Date: August 25, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 5 hours from the hours of 4 p.m. until 9 p.m. Sunday, August 25.

JUSTIFICATION: Catch during Thursday night's 5-hour opening was 2,973 chum salmon by 16 permit holders. The catch and CPUE was above average for this date. The commercial harvest for the season is over 299,000 chum salmon and with the upcoming period will likely push past 300,000 chum salmon harvested for only the tenth time in the 52-year history of the fishery. This season's Kobuk River test catch index has been the best in the 21-year project history. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-43-13    Effective Date: August 26, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 6 hours from the hours of 3 p.m. until 9 p.m. Monday, August 26.

JUSTIFICATION: Catch during Sunday night's 5-hour opening was 2,874 chum salmon by 13 permit holders. The catch was average for this date. The commercial harvest for the season is over 300,000 chum salmon and is the best since 1988. The Kobuk River test fishing crew has finished for the season and the chum salmon catch index was the highest in the 21 year project history. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-44-13    Effective Date: August 27, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 6 hours, from the hours of 6 a.m. to 12 p.m. daily, beginning on Tuesday, August 27 until 12 p.m. Saturday, August 31.

JUSTIFICATION: There is a small market buyer interested in buying salmon from a few permit holders. The major buyer is buying during a fishing period later in the day. Compared to the normal fishing effort of over 40 permit

---

-continued-

holders during some periods this season the small effort should not jeopardize making escapement goals or subsistence opportunity.

Emergency Order: 3-S-X-45-13    Effective Date: August 27, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 6 hours from the hours of 3 p.m. until 9 p.m. Tuesday, August 27.

JUSTIFICATION: Catch during Monday night's 6-hour opening was 392 chum salmon by 6 permit holders. The catch was well below average for this date, but most permit holders were kept on the beach by weather. The commercial harvest for the season is over 300,000 chum salmon and is the best since 1988. The Kobuk River test fishing crew has finished for the season and the chum salmon catch index was the highest in the 21 year project history. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-46-13    Effective Date: August 28, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 6 hours from the hours of 3 p.m. until 9 p.m. Wednesday, August 28.

JUSTIFICATION: Catch during Tuesday two 6-hour opening was 2,903 chum salmon. The catch was average for the fished. The commercial harvest for the season is over 300,000 chum salmon and is the best since 1988. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-47-13    Effective Date: August 29, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 6 hours from the hours of 12 p.m. until 6 p.m. Thursday, August 29.

JUSTIFICATION: There are two buyers registered in the Kotzebue District and one had requested a six hour morning fishing period and the department had issued a previous emergency order for a six hour fishing period starting at 6 a.m. The second buyer requested a six hour afternoon fishing period. Because the commercial fishing season is winding down and there are no longer any buyer capacity concerns the department is allowing a combined 12 hour fishing period. The commercial harvest for the season is over 300,000 chum salmon and is the best since 1988. The last scheduled day of commercial salmon fishing is August 31. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-X-48-13    Effective Date: August 30, 2013

EXPLANATION: This emergency order opens commercial salmon fishing in the Kotzebue District for 6 hours from the hours of 12 p.m. until 6 p.m. Friday, August 30.

JUSTIFICATION: There are two buyers registered in the Kotzebue District and one had requested a six hour morning fishing period and the department had issued a previous emergency order for a six hour fishing period starting at 6 a.m. The second buyer requested a six hour afternoon fishing period. Because the commercial fishing season is winding down and there are no longer any buyer capacity concerns the department is allowing a combined 12 hour fishing period. The commercial harvest for the season is over 300,000 chum salmon and is the best since 1988. The last scheduled day of commercial salmon fishing is August 31. Continuing with short duration commercial openings should not jeopardize subsistence opportunity or escapement.

---

-continued-

### **NORTON SOUND SALMON**

Emergency Order: 3-S-Z-01-13   Effective Date: June 15, 2013

EXPLANATION: This emergency order sets the subsistence salmon gillnet fishing schedule for the Nome Subdistrict and catch limits for the Nome Subdistrict (Subdistrict 1), and Pilgrim and Kuzitrin Rivers in the Port Clarence District. The subsistence salmon gillnet schedule will be from 6 p.m. Wednesday until 6 p.m. Saturday in the Nome Subdistrict marine waters west of Cape Nome and the catch limits for all locations are listed on the permits. Beach seines are allowed to be used during the salmon gillnet schedule.

JUSTIFICATION: The department forecast for 2013 is that the chum salmon run will exceed the ANS and Tier II restrictions will not be required. Because of the late spring the department is allowing beach seines to be used to increase the efficiency of the harvest and will reassess the use of beach seines when the peak of the run enters Nome Subdistrict rivers in July. Catch limits are still in effect for the various marine and fresh water subsistence areas. All catch limits are listed on the permits. The department staff will be flying frequent aerial surveys and boating some of the rivers to track the salmon migration strength and progress. The weirs and towers on the Nome, Snake, Eldorado, Solomon and Pilgrim Rivers, will also be used to track the various salmon migrations. If a stream appears to have adequate escapement, catch limits will be lifted in that area.

Emergency Order: 3-S-Z-02-13   Effective Date: June 17, 2013

EXPLANATION: This emergency order closes all marine waters in Norton Sound Subdistricts 5 and 6, the Shaktoolik and Unalakleet Subdistricts to subsistence salmon fishing effective 8:00 a.m. Monday, June 17 and reopens Subdistricts 5 and 6 to subsistence fishing schedules and gear restrictions in Subdistrict 5 until midnight Sunday evening, July 14.

Effective 8:00 a.m. Monday, June 17, subsistence salmon fishing will reopen in the marine waters of Subdistrict 5 to a subsistence fishing schedule of two 48-hour periods per week, but will be restricted to set gillnets with a mesh size no greater than 6 inches. Subsistence fishing periods in the Shaktoolik Subdistrict will be from 6:00 p.m. Mondays to 6:00 p.m. Wednesdays, and from 6:00 p.m. Thursdays to 6:00 p.m. Saturdays.

Also effective 8:00 a.m. Monday, June 17, the marine waters of Subdistrict 6, the Unalakleet Subdistrict, will reopen to subsistence salmon fishing on a schedule of two 24-hour periods per week. Subsistence fishing periods in the Unalakleet Subdistrict will be from 6:00 p.m. Mondays to 6:00 p.m. Tuesdays and from 6:00 p.m. Fridays to 6:00 p.m. Saturdays.

JUSTIFICATION: Chinook salmon runs to southeastern Norton Sound are expected to be very poor this season. The combined Subdistricts 5 and 6 Chinook salmon run could be as low as 4,500 fish this season, and this level of abundance is not sufficient to justify sport fishery harvests or support historical Chinook salmon harvest rates. These preemptive restrictions are being implemented this season to significantly reduce harvest rates of Chinook salmon and ensure that escapement goals are achieved in the Shaktoolik and Unalakleet River drainages. In addition to restrictions, subsistence users are also strongly encouraged to redirect harvest pressure on other more numerous species this season, such as chum salmon which are anticipated to be abundant in southern Norton Sound. Once the Chinook salmon run begins, inseason management will be based upon escapement counts and subsistence fishing reports. Fishing restrictions may be relaxed or further restrictions may be necessary depending on run assessment.

Emergency Order: 3-S-Z-03-13   Effective Date: June 17, 2013

EXPLANATION: This emergency order closes all fresh waters of the Shaktoolik and Unalakleet River drainages to subsistence salmon fishing effective 8:00 a.m. Monday, June 17, and immediately reopens subsistence fishing in the Shaktoolik and Unalakleet River drainages to subsistence fishing with set gillnets with a mesh size not greater than 4 ½ inches. Additionally, this emergency order establishes a subsistence fishing schedule for the Unalakleet River drainage effective 8:00 a.m. Monday, June 17, of two 36-hour periods per week.

---

-continued-



Effective 8:00 a.m. Monday, June 17, subsistence salmon fishing will reopen in the Shaktoolik River drainage but will be limited to set gillnets with a mesh size no greater than 4 ½ inches.

Also effective 8:00 a.m. Monday, June 17, the Unalakleet River drainage will reopen to subsistence salmon fishing on a schedule of two 36-hour periods per week. Subsistence fishing periods in the Unalakleet River drainage will be from 8:00 a.m. Mondays to 8:00 p.m. Tuesdays and from 8:00 a.m. Fridays to 8:00 p.m. Saturdays.

**JUSTIFICATION:** Chinook salmon runs to southeastern Norton Sound are expected to be very poor this season. The combined Subdistricts 5 and 6 Chinook salmon run could be as low as 4,500 fish this season, and this level of abundance is not sufficient to justify sport fishery harvests or support historical Chinook salmon harvest rates. These preemptive restrictions are being implemented this season to significantly reduce harvest rates of Chinook salmon and ensure that escapement goals are achieved in the Shaktoolik and Unalakleet River drainages. In addition to restrictions, subsistence users are also strongly encouraged to redirect harvest pressure on other more numerous species this season, such as chum salmon which are anticipated to be abundant in southern Norton Sound. Once the Chinook salmon run begins, inseason management will be based upon escapement counts and subsistence fishing reports. Fishing restrictions may be relaxed or further restrictions may be necessary depending on run assessment.

**Emergency Order:** 3-S-Z-04-13 **Effective Date:** June 25, 2013

**EXPLANATION:** This emergency order opens Subdistrict 4 of the Norton Sound Subdistrict, the Norton Bay Subdistrict, to two 48-hour periods from 6:00 p.m. Tuesday, June 25 to 6:00 p.m. Thursday, June 27 and from 6:00 p.m. Friday, June 28 to 6:00 p.m. Sunday, June 30. Permit holders in Subdistrict 4 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches.

**JUSTIFICATION:** An average to above average run of chum salmon is anticipated this season in Norton Sound Subdistrict 4, the Norton Bay Subdistrict. Escapement and subsistence harvest needs are expected to easily be met and additional surplus should be available for commercial harvest. Norton Sound Seafood Products has expressed interest in purchasing salmon in the Norton Bay Subdistrict. The department will allow two 48-hour commercial openings to gauge early chum salmon run strength and provide commercial harvest opportunity.

**Emergency Order:** 3-S-Z-05-13 **Effective Date:** July 1, 2013

**EXPLANATION:** Effective 12:00 a.m. Monday, July 1, the marine waters of Subdistricts 5 and 6, and all freshwaters of the Shaktoolik and Unalakleet River drainages will open to subsistence salmon fishing with beach seines 24 hours a day, seven days a week until Saturday, August 10. This emergency order also prohibits the retention of any king salmon incidentally captured in beach seines while targeting other salmon for subsistence uses. All king salmon incidentally captured in beach seines must be immediately released back into the water unharmed.

**JUSTIFICATION:** Effective July 1, subsistence salmon fishing in all fresh and marine waters of the Shaktoolik and Unalakleet Subdistricts will open to the use of beach seines 24 hours a day, seven days a week to target salmon other than Chinook salmon. The Alaska Board of Fisheries passed two new regulations in 2013: one requires that beach seines in the Shaktoolik and Unalakleet Subdistricts have a mesh size of 4 ½ inches or less; the other prohibits the retention of any Chinook salmon captured in beach seine gear. Chinook salmon incidentally captured in beach seines must be immediately released unharmed back into the water. Using beach seine gear provides additional opportunity for subsistence users to target abundant pink and chum salmon while ensuring that king salmon can reach spawning areas of the Shaktoolik and Unalakleet River drainages unharmed.

**Emergency Order:** 3-S-Z-06-13 **Effective Date:** July 2, 2013

**EXPLANATION:** This emergency order reopens Subdistrict 4 of the Norton Sound Subdistrict, the Norton Bay Subdistrict, to two 48-hour periods from 6:00 p.m. Tuesday, July 2 to 6:00 p.m. Thursday, July 4 and from 6:00 p.m. Friday, July 5 to 6:00 p.m. Sunday, July 7. Permit holders in Subdistrict 4 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches.

---

-continued-

**JUSTIFICATION:** The commercial catch of chum salmon during the most recent period in Norton Bay Subdistrict was 4,323 chum salmon by 8 permit holders. Harvest and catch per unit of effort were the most for any period in the history of the Norton Bay commercial salmon fishery. Thus far, 8 permit holders have harvested nearly 5,700 chum salmon in the fishery, which is also record setting for late June. Inglutalik River tower personnel have enumerated over 600 chum salmon as of June 28, which is similar to previous years when 30,000–65,000 chum salmon were counted for the season. Inriver abundance of Norton Bay salmon producing drainages as indexed by the Inglutalik River tower appears more than sufficient to provide for escapements and subsistence use by Koyuk residents.

**Emergency Order:** 3-S-Z-07-13    **Effective Date:** July 2, 2013

**EXPLANATION:** This emergency order opens Subdistrict 5 of the Norton Sound Subdistrict, the Shaktoolik Subdistrict, to two 24-hour periods from 6:00 p.m. Tuesday, July 2 to 6:00 p.m. Wednesday, July 3, and from 6:00 p.m. Friday, July 5 to 6:00 p.m. Saturday, July 6. Permit holders in Subdistricts 5 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches. This emergency order also prohibits the sale of king salmon incidentally harvested during these directed chum salmon openings.

**JUSTIFICATION:** This index opening will provide opportunity to gauge early run strength of chum salmon to Subdistrict 5 and utilize projected commercial harvest surpluses of chum salmon. Shaktoolik sonar estimated passage of chum salmon is over 7,600 fish, which indicates that inriver abundance of chum salmon will be sufficient to provide for subsistence uses and escapement needs in the Shaktoolik Subdistrict. Thus, these periods should not jeopardize escapement needs or subsistence uses of chum salmon in Subdistrict 5.

Although chum salmon abundance is sufficient to warrant additional fishing time in Subdistricts 5, periods will be brief for early July to minimize incidental harvest of king salmon. The department will evaluate escapement counts of king salmon this week to determine if additional chum salmon commercial harvest opportunity can be provided. This brief period will also provide an opportunity to evaluate incidental harvest of king salmon in commercial 6-inch mesh set net gear, which will factor in determining the location and extent of future openings directed on chum salmon in Subdistrict 5.

Per the Subdistricts 5 and 6 King Salmon Management Plan, permit holders may retain but cannot sell king salmon incidentally harvested in the commercial salmon fishery. King salmon retained for subsistence purposes must be recorded in the personal use section of each fish ticket at the time of delivery.

**Emergency Order:** 3-S-Z-08-13    **Effective Date:** July 1, 2013

**EXPLANATION:** This emergency order opens Norton Sound Subdistrict 6, the Unalakleet Subdistrict to commercial salmon fishing for two 24-hour periods from 6:00 p.m. Monday, July 1 to 6:00 p.m. Tuesday, July 2 and from 6:00 p.m. Friday, July 5 to 6:00 p.m. Saturday, July 6. Permit holders in Subdistricts 6 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches. This emergency order also prohibits the sale of king salmon incidentally harvested during these directed chum salmon openings.

**JUSTIFICATION:** This index opening will provide opportunity to gauge early run strength of chum salmon to Subdistrict 6 and utilize projected commercial harvest surpluses of chum salmon. Chum salmon counts at the Unalakleet River weir are 3,200 chum salmon and the Unalakleet River aerial survey goal of 2,400–4,800 chum salmon is projected to easily be reached. This indicates that inriver abundance of chum salmon will be sufficient to provide for subsistence uses and escapement needs in the Unalakleet Subdistrict. Thus, these periods should not jeopardize escapement needs or subsistence uses of chum salmon in Subdistrict 6.

Although chum salmon abundance is sufficient to warrant additional fishing time in Subdistrict 6, periods will be brief for early July to minimize incidental harvest of king salmon. The department will evaluate escapement counts of king salmon this week to determine if additional chum salmon commercial harvest opportunity can be provided. This brief period will also provide an opportunity to evaluate incidental harvest of king salmon in commercial 6-inch mesh set net gear, which will factor in determining the location and extent of future openings directed on chum salmon in Subdistrict 6.

---

-continued-

Per the Subdistricts 5 and 6 King Salmon Management Plan, permit holders may retain but cannot sell king salmon incidentally harvested in the commercial salmon fishery. King salmon retained for subsistence purposes must be recorded in the personal use section of each fish ticket at the time of delivery.

Emergency Order: 3-S-Z-09-13   Effective Date: July 8, 2013

EXPLANATION: This emergency order closes all marine waters in Norton Sound Subdistrict 6, the Unalakleet Subdistrict, to subsistence salmon fishing effective 6:00 p.m. Monday, July 8 and immediately reopens Subdistrict 6 to a subsistence fishing schedule with fishermen restricted to set gillnets with a mesh size no larger than 6 inches. The schedule will be from 6:00 p.m. Monday, July 8 to 6:00 p.m. Tuesday, July 9, and from 6:00 p.m. Friday, July 12 to 6:00 p.m. Thursday, July 13. After July 13, the Subdistrict 6 marine subsistence fishery will remain restricted to 6 inches or less until midnight Sunday evening, July 21.

JUSTIFICATION: As expected, the Chinook salmon run to the Unalakleet River drainage is showing very late timing and poor initial run strength. As of July 1, a total of 31 Chinook salmon have been enumerated at the mainstem Unalakleet River weir. North River tower has not been operational since July 1 due to high water levels. However, the Unalakleet River mainstem comprises 45–60% of the drainagewide escapement, which suggests the projected North River Chinook salmon escapement is likely to fall short of the tower-based escapement goal range of 1,200–2,600 fish. Therefore, additional conservation measures are necessary if there is to be any chance of meeting Chinook salmon escapement needs in the Unalakleet River drainage.

Emergency Order: 3-S-Z-10-13   Effective Date: July 6, 2013

EXPLANATION: This emergency order closes all fresh waters of the Unalakleet River drainage to subsistence salmon fishing with set gillnets effective 8:00 p.m. Monday, July 6 until 12:00 a.m. Monday, July 22. Additionally, this emergency order rescinds the subsistence fishing schedule for the Unalakleet River drainage effective 8:00 p.m. Saturday, July 6 until further notice. Subsistence fishermen may continue to use beach seine gear 24 hours a day, 7 days a week to target other salmon, but king salmon must be immediately released unharmed back into the water. Beach seines must have a mesh size of 4 ½ inches or less.

JUSTIFICATION: As expected, the Chinook salmon run to the Unalakleet River drainage is showing very late timing and poor initial run strength. As of July 2, a total of 31 Chinook salmon have been enumerated at the mainstem Unalakleet River weir. North River tower has not been operational since July 1 due to high water levels. However, the Unalakleet River mainstem comprises 45–60% of the drainagewide escapement, which suggests the projected North River Chinook salmon escapement is likely to fall short of the tower-based escapement goal range of 1,200–2,600 fish. Therefore, additional conservation measures are necessary if there is to be any chance of meeting Chinook salmon escapement needs in the Unalakleet River drainage.

Emergency Order: 3-S-Z-11-13   Effective Date: July 9, 2013

EXPLANATION: This emergency order reopens Subdistrict 4 of the Norton Sound Subdistrict, the Norton Bay Subdistrict, for two 48-hour periods from 6:00 p.m. Tuesday, July 9 to 6:00 p.m. Thursday, July 11 and from 6:00 p.m. Friday, July 12 to 6:00 p.m. Sunday, July 14. Permit holders in Subdistrict 4 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches.

JUSTIFICATION: The commercial catch of chum salmon during the most recent period in Norton Bay Subdistrict was 5,700 chum salmon by 8 permit holders. This was the second period in a row with a record high harvest for a single period in the Norton Bay Subdistrict commercial fishery. The total harvest to date of 16,595 chum salmon is now on track to exceed the record harvest of 21,973 chum salmon caught during the 1978 season. Inglutalik River tower personnel have enumerated over 5,600 chum salmon as of July 7, which is more than the July 7 cumulative count of 4,521 in 2012; 32,832 chum salmon were counted during the 2012 season. Current levels of inriver abundance of chum salmon for Norton Bay drainages as indexed by the Inglutalik River tower appear more than sufficient to provide for escapement and subsistence uses by Koyuk residents.

---

-continued-

Emergency Order: 3-S-Z-12-13    Effective Date: July 8, 2013

EXPLANATION: This emergency order reopens Subdistrict 5 of the Norton Sound Subdistrict, the Shaktoolik Subdistrict, for one 48-hour period from 6:00 p.m. Monday, July 8 to 6:00 p.m. Wednesday, July 10, and one 24-hour period from 6:00 p.m. Friday, July 12 to 6:00 p.m. Saturday, July 13. Permit holders in Subdistricts 5 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches. This emergency order also prohibits the sale of king salmon incidentally harvested during these directed chum salmon openings.

JUSTIFICATION: Chum salmon abundance in the Shaktoolik River drainage is more than sufficient to warrant additional fishing time in Subdistricts 5. Periods have been brief thus far in order to minimize incidental harvest of king salmon. Permit holders were largely stuck on the beach due to hazardous surf conditions arising from southerly winds. The first period is 48 hours in duration in order to mitigate foregone chum salmon harvest opportunities that were lost last week. Historical run timing information suggests that the bulk of the Shaktoolik River king salmon run is in the lower reaches of the Shaktoolik River drainage. Therefore, incidental catches of king salmon for these periods are expected to be very minimal.

Per the Subdistricts 5 and 6 King Salmon Management Plan, permit holders may retain but cannot sell king salmon incidentally harvested in the commercial salmon fishery. King salmon retained for subsistence purposes must be recorded in the personal use section of each fish ticket at the time of delivery.

Emergency Order: 3-S-Z-13-13    Effective Date: July 8, 2013

EXPLANATION: This emergency order reopens Norton Sound Subdistrict 6, the Unalakleet Subdistrict to commercial salmon fishing for two 24-hour periods from 6:00 p.m. Monday, July 8 to 6:00 p.m. Tuesday, July 9 and from 6:00 p.m. Friday, July 12 to 6:00 p.m. Saturday, July 13. Permit holders in Subdistricts 6 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches. This emergency order also prohibits the sale of king salmon incidentally harvested during these directed chum salmon openings.

JUSTIFICATION: This index opening will provide opportunity to utilize commercial harvest surpluses of chum salmon. Cumulative chum salmon passage at the Unalakleet River weir now stands at 21,208 chum salmon as of July 7. This is the highest count of chum salmon for July 7 in the project's 4-year history and the Unalakleet River aerial survey goal of 2,400–4,800 chum salmon has been exceeded. Inriver abundance of chum salmon will be sufficient to provide for subsistence uses and escapement needs in the Unalakleet Subdistrict.

Chum salmon abundance is more than sufficient to warrant longer commercial periods directed on chum salmon in Subdistrict 6. However, more aggressive fishing periods will not be scheduled until Monday, July 15 in order to allow the remainder of the king salmon run to enter the Unalakleet River drainage unharmed. Beginning Monday, July 15, 48-hour periods are expected after the bulk of the king salmon run has entered the lower Unalakleet River drainage, with extended (72-hour) periods possible to more efficiently harvest chum salmon surpluses.

Per the Subdistricts 5 and 6 King Salmon Management Plan, permit holders may retain but cannot sell king salmon incidentally harvested in the commercial salmon fishery. King salmon retained for subsistence purposes must be recorded in the personal use section of each fish ticket at the time of delivery.

Emergency Order: 3-S-Z-14-13    Effective Date: July 12, 2013

EXPLANATION: This emergency order supersedes emergency orders 3-S-Z-02-13 and 3-S-Z-09-1,3 and closes all marine waters of Norton Sound Subdistricts 5 and 6, the Shaktoolik and Unalakleet Subdistricts, to subsistence salmon fishing effective 6:00 p.m. Friday, July 12. This emergency order rescinds subsistence fishing schedules of two 48-hour openings per week in the Shaktoolik Subdistrict, and two 24-hour periods per week in the Unalakleet Subdistrict. This emergency order also immediately reopens subsistence salmon fishing in the marine waters of both Subdistricts 5 and 6 for 24 hours a day, 7 days a week, but fishing is restricted to set gillnets with a mesh size of 6 inches or less.

---

-continued-

**JUSTIFICATION:** Westerly winds and high water levels hastened the entry of king salmon into the lower reaches of the Shaktoolik and Unalakleet River drainages as indicated by a major decline in incidental catches of king salmon in the commercial chum salmon fishery this past week. All available information suggests that the bulk of the king salmon run is now protected within the lower reaches of the Shaktoolik and Unalakleet River drainages. King salmon runs to southern Norton Sound have been very weak and have exhibited later than average run timing. However, closures and restrictions are in place to conserve these king salmon so they may reach known spawning reaches and provide for future returns. Consequently, the department is increasing fishing time in the marine waters to provide additional subsistence opportunity for users to target chum salmon which are in abundance this season.

**Emergency Order:** 3-S-Z-15-13   **Effective Date:** July 11, 2013

**EXPLANATION:** This emergency order supersedes emergency order 3-S-Z-13-13 and extends the July 12 24-hour opening in Norton Sound Subdistrict 6, the Unalakleet Subdistrict, by an additional 24 hours, effectively creating a 48-hour period. This emergency order permits commercial salmon fishing for 48 hours from 6:00 p.m. Thursday, July 11 to 6:00 p.m. Saturday, July 13. Permit holders in Subdistricts 6 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches. This emergency order also prohibits the sale of king salmon incidentally harvested during these directed chum salmon openings.

**JUSTIFICATION:** Chum salmon abundance is more than sufficient to warrant longer commercial periods directed on chum salmon in Subdistrict 6. Additional opportunity to target chum salmon for commercial purposes is being provided because recent incidental harvests of king salmon have been very low. Inclement weather from westerly winds kept many fishermen on the beach but also appears to have had the dual effect of pushing the remainder of the king salmon run into the lower reach of the Unalakleet River unharmed.

Per the Subdistricts 5 and 6 King Salmon Management Plan, permit holders may retain but cannot sell king salmon incidentally harvested in the commercial salmon fishery. King salmon retained for subsistence purposes must be recorded in the personal use section of each fish ticket at the time of delivery.

**Emergency Order:** 3-S-Z-16-13   **Effective Date:** July 12, 2013

**EXPLANATION:** This emergency order supersedes emergency order 3-S-Z-12-13 and extends fishing time in the Norton Sound Subdistrict 5, the Shaktoolik Subdistrict, by 24 hours, thereby creating one 48-hour commercial salmon fishing period from 6:00 p.m. Friday, July 12 to 6:00 p.m. Sunday, July 14. Permit holders in Subdistricts 5 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches. This emergency order also prohibits the sale of king salmon incidentally harvested during these directed chum salmon openings.

**JUSTIFICATION:** Chum salmon abundance in the Shaktoolik River drainage is more than sufficient to warrant additional fishing time in Subdistricts 5. Apportioned sonar count estimated passage for the Shaktoolik River is 35,510 chum salmon which is tracking similar to the 39,000 chum salmon observed at the Unalakleet River weir. Additional opportunity to target chum salmon for commercial purposes is being provided because recent incidental harvests of king salmon have been very low. Inclement weather from westerly winds kept many fishermen on the beach in the Shaktoolik Subdistrict, but also appears to have had the dual effect of pushing the remainder of the king salmon run into the lower reach of the Shaktoolik River unharmed. Estimated apportioned Shaktoolik sonar passage as of July 10 is between 1,100–2,200 king salmon.

Per the Subdistricts 5 and 6 King Salmon Management Plan, permit holders may retain but cannot sell king salmon incidentally harvested in the commercial salmon fishery. King salmon retained for subsistence purposes must be recorded in the personal use section of each fish ticket at the time of delivery.

**Emergency Order:** 3-S-Z-17-13   **Effective Date:** July 12, 2013

**EXPLANATION:** This emergency order supersedes emergency order 3-S-Z-11-13 and extends fishing time in Norton Sound Subdistrict 4, the Norton Bay Subdistrict, by an additional 24 hours. This emergency order therefore,

---

-continued-

reopens commercial salmon fishing in Subdistrict 4 for 72 hours from 6:00 p.m. Friday, July 12 to 6:00 p.m. Monday, July 15. Permit holders in Subdistrict 4 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches.

**JUSTIFICATION:** The commercial catch of chum salmon during the most recent period in Norton Bay Subdistrict was 3,600 chum salmon by 8 permit holders. Harvests of chum salmon per period continue to be well above average for mid-July. Total harvest of chum salmon in Norton Bay Subdistrict is expected to easily surpass the record harvest of 21,973 chum salmon set in 1978 during this period. Thus far, a total of 20,254 chum salmon have been harvested in the commercial fishery. Inglutalik River tower personnel have enumerated a minimum count of over 6,500 chum salmon as of July 11. High water levels and poor water clarity have made counting operations difficult since July 7 and chum salmon passage is most likely underestimated. Regardless, current levels of inriver abundance of chum salmon for Norton Bay drainages as indexed by the Inglutalik River tower appear more than sufficient to provide for escapement needs and subsistence uses by Koyuk residents.

**Emergency Order:** 3-S-Z-18-13    **Effective Date:** July 15, 2013

**EXPLANATION:** This emergency order reopens Subdistrict 5 of the Norton Sound Subdistrict, the Shaktoolik Subdistrict, for one 72-hour period from 6:00 p.m. Monday, July 15 to 6:00 p.m. Thursday, July 18. Permit holders in Subdistricts 5 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches. This emergency order also prohibits the sale of king salmon incidentally harvested during these directed chum salmon openings.

**JUSTIFICATION:** Chum salmon abundance in the Shaktoolik River drainage is more than sufficient to warrant additional fishing time in Subdistricts 5. Apportioned sonar count estimated passage for the Shaktoolik River is 35,510 chum salmon which is tracking similar to the 39,000 chum salmon observed at the Unalakleet River weir. Additional opportunity to target chum salmon for commercial purposes is being provided because recent incidental harvests of king salmon have been very low. Inclement weather from westerly winds kept many fishermen on the beach in the Shaktoolik Subdistrict, but also appears to have had the dual effect of pushing the remainder of the king salmon run into the lower reach of the Shaktoolik River unharmed. Estimated apportioned Shaktoolik sonar passage as of July 10 is between 1,100–2,200 king salmon. This 72-hour period will provide additional opportunity to efficiently harvest commercial chum salmon harvestable surpluses during the second peak of the southern Norton Sound chum salmon run.

Per the Subdistricts 5 and 6 King Salmon Management Plan, permit holders may retain but cannot sell king salmon incidentally harvested in the commercial salmon fishery. King salmon retained for subsistence purposes must be recorded in the personal use section of each fish ticket at the time of delivery.

**Emergency Order:** 3-S-Z-19-13    **Effective Date:** July 15, 2013

**EXPLANATION:** This emergency order reopens Norton Sound Subdistrict 6, the Unalakleet Subdistrict, for one 72-hour period from 6:00 p.m. Monday, July 15 to 6:00 p.m. Thursday, July 18. Permit holders in Subdistricts 6 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches. This emergency order also prohibits the sale of king salmon incidentally harvested during these directed chum salmon openings.

**JUSTIFICATION:** Cumulative chum salmon passage at the Unalakleet River weir now stands at over 39,000 chum salmon as of July 11, which is the highest count of chum salmon for July 11 in the project's 4-year history. Escapement needs of chum salmon have been achieved and inriver abundance of chum salmon is more than sufficient to provide for subsistence uses of Unalakleet Subdistrict residents.

Given the record abundance of chum salmon, longer commercial periods directed on chum salmon are warranted in Subdistrict 6. This additional opportunity to target chum salmon for commercial purposes is being provided because recent incidental harvests of king salmon have been very low. Inclement weather from westerly winds kept many fishermen on the beach but also appears to have had the dual effect of pushing the remainder of the king salmon run into the lower reach of the Unalakleet River unharmed. Weir counts of king salmon have picked up in recent days

---

-continued-

with a total of 313 king salmon enumerated as of July 11.

Per the Subdistricts 5 and 6 King Salmon Management Plan, permit holders may retain but cannot sell king salmon incidentally harvested in the commercial salmon fishery. King salmon retained for subsistence purposes must be recorded in the personal use section of each fish ticket at the time of delivery.

Emergency Order: 3-S-Z-20-13    Effective Date: July 16, 2013

EXPLANATION: This emergency order reopens Subdistrict 4 of the Norton Sound Subdistrict, the Norton Bay Subdistrict, for one 72-hour period from 6:00 p.m. Tuesday, July 16 to 6:00 p.m. Friday, July 19. Permit holders in Subdistrict 4 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches.

JUSTIFICATION: Norton Bay fishermen have already harvested nearly 24,000 chum salmon this season, which is above the previous record harvest of 21,973 taken back in 1978. Escapement of chum salmon into Norton Bay salmon-producing drainages has also been strong as indexed by the NSEDC-operated Inglutalik River tower. Cumulative passage of chum salmon at Inglutalik River tower is over 30,000 fish, which is tracking similar to the 2011 season when over 64,000 chum salmon were counted. Current levels of inriver chum salmon abundance are more than sufficient to meet subsistence needs and provide escapements to ensure future returns of chum salmon. The increase in commercial fishing time is warranted because large harvestable surpluses of chum salmon remain in Norton Bay and there continues to be buyer interest.

Emergency Order: 3-S-Z-21-13    Effective Date: July 15, 2013

EXPLANATION: This emergency order adds 48-hours to the subsistence salmon gillnet fishing schedule for Subdistrict 1 west of Cape Nome. The subsistence salmon gillnet schedule will change from 6 p.m. Wednesday to 6 p.m. Saturday to the expanded schedule from 6 p.m. Monday until 6 p.m. Saturday in Subdistrict 1 marine waters west of Cape Nome.

JUSTIFICATION: The Nome Subdistrict escapement range goal of 23,000-35,000 chum salmon will easily be exceeded this year. At the Eldorado River, escapement is 11,000 chum salmon and that exceeds the escapement goal range of 6,000 to 9,200 chum salmon. Historically, mid-July is average midpoint of the chum salmon run at Eldorado River weir. At Snake River, 800 chum salmon have passed the weir (escapement goal range is 1,600 to 2,500 chum salmon). At Nome River, 1,150 chum salmon have passed the weir (escapement goal range is 2,900 to 4,300 chum salmon). Mid-July is average first-quarter point of the chum salmon run at both Snake and Nome weirs. Nome and Snake rivers are the index rivers for escapement west of Cape Nome and with both projected to reach escapement goal ranges the department is expanding to a 5-day a week subsistence set gillnet fishing schedule west of Cape Nome.

Emergency Order: 3-S-Z-22-13    Effective Date: July 18, 2013

EXPLANATION: This emergency order opens the marine waters from the Cape Nome jetty eastward to Topkok Head in Subdistrict 1 of the Norton Sound Subdistrict, the Nome Subdistrict, for one 24-hour period from 6:00 p.m. Thursday, July 18 to 6:00 p.m. Friday, July 19. Permit holders in Subdistrict 1 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches.

JUSTIFICATION: The Nome Subdistrict commercial chum salmon fishery was closed in 1991 due to chronically low runs and difficulty meeting escapement needs. Since the mid-2000s, runs of chum salmon to the Nome Subdistrict have been sufficient to meet escapement goals in most years and provide surpluses for subsistence harvest needs. As a result, the State of Alaska Board of Fisheries passed regulations in January, 2013 allowing for a commercial chum salmon fishery based on conservative management guidelines. Additionally, the board adopted regulations that divided Nome Subdistrict into two management areas: one from Cape Nome east to Topkok Head, and another from Cape Nome west to Cape Rodney. This regulation was adopted to provide the department with

---

-continued-

guidance to manage subsistence and commercial salmon fisheries east or west of Cape Nome based on escapement information from drainages located east or west of Cape Nome.

The chum salmon escapement goal range has already been achieved at Eldorado River, the index river for chum salmon escapement east of Cape Nome, and a limited commercial chum salmon fishery should not jeopardize subsistence uses or escapement needs. Projections of chum salmon escapement based on historical data for western Nome Subdistrict drainages will most likely approach or exceed the upper bound of established escapement goal ranges. As of July 16, cumulative escapement counts of chum salmon at the Eldorado, Nome, and Snake Rivers are 12,788, 1,263, and 872 chum salmon, respectively. Harvestable surpluses of chum salmon are available east of Cape Nome and there is buyer interest to purchase these fish.

Emergency Order: 3-S-Z-23-13    Effective Date: July 17, 2013

EXPLANATION: This emergency order opens Subdistrict 2, the Golovnin Bay Subdistrict of the Norton Sound District, to commercial salmon fishing for one 48-hour period directed at pink salmon from 6:00 p.m. Wednesday, July 17 to 6:00 p.m. Friday, July 19. Permit holders are restricted to gillnets with a mesh size of four and one-half inches or less.

JUSTIFICATION: Golovin Subdistrict salmon fisheries are being managed this season based on escapement data from the Kwiniuk River tower, located in Subdistrict 3, the nearby Elim Subdistrict. The Niukluk River tower project is no longer operational to assess escapements in season in the Golovin Subdistrict. However, the Kwiniuk River tower-based escapement goal threshold of 8,400 pink salmon is projected to be reached and there is buyer interest in pink salmon. Additionally, the former Niukluk River escapement goal threshold of 10,500 pink salmon has been achieved continually since 1998, and there are no concerns meeting subsistence and escapement needs this season.

There were no directed chum salmon openings this season in Subdistrict 2 as a result of a weak chum salmon run. The Subdistricts 2 and 3 management plan does not allow for commercial pink salmon fishing to occur before July 14 in the Golovin Subdistrict in years of low chum salmon abundance. However, the bulk of the chum salmon run is believed to be in river by this time and pink salmon commercial fishing should not have a negative impact on chum salmon escapements in the Fish River drainage. Pink salmon run strength will be indexed using comparative catch statistics from this opening and Kwiniuk River tower counts in order to determine if more commercial fishing opportunity can be provided while not jeopardizing subsistence uses of pink salmon.

Emergency Order: 3-S-Z-24-13    Effective Date: July 19, 2013

EXPLANATION: This emergency order reopens Subdistrict 5 of the Norton Sound Subdistrict, the Shaktoolik Subdistrict, for one 72-hour period from 6:00 p.m. Friday, July 19 to 6:00 p.m. Monday, July 22. Permit holders in Subdistricts 5 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches. This emergency order also prohibits the sale of king salmon incidentally harvested during these directed chum salmon openings.

JUSTIFICATION: Chum salmon abundance in the Shaktoolik River drainage is more than sufficient to warrant additional fishing time in Subdistricts 5. Apportioned sonar count estimated passage for the Shaktoolik River is 36,000 chum salmon. Additional opportunity to target chum salmon for commercial purposes is warranted due to the large chum salmon surplus.

Per the Subdistricts 5 and 6 King Salmon Management Plan, permit holders may retain but cannot sell king salmon incidentally harvested in the commercial salmon fishery. King salmon retained for subsistence purposes must be recorded in the personal use section of each fish ticket at the time of delivery.

Emergency Order: 3-S-Z-25-13    Effective Date: July 19, 2013

EXPLANATION: This emergency order reopens Norton Sound Subdistrict 6, the Unalakleet Subdistrict, for one 72-hour period from 6:00 p.m. Friday, July 19 to 6:00 p.m. Monday, July 22. Permit holders in Subdistricts 6 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6

---

-continued-



inches. This emergency order also prohibits the sale of king salmon incidentally harvested during these directed chum salmon openings.

**JUSTIFICATION:** Cumulative chum salmon passage at the Unalakleet River weir now stands at over 63,000 chum salmon as of July 18. Escapement needs of chum salmon have been achieved and inriver abundance of chum salmon is more than sufficient to provide for subsistence uses of Unalakleet Subdistrict residents.

Given the near record abundance of chum salmon, longer commercial periods directed on chum salmon are warranted in Subdistrict 6. This additional opportunity to target chum salmon for commercial purposes is being provided because recent incidental harvests of king salmon have been very low.

Per the Subdistricts 5 and 6 King Salmon Management Plan, permit holders may retain but cannot sell king salmon incidentally harvested in the commercial salmon fishery. King salmon retained for subsistence purposes must be recorded in the personal use section of each fish ticket at the time of delivery.

**Emergency Order:** 3-S-Z-26-13    **Effective Date:** July 20, 2013

**EXPLANATION:** This emergency order reopens Subdistrict 4 of the Norton Sound Subdistrict, the Norton Bay Subdistrict, for one 72-hour period from 6:00 p.m. Saturday, July 20 to 6:00 p.m. Tuesday, July 23. Permit holders in Subdistrict 4 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches.

**JUSTIFICATION:** Norton Bay fishermen have already harvested over 27,000 chum salmon this season, which is above the previous record harvest of 21,973 taken back in 1978. Escapement of chum salmon into Norton Bay salmon-producing drainages has also been strong as indexed by the NSEDC-operated Inglutalik River tower. Cumulative passage of chum salmon at Inglutalik River tower is over 42,000 fish, which is tracking similar to the 2011 season when over 64,000 chum salmon were counted. Current levels of inriver chum salmon abundance are more than sufficient to meet subsistence needs and provide escapements to ensure future returns of chum salmon. The increase in commercial fishing time is warranted because large harvestable surpluses of chum salmon remain in Norton Bay and there continues to be buyer interest.

**Emergency Order:** 3-S-Z-27-13    **Effective Date:** July 20, 2013

**EXPLANATION:** This emergency order reopens the marine waters from the Cape Nome jetty eastward to Topkok Head in Subdistrict 1 of the Norton Sound Subdistrict, the Nome Subdistrict, for one 24-hour period from 12:00 p.m. Saturday, July 20 to 12:00 p.m. Sunday, July 21. Permit holders in Subdistrict 1 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches.

**JUSTIFICATION:** The Nome Subdistrict commercial chum salmon fishery was closed in 1991 due to chronically low runs and difficulty meeting escapement needs. Since the mid-2000s, runs of chum salmon to the Nome Subdistrict have been sufficient to meet escapement goals in most years and provide surpluses for subsistence harvest needs. As a result, the State of Alaska Board of Fisheries passed regulations in January, 2013 allowing for a commercial chum salmon fishery based on conservative management guidelines. Additionally, the board adopted regulations that divided Nome Subdistrict into two management areas: one from Cape Nome east to Topkok Head, and another from Cape Nome west to Cape Rodney. This regulation was adopted to provide the department with guidance to manage subsistence and commercial salmon fisheries east or west of Cape Nome based on escapement information from drainages located east or west of Cape Nome.

Escapement needs of chum salmon in drainages east of Cape Nome have already been met. Eldorado River weir chum salmon passage is nearly 16,000 fish, which is well above the escapement goal range of 6,000–9,200 chum salmon. Harvestable surpluses of chum salmon are available for commercial utilization east of Cape Nome. Normal and late run timing projections also suggest the upper end the Nome and Snake River escapement goal ranges in the western half of the Nome Subdistrict will also be reached. Commercial openings west of Cape Nome are possible next week if escapement counts continue to improve.

---

-continued-

Emergency Order: 3-S-Z-28-13    Effective Date: July 20, 2013

EXPLANATION: This emergency order reopens Subdistrict 2, the Golovnin Bay Subdistrict of the Norton Sound District, to commercial salmon fishing for 48 hours from 6:00 p.m. Saturday, July 20 to 6:00 p.m. Monday, July 22. This period is directed at chum salmon and permit holders will be restricted to gillnets with a mesh size of 6 inches or less.

JUSTIFICATION: Commercial chum salmon fishing was not allowed in Golovin Bay this season until now. This period is being established because department biologists observed an estimated 16,000 chum salmon in the Boston Creek tributary alone, as well as several thousand chum salmon at the confluence of the Niukluk and Fish Rivers during an aerial survey July 18. The chum salmon run to the Fish River appears to be very late this season, but escapement of chum salmon into the Fish River has improved dramatically from counts from a July 9 survey. July 18 survey results indicate chum salmon escapement to the Niukluk River has a good chance of nearing the former tower-based escapement goal threshold of 23,000 chum salmon. Additionally, the bulk of the Niukluk River chum salmon run is now considered to be in river by this date. Thus, Niukluk River chum salmon should comprise a miniscule proportion of the commercial harvest during the upcoming opening. Niukluk River tower is no longer operational, but department will fly a peak spawning ground survey at the end of the month to index overall escapement of chum salmon into the Niukluk River. Incidental coho salmon harvests from this period will also provide an early index of coho salmon run strength to Golovin Subdistrict.

Emergency Order: 3-S-Z-29-13    Effective Date: July 20, 2013

EXPLANATION: This emergency order opens all marine waters from Carson Creek eastward to Iron Creek in Norton Sound Subdistrict 3, the Elim Subdistrict, to commercial salmon fishing for 48 hours from 6:00 p.m. Saturday, July 20 to 6:00 p.m. Monday, July 22.

JUSTIFICATION: Elim Subdistrict has not been opened this season because of conservation concerns with chum salmon based on Kwiniuk River tower escapement falling short of the escapement goal. Interestingly, however, the nearby Tubutulik River escapement goal range is projected to be reached based on an early July survey of 4,600 chum salmon. A commercial fishing period is being allowed for 48 hours, but commercial fishing will be limited to the western half of the Elim Subdistrict from Carson Creek eastward to Iron Creek. This will provide commercial permit holders with some late opportunity to target chum salmon, while minimizing the impact to the Kwiniuk River chum salmon escapement. Spawning ground surveys will also be flown later this month to index this season's Tubutulik River chum salmon escapement. The department will evaluate catch statistics and escapement information following these openings to determine if additional commercial opportunity is warranted.

Emergency Order: 3-S-Z-30-13    Effective Date: July 19, 2013

EXPLANATION: This emergency order raises the sockeye salmon subsistence catch limit to 50 fish at Pilgrim River.

JUSTIFICATION: The Pilgrim River weir count for sockeye salmon is 9,309 fish through July 18. This year's sockeye salmon run is already the best since 2008 and run timing projections give a final count of between 15,000 and 20,000 fish. Raising the limit to 50 sockeye salmon should not jeopardize the escapement goal of 4,000 to 8,000 sockeye salmon observed by aerial survey in Salmon Lake.

Emergency Order: 3-S-Z-31-13    Effective Date: July 22, 2013

EXPLANATION: This emergency order closes all fresh waters of the Shaktoolik and Unalakleet River drainages to subsistence salmon fishing effective 12:01 a.m. Monday, July 22, and immediately reopens subsistence salmon fishing in the Shaktoolik and Unalakleet River drainages 24 hours a day, 7 days a week. However, this emergency limits subsistence salmon fishing to set gillnets with a mesh size not greater than 6 inches.

JUSTIFICATION: Current cumulative escapement counts of Chinook salmon on the North and Unalakleet River

---

-continued-

projects are 380 and 690 Chinook salmon, respectively. Projected escapement at the North River is expected to range between 530–850 Chinook salmon, whereas the Unalakleet River mainstem projected escapement is expected to range from 800 to 1,000 Chinook salmon. The latest assessment indicates that the North River tower-based escapement goal range of 1,200–2,600 Chinook salmon will not be achieved this season. Additionally, cumulative Unalakleet River weir passage to date also suggests that the Unalakleet River aerial survey goal range of 550–1,100 Chinook salmon will also not be reached this season. Normally, subsistence fishing for salmon would be allowed with unrestricted mesh size in freshwater areas at this time of year. However, the exceptionally late, but weak run of Chinook salmon warrants additional restrictions to protect Chinook salmon that will contribute to the spawning escapement and provide future returns.

Emergency Order: 3-S-Z-32-13    Effective Date: July 23, 2013

EXPLANATION: This emergency order reopens the marine waters from the Cape Nome jetty eastward to Topkok Head in Subdistrict 1 of the Norton Sound Subdistrict, the Nome Subdistrict, for one 24-hour period from 12:00 p.m. Tuesday, July 23 to 12:00 p.m. Wednesday, July 24. Permit holders in Subdistrict 1 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches.

JUSTIFICATION: The Nome Subdistrict commercial chum salmon fishery was closed in 1991 due to chronically low runs and difficulty meeting escapement needs. Since the mid-2000s, runs of chum salmon to the Nome Subdistrict have been sufficient to meet escapement goals in most years and provide surpluses for subsistence harvest needs. As a result, the State of Alaska Board of Fisheries passed regulations in January, 2013 allowing for a commercial chum salmon fishery based on conservative management guidelines. Additionally, the board adopted regulations that divided Nome Subdistrict into two management areas: one from Cape Nome east to Topkok Head, and another from Cape Nome west to Cape Rodney. This regulation was adopted to provide the department with guidance to manage subsistence and commercial salmon fisheries east or west of Cape Nome based on escapement information from drainages located east or west of Cape Nome.

The Nome Subdistrict chum salmon escapement goal range of 23,000 to 35,000 fish has been achieved and will easily be exceeded this year. There are also three rivers that have escapement goals. The escapement goal range at Eldorado River is 6,000 to 9,200 chum salmon and the escapement this season is over double the high end of the range. At Snake River the escapement goal range is 1,600 to 2,500 chum salmon and has been met and will likely be exceeded. At Nome River the escapement goal range is 2,900–4,300 chum salmon and the escapement is 1,700 chum salmon through Sunday, July 21. The average midpoint of chum passage at the weir is July 23 so Nome River is projected to meet escapement.

Emergency Order: 3-S-Z-33-13    Effective Date: July 23, 2013

EXPLANATION: This emergency order reopens Subdistrict 5 of the Norton Sound Subdistrict, the Shaktoolik Subdistrict, for one 72-hour period from 12:00 p.m. Tuesday, July 23 to 12:00 p.m. Friday, July 26. Permit holders in Subdistricts 5 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches. This emergency order also prohibits the sale of king salmon incidentally harvested during these directed chum salmon openings.

JUSTIFICATION: Chum salmon abundance in the Shaktoolik River drainage is more than sufficient to warrant additional fishing time in Subdistricts 5. Apportioned sonar count estimated passage for the Shaktoolik River is 52,000 chum salmon. Additional opportunity to target chum salmon for commercial purposes is warranted due to the large chum salmon surplus.

Per the Subdistricts 5 and 6 King Salmon Management Plan, permit holders may retain but cannot sell king salmon incidentally harvested in the commercial salmon fishery. King salmon retained for subsistence purposes must be recorded in the personal use section of each fish ticket at the time of delivery.

Emergency Order: 3-S-Z-34-13    Effective Date: July 23, 2013

---

-continued-

**EXPLANATION:** This emergency order reopens Norton Sound Subdistrict 6, the Unalakleet Subdistrict, for one 72-hour period from 6:00 p.m. Tuesday, July 23 to 6:00 p.m. Friday, July 26. Permit holders in Subdistricts 6 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches. This emergency order also prohibits the sale of king salmon incidentally harvested during these directed chum salmon openings.

**JUSTIFICATION:** Cumulative chum salmon passage at the Unalakleet River weir now stands at over 76,000 chum salmon as of July 18. Escapement needs of chum salmon have been achieved and inriver abundance of chum salmon is more than sufficient to provide for subsistence uses of Unalakleet Subdistrict residents. Given the above average abundance of chum salmon, longer commercial periods directed on chum salmon are warranted in Subdistrict 6.

Per the Subdistricts 5 and 6 King Salmon Management Plan, permit holders may retain but cannot sell king salmon incidentally harvested in the commercial salmon fishery. King salmon retained for subsistence purposes must be recorded in the personal use section of each fish ticket at the time of delivery.

**Emergency Order:** 3-S-Z-35-13    **Effective Date:** July 23, 2013

**EXPLANATION:** This emergency order reopens Subdistrict 2, the Golovnin Bay Subdistrict of the Norton Sound District, to commercial salmon fishing for 24 hours from 6:00 p.m. Tuesday, July 23 to 6:00 p.m. Wednesday, July 24. This period is directed at chum salmon and permit holders will be restricted to gillnets with a mesh size of 6 inches or less.

**JUSTIFICATION:** Commercial chum salmon fishing was not allowed in Golovin Bay this season until now. This period is being established because department biologists observed an estimated 16,000 chum salmon in the Boston Creek tributary alone, as well as several thousand chum salmon at the confluence of the Niukluk and Fish Rivers during an aerial survey July 18. The chum salmon run to the Fish River appears to be very late this season, but escapement of chum salmon into the Fish River has improved dramatically from counts from a July 9 survey. July 18 survey results indicate chum salmon escapement to the Niukluk River has a good chance of nearing the former tower-based escapement goal threshold of 23,000 chum salmon. Additionally, the bulk of the Niukluk River chum salmon run is now considered to be in river by this date. Thus, Niukluk River chum salmon should comprise a miniscule proportion of the commercial harvest during the upcoming opening. Niukluk River tower is no longer operational, but department will fly a peak spawning ground survey at the end of the month to index overall escapement of chum salmon into the Niukluk River. Incidental coho salmon harvests from this period will also provide an early index of coho salmon run strength to Golovin Subdistrict.

**Emergency Order:** 3-S-Z-36-13    **Effective Date:** July 24, 2013

**EXPLANATION:** This emergency order opens all marine waters from Carson Creek eastward to Iron Creek in Norton Sound Subdistrict 3, the Elim Subdistrict, to commercial salmon fishing for 24 hours from 6:00 p.m. Wednesday July 24 to 6:00 p.m. Thursday July 25.

**JUSTIFICATION:** Elim Subdistrict has not been opened this season because of conservation concerns with chum salmon based on Kwiniuk River tower escapement falling short of the escapement goal. Interestingly, however, the nearby Tubutulik River escapement goal range is projected to be reached based on an early July survey of 4,600 chum salmon. A commercial fishing period is being allowed for 48 hours, but commercial fishing will be limited to the western half of the Elim Subdistrict from Carson Creek eastward to Iron Creek. This will provide commercial permit holders with some late opportunity to target chum salmon, while minimizing the impact to the Kwiniuk River chum salmon escapement. Spawning ground surveys will also be flown later this month to index this season's Tubutulik River chum salmon escapement. The department will evaluate catch statistics and escapement information following these openings to determine if additional commercial opportunity is warranted.

**Emergency Order:** 3-S-Z-37-13    **Effective Date:** July 25, 2013

**EXPLANATION:** This emergency order reopens Subdistrict 4 of the Norton Sound Subdistrict, the Norton Bay

---

-continued-

Subdistrict, for one 64-hour period from 8:00 a.m. Thursday, July 25 to 12:00 midnight Saturday evening, July 27. Permit holders in Subdistrict 4 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches.

**JUSTIFICATION:** Norton Bay fishermen have already harvested nearly 30,000 chum salmon this season, which is above the previous record harvest of 21,973 taken back in 1978. Escapement of chum salmon into Norton Bay salmon-producing drainages has also been strong as indexed by the NSEDC-operated Inglutalik River tower. Cumulative passage of chum salmon at Inglutalik River tower is over 42,000 fish, which is more than sufficient to meet subsistence needs and provide escapements to ensure future returns of chum salmon. The increase in commercial fishing time is warranted because large harvestable surpluses of chum salmon remain in Norton Bay and there continues to be buyer interest. The department will switch to coho salmon management once coho salmon catches exceed chum salmon catches.

**Emergency Order:** 3-S-Z-38-13 **Effective Date:** July 26, 2013

**EXPLANATION:** This emergency order reopens the all marine waters in Subdistrict 1 of the Norton Sound Subdistrict, the Nome Subdistrict, for one 24-hour period from 12:00 p.m. Friday, July 26 to 12:00 p.m. Saturday, July 27. Permit holders in Subdistrict 1 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches.

**JUSTIFICATION:** The Nome Subdistrict commercial chum salmon fishery was closed in 1991 due to chronically low runs and difficulty meeting escapement needs. Since the mid-2000s, runs of chum salmon to the Nome Subdistrict have been sufficient to meet escapement goals in most years and provide surpluses for subsistence harvest needs. As a result, the State of Alaska Board of Fisheries passed regulations in January, 2013 allowing for a commercial chum salmon fishery based on conservative management guidelines. Additionally, the board adopted regulations that divided Nome Subdistrict into two management areas: one from Cape Nome east to Topkok Head, and another from Cape Nome west to Cape Rodney. This regulation was adopted to provide the department with guidance to manage subsistence and commercial salmon fisheries east or west of Cape Nome based on escapement information from drainages located east or west of Cape Nome.

The Nome Subdistrict chum salmon escapement goal range of 23,000 to 35,000 fish has been achieved and will easily be exceeded this year. There are also three rivers that have escapement goals. The escapement goal range at Eldorado River is 6,000 to 9,200 chum salmon and the escapement this season is over double the high end of the range. The Snake and Nome River escapement goals have also been achieved and escapements at these rivers will likely approach or exceed the upper bound of their respective escapement goal ranges. This brief commercial period will also not jeopardize subsistence uses or escapement needs of chum salmon in the Nome Subdistrict.

**Emergency Order:** 3-S-Z-39-13 **Effective Date:** July 27, 2013

**EXPLANATION:** This emergency order reopens Subdistrict 5 of the Norton Sound Subdistrict, the Shaktoolik Subdistrict, for one 72-hour period from 6:00 p.m. Saturday, July 27 to 6:00 p.m. Tuesday, July 30. Permit holders in Subdistricts 5 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches. This emergency order also prohibits the sale of king salmon incidentally harvested during these directed chum salmon openings.

**JUSTIFICATION:** Chum salmon abundance in the Shaktoolik River drainage is more than sufficient to warrant additional fishing time in Subdistricts 5. Apportioned sonar count estimated passage for the Shaktoolik River is 52,000 chum salmon. Additional opportunity to target chum salmon for commercial purposes is warranted due to the large chum salmon surplus. The department will transition to coho salmon management next week.

Per the Subdistricts 5 and 6 King Salmon Management Plan, permit holders may retain but cannot sell king salmon incidentally harvested in the commercial salmon fishery. King salmon retained for subsistence purposes must be recorded in the personal use section of each fish ticket at the time of delivery.

**Emergency Order:** 3-S-Z-40-13 **Effective Date:** July 27, 2013

---

-continued-

**EXPLANATION:** This emergency order reopens Norton Sound Subdistrict 6, the Unalakleet Subdistrict, from 6:00 p.m. Saturday, July 27 to 6:00 p.m. Tuesday, July 30. Permit holders in Subdistricts 6 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches. This emergency order also prohibits the sale of king salmon incidentally harvested during these directed chum salmon openings.

**JUSTIFICATION:** Cumulative chum salmon passage at the Unalakleet River weir is nearly 84,000 chum salmon as of July 25. Escapement needs of chum salmon have been achieved and inriver abundance of chum salmon is more than sufficient to provide for subsistence uses of Unalakleet Subdistrict residents. Given the above average abundance of chum salmon, longer commercial periods directed on chum salmon are warranted in Subdistrict 6. However, the department will switch to coho salmon management next week and use shorter index periods of 48 hours in duration in order to assess coho salmon run strength.

Per the Subdistricts 5 and 6 King Salmon Management Plan, permit holders may retain but cannot sell king salmon incidentally harvested in the commercial salmon fishery. King salmon retained for subsistence purposes must be recorded in the personal use section of each fish ticket at the time of delivery.

**Emergency Order:** 3-S-Z-41-13 **Effective Date:** July 28, 2013

**EXPLANATION:** This emergency order reopens Subdistrict 4 of the Norton Sound Subdistrict, the Norton Bay Subdistrict, for 72 hours from 6:00 p.m. Sunday, July 28 to 6:00 p.m. Wednesday, July 31. Permit holders in Subdistrict 4 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches.

**JUSTIFICATION:** Norton Bay fishermen have already harvested 30,000 chum salmon this season, which is well above the previous record harvest of 21,973 taken back in 1978. Escapement of chum salmon into Norton Bay salmon-producing drainages has also been strong as indexed by the NSEDC-operated Inglutalik River tower. Cumulative passage of chum salmon at Inglutalik River tower is over 46,000 fish, which is more than sufficient to meet subsistence needs and provide escapements to ensure future returns of chum salmon. The increase in commercial fishing time is warranted because large harvestable surpluses of chum salmon remain in Norton Bay and there continues to be buyer interest. The department will switch to coho salmon management next week and use historical catch statistic comparisons with current year catch statistics to adjust fishing time in Norton Bay.

**Emergency Order:** 3-S-Z-42-13 **Effective Date:** July 31, 2013

**EXPLANATION:** This emergency order reopens the all marine waters in Subdistrict 1 of the Norton Sound Subdistrict, the Nome Subdistrict, for one 24-hour period from 12:00 p.m. Wednesday, July 31 to 12:00 p.m. Thursday, August 1. Permit holders in Subdistrict 1 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches.

**JUSTIFICATION:** The Nome Subdistrict chum salmon escapement goal range of 23,000 to 35,000 fish has been exceeded this year. There are also three rivers that have escapement goals. The Snake and Nome River escapement goals have also been achieved and escapements at these rivers will likely approach or exceed the upper bound of their respective escapement goal ranges. This brief commercial period will also not jeopardize subsistence uses or escapement needs of chum salmon in the Nome Subdistrict.

**Emergency Order:** 3-S-Z-43-13 **Effective Date:** August 1, 2013

**EXPLANATION:** This emergency order reopens Subdistrict 2, the Golovnin Bay Subdistrict of the Norton Sound District, to commercial salmon fishing for 24 hours from 6:00 p.m. Thursday, August 1 to 6:00 p.m. Friday, August 2. This period is directed at chum salmon and permit holders will be restricted to gillnets with a mesh size of 6 inches or less.

**JUSTIFICATION:** This period is being established to index early run strength of coho salmon. Catch statistics from this period will be compared to historical catch statistics to determine if additional commercial fishing is justified. Additional periods may be allowed if catch statistics indicate that continued commercial fishing will not jeopardize

---

-continued-

subsistence uses or escapement needs of coho salmon in the Fish River drainage and other drainages emptying into Golovin Lagoon. Aerial surveys will be flown on the spawning grounds in late August or early September to index spawning escapement of coho salmon in the Fish River drainage.

Emergency Order: 3-S-Z-44-13   Effective Date: August 1, 2013

EXPLANATION: This emergency order opens all marine waters from Carson Creek eastward to Iron Creek in Norton Sound Subdistrict 3, the Elim Subdistrict, to commercial salmon fishing for 24 hours from 6:00 p.m. Thursday, August 1 to 6:00 p.m. Friday August 2.

JUSTIFICATION: This will provide commercial permit holders with some early opportunity to target coho salmon and provide an early index of coho salmon run strength. Kwiniuk River tower personnel have enumerated 100 coho salmon as of July 28, which is above counts from the 2010–2012 seasons for this date. In all years with counts this high for July 28, escapements of coho salmon have been more than sufficient to meet the aerial survey escapement goal (650–1,300 cohos), provide for inriver subsistence harvests, and allow for commercial harvests. However, it is too early to ascertain if the run is merely early or if the good initial counts indicate good early run strength. Tower counts and catch statistics from this index opening will be evaluated to determine if additional commercial harvest opportunities can be allowed. Spawning ground surveys will also be flown later this month to index this season's Tubutulik River coho salmon escapement.

Emergency Order: 3-S-Z-45-13   Effective Date: July 31, 2013

EXPLANATION: This emergency order reopens Subdistrict 5 and 6 of the Norton Sound Subdistrict, the Shaktoolik and Unalakleet Subdistricts, for one 48-hour period from 6:00 p.m. Wednesday, July 31 to 6:00 p.m. Friday, August 2. Permit holders in Subdistricts 5 and 6 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches. This emergency order also prohibits the sale of king salmon incidentally harvested during this directed coho salmon opening.

JUSTIFICATION: Coho salmon abundance in southern Norton Sound as indexed by the Unalakleet River weir passage and North River tower passage is tracking ahead of counts during the 2011 and 2012 seasons. This initial opportunity to target coho salmon for commercial purposes is warranted to index early coho salmon run strength and recent run performance history suggests that this period will not negatively impact subsistence uses of coho salmon or jeopardize escapement needs of coho salmon in the Shaktoolik and Unalakleet River drainages. North River tower counts and historical radiotelemetry data will be analyzed to estimate inriver abundance of coho salmon to ensure that escapement and inriver subsistence uses of coho salmon will be met before committing to longer periods or a set schedule for the remainder of the season.

Per the Subdistricts 5 and 6 King Salmon Management Plan, permit holders may retain but cannot sell king salmon incidentally harvested in the commercial salmon fishery. King salmon retained for subsistence purposes must be recorded in the personal use section of each fish ticket at the time of delivery.

Emergency Order: 3-S-Z-46-13   Effective Date: August 2, 2013

EXPLANATION: This emergency order reopens Subdistrict 4 of the Norton Sound Subdistrict, the Norton Bay Subdistrict, for 48 hours from 6:00 p.m. Friday, August 2 to 6:00 p.m. Sunday, August 4. Permit holders in Subdistrict 4 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches.

JUSTIFICATION: Coho salmon management begins this week and the department will use historical catch statistic comparisons with current year catch statistics to adjust fishing time in Norton Bay. Spawning ground surveys will be flown in late August to index the coho salmon spawning escapement to the Ungalik and Inglutalik Rivers, the major salmon producing drainages in Norton Bay.

Emergency Order: 3-S-Z-47-13   Effective Date: August 5, 2013

---

-continued-

**EXPLANATION:** This emergency order reopens Subdistrict 2, the Golovnin Bay Subdistrict of the Norton Sound District, to commercial salmon fishing for 36 hours from 12:00 p.m. Monday, August 5 to 12:00 a.m. Wednesday, August 7. This period is directed at coho salmon and permit holders will be restricted to gillnets with a mesh size of 6 inches or less.

**JUSTIFICATION:** This period is being established to provide additional commercial harvest opportunity directed on coho salmon. Catch statistics from this period will be compared to historical catch statistics to determine if additional commercial fishing is justified. The August 1<sup>st</sup> opening catch of 527 coho salmon in Golovin Subdistrict by 7 permit holders was 92% above the long-term average catch of 274 coho salmon for the first week of August. The latest catch statistics indicate that abundance of coho salmon will be sufficient to achieve escapement needs and provide for inriver subsistence uses of coho salmon. Catch statistics will be evaluated from this 36-hour opening to determine if additional fishing or periods of longer duration can be allowed without jeopardizing escapement or subsistence needs.

Additional periods may be allowed if catch statistics indicate that continued commercial fishing will not jeopardize subsistence uses or escapement needs of coho salmon in the Fish River drainage and other drainages emptying into Golovin Lagoon. Aerial surveys will be flown on the spawning grounds in late August or early September to index spawning escapement of coho salmon in the Fish River drainage.

**Emergency Order:** 3-S-Z-48-13 **Effective Date:** August 5, 2013

**EXPLANATION:** This emergency order opens all marine waters from Carson Creek eastward to Iron Creek in Norton Sound Subdistrict 3, the Elim Subdistrict, to commercial salmon fishing for 36 hours from 12:00 p.m. Monday, August 5 to 12:00 a.m. Wednesday August 7.

**JUSTIFICATION:** This brief opening will provide commercial permit holders with harvest opportunity to target coho salmon and provide an additional index of coho salmon run strength. In the Elim Subdistrict, 12 permit holders caught 457 cohos during the most recent 24 hours opening. This catch was slightly above average for the first week of August.

There have been 657 coho salmon enumerated at the Kwiniuk River as of midnight August 3, which is 21% below the 2008–2012 average count of 797 cohos for August 2. However, projected escapement estimates of coho salmon based on this count range from 3,900–6,800 coho salmon for the season for runs with early and normal migration timing, respectively.

Current comparative catch statistics and escapement information indicate that abundance of coho salmon is on track achieve escapement needs and provide for inriver subsistence uses of coho salmon in the Elim Subdistrict. Escapements and catch statistics will be evaluated from this 36-hour opening to determine if additional fishing or periods of longer duration can be allowed without jeopardizing escapement or subsistence needs. Escapement of coho salmon to Subdistrict 3 will be evaluated using tower counts at the Kwiniuk River and the Kwiniuk River aerial survey goal range of 650–1,300 fish. Spawning ground surveys will also be flown later this month to index this season's Tubutulik River coho salmon escapement.

**Emergency Order:** 3-S-Z-49-13 **Effective Date:** August 5, 2013

**EXPLANATION:** This emergency order reopens Subdistrict 4 of the Norton Sound Subdistrict, the Norton Bay Subdistrict, for 48 hours from 6:00 p.m. Monday, August 5 to 6:00 p.m. Wednesday, August 7. Permit holders in Subdistrict 4 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches.

**JUSTIFICATION:** Coho salmon season is underway. Recent catches of coho salmon from the Norton Bay Subdistrict this past week were average to above average for early August. Additionally, 2,700 cohos have been harvested to date in Norton Bay which is on track to establish a new record harvest for Norton Bay Subdistrict.

During a July 30 aerial survey of the Ungalik River, department biologists observed over 1,000 coho salmon in the lower reach of the Ungalik. The good early survey count of coho salmon suggests that inriver abundance of coho salmon will be more than sufficient to provide for future returns and meet the subsistence priority. Peak spawning

---

-continued-



ground surveys will be flown in late August to index the coho salmon spawning escapement to the Ungalik River, the major coho salmon producing drainage in Norton Bay.

Emergency Order: 3-S-Z-50-13    Effective Date: August 4, 2013

EXPLANATION: This emergency order reopens Subdistrict 5 and 6 of the Norton Sound Subdistrict, the Shaktoolik and Unalakleet Subdistricts, to a commercial fishing schedule of two 48-hour periods per week until the end of the Subdistricts 5 and 6 season. Periods will be from 6:00 p.m. Sundays to 6:00 p.m. Tuesdays and from 6:00 p.m. Wednesdays to 6:00 p.m. Fridays. Permit holders in Subdistricts 5 and 6 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches. This emergency order also prohibits the sale of king salmon incidentally harvested during the directed coho salmon fishery.

JUSTIFICATION: Coho salmon abundance in southern Norton Sound as indexed by the North River tower passage is tracking well ahead of counts during the 2011 and 2012 seasons. This schedule will provide fishermen and the buyer with an orderly fishery. The schedule is warranted based on current projections of North River coho salmon escapement, projected Unalakleet River inriver coho salmon abundance, and recent run performance trends. This schedule is not expected to negatively impact subsistence uses of coho salmon or jeopardize escapement needs of coho salmon in the Shaktoolik and Unalakleet River drainages.

Escapement to the North River tributary is at 1,602 coho salmon as of August 3. Drainagewide escapement projections based on the North River tower count and historical radiotelemetry data range from 38,000–54,000 coho salmon for runs with early and normal migration timing, respectively. More importantly, this level of inriver abundance will be more than sufficient to achieve the North River aerial survey goal range of 550–1,100 fish and provide for inriver subsistence and sport fish harvest needs. Shaktoolik and Unalakleet Subdistricts will remain on this schedule for the remainder of the season unless projections decrease dramatically and reductions in fishing time are needed to ensure that subsistence harvest needs will not be jeopardized by commercial fishing.

Per the Subdistricts 5 and 6 King Salmon Management Plan, permit holders may retain but cannot sell king salmon incidentally harvested in the commercial salmon fishery. King salmon retained for subsistence purposes must be recorded in the personal use section of each fish ticket at the time of delivery.

Emergency Order: 3-S-Z-51-13    Effective Date: August 8, 2013

EXPLANATION: This emergency order reopens Subdistrict 2, the Golovnin Bay Subdistrict of the Norton Sound District, to commercial salmon fishing for 36 hours from 12:00 p.m. Thursday, August 8 to 12:00 a.m. Saturday, August 10. This period is directed at coho salmon and permit holders will be restricted to gillnets with a mesh size of 6 inches or less.

JUSTIFICATION: This period is being established to provide additional commercial harvest opportunity directed on coho salmon. The August 5<sup>th</sup> opening catch of 861 coho salmon in Golovin Subdistrict by 8 permit holders was 192% above the long-term average catch of 282 coho salmon for the first week of August. The August 5<sup>th</sup> catch was also the fifth best ever for a single period and the 2013 cumulative harvest of 1,700 cohos lags only behind the record 2010 season's catch for this date.

Additional periods may be allowed if catch statistics indicate that continued commercial fishing will not jeopardize subsistence uses or escapement needs of coho salmon in the Fish River drainage and other drainages emptying into Golovin Lagoon. Aerial surveys will be flown on the spawning grounds in late August or early September to index spawning escapement of coho salmon in the Fish River drainage.

Emergency Order: 3-S-Z-52-13    Effective Date: August 8, 2013

EXPLANATION: This emergency order opens all marine waters of Norton Sound Subdistrict 3, the Elim Subdistrict, to commercial salmon fishing for 36 hours from 12:00 p.m. Thursday, August 8 to 12:00 a.m. Saturday August 10.

JUSTIFICATION: This brief opening will provide commercial permit holders with harvest opportunity to target coho salmon and provide an additional index of coho salmon run strength. In the Elim Subdistrict, 9 permit holders

---

-continued-

caught 511 cohos during the most recent 36-hour opening. This catch was 19% above the average catch of 421 cohos for the first week of August.

There have been nearly 1,200 coho salmon enumerated at the Kwiniuk River as of 8:00 a.m. August 8, which is 18% below the 2008–2012 average count of 1,448 cohos for August 8. However, projected escapement estimates of coho salmon based on this count range from 4,200–5,700 coho salmon for runs with early and normal migration timing, respectively. Current catch statistics and escapement information indicate that abundance of coho salmon continues to build in Subdistrict 3 and will be sufficient to achieve escapement needs and provide for subsistence uses. Escapements and catch statistics will be evaluated from this 36-hour opening to determine if additional fishing or periods of longer duration can be allowed without jeopardizing escapement or subsistence needs. Escapement of coho salmon to Subdistricts 2 and 3 will be evaluated using tower counts at the Kwiniuk River and the Kwiniuk River aerial survey goal range of 650–1,300 fish.

Emergency Order: 3-S-Z-53-13 Effective Date: August 9, 2013

EXPLANATION: This emergency order reopens Subdistrict 4 of the Norton Sound Subdistrict, the Norton Bay Subdistrict, for 48 hours from 6:00 p.m. Friday, August 9 to 6:00 p.m. Sunday, August 11. Permit holders in Subdistrict 4 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches.

JUSTIFICATION: Coho salmon season is underway. Recent catches of coho salmon from the Norton Bay Subdistrict this past week were average to above average for early August. Additionally, nearly 3,500 cohos have been harvested to date in Norton Bay which is on track to establish a new record harvest for Norton Bay Subdistrict this season.

During a July 30 aerial survey of the Ungalik River, department biologists observed over 1,000 coho salmon in the lower reach of the Ungalik. The good early survey count of coho salmon suggests that inriver abundance of coho salmon will be more than sufficient to provide for future returns and meet the subsistence priority. Peak spawning ground surveys will be flown in late August to index the coho salmon spawning escapement to the Ungalik River, the major coho salmon producing drainage in Norton Bay.

Emergency Order: 3-S-Z-54-13 Effective Date: August 11, 2013

EXPLANATION: This emergency order reopens Subdistricts 2 and 3 of the Norton Sound District, to commercial salmon fishing for 36 hours from 12:00 p.m. Sunday, August 11 to 12:00 a.m. Tuesday, August 13. This period is directed at coho salmon and permit holders will be restricted to gillnets with a mesh size of 6 inches or less.

JUSTIFICATION: This period is being established to provide additional commercial harvest opportunity directed on coho salmon. The August 8 opening catch of 1,476 coho salmon in Golovin Subdistrict by 7 permit holders was a record catch for one fishing period. Department staff plan to fly an aerial survey next week of the Niukluk River drainage to estimate early escapement numbers. In Elim Subdistrict the catch was 886 coho salmon by 14 permit holders and was average for this date. Escapement at Kwiniuk River counting tower is 1,254 coho salmon with the average first quarter point of passage at the tower on August 10. Projections show that escapement will be met. Continuing with short duration openings should not jeopardize reaching escapement goals or subsistence fishing opportunity.

Emergency Order: 3-S-Z-55-13 Effective Date: August 12, 2013

EXPLANATION: This emergency order opens Subdistrict 4 of the Norton Sound District to a fishing schedule of two 48-hour commercial fishing periods per week from 6:00 p.m. Monday to 6:00 p.m. Wednesday and from 6 p.m. Thursday until 6 p.m. Saturday through September 7. Permit holders in Subdistrict 4 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches.

JUSTIFICATION: Catches of coho salmon from the Norton Bay Subdistrict during recent 48 hour fishing periods have been average for mid-August. Additionally, nearly 4,200 coho salmon have been harvested to date in Norton

---

-continued-

Bay which is on track to establish a new record harvest for Norton Bay Subdistrict this season.

During a July 30 aerial survey of the Ungalik River, department biologists observed over 1,000 coho salmon in the lower reach of the Ungalik. The tower crew has enumerated over 5,000 coho salmon through August 11, but water has been high and turbid so there may some doubt as to accurate speciation. The good early survey count of coho salmon suggests that inriver abundance of coho salmon will be more than sufficient to provide for future returns and meet the subsistence priority. Peak spawning ground surveys will be flown in late August to index the coho salmon spawning escapement to the Ungalik River, the major coho salmon producing drainage in Norton Bay.

Emergency Order: 3-S-Z-56-13    Effective Date: August 14, 2013

EXPLANATION: This emergency order reopens Subdistricts 2 and 3 of the Norton Sound District, the Golovin and Elim Subdistricts, to commercial salmon fishing for 48 hours from 6:00 p.m. Wednesday, August 14 to 6:00 p.m. Friday, August 16. This period is directed at coho salmon and permit holders will be restricted to gillnets with a mesh size of 6 inches or less.

JUSTIFICATION: This period is being established to maximize commercial harvest opportunity of coho salmon at the historical peak of the coho salmon run. The August 8 opening catch of 1,476 coho salmon in the Golovin Subdistrict was a record catch for one fishing period, and the August 11<sup>th</sup> catch of 940 was more than double the historical average catch for mid-August. On August 13<sup>th</sup>, department staff flew an early aerial survey of the Niukluk River drainage to estimate initial escapement numbers. There is no longer an escapement counting project at Niukluk River in the Golovin Subdistrict. However, 630 coho salmon were counted in the Niukluk River from Ophir Creek downstream to the Fish River confluence during an August 13 aerial survey conducted by department personnel under fair to poor viewing conditions. These early survey results suggest that coho salmon escapement will be achieved as the old aerial survey goal of 950–1,900 coho salmon correlated well with the tower-based goal of 2,400–7,200 coho salmon. The survey count of 630 fish is also considered to be very conservative because of the difficult viewing conditions. To ensure that escapement and subsistence needs of coho salmon are met, the department may need to reduce commercial fishing opportunity in Golovin Subdistrict if catch statistics drop below historical average catch statistics during this opening.

In Elim Subdistrict the catch was 734 coho salmon by 15 permit holders and was below average for this date. However, escapement at Kwiniuk River counting tower is nearly 1,800 coho salmon with the average first quarter point of passage at the tower on August 10. Projections show that the Kwiniuk River aerial survey escapement goal of 650–1,300 will be easily achieved. This 48-hour opening should also not jeopardize reaching escapement goals or subsistence fishing opportunity.

Emergency Order: 3-S-Z-57-13    Effective Date: August 19, 2013

EXPLANATION: This emergency order reopens Subdistricts 2 and 3 of the Norton Sound District, the Golovin and Elim Subdistricts, to commercial salmon fishing for 48 hours from 6:00 p.m. Monday, August 19 to 6:00 p.m. Wednesday, August 21. This period is directed at coho salmon and permit holders will be restricted to gillnets with a mesh size of 6 inches or less.

JUSTIFICATION: This period is being established to provide additional commercial harvest opportunity of coho salmon. An estimated 609 and 880 coho salmon were harvested in the Golovin and Elim Subdistricts during the most recent 48-hour opening ending Friday, August 16. Catches were above average in Golovin and average in Elim for the third week of August.

On August 13<sup>th</sup>, department staff flew an early aerial survey of the Niukluk River drainage to estimate initial escapement numbers. There is no longer an escapement counting project at Niukluk River in the Golovin Subdistrict. However, 630 coho salmon were counted in the Niukluk River from Ophir Creek downstream to the Fish River confluence during an August 13 aerial survey conducted by department personnel under fair to poor viewing conditions. These early survey results suggest that coho salmon escapement will be achieved as the old aerial survey goal of 950–1,900 coho salmon correlated well with the tower-based goal of 2,400–7,200 coho salmon.

---

-continued-

The survey count of 630 fish is also considered to be very conservative because of the difficult viewing conditions. Peak spawning ground surveys will be flown later this month.

Escapement of coho salmon in the Elim Subdistrict as indexed by Kwiniuk River counting tower is 2,250 coho salmon with the average midpoint of passage at the tower on August 20. Projections show that the Kwiniuk River aerial survey escapement goal of 650–1,300 will be easily achieved. Furthermore, continued commercial fishing with 48-hour openings and escapement windows between periods should also not jeopardize subsistence uses of coho salmon for Elim residents.

Emergency Order: 3-S-Z-58-13   Effective Date: August 22, 2013

EXPLANATION: This emergency order supersedes 3-S-Z-55-13 and closes the fishing period beginning at 6 p.m. August 22 and moves that scheduled 48-hour fishing period 24 hours later to now start at 6 p.m. Friday, August 23 and conclude at 6 p.m. Sunday, August 25. Permit holders in Subdistrict 4 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches.

JUSTIFICATION: The buyer has requested that the scheduled fishing period be shifted to one day later. Because Subdistrict 4 is on a schedule of two 48-hour commercial salmon fishing periods per week there is no concern with shifting the fishing period to one day later.

Emergency Order: 3-S-Z-59-13   Effective Date: August 23, 2013

EXPLANATION: This emergency order reopens Subdistricts 2 and 3 of the Norton Sound District, the Golovin and Elim Subdistricts, to commercial salmon fishing for 48 hours from 6:00 p.m. Friday, August 23 to 6:00 p.m. Sunday, August 25. This period is directed at coho salmon and permit holders are restricted to gillnets with a mesh size of 6 inches or less.

JUSTIFICATION: The recent 48-hour fishing period in Subdistricts 2 and 3 had average catches when compared to the historical averages. The escapement goal is projected to be made in each subdistrict. The escapement goal is determined by aerial survey in Subdistrict 2 and is expected to be made based on an earlier aerial survey projection. The escapement goal in Subdistrict 3 is also determined by aerial survey and Kwiniuk River counting crew has enumerated four times the lower end of the coho salmon escapement goal needed by aerial survey. The upcoming fishing period should not jeopardize subsistence opportunity or escapement.

Emergency Order: 3-S-Z-60-13   Effective Date: August 27, 2013

EXPLANATION: This emergency order opens all marine waters of Norton Sound Subdistrict 3, the Elim Subdistrict, to commercial salmon fishing for one 48-hour period from 6:00 p.m. Tuesday, August 27 to 6:00 p.m. Thursday, August 29, and one 40-hour period from 8:00 a.m. Friday, August 30 to 12:00 midnight Saturday evening, August 31. Permit holders are limited to 100 fathoms of net with a mesh size no greater than 6 inches. This emergency order also closes the commercial salmon season in the Elim Subdistrict effective 12:00 midnight.

JUSTIFICATION: Escapement of coho salmon to Subdistrict 3 as indexed by Kwiniuk River tower counts is over 3,200 coho salmon. Current projections of inriver abundance based on these tower counts range from 4,100 and 5,300 cohos. This level of inriver abundance is sufficient to provide for upriver subsistence harvests of coho salmon and enough fish to achieve the Kwiniuk River aerial survey goal range of 650–1,300 fish. Catches of coho salmon in the commercial fishery continue to be average to above average for late August and these additional periods are warranted to utilize remaining harvestable surpluses.

Emergency Order: 3-S-Z-61-13   Effective Date: August 27, 2013

EXPLANATION: This emergency order supersedes emergency order 3-S-Z-55-13 and reopens Subdistrict 4 of the Norton Sound Subdistrict, the Norton Bay Subdistrict, to a new commercial fishing schedule of two 48-hour periods per week until the end of the Subdistrict 4 season, which closes by emergency order effective 6:00 p.m. Thursday,

---

-continued-

September 5. Periods will now be from 6:00 p.m. Tuesdays to 6:00 p.m. Thursdays and from 6:00 p.m. Fridays to 6:00 p.m. Sundays. Permit holders in Subdistrict 4 are limited to 100 fathoms of net in aggregate length and set gillnets must have a stretched-mesh size no greater than 6 inches.

**JUSTIFICATION:** Norton Bay fishermen have harvested a record 5,400 cohos this season. Commercial fishing effort has been minimal during the coho salmon season. Additionally, there were early indications in late July that inriver abundance of coho salmon would be sufficient to provide for subsistence uses and escapement needs of coho salmon. A July 30 aerial survey resulted in over 1,000 cohos counted in the lower Ungalik River. Additionally, previous years with similar fishing effort have had a limited impact on inriver abundance of coho salmon. Therefore, the limited level of participation in the commercial fishery coupled with escapement windows between periods should not jeopardize subsistence needs or have an adverse impact on escapements of coho salmon in Norton Bay drainages.

**Emergency Order:** 3-S-Z-62-13 **Effective Date:** September 9, 2013

**EXPLANATION:** : This emergency order opens the eastern end of Salmon Lake to subsistence salmon fishing. The catch limit is 100 sockeye salmon.

**JUSTIFICATION:** The escapement past Pilgrim River weir was over 12,000 sockeye salmon and the aerial survey goal of 4,000 to 8,000 sockeye salmon in Salmon Lake was easily reached. The majority of sockeye salmon spawn in the western half of Salmon Lake and that area remains closed by regulation. Because there are sufficient numbers of spawning salmon the department will allow subsistence salmon fishing in that area of the lake where fishing will allow some take of salmon to provide subsistence opportunity.

#### **NORTON SOUND SALMON – SPORT FISH**

**Emergency Order:** 3-KS-04-13 **Effective Date:** June 17, 2013

**EXPLANATION:** This emergency order prohibits the retention of king salmon in all waters of the Unalakleet and Shaktoolik river drainages, effective 12:01 a.m., Monday, June 17, 2013, and prohibits the use of bait while sport fishing in these rivers.

**Emergency Order:** 3-KS-08-13 **Effective Date:** July 11, 2013

**EXPLANATION:** This emergency order supersedes Emergency Order No. 3-KS-04-13. This emergency order closes all waters to sport fishing for king salmon, and prohibits the use of bait while sport fishing, in the Unalakleet and Shaktoolik river drainages effective 12:01 A.M. Thursday, July 11, 2013.

---



## **APPENDIX H: ARCTIC FISHERIES**

Appendix H1.—Commercial freshwater finfish harvest and sales, Colville River, Arctic Area, 1990–2007.

Year	Number of fish harvested intended for commercial sale <sup>a</sup>					Estimated commercial sales based on fish tickets	
	Broad whitefish	Humpback whitefish	Least Cisco ("herring")	Arctic Cisco ("kaktok")	Total harvest	Arctic Cisco	Whitefish species <sup>b</sup>
1990	0	5,694	21,003	19,374	46,071	12,571 <sup>c</sup>	14,249 <sup>c</sup>
1991	0	1,240	5,697	13,805	20,742	1,970 <sup>d</sup>	3,307 <sup>d</sup>
1992	126	5,209	6,962	20,939	33,236	<sup>e</sup>	10,200 <sup>f</sup>
1993	20	5,339	6,037	31,310	42,706	11,291 <sup>d</sup>	6,170 <sup>d</sup>
1994	—	6,056 <sup>g</sup>	10,176	8,958	25,190	7,434 <sup>d</sup>	4,121 <sup>d</sup>
1995	—	33,794 <sup>h</sup>	—	—	33,794	13,921	6,000
1996	—	6,425 <sup>g</sup>	7,796	21,817	36,038	9,076	4,127
1997	—	1,721 <sup>g</sup>	10,754	9,403	21,878	9,403	4,760
1998	—	4,881 <sup>g</sup>	9,936	7,019	21,836	5,648	7,105
1999	—	6,875 <sup>g</sup>	7,430	8,832	23,137	7,095	6,170
2000	—	3,706 <sup>g</sup>	5,758	2,619	12,083	2,809	6,569
2001	—	6,078 <sup>g</sup>	2,839	1,740	10,657	1,779	7,306
2002	—	4,183 <sup>g</sup>	5,503	3,935	13,621	899	4,093
2003	—	6,463 <sup>g</sup>	4,777	5,627	16,867	0	1,292
2004	—	1,145 <sup>g</sup>	3,061	3,061	7,267	2,412 <sup>f</sup>	476
2005	—	490 <sup>g</sup>	2,870	9,343	12,703	2,975 <sup>f</sup>	2,170
2006	—	1,188 <sup>g</sup>	4,995	3,293	9,476	1,482 <sup>f</sup>	3,655
2007	—	462 <sup>g</sup>	2,265	390	3,117	<sup>e</sup>	<sup>e</sup>
2002-2006							
Average	—	2,694	4,241	5,052	11,987	1,554	2,337

<sup>a</sup> Reported on daily catch form returned to ADF&G. Catch reports were returned to the department following the fishing season. All fish reported on the catch report were harvested with the intent to sell. Dashes indicate information is not available.

<sup>b</sup> Whitefish species include mostly humpback whitefish and least cisco, with occasional broad whitefish.

<sup>c</sup> Commercial harvest estimate based on one fish ticket average weights of 0.89 lb (900 Arctic cisco at 800 lb) and 0.61 lb (1,400 whitefish species at 850 lb).

<sup>d</sup> Estimated commercial harvest sales based on 1995 to 2001 average weight of 0.92 lb for Arctic cisco and 0.89 lb for whitefish species (humpback and broad whitefish, and least cisco).

<sup>e</sup> No information is available from fish tickets indicating that harvested fish were sold commercially.

<sup>f</sup> Mixed commercial harvest of mostly Arctic cisco along with humpback whitefish, broad whitefish, and least cisco. Estimated commercial harvest sales based on 1995 to 2001 combined average of \$1.07/lb. for whitefish species and Arctic cisco.

<sup>g</sup> Humpback whitefish harvest includes undetermined amounts of broad whitefish.

<sup>h</sup> Humpback whitefish harvest includes undetermined amounts of broad whitefish, least cisco, and Arctic cisco.