

A detailed microscopic image of green algae. The image shows numerous long, thin, filamentous structures (likely diatoms or similar algae) and several spherical, radiating colonies. The filaments are dark green and appear to be intertwined, while the spherical colonies are also dark green and have a distinct radial pattern. The background is a light, almost white color, which makes the green structures stand out prominently.

H. B. N. HYNES

# The Ecology of Running Waters

UNIVERSITY OF TORONTO PRESS

**This document is copyrighted material.**

Alaska Resources Library and Information Services (ARLIS) is providing this excerpt in an attempt to identify and post all documents from the Susitna Hydroelectric Project.

This book is identified as APA no. 2158 in the Susitna Hydroelectric Project Document Index (1988), compiled by the Alaska Power Authority.

It is unable to be posted online in its entirety. Selected pages are displayed here to identify the published work.

The book is available at call number QH541.5.S7H9 1970 in the ARLIS Susitna collection.

WTT  
S41.5  
.S7  
H9  
1974

# The Ecology of Running Waters

---

H. B. N. HYNES

*Professor of Biology,  
University of Waterloo, Ontario, Canada*

**ARLIS**  
Alaska Resources  
Library & Information Services  
Anchorage, Alaska

UNIVERSITY OF TORONTO PRESS

# *Contents*

PREFACE	vii
LIST OF FIGURES	xv
LIST OF TABLES	xxi
I Water flow and stream channels	I
II Physical characteristics of flowing water	28
III Chemical characteristics of flowing water	36
IV Attached algae	53
v Higher plants	78
VI Plankton	94
VII Composition of the benthic invertebrate fauna	112
VIII Anatomical and behavioural adaptations of benthic invertebrates	121
IX Further adaptations of benthic invertebrates	161
x Feeding mechanisms and food of benthic invertebrates	183
XI Factors controlling benthic invertebrates	196
XII Quantitative study of benthic invertebrates	236
XIII Effects of downstream movements of organisms on the benthos	256
XIV Life histories and seasonal cycles of benthic invertebrates	271

xiv CONTENTS

xv	The fishes of running water	300
xvi	Ecological factors affecting fishes	319
xvii	Movements and breeding of fishes	342
xviii	Feeding habits of fishes	364
xix	Other vertebrates	377
xx	Longitudinal zonation	383
xxi	Special habitats	398
xxii	The ecosystem	411
xxiii	Effects of man on watercourses	443
	BIBLIOGRAPHY AND INDEX OF CITATIONS	451
	INDEX OF ORGANISMS	519
	SUBJECT INDEX	543

# Figures

I, 1. Diagram of the routes taken by the products of precipitation	3
I, 2. Two types of notched weir used to measure the discharge of streams	4
I, 3. A simple current meter	6
I, 4. The rate of flow of water at different depths in an open channel	6
I, 5. The patterns of flow in cross-sections of open channels	7
I, 6. Photograph of particles carried by flowing water over a stone	<i>facing page</i> 8
I, 7. Diagrams of the patterns of flow below a waterfall and at the foot of a slope	10
I, 8. A sketch-map showing stream orders	13
I, 9. Scour and refill during a flood	17
I, 10. A rating curve for gauging stations in the eastern United States	18
I, 11. Diagrams of the patterns of flow near small irregularities in a channel and in a meander	20
II, 1. The levels of light at the surface of a small stream throughout the year	34
IV, 1. Diatoms of the attached algal community	54
IV, 2. Red, blue-green, and yellow algae of the attached community	55
IV, 3. Green algae of the attached community	56
IV, 4. Diagram of the distribution of certain algae on submerged and emergent stones	75

xvi FIGURES

iv, 5. Photographs of travertine and diagrams of some of the algae which make it	<i>facing page 76</i>
v, 1. The position which Podostemaceae occupy on emergent rocks	82
v, 2. Diagram of formation and erosion of banks of sediment formed by rooted plants	89
v, 3. Changes in the rooted vegetation in the River Lark during a ten-month period	90
v, 4. Changes in the vegetation in the River Acher during three years	91
vi, 1. Common phytoplankters from running water	95
vi, 2. Common zooplankters from running water	96
vi, 3. The abundance of plankton in the Sacramento River at different points and times of year	97
vii, 1. Survivors of ancient invertebrate stocks which now occur mainly in running water	113
vii, 2. Two members of the Gerridae which occur primarily on running water	116
vii, 3. Comparison between the nymphs of some European stoneflies and their morphological equivalents among the Gripopterygidae of South America	118
viii, 1. Two flattened mayflies of the family Ecdyonuridae which live exposed on stones	123
viii, 2. Flattened mayflies which live in sheltered places under stones	124
viii, 3. Two flattened animals which are common under stones in rapid water in Central Africa	124
viii, 4. Flatworms and a leech which occur under stones in rapid water	125
viii, 5. Streamlined nymphs of the family Baetidae	126
viii, 6. A caddis-worm with a long tail to its case and several animals with lateral projections	127
viii, 7. Larvae of Blepharoceridae	129
viii, 8. Broad-footed snails	131
viii, 9. Larvae with peripheral bristles which fit the surfaces of stones	132
viii, 10. Two mayflies with friction-pads of bristles	133

VIII, 11. Ventral views of the abdomens of a dragonfly and a mayfly with friction-pads	134
VIII, 12. The hooks at the rear ends of insect larvae from rapid water	135
VIII, 13. Larvae of <i>Deuterophlebia</i> and larva and pupa of <i>Limnophora</i>	135
VIII, 14. Larvae of pyralid moths and <i>Rheotanytarsus</i> , and adult <i>Corophium</i>	137
VIII, 15. Larva of <i>Simulium</i>	138
VIII, 16. Caddis-worms with heavy stone cases	140
VIII, 17. The shells of several large bivalves	141
VIII, 18. Prosobranch Gastropoda and Sphaeriidae which occur in streams and rivers	142
VIII, 19. Lateral views of <i>Taeniopteryx</i> and <i>Araucanioperla</i>	144
VIII, 20. Larvae of Diptera which burrow in the substratum	147
VIII, 21. Elongate stonefly and mayfly nymphs which burrow in gravel	148
VIII, 22. The mitten-crab	154
VIII, 23. The mandibles of a caddis-fly pupa and the pupae of three families of Diptera which emerge under water	157
VIII, 24. Adult insects which creep down into quite rapid water to oviposit	158
IX, 1. Graphs showing the effect of current speed on the oxygen consumption of some animals	164
IX, 2. Oxygen concentrations which are just too low for the survival of experimental animals at different current speeds	165
IX, 3. Insects which use plastron respiration	171
IX, 4. Animals which are associated with the water surface in streams	174
IX, 5. Mayfly nymphs which burrow in sand in rivers	176
IX, 6. Mayflies and a dragonfly which burrow in soft substrata	177
IX, 7. Very flattened animals which live in silty substrata	178
IX, 8. Nymphs of mayflies which occur in silty places in streams	179
IX, 9. Phoretic associations between <i>Simulium</i> and mayflies and crabs	181

xviii FIGURES

x, 1.	The head of <i>Brachyptera</i> and comparison of its maxilla with that of an unspecialized stonefly	183
x, 2.	Ventral views of the heads of mayfly nymphs and details of mouth parts	184
x, 3.	Insects which strain particles from the water with fringes of hairs	185
x, 4.	Head of the larva of <i>Simulium</i> and a position commonly assumed during feeding	186
x, 5.	Photographs of nets of <i>Hydropsyche</i> and <i>Neureclipsis</i> facing page	188
x, 6.	Cases and nets of <i>Diplectrona</i> and <i>Hydropsyche</i>	189
x, 7.	The larva of <i>Hydropsyche</i> and the head of an African <i>Cheumatopsyche</i>	189
x, 8.	The larval case of a Brazilian species of <i>Macronema</i>	190
x, 9.	Nymphs of Zygoptera	191
x, 10.	The carnivorous mayfly <i>Centroptiloides</i>	194
xI, 1.	The larva of <i>Helicopsyche</i>	198
xI, 2.	The numbers of species of three insect orders emerging at four points along a stream	202
xI, 3.	The oxygen consumption of two species of the leech <i>Erpobdella</i>	217
xIII, 1.	The numbers of <i>Gammarus</i> caught in drift samples at different times of day and year	263
xIII, 2.	The numbers of <i>Baetis</i> caught in drift samples at different times of day and year	264
xIV, 1.	The size-distributions in monthly collections of three species from a stream in Wales	272
xIV, 2.	A diapausing nymph of <i>Capnia bifrons</i>	281
xIV, 3.	Diagrams of the various types of life cycle commonly found in stream-dwelling insects	295
xIV, 4.	Diagrams of unusual types of life cycle found in stream-dwelling insects and of typical cycles of Peracarida and Mollusca	296
xIV, 5.	Diagrams of the seasonal fluctuations in numbers and biomass of invertebrates in streams	298
xv, 1.	Outlines of fishes which live in fast-flowing water	303
xv, 2.	Outlines of fishes which live in slow-flowing water	303

xv, 3. Outlines of small fishes which live on the bottom in fast-flowing water	304
xv, 4. Ventral views of fishes with friction devices and suckers	306
xv, 5. Tadpoles from rapid streams	307
xvi, 1. The relationship between the lethal temperature and the temperature to which specimens of <i>Salvelinus fontinalis</i> have been acclimated	323
xvi, 2. Changes in the instantaneous growth-rate of <i>Salvelinus fontinalis</i>	325
xx, 1. Diagram of the occurrence and abundance of common species of Ephemeroptera in rivers in north-western Europe	389
xx, 2. The altitudinal distribution of the Elminthidae in streams on Mount Elgon	390
xx, 3. The altitudinal distribution of the important members of the fauna in streams on Mount Elgon	395
xxi, 1. Madicolous animals which may occur in collections made in streams	409
xxii, 1. Diagram of the trophic relationships of the ecosystem in the rhithron	439
xxii, 2. Diagram of the trophic relationships of the ecosystem in the potamon	440

# Tables

I, 1.	Mean velocities of clear and muddy water required to initiate movement of various types of particles	11
I, 2.	Speeds of flow required to move mineral particles of different sizes	11
I, 3.	Calculated mean velocities of water in three channels of differing dimensions at various slopes	19
I, 4.	Suggested terminology, categories and methods for particle-size analysis	24
III, 1.	Chemical content of the water of the River Dee at various points below its source	45
IV, 1.	Some common attached algae of various types of habitat in running water	59
IV, 2.	The rate of oxygen production by algal communities in laboratory streams at different temperatures and illuminations	62
IV, 3.	The effect of current on the growth of algae in the laboratory	64
V, 1.	The dominant submerged plants in various types of river in Great Britain	84
VI, 1.	The numbers of plankton organisms on various dates at two stations on the Montreal River	102
VI, 2.	The numbers of plankters at various distances below the Lunzer Untersee	103
VI, 3.	The numbers of planktonic organisms at different seasons in the River Danube	109
VIII, 1.	The maximum current speeds resisted, moved against, and inhabited in nature by a number of European stream animals	149

xxii TABLES

IX, 1.	The oxygen consumption of the nymphs of various European mayflies	161
XI, 1.	The percentage composition of adults of various groups of insects emerging over different types of substratum in streams in Algonquin Park	208
XI, 2.	The percentage composition of the fauna on various types of stream bed in north-eastern England	209
XI, 3.	The percentage composition of the faunas of riffles and pools in the Black River	211
XI, 4.	The numbers of various animals in ten samples taken in April 1962 from Shelligan Burn	212
XI, 5.	The percentage composition and mean number of animals per square metre on various types of substratum in Doe Run	214
XI, 6.	The numbers and weights of animals in different species of plant in the River Angara	215
XI, 7.	The average number of animals per square foot taken from three streams in Utah with a Surber sampler	225
XI, 8.	The average amounts of invertebrate fauna in three small rivers and their tributaries in Transcarpathia	227
XI, 9.	The average number of invertebrates per square foot at three altitudes in the Sagana River	228
XII, 1.	The numbers of invertebrates collected in six standardized samples from Allt Leathan	240
XII, 2.	The numbers and weights of animals per one tenth square metre collected by three methods from the River Polenz	245
XII, 3.	The weights of invertebrates collected with a Surber sampler at the sides and centres of various streams in New York State	248
XII, 4.	The numbers and weights of organisms per square metre at different distances from the bank in the River Plane	249
XII, 5.	The average numbers of animals collected by a standardized method in spring and summer from streams in the Scottish Highlands	252
XII, 6.	The numbers and weights of invertebrates at various points on the River Hæelva	254
XIII, 1.	The numbers of animals per square foot collected on various dates above and below four lakes on the Montreal River	258

XIII, 2.	The numbers and weights of benthic organisms per 1000 sq. cm. at two stations below a lake on Norvijokk	259
XIII, 3.	The percentage composition of the benthic fauna and the drift in the Missouri River at Boonville	260
XIII, 4.	Percentages of total numbers of Ephemeroptera, Plecoptera and Simuliidae, which were longer than 3 mm., caught in drift samples in each three-hour period during two days from Berry Creek	265
XIV, 1.	The numbers of species and genera of midges occurring in different types of stream in Europe	274
XIV, 2.	The mean lengths of specimens of <i>Capnia bifrons</i> emerging in successive half-monthly periods	285
XIV, 3.	The percentage composition, by species, of larvae of <i>Simulium</i> from eight samples collected at various altitudes on the River Manafwa	290
XVI, 1.	The ecological distribution of various species of fish in the Ohio River, and in small streams flowing into it, in Kentucky	330
XVIII, 1.	The average total volumes and numbers of invertebrates in stomachs of rainbow trout of different sizes	369
XVIII, 2.	The percentage composition by volume of the stomach contents of young salmon and trout from the River Endrick	370
XVIII, 3.	The percentage composition by volume of the stomach contents of burbot of different sizes from the White River	370
XVIII, 4.	The percentage composition by volume of the stomach contents of four species of fishes from the River Endrick	374
XVIII, 5.	The percentage composition by numbers of the gut contents of three species of fishes from the Morávka River	375
xx, 1.	The communities of fishes in the four fish-zones of Western Europe	384
xx, 2.	Huet's 'slope-rule' for western European streams	384
xx, 3.	The distribution of fish species along the length of two adjacent tributaries of New River	386
xx, 4.	The distribution of Unionidae in the Huron River system	388
xx, 5.	The percentage composition of the fauna of the Afon Hirnant at four altitudes	394

xxiv TABLES

xxii, 1.	The estimated numbers at different stages in their life histories of two species of stonefly during two years in a reach of Walla Brook	421
xxii, 2.	The original numbers and percentage survivors of two species of stonefly at various stages in their growth in Walla Brook and Afon Hirnant	422
xxii, 3.	The percentage composition of the invertebrate fauna of the upper River Derwent in the spring of nine successive years	423
xxii, 4.	The average biomass and total production of trout in three stations on Walla Brook during two years	429
xxii, 5.	The biomass and production of <i>Salmo clarkii</i> in four sections of Berry Creek	431
xxii, 6.	The percentages of taxa which occur in different numbers of nine rivers in south-eastern United States	442