

Table 6. Comparison of age composition, weight, and length (AWL) data from the sac roe herring fishery, Kodiak Management Area, 1989-1996.

YEAR	Age Class										Average	N
	2	3	4	5	6	7	8	9	10	11+		
<i>Percent at Age</i>												
1989	0.3	7.1	6.2	42.0	19.3	1.0	10.0	4.5	8.5	1.1		3,026
1990	0.7	52.0	3.7	3.3	20.4	8.5	0.6	3.0	2.6	5.2		7,672
1991	<0.1	26.3	49.4	2.6	1.4	8.7	5.2	0.6	2.7	3.0		5,498
1992	0.5	2.9	62.3	26.9	2.0	0.7	2.0	1.2	<0.1	1.4		9,325
1993	0.3	8.7	4.5	63.5	18.2	1.2	0.2	1.6	0.8	1.0		7,396
1994	0.1	11.5	11.6	3.8	42.4	24.5	1.9	0.7	2.0	1.5		6,965
1995	0.3	3.2	16.0	8.9	2.2	37.4	28.3	0.7	0.4	2.3		3,775
1996	0.6	38.6	6.2	11.4	5.0	1.8	23.3	10.4	0.5	1.7		6,890
<i>Average Weights (Grams)</i>												
1989	60	85	130	176	224	251	269	271	276	284	199	2,569
1990	52	95	141	167	191	248	229	280	283	290	152	4,885
1991	53	81	137	169	191	221	255	261	301	292	153	4,239
1992	40	87	114	176	201	226	247	284	283	303	140	8,139
1993	58	94	134	144	203	226	226	250	281	310	156	6,852
1994	78	99	129	166	182	251	251	256	282	316	188	6,836
1995	49	109	146	166	200	208	264	270	270	310	209	3,763
1996	52	93	132	187	193	227	230	294	284	305	171	6,768
<i>Average Lengths (Millimeters)</i>												
1989	168	185	210	227	247	254	259	259	260	263	235	3,026
1990	158	188	214	226	232	250	245	261	258	260	212	7,671
1991	166	184	215	225	235	244	251	257	266	263	215	5,497
1992	162	194	202	231	240	250	254	264	265	268	214	9,323
1993	161	191	211	216	240	247	244	254	261	264	219	7,394
1994	175	194	211	227	231	253	252	253	259	267	231	6,965
1995	163	201	220	228	241	243	261	264	265	272	242	3,774
1996	158	192	212	235	236	248	247	266	261	268	223	6,803

