

Alaska Subsistence and Personal Use Salmon Fisheries 2012 Annual Report

by

James A. Fall

Nicole M. Braem

Caroline L. Brown

Sarah S. Evans

Lisa Hutchinson-Scarbrough

Bronwyn Jones

Robbin La Vine

Terri Lemons

Meredith Ann Marchioni

Elizabeth Mikow

Joshua T. Ream

and

Lauren A. Sill

September 2014

Alaska Department of Fish and Game

Division of Subsistence



Symbols and Abbreviations

The following symbols and abbreviations, and others approved for the Système International d'Unités (SI), are used without definition in the reports by the Division of Subsistence. All others, including deviations from definitions listed below, are noted in the text at first mention, as well as in the titles or footnotes of tables, and in figure or figure captions.

Weights and measures (metric)		General		Mathematics, statistics	
centimeter	cm	Alaska Administrative Code	AAC	<i>all standard mathematical signs, symbols and abbreviations</i>	
deciliter	dL	all commonly-accepted abbreviations	e.g., Mr., Mrs., AM, PM, etc.	alternate hypothesis	H _A
gram	g			base of natural logarithm	e
hectare	ha			catch per unit effort	CPUE
kilogram	kg			coefficient of variation	CV
kilometer	km	all commonly-accepted professional titles	e.g., Dr., Ph.D., R.N., etc.	common test statistics (F, t, χ^2 , etc.)	
liter	L			confidence interval	CI
meter	m			correlation coefficient (multiple)	R
milliliter	mL	at	@	correlation coefficient (simple)	r
millimeter	mm	compass directions:		covariance	cov
		east	E	degree (angular)	°
		north	N	degrees of freedom	df
		south	S	expected value	E
		west	W	greater than	>
		copyright	©	greater than or equal to	≥
		corporate suffixes:		harvest per unit effort	HPUE
		Company	Co.	less than	<
		Corporation	Corp.	less than or equal to	≤
		Incorporated	Inc.	logarithm (natural)	ln
		Limited	Ltd.	logarithm (base 10)	log
		District of Columbia	D.C.	logarithm (specify base)	log ₂ , etc.
		et alii (and others)	et al.	minute (angular)	'
		et cetera (and so forth)	etc.	not significant	NS
		exempli gratia (for example)	e.g.	null hypothesis	H ₀
		Federal Information Code	FIC	percent	%
		id est (that is)	i.e.	probability	P
		latitude or longitude	lat. or long.	probability of a type I error (rejection of the null hypothesis when true)	α
		monetary symbols (U.S.)	\$, ¢	probability of a type II error (acceptance of the null hypothesis when false)	β
		months (tables and figures)	first three letters (Jan.,...,Dec)	second (angular)	"
		registered trademark	®	standard deviation	SD
		trademark	™	standard error	SE
		United States (adjective)	U.S.	variance	
		United States of America (noun)	USA	population	Var
		U.S.C.	United States Code	sample	var
		U.S. state	two-letter abbreviations (e.g., AK, WA)		
Weights and measures (English)					
cubic feet per second	ft ³ /s				
foot	ft				
gallon	gal				
inch	in				
mile	mi				
nautical mile	nmi				
ounce	oz				
pound	lb				
quart	qt				
yard	yd				
Time and temperature					
day	d				
degrees Celsius	°C				
degrees Fahrenheit	°F				
degrees kelvin	K				
hour	h				
minute	min				
second	s				
Physics and chemistry					
<i>all atomic symbols</i>					
alternating current	AC				
ampere	A				
calorie	cal				
direct current	DC				
hertz	Hz				
horsepower	hp				
hydrogen ion activity (negative log of)	pH				
parts per million	ppm				
parts per thousand	ppt, ‰				
volts	V				
watts	W				
		Measures (fisheries)			
		fork length	FL		
		mid-eye-to-fork	MEF		
		mid-eye-to-tail-fork	METF		
		standard length	SL		
		total length	TL		

TECHNICAL PAPER NO. 406

**ALASKA SUBSISTENCE AND PERSONAL USE SALMON FISHERIES
2012 ANNUAL REPORT**

by

James A. Fall, Lisa Hutchinson-Scarbrough, Bronwyn Jones,
Robbin La Vine, Terri Lemons, and Joshua T. Ream
Alaska Department of Fish and Game, Division of Subsistence, Anchorage

Nicole M. Braem, Caroline L. Brown,
Hiroko Ikuta, and Elizabeth Mikow
Alaska Department of Fish and Game, Division of Subsistence, Fairbanks

Sarah S. Evans
Alaska Department of Fish and Game, Division of Subsistence, Dillingham

and

Meredith Ann Marchioni and Lauren A. Sill
Alaska Department of Fish and Game, Division of Subsistence, Douglas

Alaska Department of Fish and Game
Division of Subsistence
333 Raspberry Road, Anchorage, AK 99518

September 2014

The Division of Subsistence Technical Paper series was established in 1979 and represents the most complete collection of information about customary and traditional uses of fish and wildlife resources in Alaska. The papers cover all regions of the state. Some papers were written in response to specific fish and game management issues. Others provide detailed, basic information on the subsistence uses of particular communities which pertain to a large number of scientific and policy questions.

Technical Paper series reports are available through the Alaska Resources Library and Information Services (ARLIS), the Alaska State Library and on the Internet: <http://www.adfg.alaska.gov/sf/publications/>. This publication has undergone editorial and professional review.

*James A. Fall, Lisa Hutchinson-Scarborough, Bronwyn Jones,
Robbin La Vine, Terri Lemons, and Joshua T. Ream
Alaska Department of Fish and Game, Division of Subsistence
333 Raspberry Road, Anchorage, AK 99518-1565*

*Nicole M. Braem, Caroline L. Brown,
Hiroko Ikuta, and Elizabeth Mikow
Alaska Department of Fish and Game, Division of Subsistence
1300 College Road, Fairbanks, AK 99701-1551*

*Sarah S. Evans
Alaska Department of Fish and Game, Division of Subsistence
546 Kenny Wren Road, PO Box 230, Dillingham, AK 99576-0230*

*and
Meredith Ann Marchioni and Lauren A. Sill
Alaska Department of Fish and Game, Division of Subsistence
PO Box 110024, Juneau, AK 99811-0024 USA*

This document should be cited as:

Fall, J. A., N. M. Braem, C. L. Brown, S. S. Evans, L. Hutchinson-Scarborough, B. Jones, R. La Vine, T. Lemons, M. A. Marchioni, E. Mikow, J. T. Ream, and L. A. Sill. 2014. Alaska subsistence and personal use salmon fisheries 2012 annual report. Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 406, Anchorage.

The Alaska Department of Fish and Game (ADF&G) administers all programs and activities free from discrimination based on race, color, national origin, age, sex, religion, marital status, pregnancy, parenthood, or disability. The department administers all programs and activities in compliance with Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act (ADA) of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972.

If you believe you have been discriminated against in any program, activity, or facility please write:

ADF&G ADA Coordinator, P.O. Box 115526, Juneau, AK, 99811-5526

U.S. Fish and Wildlife Service, 4040 N. Fairfax Drive, MS 2042, Arlington, VA, 22203

Office of Equal Opportunity, U.S. Department of the Interior, 1849 C Street NW, MS 5230, Washington DC 20240

The department's ADA Coordinator can be reached via phone at the following numbers:

(VOICE) 907-465-6077, (Statewide Telecommunication Device for the Deaf) 1-800-478-3648, (Juneau TDD) 907-465-3646, or (FAX) 907-465-6078

For information on alternative formats and questions on this publication, please contact:

ADF&G Division of Subsistence at <http://www.adfg.alaska.gov/index.cfm?adfg=contacts.anchorage>.

ERRATA

Originally published in September 2014, Technical Paper No. 406, *Alaska subsistence and personal use salmon fisheries 2012 annual report*, has been revised and updated to reflect the following changes:

- Table 2-2, which was omitted in the original copy, has been added at page 12.

Table of Contents

	Page
LIST OF TABLES.....	v
TABLE OF FIGURES.....	vii
ABSTRACT	viii
CHAPTER 1: INTRODUCTION	1
CHAPTER 2: OVERVIEW OF SUBSISTENCE FISHERIES IN ALASKA	7
Subsistence Harvests in Rural Alaska	7
Subsistence Salmon Harvests in 2012	7
Personal Use Salmon Harvests in 2012	8
Statewide Subsistence and Personal Use Salmon Harvests, 1994–2012	8
CHAPTER 3: NORTON SOUND-PORT CLARENCE AREA AND ARCTIC-KOTZEBUE AREA ALASKA	27
Introduction	27
Norton Sound–Port Clarence Area Salmon	27
Background.....	27
Regulations	27
Subsistence Salmon Harvest Data Collection Methods	29
Subsistence Salmon Harvests in 2012	31
Arctic-Kotzebue Area Salmon.....	35
Introduction	35
Background.....	35
Regulations	35
Subsistence Salmon Harvest Data Collection Methods	36
Arctic-Kotzebue Area Salmon, Sheefish, Whitefishes, and Arctic Char/Dolly Varden	36
CHAPTER 4: YUKON AREA.....	51
Background.....	51
Regulations	51
Subsistence Harvest Assessment Methods	55
Subsistence Salmon Harvests in 2012	56
Nonsalmon Fish Harvests	58
The Role of Salmon within Annual Subsistence Harvests	59
CHAPTER 5: KUSKOKWIM AREA	81
Background.....	81
Regulations	82
Subsistence Fishery	83
Commercial Fishery.....	84
Subsistence Salmon Harvest Assessment Methods	85
Household Harvest Surveys.....	85
Harvest Calendars	86
Data Correction and Archiving	87
Data Analysis.....	87
2012 Sampling Summary	87
2012 Subsistence Salmon Harvest Summary	88
Use of Salmon for Dog Food.....	88

Gear Types.....	89
Salmon Retained from Commercial Fishing for Subsistence Uses.....	89
Other Fish	89
The Role of Salmon within Annual Subsistence Harvests	90
CHAPTER 6: BRISTOL BAY AREA	103
Background.....	103
Regulations	103
Inseason Management in 2012	104
Salmon Harvest Assessment Program	104
Subsistence Salmon Harvests in 2012	104
Other Subsistence Fisheries.....	106
Subsistence Regulations	106
Subsistence Harvests and Uses	106
CHAPTER 7: CHIGNIK MANAGEMENT AREA.....	117
BACKGROUND	117
REGULATIONS	117
Recent Regulatory History	118
Harvest Assessment Program	120
CMA Subsistence Salmon Harvests	121
Gear Type	123
Federal Subsistence Fishery in CMA	123
Salmon Removal From Commercial Harvests for Home Use (“Home Pack”)	124
Other Chignik Area Subsistence Fisheries	124
Discussion.....	124
CHAPTER 8: ALASKA PENINSULA AREA	141
Background.....	141
Regulations	141
Harvest Assessment Program	141
Subsistence Salmon Harvests in 2012	141
Other Subsistence Fisheries.....	142
CHAPTER 9: ALEUTIAN ISLANDS AREA	149
Introduction	149
Salmon Harvests in the Unalaska District	150
Salmon Harvest Regulations.....	150
Salmon Harvest Assessment Program	150
Subsistence Salmon Harvests in 2012	150
Salmon Harvests in the Adak District.....	151
Salmon Harvest Regulations.....	152
Salmon Harvest Assessment Program	152
Subsistence Salmon Harvests in 2012, Adak District.....	152
Salmon Harvests at Akutan, Nikolski, and Atka	152
Other Subsistence Fisheries in the Aleutian Islands Area	153
Finfishes.....	153
Shellfish	153

CHAPTER 10: KODIAK AREA	159
Introduction	159
Salmon Harvest in the Kodiak Area	159
Salmon Harvest Regulations.....	159
Salmon Harvest Assessment Program	160
Subsistence Salmon Harvests in 2012	160
Other Subsistence Fisheries in the Kodiak Area.....	162
Finfishes.....	162
Shellfish	162
CHAPTER 11: COOK INLET AREA	169
Introduction	169
Port Graham and Koyuktoilik Subdistricts	169
History and Regulations.....	169
Harvest Assessment Methods	170
Harvest Estimates for 2012.....	170
Seldovia Subsistence Fishery	171
History and Regulations.....	171
Harvest Assessment Methods	171
The 2012 Season.....	171
Tyonek Subdistrict.....	172
History and Regulations.....	172
Harvest Assessment Methods	172
The 2012 Season.....	172
Upper Yentna River Fish Wheel Fishery.....	173
History and Regulations.....	173
Harvest Assessment Methods	173
Harvests in 2012	173
Federal Subsistence Salmon Fisheries in Cook Inlet.....	173
Cook Inlet Personal Use Salmon Fisheries.....	174
Background.....	174
Upper Cook Inlet Personal Use Salmon Fisheries	174
Lower Cook Inlet Personal Use Salmon Fisheries.....	176
Other Subsistence Fisheries in Cook Inlet.....	176
CHAPTER 12: PRINCE WILLIAM SOUND AREA.....	201
Introduction	201
Upper Copper River State and Federal Subsistence Fisheries: Glennallen Subdistrict	202
Regulations	202
Harvest Assessment Program	202
Subsistence Salmon Harvests in 2012	203
Upper Copper River State Personal Use Fishery: Chitina Subdistrict.....	203
Background and History	203
Regulations	204
Harvest Assessment Program	204
Personal Use Salmon Harvests in 2012	204
Upper Copper River Federal Subsistence Fishery: Chitina Subdistrict	204
Regulations	204
Federal Subsistence Harvests in 2012.....	205
Native Village of Batzulnetas Subsistence Fishery	205

Copper River District Subsistence Fishery	205
Background and Regulations	205
Harvest Assessment Program	205
Subsistence Salmon Harvests in 2012	206
Eastern District Subsistence Salmon Fishery	206
Southwestern District Subsistence Salmon Fishery.....	206
Prince William Sound: General Districts.....	207
Other Subsistence Fisheries in the Prince William Sound Area	207
CHAPTER 13: THE SOUTHEAST REGION	227
Introduction	227
Harvest Assessment Programs.....	227
Subsistence Salmon Harvests in 2012	228
Yakutat Management Area	228
Yakutat Area Subsistence Fisheries.....	228
Haines Management Area.....	229
Haines Area Subsistence Fisheries	229
Juneau Management Area.....	231
Angoon Subsistence Area.....	231
Hoonah Subsistence Area	232
Elfin Cove, Gustavus, Pelican, and Tenakee Springs Subsistence and Personal Use Salmon Fishing.....	233
Juneau Personal Use Area.....	234
Sitka Management Area	235
Sitka Subsistence and Personal Use Salmon Fisheries	235
Petersburg Management Area.....	236
Kake Subsistence Area	236
Petersburg–Wrangell Subsistence–Personal Use Area	237
2012 Federal Stikine River Subsistence Salmon Fishery: Regulations	239
Point Baker–Port Protection Subsistence Fisheries	241
Ketchikan Management Area	241
Craig, Klawock, and Hydaburg Subsistence Fisheries	242
Kasaan Subsistence Area.....	243
Ketchikan Personal Use Area	244
ACKNOWLEDGEMENTS.....	254
REFERENCES CITED	255

LIST OF TABLES

Table 2-1.–Alaska subsistence and personal use salmon harvests, 2012.....	10
Table 2-2.–Historical Alaska subsistence salmon harvests, 1994–2012.....	12
Table 2-3.–Historical Alaska personal use salmon harvests, 1994–2012.....	13
Table 2-4.–Historical Alaska subsistence and personal use salmon harvests, 1994–2012.....	14
Table 2-5.–Alaska subsistence salmon harvests by species and place of residence, 2012.....	15
Table 3-1.–Subsistence salmon harvests by Norton Sound subdistricts, Northwest Alaska, 2012.....	40
Table 3-2.–Subsistence salmon harvests by community, Norton Sound-Port Clarence Area, Alaska, 2012.....	41
Table 3-3.–Historical subsistence salmon harvests by district, Northwest Alaska, 1994–2012.....	42
Table 3-4.–Subsistence salmon harvests by district, Northwest Alaska, 2012.....	44
Table 3-5.–Historical subsistence salmon harvests, Northwest Alaska, 1975–2012.....	45
Table 3-6.–Subsistence salmon harvests by Kotzebue District ^a communities.....	46
Table 3-7.–Subsistence nonsalmon harvests by Kotzebue District ^a communities.....	47
Table 3-8.–Subsistence salmon harvests by Arctic District communities.....	48
Table 3-9.–Subsistence nonsalmon harvests by Arctic District communities.....	48
Table 4-1.–Subsistence fishing schedule by district, Lower Yukon Area, 2012.....	61
Table 4-2.–Subsistence fishing schedule by district, Upper Yukon Area, 2012.....	64
Table 4-3.–Household subsistence and personal use permits, listed by fishery and community of residence, Yukon Area, 2012.....	70
Table 4-4.–Estimated number of subsistence fishing households in surveyed communities, with community and district totals, Yukon Area, 2012.....	71
Table 4-5.–Estimated subsistence salmon harvests by community, Yukon Area, 2012.....	72
Table 4-6.–Historical subsistence salmon harvests, Yukon Area, 1976–2012.....	73
Table 4-7.–Comparison of amounts necessary for subsistence (ANS) and estimated subsistence salmon harvests, Yukon Area, 1998–2012.....	74
Table 4-8.–Estimated subsistence harvest of nonsalmon fish by community, Yukon Area, 2012.....	75
Table 5-1.–Subsistence salmon harvests by community, Kuskokwim Area, 2012.....	91
Table 5-2.–Subsistence salmon harvests in 7 coastal Kuskokwim communities, 2011.....	93
Table 5-3.–Historical subsistence salmon harvests, Kuskokwim Area, 1989–2012.....	94
Table 5-4.–Number of households that own dogs, fed salmon to dogs, and total number of salmon fed to dogs, Kuskokwim Area, 2012.....	95
Table 5-5.–Gear types used for subsistence fishing, Kuskokwim Area, 2012.....	97
Table 5-6.–Reported number of salmon retained from commercial harvest for subsistence use, Kuskokwim Area, 2012.....	98
Table 5-7.–Subsistence nonsalmon fish harvests by community, Kuskokwim Area, 2012.....	100
Table 6-1.–Estimated subsistence salmon harvests by district and location fished, Bristol Bay Area, 2012.....	109
Table 6-2.–Estimated historical subsistence salmon harvests, Bristol Bay Area, 1983–2012.....	110
Table 6-3.–Estimated subsistence salmon harvests by community, Bristol Bay Area, 2012.....	111
Table 6-4.–Uses and harvests of fish other than salmon, Bristol Bay communities.....	113
Table 6-5.–Nonsalmon finfish used for subsistence purposes in the Bristol Bay Area.....	114
Table 7-1.–Historical subsistence salmon harvests, Chignik Area, 1977–2012.....	127
Table 7-2.–Participation in Chignik area subsistence salmon fishery and estimated total salmon harvest by place of residence, 1980–2012.....	129
Table 7-3.–Estimated subsistence salmon harvests by community, Chignik Area, 2012.....	129
Table 7-4.–Subsistence salmon harvests by species and subarea of harvest, Chignik Area, 2012.....	131
Table 7-5.–2012 Chignik area subsistence salmon harvests by species, fishing location, and date.....	132
Table 7-6.–Chignik area salmon removed from commercial catch for home use, 1994–2012.....	133
Table 8-1.–Historical subsistence salmon harvests, Alaska Peninsula Area, 1985–2012.....	144
Table 8-2.–Subsistence salmon harvest estimates by community, Alaska Peninsula Area, 2012.....	145
Table 8-3.–Percentage of households using selected nonsalmon finfishes, Alaska Peninsula Area communities.....	146
Table 9-1.–Historical subsistence salmon harvests, Unalaska District, 1985–2012.....	154
Table 9-2.–Estimated subsistence salmon harvests by community, Unalaska District, 2012.....	155
Table 9-3.–Historical subsistence and personal use salmon harvests, Adak District, 1988–2012.....	156
Table 9-4.–Estimated subsistence salmon harvests by community, Adak District, 2012.....	157

Table 9-5.–Estimated subsistence harvests of salmon by residents of Akutan, Atka, and Nikolski.....	157
Table 10-1.–Historical subsistence salmon harvests, Kodiak Area, 1986–2012.....	163
Table 10-2.–Reported subsistence salmon harvests by community and species, Kodiak Area, 2012.....	164
Table 10-3.–Permits returned and salmon harvests reported by the villages of Akhiok, Karluk, Larsen Bay, Old Harbor, Ouzinkie, and Port Lions.....	165
Table 11-1.–Historical subsistence salmon harvests, Port Graham and Koyuktolik subdistricts, 1981–2012.....	177
Table 11-2.–Subsistence salmon harvests by community, Port Graham and Koyuktolik subdistricts, 2012.....	178
Table 11-3.–Subsistence salmon harvests by community, Seldovia, 2012.....	178
Table 11-4.–Historical subsistence salmon harvests, Seldovia, 1996–2012.....	178
Table 11-5.–Subsistence salmon harvests by community, Tyonek Subdistrict, 2012.....	179
Table 11-6.–Historical subsistence salmon harvests, Tyonek Subdistrict, 1981–2012.....	179
Table 11-7.–Subsistence salmon harvests by community, Upper Yentna River, 2012.....	180
Table 11-8.–Historical subsistence and personal use salmon harvests, Upper Yentna River, 1996–2012.....	181
Table 11-9.–Federal subsistence salmon harvests by community, Kenai and Kasilof rivers, 2012.....	181
Table 11-10.–Historical federal subsistence salmon harvests, Kenai and Kasilof rivers, 2007–2012.....	182
Table 11-11.–Miscellaneous Upper Cook Inlet personal use and subsistence salmon harvests, 1981–1995.....	182
Table 11-12.–Cook Inlet personal use salmon fisheries, 2012.....	183
Table 11-13.–Estimated personal use salmon harvests, Upper Cook Inlet personal use fishery total, 1996–2012.....	184
Table 11-14.–Personal use salmon harvest estimates by community, Upper Cook Inlet, 2012.....	185
Table 11-15.–Estimated personal use salmon harvests, Kasilof River setnet fishery, 1982–2012.....	188
Table 11-16.–Estimated personal use salmon harvests, Kasilof River dip net fishery, 1981–2012.....	189
Table 11-17.–Estimated personal use salmon harvests, Kenai River dip net fishery, 1981–2012.....	191
Table 11-18.–Estimated personal use salmon harvests, Fish Creek dip net fishery, 1987–2012.....	192
Table 11-19.–Estimated personal use salmon harvests, unknown fishery, 1996–2012.....	193
Table 11-20.–Beluga River senior personal use dip net fishery summary, 2008–2012.....	193
Table 11-21.–Personal use/subsistence salmon harvests, Kachemak Bay setnet fishery (excluding the Port Graham/Nanwalek subsistence fishery and the Seldovia subsistence fishery), Lower Cook Inlet, 1969–2012.....	194
Table 11-22.–Estimated personal use salmon harvests, China Poot dip net fishery, 1980–1995.....	195
Table 12-1.–Subsistence harvests by village fish wheel permits, Glennallen Subdistrict, 1997–2012.....	209
Table 12-2.–Historical subsistence salmon harvests, Glennallen Subdistrict, 1989–2012.....	210
Table 12-3.–Subsistence salmon harvests by community, Glennallen Subdistrict, 2012.....	211
Table 12-4.–Historical subsistence and personal use salmon harvests, state Chitina Subdistrict permits, 1989–2012.....	213
Table 12-5.–Personal use salmon harvests by community, state Chitina Subdistrict permits, 2012.....	214
Table 12-6.–Historical subsistence salmon harvests, federal Chitina Subdistrict permits, 2003–2012.....	217
Table 12-7.–Subsistence salmon harvests by community, federal Chitina Subdistrict permits, 2012.....	217
Table 12-8.–Historical subsistence salmon harvests, Batzulnetas fishery, 1987–2012.....	218
Table 12-9.–Historical subsistence salmon harvests, Copper River District (Copper River Flats), 1965–2012.....	219
Table 12-10.–Subsistence salmon harvests by community, Copper River District (Copper River Flats), 2012.....	220
Table 12-11.–Historical subsistence salmon harvests, Prince William Sound, Eastern District, 1988–2012.....	221
Table 12-12.–Estimated harvests of salmon for home use, Tatitlek, 2003.....	222
Table 12-13.–Historical subsistence salmon harvests, Prince William Sound, Southwestern District, 1988–2012.....	222
Table 12-14.–Estimated harvests of salmon for home use, Chenega Bay, 2003.....	223
Table 12-15.–Historical subsistence salmon harvests, Prince William Sound general, 1960–2012.....	224
Table 12-16.–Subsistence salmon harvests by community, Prince William Sound general, 2012.....	225
Table 13-1.–Subsistence and personal use salmon harvests by district, Southeast region, 2012.....	246
Table 13-2.–Historical subsistence and personal use salmon harvests, Southeast region, 1985–2012.....	247
Table 13-3.–Estimated subsistence and personal use salmon harvests by management and use areas, Southeast region, 2012.....	248
Table 13-4.–Subsistence and personal use salmon harvests by community, Southeast region, 2012.....	249
Table 13-5.–Subsistence salmon harvests by community for the Federal Stikine River subsistence salmon fishery, Southeast region, 2012.....	250
Table 13-6.–Historical subsistence salmon harvests by community for the Federal Stikine River subsistence salmon fishery, Southeast region, 2004–2012.....	250

TABLE OF FIGURES

Figure 1-1.—Alaska subsistence fisheries areas.....	5
Figure 2-1.—Composition of subsistence harvest by rural Alaska residents, 2012.....	22
Figure 2-2.—Alaska subsistence salmon harvest by species, 2012.....	22
Figure 2-3.—Alaska subsistence salmon harvest by area, 2012.....	23
Figure 2-4.—Subsistence Chinook salmon harvest by area, 2012.....	23
Figure 2-5.—Subsistence sockeye salmon harvest by area, 2012.	24
Figure 2-6.—Subsistence chum salmon harvest by area, 2012.	24
Figure 2-7.—Subsistence coho salmon harvest by area, 2012.....	25
Figure 2-8.—Subsistence pink salmon harvest by area, 2012.	25
Figure 2-9.—Alaska personal use salmon harvest by species, 2012.	26
Figure 2-10.—Alaska subsistence and personal use salmon harvest by species, 2012.....	26
Figure 3-1.—Species composition of estimated subsistence salmon harvests, Norton Sound District, 2012.	49
Figure 3-2.—Species composition of estimated subsistence salmon harvests, Port Clarence District, 2012.	49
Figure 3-3.—Species composition of estimated subsistence salmon harvests, Kotzebue District, 2012.....	50
Figure 3-4.—Species composition of estimated subsistence salmon harvests, Arctic District, 2012.....	50
Figure 4-1.—Map of the Alaska portion of the Yukon River drainage, showing communities and districts.	76
Figure 4-2.—Yukon Area estimated subsistence salmon harvests, 2012.	77
Figure 4-3.—Estimated subsistence salmon harvests by species, Yukon Area, 1988–2012.	78
Figure 4-4.—Estimated number of dogs by district, Yukon Area, 2012.	79
Figure 4-5.—Primary gear type utilized for subsistence salmon fishing, Yukon Area, 2012.	79
Figure 5-1.—Kuskokwim subsistence salmon harvest composition, 2012.	102
Figure 6-1.—Bristol Bay Area subsistence salmon harvest composition, 2012.....	115
Figure 6-2.—Bristol Bay Area subsistence salmon harvests by district, 2012.....	115
Figure 7-1.—Location of Chignik Management Area (CMA) and communities within the CMA on Alaska Peninsula.	136
Figure 8-1.—Composition of Alaska Peninsula Area subsistence salmon harvest by species, 2012.	147
Figure 8-2.—Subsistence salmon harvests by community, Alaska Peninsula Area, 2012.	147
Figure 9-1.—Composition of Unalaska District estimated subsistence salmon harvest by species, 2012.	158
Figure 10-1.—Kodiak Area map, 2012.	166
Figure 10-2.—Subsistence salmon harvests by community, Kodiak Area, 2012.....	167
Figure 10-3.—Composition of Kodiak Area subsistence salmon harvest by species, 2012.....	167
Figure 10-4.—Salmon retained from commercial harvests for home use, Kodiak Area, 2012.	168
Figure 11-1.—Anchorage Nonsubsistence Area map.....	196
Figure 11-2.—Subsistence salmon harvests in the Port Graham and Koyuktolik subdistricts, 2012.	197
Figure 11-3.—Subsistence salmon harvests in Seldovia, 2012.	197
Figure 11-4.—Subsistence salmon harvests in the Tyonek Subdistrict, 2012.	198
Figure 11-5.—Permits issued, by place of residence, for the Upper Yentna River fishery, 2012.	198
Figure 11-6.—Subsistence salmon harvests in the Upper Yentna River, 2012.	199
Figure 13-1.—Juneau Nonsubsistence Area map, 2012.....	251
Figure 13-2.—Ketchikan Nonsubsistence Area map, 2012.....	252
Figure 13-3.—Southeast region subsistence and personal use harvests by species, 2012.....	253

ABSTRACT

Each year thousands of Alaskans participate in subsistence activities including the harvest of wild resources from Alaska's fisheries. Subsistence fishing is an important element of Alaska's social and cultural heritage, as well as a crucial component of the subsistence sector of the state's economy. In addition, personal use fisheries provide opportunities for Alaska residents to harvest salmon for home use using efficient methods in areas closed to subsistence fishing. This report summarizes Alaska's 2012 subsistence and personal use salmon fisheries based upon subsistence and personal use permit data and harvest assessment surveys from across the state. New information is compared to findings from previous years and the results are discussed. Where available, information about other subsistence finfish fisheries is included. Additional information from federal agencies regulating and administering certain subsistence fisheries, beginning in 1997, is included where available.

Key words: Pacific salmon, *Oncorhynchus* spp., sheefish, whitefish, rainbow/steelhead trout, Arctic char/Dolly Varden, northern pike, Chinook salmon, coho salmon, sockeye salmon, pink salmon, chum salmon, Norton Sound, Port Clarence, Kotzebue, Yukon, Kuskokwim, Bristol Bay, Chignik, Alaska Peninsula, Aleutian Islands, Kodiak, Cook Inlet, Prince William Sound, Southeast Alaska, Yakutat, subsistence salmon fisheries, personal use salmon fisheries

CHAPTER 1: INTRODUCTION

This is the fourteenth report in a series of annual reports on Alaska's subsistence and personal use fisheries. It was prepared by the Alaska Department of Fish and Game (ADF&G) Division of Subsistence.

Alaska state law defines subsistence fishing as the taking of fish, shellfish, or other fisheries resources by Alaska residents for subsistence uses (AS 16.05.940 (31)). Subsistence uses of wild resources are defined as "noncommercial, customary and traditional uses" for a variety of purposes. These include:

... direct personal or family consumption as food, shelter, fuel, clothing, tools, or transportation, for the making and selling of handicraft articles out of nonedible by-products of fish and wildlife resources taken for personal or family consumption, and for the customary trade, barter, or sharing for personal or family consumption ... (AS 16.05.940 (33))

Under Alaska's subsistence statute, the Alaska Board of Fisheries (BOF) must identify fish stocks that support subsistence fisheries. The BOF applies the Joint Board of Fisheries and Game Subsistence Procedures (5 AAC 99.010) to make these determinations, which are called "customary and traditional (C&T) findings." If there is a harvestable surplus of these stocks with C&T uses, the BOF must adopt regulations that provide reasonable opportunities for subsistence uses. When it is necessary to restrict harvests, the statute directs the BOF to assign a preference to subsistence uses (AS 16.05.258).

The Joint Board of Fisheries and Game (Joint Board) is required to identify "nonsubsistence areas," where "dependence upon subsistence is not a principal characteristic of the economy, culture, and way of life of the area or community" (AS 16.05.258 (c)). The Joint Board has identified 5 nonsubsistence areas: the Ketchikan Nonsubsistence Area, the Juneau Nonsubsistence Area, the Anchorage–Matsu–Kenai Nonsubsistence Area, the Fairbanks Nonsubsistence Area, and the Valdez Nonsubsistence Area (5 AAC 99.015). The BOF may not authorize subsistence fisheries in nonsubsistence areas.

Alaska state law recognizes 3 additional categories of fishing: commercial, sport, and personal use. Commercial fishing is the taking of fish "with the intent of disposing of them for profit, or by sale, barter, trade, or in commercial channels" (AS 16.05.940 (5)).

Sport fishing is defined as the taking "for personal use, and not for sale or barter, any fresh water, marine, or anadromous fish by hook and line held in the hand, or by hook and line with the line attached to a pole or rod which is held in the hand or closely attended, or by other means defined by the Board of Fisheries" (AS 16.05.940 (30)).

Personal use fishing is defined as the taking of fish "by Alaska residents for personal use and not for sale or barter, with gill or dip net, seine, fish wheel, long line, or other means defined by the Board of Fisheries" (AS 16.05.940 (25)). Personal use fisheries differ from subsistence fisheries in that they do not meet the criteria for customary and traditional fisheries as established by the Joint Board (5 AAC 99.010), or because they occur within nonsubsistence areas; in addition, a sport fishing license is required. This type of fishery provides Alaska residents with opportunities to harvest fish for noncommercial purposes, utilizing gear other than rod and reel, within nonsubsistence areas.

Every year, the ADF&G Division of Commercial Fisheries prepares Fishery Management Reports (FMRs, formerly "annual management reports," AMRs) for most fishery management areas in the state. Figure 1-1 shows the location of these management areas. Although the FMRs focus primarily on commercial fisheries, most also routinely summarize basic data for programs that collect harvest information for subsistence fisheries and for those personal use fisheries that the Division of Commercial Fisheries administers. Detailed annual reports about subsistence fisheries harvest assessment programs are prepared in the Northwest Alaska, Yukon River, and Kuskokwim River areas. Additionally, the Division of Sport Fish prepares summaries for the personal use salmon fisheries it administers in the Cook Inlet

and Prince William Sound (Upper Copper River) areas. However, until the Division of Subsistence annual subsistence fisheries report series began in 1999, there was no single source that compiled subsistence and personal use fisheries harvest data from all management areas. That is the purpose of this 2012 annual report.

The Federal Subsistence Board (FSB) adopts subsistence fishing regulations for federal waters in Alaska, in compliance with the Alaska National Interest Lands Conservation Act (ANILCA). Only eligible Alaska rural residents may participate in federal subsistence fisheries. For most subsistence fisheries, a single program administered by ADF&G provides harvest estimates for all participants regardless of the location of effort. However, for some fisheries (such as the Pacific salmon *Oncorhynchus* fisheries of the Upper Copper River District), FSB regulations require a federal subsistence fishing permit. The following chapters on each management area note where separate state and federal harvest monitoring programs operate. Subsistence harvest estimates in this report include data from both state and federal permit programs.

It is important to recognize the limitations associated with the effort to present a comprehensive annual report on Alaska's subsistence and personal use fisheries. These limitations include:

- Annual harvest assessment programs do not take place for all subsistence fisheries. Programs are in place for most salmon fisheries, but few other finfish fisheries or shellfish fisheries have annual harvest monitoring programs.
- Annual harvest data summarized in this report are limited to fisheries classified as subsistence or personal use by regulation, which, especially for salmon, generally means fish taken with gillnets, beach seines, dip nets, or fish wheels. In some parts of Alaska, substantial numbers of fish for home uses are taken with rod and reel (considered sport gear by most state area regulations) or are retained from commercial harvests. With noted exceptions, these harvests are not included in the analysis of subsistence harvest data in this report because they are not collected by annual subsistence fisheries harvest programs. Therefore, the harvest data in this report are a conservative estimate of the number of salmon being taken for subsistence uses in Alaska. Underestimations of subsistence salmon harvests are a particular issue in the Southeast region.
- Between management areas, and sometimes between districts within management areas, there is inconsistency in how subsistence and personal use harvest data are collected, analyzed, and reported.
- In some areas there are no routine mechanisms for evaluating the quality of subsistence harvest data. For example, in some areas it is not known if all subsistence fishers are obtaining permits and providing accurate harvest reports. This can result in a significant underestimation of harvests.
- There are also few programs for contextualizing annual subsistence harvest data so as to interpret changes in harvests. In some cases, however, FMRs do contain discussions of data limitations and harvest trends.

Despite these limitations, it is nonetheless possible to present an informative, conservative statewide overview of subsistence and personal use harvests of salmon. Information for all areas of the state where subsistence and personal use salmon fisheries occur is covered in this report. In past reports, we only included data for personal use salmon fisheries in the Yukon Management Area, the Prince William Sound Management Area (specifically, the Chitina Subdistrict of the Upper Copper River District), and the Southeast region because these fisheries were classified as subsistence fisheries in the past, and are administered in programs that collect subsistence harvest data. We did not include data from the Cook Inlet Management Area personal use salmon fisheries in past statewide overviews, primarily because most of these fisheries have relatively short histories. However, beginning in the report for 2010, we

added harvest data from the Cook Inlet personal use salmon fisheries so as to provide a complete statewide summary for all subsistence and personal use salmon harvests.

The quality and quantity of subsistence harvest data for finfish other than salmon and for shellfish are very uneven. For other finfish, if annual subsistence harvest information is collected, it is reflected in this report if the summary data were available to the Division of Subsistence. Otherwise, we have usually noted which species are primarily used for subsistence, relying in general on baseline studies conducted by the Division of Subsistence. In a small number of instances we have drawn from reports prepared for the BOF. This annual report does not attempt to provide a comprehensive overview of subsistence shellfish harvests.

In 1988, the Division of Subsistence prepared the first version of the Historical Subsistence Salmon Harvest Database (HSSHDB). As part of the cooperative agreement that supported the development of this annual report series, this database was updated, upgraded, and renamed the Alaska Subsistence Fisheries Database (ASFDB).¹ The database is written for Microsoft Access software.² It is organized by 21 subsistence fisheries and is generally reflective of unique harvest assessment programs and regulatory structures. It contains harvest data organized by species, year, community of residence of permit holder, and gear type. The number of permits issued and returned each year is included as well. The most complete data sets are sought; data sets which, in some cases, are more up-to-date than those reported in FMRs.

In 2008, the division received funding from the Alaska State Legislature to develop and annually update a web-based version of the ASFDB. This version of the database was developed using Microsoft SQL Server to store the data and Adobe ColdFusion 8 to create the user interface. The final product, projected to be available to the public in 2015, will contain all historical information from the HSSHDB along with contemporary data from the ASFDB stored in Microsoft Access, and will be updated periodically with new subsistence and personal salmon fisheries data.

The historical ASFDB is not currently available online. Upon request, the Division of Subsistence distributes the database on CD-ROM, along with the Community Subsistence Information System³, formerly the Community Profile Database (Scott et al. 2001). The CSIS includes the results of Division of Subsistence systematic household harvest surveys and is the primary source for subsistence harvest data for shellfish and for finfishes other than salmon.

In most fisheries data analysis, the Division of Subsistence expands harvest estimates from reported harvests in order to account for unreturned permits. In a few cases, this results in a larger estimate than is found in those FMRs that routinely only summarize data from returned permits. Also, the ASFDB calculates harvest estimates first for all permit holders by community represented in the fishery, and then sums these community estimates for a fishery total. This method is in contrast to the expansion method used by other divisions to analyze data from certain fisheries, such as the subsistence fishery in the Glennallen Subdistrict of the Prince William Sound Area. The harvest data analysis for this fishery presented in the FMR only considers the total number of issued and returned permits in expansion, resulting in slightly different estimates of total harvests than those in this report.

It is important to note that the preparation of this annual report and the supporting database were 2 objectives of the Statewide Subsistence Fisheries Harvest Monitoring Strategy project funded by the U.S. Fish and Wildlife Service (USFWS) Office of Subsistence Management (OSM) and implemented jointly by the Division of Subsistence and the Alaska Inter-Tribal Council (AITC). A central goal of the project was to develop recommendations for a unified subsistence harvest assessment program for Alaska's subsistence fisheries. A working group composed of state, federal, and tribal members developed these

1. David A. Caylor and Louis A. Brown. 2006. ASFDB. Alaska Department of Fish and Game Division of Subsistence, Juneau.

2. Product names are given for scientific completeness; they do not constitute product endorsement.

3. ADF&G Division of Subsistence, Community Subsistence Information System (CSIS): <http://www.subsistence.adfg.state.ak.us/CSIS/>.

recommendations. The recommendations are available as a separate document (ADF&G and AITC 2000) a final report with an overview of all the project activities is also available (Fall and Shanks 2000). The final report also includes comments on existing subsistence harvest assessment programs, based on working group discussions as well as interviews of ADF&G staff conducted by the Division of Subsistence. We have drawn on these comments for most of the evaluations of harvest data in this annual report. As background for the efforts of the working group, Division of Subsistence staff prepared detailed overviews of current subsistence fisheries harvest assessment programs. These are the basis of the program descriptions that appear in this report, with updates as necessary.

A final note regarding data ranges and averages: except where otherwise noted, averages in this report do not include the current data year (2012). Both date and numeric ranges are inclusive. The following list illustrates named-ranges used in this report and their meanings.

- 5-year average: 2007–2011
- 10-year average: 2002–2011
- 15-year average: 1997–2011
- Historical average: yyyy–2011, beginning of range varies depending on available data




Figure 1-1.—Alaska subsistence fisheries areas.

CHAPTER 2: OVERVIEW OF SUBSISTENCE FISHERIES IN ALASKA

SUBSISTENCE HARVESTS IN RURAL ALASKA

Of the estimated 36.9 million pounds of wild foods annually harvested for subsistence purposes in rural Alaska communities, subsistence fisheries contribute about 32% from salmon, 21% from other finfish and 3% from shellfish (Fall 2014:2, 3) (Figure 2-1). On average, the subsistence fisheries harvest provides about 167 lb of food per person annually in rural Alaska (Fall 2014:2). Although they constitute a major portion of the food supply, subsistence harvests represent just a small part of the annual harvest of wild resources in Alaska: about 1.1% (fish, game, and other resources combined). Commercial fisheries take 98.2% of the wild resource harvest, personal use fishing and general hunting by Alaskans take 0.2%, and sport fisheries and hunts take about 0.6% of the fish and game harvest.

SUBSISTENCE SALMON HARVESTS IN 2012

The estimated total subsistence harvest of salmon in Alaska in 2012, based on annual harvest assessment programs, was 935,470 fish (Table 2-1).⁴ The estimated statewide harvest by species was as follows: 344,071 sockeye salmon *O. nerka* (37%), 367,692 chum salmon *O. keta* (39%), 74,381 Chinook salmon *O. tshawytscha* (8%), 80,275 coho salmon *O. kisutch* (9%), and 69,051 pink salmon *O. gorbuscha* (7%) (Figure 2-2).

In 2012, fisheries in 8 management areas accounted for 95% of the total estimated statewide subsistence salmon harvest (Table 2-1; Figure 2-3). These were the Yukon Management Area (284,301 salmon; 30% of the statewide total); the Kuskokwim Management Area (190,245 salmon; 20%); the Bristol Bay Management Area (122,582 salmon; 13%); the Glennallen Subdistrict of the Prince William Sound Management Area (98,110 salmon; 11%); the Norton Sound-Port Clarence Area⁵ (91,696 salmon; 10%); Southeast Region⁶ (including the Stikine River federal fishery) (7,725 salmon; 5%); the Kotzebue District (29,092 salmon; 3%); and the Kodiak Management Area (28,159; 3%).

The largest estimated subsistence harvests of Chinook salmon in 2012 occurred in the Yukon Management Area (30,486 salmon; 41%), followed by the Kuskokwim Management Area (25,336 salmon; 34%), Bristol Bay Management Area (12,136 salmon; 16%), the Glennallen Subdistrict (2,649 salmon; 4%); and the Norton Sound-Port Clarence Area (1,335 salmon; 2%) (Figure 2-4). For sockeye salmon, the largest estimated subsistence harvests in 2012 were in the Bristol Bay Management Area (100,728 salmon; 29%), followed by the Glennallen Subdistrict (94,991 salmon; 27%), the Kuskokwim Management Area (50,616 salmon; 15%), the Southeast Region (40,007 salmon; 12%), and the Kodiak Management Area (23,865 salmon; 7%) (Figure 2-5).

In 2012, as in past recent years, 4 areas dominated the subsistence chum salmon estimated harvest: the Yukon Management Area (227,032 salmon; 62% of the statewide harvest), Kuskokwim Management Area (81,912 salmon; 22%), Kotzebue District (26,694 salmon; 7%, and the Norton Sound-Port Clarence

4. Annual reports prior to 2010 included personal use salmon harvests from Southeast Alaska and the Chitina Subdistrict of the Upper Copper River in the discussion of subsistence harvests. Beginning with the 2010 report, personal use salmon fisheries are discussed separately. One exception is the small personal use harvest that occurs in those portions of the Yukon Management Area that are within the Fairbanks Nonsubsistence Area. Also, as noted in Chapter 1, Cook Inlet Area personal use salmon harvest data have been added to the annual report.

5. Subsistence harvest estimates for Northwest Alaska (Norton Sound-Port Clarence Area and the Arctic-Kotzebue Area for 2003, 2004, and 2012 do not include the regional center of Kotzebue, which was included in the harvest assessment program for 1994–2002. No subsistence fisheries harvest data were collected in the Kotzebue District for 2005 through 2011. Therefore, the estimated harvest totals for Northwest Alaska as reported for 2003–2011 are incomplete. See also Chapter 3.

6. As discussed further in Chapter 13, state subsistence regulations for the Southeast Region focus on sockeye salmon. Small harvests of Chinook and coho salmon are reported on permit returns as incidental to sockeye salmon harvests. The major portion of coho and Chinook salmon harvests for home uses in Southeast is taken with rod and reel (sport gear). Thus the Southeast Region is particularly underrepresented in statewide overviews based on permit data.

Area (24,049 salmon; 7%) (the latter two areas were combined as “Northwest Alaska” in previous annual reports) (Figure 2-6). Of the statewide estimated subsistence harvest of coho salmon in 2012, the greatest share was taken in the Kuskokwim Management Area (30,221 salmon; 38%), followed by the Yukon Management Area (21,633 salmon; 27%), Norton Sound-Port Clarence Area (12,203 salmon; 15%), Bristol Bay Management Area (3,837 salmon; 5%), the Southeast Region (2,639 salmon; 3%), the Kodiak Management Area (2,920 salmon; 4%), and the Alaska Peninsula Management Area (1,936 salmon; 2%) (Figure 2-7). Finally, the largest portion by far of the statewide estimated pink salmon subsistence harvest in 2012 occurred in the Norton Sound-Port Clarence Area (52,250 salmon; 76%), followed by the Yukon Area (5,150 salmon; %), and the Kuskokwim Area (2,160 salmon; 3%) (Figure 2-8).

Table 2-2 reports historical estimated subsistence salmon harvests for 1994 through 2012 based on annual harvest assessment programs. While earlier estimates for many of the fisheries are available, 1994 marks the first year that data from all of the included fisheries were available and collected with methods comparable to those currently in use.

The 19-year period reflected in Table 2-2 shows a general downward trend. Estimates from 2000 through 2008 suggested this trend might have been stabilizing. However, the 2011 estimate of 839,903 salmon was the fourth-lowest within the 19-year period, with the 775,642 salmon harvested in 2009 being the lowest estimate, and the 834,627 salmon harvested in 2010 the third-lowest, since 1994. The estimate for 2012 of 935,470 was the highest since 2008. The 2012 estimate was higher than the recent 5-year average (853,958 salmon) and the recent 10-year average (877,813 salmon), but lower than the historical average since 1994 (942,759 salmon). The collection of harvest data in the Kotzebue District for the first time since 2003 may account for this slight rise in the statewide subsistence harvest estimate compared to recent years. It should also be noted that the estimate of 74,381 Chinook salmon harvested in subsistence fisheries in 2012 is the lowest on record, and is just 47% of the annual average since 1994 and 42% below the next-lowest annual estimate (128,657 Chinook salmon in 2011).

PERSONAL USE SALMON HARVESTS IN 2012

In 2012, personal use fisheries produced an estimated harvest of 793,345 salmon (Table 2-1). The Kenai River dip net fishery accounted for 67% of the statewide personal use salmon harvest (535,235 fish), followed by the Chitina Subdistrict dip net fishery (17%; 138,465 salmon), the Kasilof River dip net fishery (10%; 75,648 salmon), the Kasilof River setnet fishery (2%; 15,970 salmon), the Southeast Region (2%; 12,213 salmon), and the Kachemak Bay set net fishery (<1%; 1,894 salmon). Sockeye salmon composed 98% of the Alaska personal use salmon harvest in 2012 (Figure 2-9).

The personal use harvest of 793,345 salmon in 2012 was the second-largest total since comprehensive records became available in 1994, just 402 fish less than the record of 793,747 salmon set in 2011 (Table 2-3). The average annual personal use harvest since 1994 of 446,132 salmon is 56% of the 2012 total. Increased harvests in the Upper Cook Inlet personal use dip net fisheries accounts for most of the growth of personal use harvests since 1994 (see Chapter 11).

STATEWIDE SUBSISTENCE AND PERSONAL USE SALMON HARVESTS, 1994–2012

Table 2-4 reports historical estimated subsistence and personal use salmon harvests for 1994 through 2012 based on annual harvest assessment programs. As noted above, 1994 marks the first year that comparable data from all of the included fisheries are available.

The 19-year period reflected in Table 2-4 shows generally stable statewide harvest totals: the recent (2007–2011) 5-year average harvest was 1,453,999 salmon compared to a 19-year annual average of 1,358,061 salmon. The total harvest estimate for 2012 of 1,728,815 salmon is the highest within the 19-year period. As noted above, however, harvests in subsistence fisheries have generally declined since 1994 while personal use harvests have increased. In 2012, sockeye salmon made up 65% of the combined

subsistence and personal use salmon harvests, followed by chum (21%), coho (5%), Chinook (4%), and pink salmon (5%) (Figure 2-10).

Table 2-5 reports subsistence and personal use harvests in 2012 by species and participants' place of residence, with harvests from all subsistence and personal use fisheries combined.

Table 2-1.—Alaska subsistence and personal use salmon harvests, 2012.

Fishery	Households or permits		Estimated salmon harvest					
	Total ^a	Surveyed or returned	Chinook	Sockeye	Coho	Chum	Pink	Total
<i>Subsistence</i>								
Adak District	1	1	0	0	0	0	0	0
Alaska Peninsula Management Area	172	138	287	9,429	1,936	1,637	941	14,230
Arctic District ^b	219	120	34	79	477	710	1,256	2,556
Batzulnetas Fishery	3	3	1	136	0	0	0	137
Bristol Bay Management Area	1,107	932	12,136	100,728	3,837	4,007	1,874	122,582
Chignik Management Area	106	87	116	5,607	1,488	220	810	8,241
Chitina Subdistrict: Federal	90	80	5	981	9	0	0	995
Copper River Flats	378	359	248	4,499	0	19	0	4,766
Glennallen Subdistrict	1,805	1,557	2,649	94,991	470	0	0	98,110
Kenai and Kasilof Rivers: Federal	133	121	0	1,438	0	0	0	1,438
Kodiak Management Area ^a	1,866	1,866	54	23,865	2,920	166	1,154	28,159
Kotzebue District ^b	545	360	16	455	1,230	26,694	697	29,092
Kuskokwim Management Area	4,294	1,569	25,336	50,616	30,221	81,912	2,160	190,245
Norton Sound - Port Clarence Area ^b	1,270	1,234	1,335	1,859	12,203	24,049	52,250	91,696
Port Graham & Koyuktolik Subdistricts ^a	8	8	24	961	414	31	482	1,912
Prince William Sound (General)	14	12	0	67	0	32	0	99
PWS Eastern District (Tatitlek)	16	8	15	954	75	8	0	1,052
PWS Southwestern District (Chenega Bay)	23	14	0	603	20	77	0	700
Seldovia Fishery	20	7	8	79	0	0	54	141
Southeast Region	2,944	2,530	718	40,007	2,639	987	1,828	46,179
Stikine River Federal Fishery	130	130	53	1,302	112	47	32	1,546
Tyonek Fishery	89	69	840	176	138	2	4	1,160
Unalaska District	211	169	20	4,960	429	43	338	5,790
Upper Yentna Fishery	21	21	0	279	24	19	21	343
Yukon Management Area ^c	3,133	1,575	30,486	0	21,633	227,032	5,150	284,301
Subtotal, Subsistence	18,598	12,970	74,381	344,071	80,275	367,692	69,051	935,470

-continued-

Table 2-1.—Page 2 of 2.

Table 2-1. Page 2 of 2.

Fishery	Households or permits		Fishery	Estimated salmon harvest				
	Total ^a	Surveyed or returned		Total ^a	Surveyed or returned	Fishery	Total ^a	
<i>Personal use</i>								
Chitina Subdistrict: State ^d	10,016	8,030	613	136,441	1,411	0	0	138,465
Kachemak Bay set net ^e	98	95	5	137	1,471	6	275	1,894
Kasilof River set net ^e	NA	NA	103	15,638	161	15	53	15,970
Kasilof River dip net ^e	NA	NA	16	73,419	1,170	147	896	75,648
Kenai River dip net ^e	NA	NA	40	526,992	4,008	425	3,770	535,235
Fish Creek dip net ^e	NA	NA	--	--	--	--	--	--
Unknown Upper Cook Inlet ^e	NA	NA	8	13,548	173	40	135	13,904
Beluga River dip net	7	7	0	9	7	0	0	16
Southeast Region	323	323	45	10,420	571	199	978	12,213
Subtotal, Personal use^e	44,759	35,535	830	776,604	8,972	832	6,107	793,345
Total	63,357	48,505	75,211	1,120,675	89,247	368,524	75,158	1,728,815

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Note Included in this table are all harvest estimates based upon annual harvest monitoring programs.

- Because the numbers of permits issued for the Kodiak and Port Graham/Koyuktolik fisheries are unknown, the numbers of permits returned are used in place of these values.
- Formerly included within Northwest Alaska. Partial coverage for Arctic and Kotzebue Districts; see Chapter 3 for details.
- Includes a small personal use harvest that occurs within the Fairbanks Nonsubsistence Area.
- Reclassified as a personal use fishery in 2003.
- A single permit is issued for the Kasilof set net, Kasilof dip net, Kenai dip net, and Fish Creek dip net fisheries. In some cases, returned permits did not indicate the area fished.

NA = Data not available.

Table 2-2.—Historical Alaska subsistence salmon harvests, 1994–2012.

Year	Households or permits		Estimated salmon harvest					
	Total	Surveyed or returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1994	15,493	10,553	183,936	338,946	135,896	417,199	94,469	1,170,446
1995	15,596	10,328	180,805	291,539	120,048	499,992	54,908	1,147,292
1996	16,512	11,789	158,369	320,821	121,381	498,525	80,928	1,180,026
1997	17,668	12,863	176,703	376,397	98,883	347,808	41,543	1,041,335
1998	17,772	12,513	170,271	328,857	93,055	302,037	74,216	968,436
1999	17,290	12,763	155,088	358,866	89,627	338,351	32,402	974,334
2000	16,678	12,765	130,822	296,875	99,338	247,337	51,714	826,087
2001	18,693	13,061	161,632	340,411	98,517	240,581	42,435	883,576
2002	17,266	13,026	142,459	299,182	92,192	229,179	85,431	848,443
2003	18,131	13,211	164,555	324,539	106,488	238,582	66,794	900,958
2004	18,374	13,549	173,746	332,543	100,860	239,811	91,597	938,557
2005	16,256	11,013	153,431	323,218	97,993	257,200	76,071	907,912
2006	16,988	11,400	139,815	314,435	93,478	291,510	73,234	912,473
2007	17,068	10,374	154,974	319,885	78,704	273,802	33,513	860,877
2008	17,226	11,248	174,115	315,040	113,242	270,502	85,842	958,741
2009	16,989	11,607	141,302	296,104	86,363	213,835	38,038	775,642
2010	16,020	11,381	133,252	326,363	80,217	235,763	59,031	834,627
2011	17,181	12,155	128,657	341,388	77,180	257,032	35,646	839,903
2012	18,598	11,970	74,381	344,071	80,275	367,692	69,051	935,470
5-year average (2007–2011)	16,897	11,353	146,460	319,756	87,141	250,187	50,414	853,958
10-year average (2002–2011)	17,150	11,896	150,630	319,270	92,672	250,722	64,520	877,813
Historical average (1994–2011)	17,067	11,978	156,885	324,745	99,081	299,947	62,101	942,759

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Note Included in this table are all harvest estimates based upon annual harvest monitoring programs.

Table 2-3.—Historical Alaska personal use salmon harvests, 1994–2012.

Year	Households or permits		Estimated salmon harvest					
	Total	Surveyed or returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1994	7,346	6,223	5,524	142,944	15,810	1,619	2,831	168,729
1995	6,997	5,674	7,029	139,861	18,455	1,672	1,579	168,596
1996	22,071	20,707	4,360	241,293	11,562	374	3,995	261,585
1997	24,281	22,939	6,318	298,151	2,753	100	1,101	308,424
1998	25,764	23,155	7,430	314,131	6,302	225	2,100	330,187
1999	27,907	24,587	7,630	360,885	5,485	1,062	3,097	378,159
2000	25,007	22,006	4,653	274,422	9,576	1,555	3,782	293,988
2001	27,017	23,392	4,631	365,875	6,990	1,746	4,037	383,279
2002	24,921	20,560	3,449	358,608	6,965	1,512	10,044	380,578
2003	26,101	21,707	3,766	394,928	6,004	1,446	3,387	409,532
2004	30,673	25,205	3,775	470,804	8,220	1,729	3,571	488,100
2005	30,817	26,677	3,367	508,419	6,350	1,218	3,776	523,130
2006	27,545	23,772	4,263	354,130	7,600	1,212	13,741	380,946
2007	31,855	27,922	4,773	496,317	6,139	797	4,267	512,294
2008	32,582	27,935	3,646	410,298	7,991	927	13,051	435,913
2009	38,443	32,800	1,654	558,352	6,872	873	7,705	575,456
2010	41,505	33,580	1,826	660,892	11,475	1,212	7,393	682,797
2011	44,208	35,265	2,661	773,540	9,714	1,461	6,371	793,747
2012	44,759	35,535	830	776,604	8,972	832	6,107	793,345
5-year average (2007–2011)	37,719	31,500	2,912	579,880	8,438	1,054	7,757	600,041
10-year average (2002–2011)	32,865	27,542	3,318	498,629	7,733	1,239	7,331	518,249
Historical average (1996–2011)	30,044	25,763	4,263	427,565	7,500	1,091	5,714	446,132

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Note Included in this table are all harvest estimates based upon annual harvest monitoring programs.

Table 2-4.—Historical Alaska subsistence and personal use salmon harvests, 1994–2012.

Year	Households or permits		Estimated salmon harvest					
	Total	Surveyed or returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1994	22,839	16,776	189,460	481,890	151,707	418,818	97,300	1,339,175
1995	22,593	16,002	187,834	431,401	138,503	501,664	56,487	1,315,888
1996	38,583	32,496	162,730	562,114	132,944	498,900	84,923	1,441,611
1997	41,949	35,802	183,022	674,548	101,637	347,909	42,644	1,349,759
1998	43,536	35,668	177,701	642,987	99,357	302,262	76,316	1,298,623
1999	45,197	37,350	162,717	719,752	95,112	339,413	35,499	1,352,493
2000	41,685	34,771	135,476	571,297	108,914	248,892	55,496	1,120,074
2001	45,710	36,453	166,263	706,285	105,507	242,327	46,472	1,266,854
2002	42,187	33,586	145,908	657,790	99,157	230,691	95,475	1,229,021
2003	44,232	34,918	168,321	719,467	112,493	240,028	70,181	1,310,489
2004	49,047	38,754	177,521	803,348	109,080	241,540	95,168	1,426,657
2005	47,073	37,690	156,798	831,637	104,343	258,418	79,847	1,431,042
2006	44,533	35,172	144,078	668,565	101,078	292,722	86,975	1,293,419
2007	48,923	38,296	159,747	816,202	84,843	274,599	37,780	1,373,171
2008	49,808	39,183	177,761	725,338	121,233	271,429	98,893	1,394,654
2009	55,432	44,407	142,956	854,456	93,235	214,708	45,743	1,351,098
2010	57,525	44,961	135,078	987,255	91,692	236,975	66,424	1,517,424
2011	61,389	47,420	131,318	1,114,928	86,894	258,493	42,017	1,633,650
2012	63,357	47,505	75,211	1,120,675	89,247	368,524	75,158	1,728,815
5-year average (2007–2011)	54,615	42,853	149,372	899,636	95,579	251,241	58,171	1,453,999
10-year average (2002–2011)	50,015	39,439	153,949	817,899	100,405	251,960	71,850	1,396,063
Historical average (1994–2011)	44,569	35,539	161,372	720,514	107,652	301,099	67,424	1,358,061

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Note Included in this table are all harvest estimates based upon annual harvest monitoring programs.

Table 2-5.—Alaska subsistence salmon harvests by species and place of residence, 2012.

Community	Households or permits		Estimated salmon harvest					
	Total	Included ^a	Chinook	Sockeye	Coho	Chum	Pink	Total
Adak	1	1	0	0	0	0	0	0
Akhiok	3	3	2	128	21	0	7	158
Akiachak	158	75	2,862	3,468	714	4,150	53	11,247
Akiak ^d	79	16	856	1,820	474	2,416	0	5,566
Alakanuk	158	54	1,081	0	252	9,461	174	10,968
Alatna	10	4	0	0	0	118	0	118
Aleknagik	29	21	696	1,548	108	86	19	2,457
Allakaket	65	22	5	0	38	4,358	0	4,401
Ambler	76	53	1	126	11	1,621	9	1,769
Anaktuvuk Pass	1	1	0	36	0	0	0	36
Anchor Point	316	269	6	4,958	45	1	22	5,031
Anchorage	19,595	15,314	1,033	347,847	3,413	508	3,114	355,915
Anderson	8	6	0	116	0	0	2	118
Angoon	98	77	0	1,308	51	0	51	1,410
Aniak	189	157	994	1,404	3,365	5,667	940	12,370
Anvik	35	27	435	0	214	1,940	0	2,589
Atmautluak	61	35	234	1,623	383	2,701	22	4,963
Atkasuk	2	0	0	28	0	0	0	29
Auke Bay	9	9	0	144	2	24	0	170
Barrow	72	38	86	1,754	6	7	12	1,864
Beaver	31	24	71	0	2	201	0	274
Bethel	2,147	462	7,321	18,596	13,281	26,872	310	66,380
Bettles	24	18	3	14	0	7	0	25
Big Lake	246	203	11	4,217	33	1	14	4,276
Birch Creek	16	12	0	0	0	0	0	0
Brevig Mission	45	45	11	388	597	3,321	3,095	7,412
Cantwell	17	15	0	300	0	0	2	303
Central	8	7	66	3	0	0	0	69
Chalkyitsik	28	18	0	0	0	162	0	162
Chefornak	1	0	0	14	0	0	0	15
Chenega Bay	16	15	0	604	20	77	0	701
Chevak	5	3	0	28	2	0	0	31
Chickaloon	30	25	1	625	0	0	4	631
Chignik Bay	7	7	11	385	105	12	19	532
Chignik Lagoon	20	18	61	1,771	12	2	37	1,884
Chignik Lake	16	13	16	1,338	45	3	41	1,442
Chiniak	28	28	0	213	57	1	10	281
Chistochina	5	5	23	711	0	0	0	734
Chitina	36	34	85	2,336	11	0	0	2,432
Chuathbaluk	34	28	103	297	179	796	2	1,377
Chugiak	888	763	50	16,833	127	5	69	17,084
Circle	19	19	280	0	5	161	0	446
Clam Gulch	54	47	0	902	12	10	7	932
Clarks Point	13	13	99	365	189	80	149	882

-continued-

Table 2-5–Page 2 of 7

Community	Households or permits		Estimated salmon harvest					
	Total	Included ^a	Chinook	Sockeye	Coho	Chum	Pink	Total
Clear	12	10	0	220	0	0	0	221
Coffman Cove	10	7	0	35	0	0	0	35
Cold Bay	24	21	2	838	21	5	0	866
Cooper Landing	112	105	0	1,501	0	0	6	1,508
Copper Center	151	123	387	10,640	82	0	0	11,109
Copperville	5	5	12	333	0	0	0	345
Cordova	305	292	207	3,738	0	0	0	3,946
Craig	162	118	0	1,159	67	27	105	1,358
Crooked Creek	37	31	124	234	149	610	2	1,119
Deering	1	1	0	63	0	0	0	63
Delta Junction	496	454	94	9,148	123	0	3	9,369
Denali National Park	53	47	5	781	4	0	10	801
Dillingham	332	279	5,055	12,959	1,420	1,331	652	21,417
Diomedes	1	1	0	0	0	0	0	0
Dot Lake	4	3	0	32	0	0	0	32
Douglas	57	54	5	385	35	1	42	469
Dutch Harbor	112	94	14	2,847	136	7	75	3,079
Eagle	34	32	167	0	0	18,731	0	18,898
Eagle River	2,390	2,051	184	44,662	309	51	281	45,486
Eek	86	45	1,004	1,490	612	1,552	50	4,708
Egegik	9	6	0	66	104	0	0	170
Eielson AFB	105	68	3	1,546	4	0	0	1,554
Ekwok	15	13	681	167	59	234	112	1,253
Elfin Cove	1	0	0	0	0	0	0	0
Elim	55	55	41	45	1,281	1,465	10,379	13,211
Elmendorf AFB	31	25	2	213	0	0	0	216
Emmonak	180	92	1,864	0	2,660	21,719	199	26,442
Ester	105	86	20	2,071	15	0	0	2,107
Excursion Inlet	3	3	0	1	18	0	0	19
Fairbanks	4,074	3,322	1,094	70,380	2,491	5,689	86	79,741
False Pass	4	4	10	288	29	42	27	396
Fort Greely	27	20	1	407	0	0	2	411
Fort Richardson	9	4	0	0	0	0	0	0
Fort Wainwright	113	78	5	1,498	16	0	7	1,526
Fort Yukon	214	89	2,141	23	4	12,659	0	14,827
Fox	3	2	0	0	0	0	0	0
Fritz Creek	72	62	0	1,326	2	1	4	1,334
Funny River	1	1	0	0	0	0	0	0
Gakona	40	38	41	2,818	59	0	0	2,919
Galena	173	55	742	96	276	3,666	3	4,783
Gambell	4	3	0	14	0	0	0	15
Girdwood	304	248	5	5,003	98	3	38	5,148
Glacier View	2	2	6	84	0	0	0	90
Glennallen	114	103	203	5,809	57	0	0	6,070

-continued-

Table 2-5–Page 3 of 7

Community	Households or permits		Estimated salmon harvest					
	Total	Included ^a	Chinook	Sockeye	Coho	Chum	Pink	Total
Golovin	29	29	39	44	246	775	2,415	3,519
Goodnews Bay	69	38	389	1,217	382	322	72	2,382
Grayling	48	18	1,081	0	26	3,420	0	4,527
Gulkana	2	1	32	600	0	0	0	632
Gustavus	22	21	0	84	0	1	2	88
Haines	433	412	235	7,353	178	349	679	8,794
Healy	80	68	4	1,360	803	595	1	2,763
Hollis	42	36	0	610	81	2	361	1,054
Holy Cross	55	31	576	0	237	1,486	0	2,299
Homer	972	837	37	17,561	120	42	200	17,960
Hoonah	124	89	0	1,582	45	103	326	2,055
Hooper Bay	222	81	1,090	14	7	15,800	1,101	18,013
Hope	47	40	5	574	1	0	2	582
Houston	44	35	8	777	2	0	5	792
Hughes	31	25	0	0	0	430	0	430
Huslia	97	35	165	96	165	9,215	101	9,742
Hydaburg	55	43	0	1,761	27	0	9	1,797
Igiugig	13	9	0	2,711	0	0	0	2,711
Iliamna	31	24	3	8,208	0	0	0	8,211
Indian	7	7	0	118	0	0	0	118
Ivanof Bay	2	2	1	70	182	27	32	312
Joint Base Elmendorf Richardson	297	216	0	5,190	30	3	44	5,267
Juneau	646	592	47	8,855	340	27	254	9,522
Kake	142	115	48	1,001	16	33	136	1,235
Kaktovik	2	1	0	0	0	0	0	0
Kaltag	58	15	1,346	0	928	3,016	0	5,290
Karluk	2	2	0	33	6	0	0	39
Kasaan	15	12	0	153	28	1	0	181
Kasigluk	104	51	552	1,451	303	3,261	0	5,567
Kasilof	469	393	19	7,911	59	16	47	8,052
Kenai	1,760	1,425	37	30,164	161	17	140	30,519
Kennicott	3	2	0	118	0	0	0	118
Kenny Lake	47	46	69	2,487	42	0	0	2,598
Ketchikan	273	225	5	3,032	54	303	297	3,690
Kiana	104	65	3	78	240	2,442	320	3,083
King Cove	57	46	52	3,643	1,100	438	28	5,260
King Salmon	81	74	173	5,329	49	17	100	5,667
Kipnuk	1	1	5	25	0	0	0	30
Klawock	123	91	0	2,597	205	33	65	2,900
Klukwan	10	9	1	414	39	72	20	547
Kobuk	37	31	4	40	14	2,637	4	2,699
Kodiak (city)	1,462	1,455	114	19,934	1,910	109	531	22,597
Kokhanok	27	20	161	16,593	0	0	1	16,755
Koliganek	15	13	852	835	361	579	207	2,834

-continued-

Table 2-5–Page 4 of 7

Community	Households or permits		Estimated salmon harvest					
	Total	Included ^a	Chinook	Sockeye	Coho	Chum	Pink	Total
Kongiganak ^b	90	0	571	1,211	458	1,901	0	4,141
Kotlik	111	38	1,173	0	420	9,625	195	11,413
Kotzebue	17	13	0	337	10	0	4	351
Koyuk	83	82	104	0	373	2,731	2,837	6,045
Koyukuk	49	22	614	0	62	2,159	0	2,835
Kwethluk	165	84	1,709	2,886	1,013	3,849	91	9,548
Lake Louise	1	1	0	44	0	0	0	44
Lake Minchumina	1	1	0	30	0	0	0	30
Larsen Bay	17	17	1	387	27	0	16	431
Levelock	4	3	0	841	0	0	0	841
Lime Village	14	10	29	780	117	419	129	1,474
Lower Kalskag	79	41	459	891	1,107	3,284	25	5,766
Manley Hot Springs	19	16	174	25	1,374	2,222	0	3,795
Manokotak	13	9	143	952	106	20	7	1,227
Marshall	69	26	1,409	0	567	6,087	5	8,068
McCarthy	34	26	0	192	9	2	0	203
McGrath	143	50	68	314	2,269	885	14	3,551
Meadow Lakes	2	1	0	20	0	0	0	20
Mekoryuk	1	1	0	25	0	0	0	25
Mendeltna	3	3	2	174	0	0	0	176
Mentasta Lake	4	4	1	163	0	0	0	164
Metlakatla	3	2	0	92	0	14	4	110
Minto	40	35	99	26	0	66	0	192
Moose Creek	1	1	0	0	0	0	0	0
Moose Pass	28	25	0	481	1	0	1	483
Mountain Village	152	57	1,789	0	256	9,716	207	11,968
Nabesna	3	3	0	91	0	0	0	91
Naknek	107	85	273	10,343	227	49	207	11,099
Nanwalek	3	2	0	338	400	5	200	944
Napakiak	100	46	457	1,155	402	1,711	0	3,726
Napaskiak	97	42	1,108	2,065	269	3,216	122	6,780
Naukati Bay	3	3	0	0	0	0	0	0
Nelchina	5	5	0	358	0	0	0	358
Nelson Lagoon	9	4	9	788	97	11	2	907
Nenana	84	76	300	919	5,913	9,041	0	16,174
New Stuyahok	39	26	2,439	1,778	345	677	137	5,375
Newhalen	15	12	0	5,067	0	0	0	5,067
Newtok	1	0	0	14	0	0	0	15
Nikiski	268	215	3	4,334	62	8	22	4,430
Nikolaevsk	18	16	0	425	0	0	2	428
Nikolai	34	30	276	0	214	1,044	0	1,534
Ninilchik	227	193	0	3,123	7	3	6	3,138
Noatak	126	83	2	94	612	7,814	80	8,601
Nome	493	492	16	1,312	1,724	3,168	10,387	16,607

-continued-

Table 2-5–Page 5 of 7

Community	Households or permits		Estimated salmon harvest					
	Total	Included ^a	Chinook	Sockeye	Coho	Chum	Pink	Total
Nondalton	31	30	0	9,327	0	0	0	9,327
Noorvik	137	83	7	110	338	9,584	275	10,314
North Pole	1,140	875	190	21,910	130	1	15	22,247
Northway	11	8	1	204	0	0	0	205
Nuiqsut	3	1	0	28	0	0	0	29
Nulato	74	25	1,955	2	41	2,983	0	4,981
Nunam Iqua (Sheldon Point)	42	34	195	0	18	2,187	1,051	3,451
Nunapitchuk	111	61	845	2,396	319	5,312	32	8,904
Old Harbor	20	20	3	604	248	77	253	1,185
Oscarville ^d	14	14	51	323	38	599	1	1,012
Other USA	7	7	0	116	36	1	2	155
Ouzinkie	32	31	1	954	438	5	71	1,469
Palmer	2,285	1,930	240	44,475	329	47	199	45,291
Paxson	1	1	1	114	0	0	0	115
Pedro Bay	15	14	0	4,028	0	0	0	4,028
Pelican	3	3	0	14	0	0	0	14
Perryville	35	27	26	1,607	1,094	172	679	3,578
Peters Creek	2	2	4	33	0	0	0	37
Petersburg	148	138	23	1,373	408	44	38	1,886
Pilot Point	6	5	18	307	60	24	0	409
Pilot Station	118	57	1,078	0	329	6,747	23	8,177
Pitkas Point	27	23	261	0	53	1,162	2	1,478
Platinum	19	16	24	173	124	76	16	413
Point Baker	3	3	0	0	0	0	0	0
Point Hope	4	2	0	46	0	0	0	47
Point Lay	67	42	14	13	372	659	1,120	2,178
Port Alexander	7	7	1	270	27	0	0	298
Port Alsworth	52	49	2	4,445	0	0	0	4,447
Port Graham	7	7	24	661	14	26	282	1,007
Port Heiden	5	4	29	193	64	55	0	340
Port Lions	40	40	0	1,277	160	0	236	1,673
Portage Creek	1	1	31	2	0	2	0	35
Prudhoe Bay	1	0	0	0	0	0	0	0
Quinhagak	163	77	2,396	2,029	1,380	2,001	70	7,877
Rampart	7	6	190	0	0	261	0	451
Red Devil	13	10	225	511	238	516	42	1,532
Ruby	66	21	1,316	0	1,806	8,299	0	11,421
Russian Mission	73	27	1,711	45	319	2,790	76	4,941
Saint Marys	129	50	2,344	0	141	12,186	643	15,314
Saint Michael	82	82	80	20	911	2,172	457	3,640
Saint Paul Island	3	3	0	134	0	0	0	134
Salcha	74	54	7	1,157	44	0	0	1,208
Sand Point	54	42	178	3,248	622	1,073	806	5,926
Savoonga	3	3	0	0	0	0	19	19

-continued-

Table 2-5–Page 6 of 7

Community	Households or permits		Estimated salmon harvest					
	Total	Included ^a	Chinook	Sockeye	Coho	Chum	Pink	Total
Saxman	21	18	0	335	0	23	13	371
Scammon Bay	99	44	1,014	0	86	7,452	1,343	9,895
Seldovia	34	19	10	302	4	11	55	382
Seward	276	228	2	4,658	20	7	14	4,702
Shageluk	29	11	75	0	0	5,051	24	5,150
Shaktolik	64	63	213	9	1,043	624	4,401	6,290
Shishmaref	3	3	0	5	0	0	0	5
Shungnak	70	47	0	90	15	2,595	9	2,709
Sitka	663	605	17	13,043	475	71	154	13,761
Skagway	21	20	7	203	0	0	23	234
Skwentna	10	10	0	148	18	14	16	196
Slana	22	20	1	707	0	0	0	708
Sleetmute	40	35	132	715	784	1,004	120	2,755
Soldotna	2,101	1,798	54	35,598	197	12	136	35,998
South Naknek	18	15	20	778	79	11	54	942
Stebbins	119	106	121	17	1,266	3,476	3,759	8,640
Sterling	481	393	5	7,983	35	8	29	8,060
Stevens Village	21	14	330	0	0	465	0	795
Stony River ^d	16	3	212	398	372	619	0	1,601
Sutton	130	106	5	2,187	2	6	12	2,212
Takotna ^b	23	0	0	2	22	0	0	24
Talkeetna	107	84	18	1,893	4	2	12	1,929
Tanacross	2	2	0	66	0	0	0	66
Tanana	104	52	2,100	0	3,060	24,798	3	29,961
Tatitlek	14	7	3	880	0	8	0	891
Tazlina	53	49	129	3,686	26	0	0	3,842
Telida ^c	2	--	--	--	--	--	--	--
Teller	45	45	26	342	100	3,864	1,951	6,283
Tenakee Springs	3	3	1	47	1	0	0	49
Thorne Bay	27	23	1	132	0	0	2	136
Togiak	54	38	951	5,379	299	779	85	7,492
Tok	80	72	17	2,353	1	3	1	2,376
Toksook Bay	2	2	0	32	0	0	3	35
Tolsona	7	7	5	342	0	0	0	347
Tonsina	10	10	7	125	0	0	0	132
Trapper Creek	33	24	2	735	1	0	1	740
Tuluksak	89	53	651	1,380	341	2,585	8	4,965
Tuntutuliak	90	53	1,123	1,516	565	2,614	15	5,833
Twin Hills	2	1	0	25	0	0	0	25
Two Rivers	25	20	1	445	0	0	0	447
Tyonek	63	47	745	121	76	0	0	942
Ugashik	10	9	7	588	168	1	0	764
Unalakleet	228	204	661	327	4,324	2,144	8,743	16,199
Unalaska	104	78	5	2,266	282	36	263	2,854

-continued-

Table 2-5–Page 7 of 7

Community	Households or permits		Estimated salmon harvest					
	Total	Included ^a	Chinook	Sockeye	Coho	Chum	Pink	Total
Unknown community	1,346	674	494	18,657	1,668	639	450	21,908
Upper Kalskag	62	31	562	770	360	1,930	30	3,652
Valdez	275	220	61	5,853	27	0	1	5,942
Venetie	76	25	86	55	0	295	0	436
Wainwright	154	78	20	80	105	51	137	393
Wales	2	2	0	5	0	0	0	5
Ward Cove	5	5	0	23	0	6	0	29
Wasilla	4,609	3,738	403	96,561	1,019	62	509	98,554
Whale Pass	1	0	0	0	0	0	0	0
White Mountain	42	41	18	44	300	237	3,662	4,262
Whittier	10	6	0	155	1	0	1	156
Willow	221	177	9	3,811	37	1	10	3,866
Wiseman	1	1	0	0	0	0	0	0
Wrangell	183	179	40	1,439	110	88	30	1,707
Yakutat	143	113	369	5,342	1,116	11	202	7,040
Total	63,357	48,505	75,211	1,120,675	89,247	368,524	75,158	1,728,815

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

- a. “Included” is the sample size or the number of permits returned.
 - b. These communities were not contacted during the 2012 study period. Harvests were estimated using historical average household harvest expanded by the number of households.
 - c. These communities were not contacted during the 2012 study period. Not enough data was available to estimate harvest.
 - d. Communities were contacted, but numbers of selected households or total number of surveyed households were insufficient. Harvests were estimated using historical average household harvest expanded by the number of households.
- Data not available.




Figure 2-1.—Composition of subsistence harvest by rural Alaska residents, 2012.




Figure 2-2.—Alaska subsistence salmon harvest by species, 2012.




Figure 2-3.—Alaska subsistence salmon harvest by area, 2012.




Figure 2-4.—Subsistence Chinook salmon harvest by area, 2012.




Figure 2-5.—Subsistence sockeye salmon harvest by area, 2012.




Figure 2-6.—Subsistence chum salmon harvest by area, 2012.




Figure 2-7.—Subsistence coho salmon harvest by area, 2012.




Figure 2-8.—Subsistence pink salmon harvest by area, 2012.




Figure 2-9.—Alaska personal use salmon harvest by species, 2012.




Figure 2-10.—Alaska subsistence and personal use salmon harvest by species, 2012.

CHAPTER 3: NORTON SOUND-PORT CLARENCE AREA AND ARCTIC-KOTZEBUE AREA ALASKA

INTRODUCTION

In 2012, the fisheries management district for the North Slope, called the Northern Area, was renamed the Arctic Area. At the same time, management was separated from the Yukon Area and combined with the Kotzebue area. The new fisheries management area is called the Arctic-Kotzebue Area. Previous statewide subsistence fisheries reports have not included information regarding subsistence fisheries on the North Slope, although ongoing division research is attempting to fill this information gap. This chapter has been reorganized and expanded to reflect these changes to the subsistence fisheries management area. It has been expanded to include the results of recent subsistence research conducted in the area, focusing on subsistence fisheries harvest information to supplement the existing annual harvest monitoring program.

NORTON SOUND–PORT CLARENCE AREA SALMON

Background

The archaeological record of the Norton Sound–Port Clarence region provides physical evidence of subsistence fishing dating back to the Arctic Small Tool/Norton Tradition, ca. 1500–1000 B.C.E. (Harritt 2010; Smith and Vreeman 1995). The area includes the regional center of Nome, with a 2012 population of 3,756, and 13 smaller communities ranging in size from 121 (Diomedes) to 712 (Savoonga).⁷ Overall, 76% of the residents of the Nome Census Area are Alaska Native, with an additional 6% reporting 2 or more racial backgrounds. More than 90% of the region’s population outside of Nome is Alaska Native, with Inupiaq, Yupik, and Siberian Yupik peoples present. Most residents of the region continue to participate in a mixed subsistence-cash economy and depend on wild foods for cultural and nutritional sustenance. While more opportunities for wage work exist in Nome itself, subsistence activities are still an important facet of life to many of its inhabitants.

In summer, subsistence fishers harvest salmon with gillnets or seines in the main Seward Peninsula rivers and coastal marine waters. Beach seines are used near the spawning grounds to harvest schooling or spawning salmon and other species of fish. A major portion of fish taken during the summer months is air dried or smoked for later consumption by residents. Chum and pink salmon are the most abundant salmon species districtwide; Chinook and coho salmon are present throughout the area but are more common in eastern and southern Norton Sound. Sockeye salmon are found in a few Seward Peninsula streams.

Regulations

The Port Clarence District includes all waters from Cape Douglas north to Cape Prince of Wales, including Salmon Lake and the Pilgrim River drainage. In most of the district, subsistence salmon fishing has few restrictions other than the general statewide provisions. Standard permit conditions include prohibition of fishing within 300 ft of a dam, fish ladder, weir, culvert, or other artificial obstruction. Salmon may be taken in most areas at any time, with no harvest limits. This area includes fishing areas used by residents of Teller, Brevig Mission, and Nome (the area is accessible via road from Nome) (Magdanz 1992:27). Since 2004, subsistence salmon permits have been required in all Port Clarence waters. In addition, in the Pilgrim River drainage, including Salmon Lake and the Kuzitrin drainage, harvests are limited, and specified areas are closed to subsistence salmon fishing. No fishing occurred in

7. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. “Research and Analysis Homepage.” Accessed July 2014. <http://labor.alaska.gov/research/pop/popest.htm>

Salmon Lake in 2009–2011 due to a crash of the sockeye population—managers had opened a portion of the lake to fishing for the previous 3 years. Prior to that, it had been closed since 1972.

The Norton Sound District encompasses all waters from Point Romanof north to Cape Douglas. It is divided into 6 subdistricts: 1) Nome, 2) Golovin, 3) Moses Point/Elim, 4) Norton Bay, 5) Shaktoolik, and 6) Unalakleet. In subdistricts 1 and 6, restrictions exist on gear, fishing periods, and areas open to fishing. In 2001, a regulatory change by the BOF made rod and reel a legal subsistence fishing gear type in the area from Cape Espenberg on northern Seward Peninsula to Bald Head, which is between Elim and Koyuk. This area includes subsistence fishing areas used by the residents of Nome, White Mountain, Golovin, Elim, Koyuk, Shaktoolik, and Unalakleet. Sport fishing bag and possession limits still apply, except when a subsistence salmon permit is required or fishing through the ice. In the former case, the harvest limits (if any exist) specified on the permit for each river apply. When fishers catch their limit in one drainage, they can fish in another. Subsistence fishing regulations are most restrictive in Subdistrict 1 (Nome) and Subdistrict 6 (Unalakleet), where the 2 largest communities in the area are located.

In Subdistrict 1 (Nome), subsistence harvests consist primarily of chum salmon and pink salmon. Chum salmon runs have been depressed since the mid-1980s, leading to increasing restrictions on all types of harvest. Upstream portions of most rivers are closed to protect spawning salmon, and harvests are limited in all subdistrict rivers. For 16 years, subsistence fishing was prosecuted primarily by emergency order, with openings much less frequent than in regulation. Fishing periods in marine waters were also limited.

From 1991–2005, the Nome Subdistrict was managed primarily by emergency order and was frequently closed to subsistence fishing for chum salmon each year on June 15 until ADF&G judged escapement goals were likely to be met. These closures, even when they were of short duration, impacted subsistence fishing because fishing often reopened during a wetter part of the summer, which made it difficult, if not impossible, to dry and process fish harvested for subsistence uses. From 1999 through 2005, chum salmon fishing in Subdistrict 1 was managed as Tier II, the only such fishery to exist in the state (5 AAC 01.182, 5 AAC 01.184).⁸ In 1999, the chum salmon return was so poor that even Tier II fishing was closed; in 2000, only 10 permits were awarded (Soong et al. 2008:10). Chum salmon returns since then have gradually improved, allowing ADF&G to manage the fishery as Tier I between 2006 and 2012, and generally to observe the fishing schedule provided for by regulation.⁹ In subdistricts 2 through 4, salmon may be taken at any time, with no harvest limits. However, restrictions exist on commercial fishers' participation in subsistence salmon fishing. Both the escapement and the commercial harvest of chum salmon experienced sharp declines starting in 1990 (Menard and Bergstrom 2006:2); however, the runs have been rebounding in recent years in the Norton Sound District (Menard et al. 2012). In Subdistrict 2 (communities of Golovin and White Mountain), both commercial and subsistence chum salmon harvests have dropped significantly since the 1990s; subsistence restrictions were in place in 2003. Chum salmon stocks in subdistricts 2 and 3 have been classified as stocks of “yield concern” since 2000, but chum salmon runs greatly improved in the late 2000s (Menard et al. 2012:8).

In subdistricts 5 and 6 (Shaktoolik and Unalakleet, respectively), continuing poor Chinook salmon runs have led to restrictions on commercial, sport, and subsistence fishing. The Shaktoolik and Unalakleet subdistricts are typically managed together because actions in one subdistrict are believed to affect the movement of fish in the other. Only 1 commercial Chinook salmon directed fishery has occurred since 2001. Restrictions were placed upon the subsistence and sport fisheries in 2003, 2004, and 2006–2012

8. A “Tier II” subsistence permit program is necessary when the number of participants in a subsistence fishery or hunt must be limited because the harvestable surplus of the fish stock or wildlife population is less than the lower bounds of the amount necessary to provide for subsistence uses. Individual Alaskans are distinguished from one another through the submission of Tier II applications, which are scored based on their history of uses of the particular resource and the ability to obtain alternative food; those with the highest scores receive Tier II permits, the others do not. Tier II provides a process that ensures that subsistence opportunities are provided to those most dependent upon the resource. Tier II implies that there is an insufficient harvestable surplus to provide for all subsistence uses (AS 16.05.258).

9. In a “Tier I” subsistence fishery, all interested Alaska residents may participate. Other fishers (commercial, sport, and personal use) are prohibited or restricted because the harvestable surplus is sufficient only to provide for customary and traditional subsistence uses (AS 16.05.258).

(Menard 2010; Menard et al. 2011, 2012, 2013). The Chinook salmon management plan adopted by the BOF in February 2007 (5 AAC 04.395) limits subsistence gillnet salmon fishing to two 48-hour fishing periods per week in marine waters from mid-June to mid-July. On the Unalakleet River, subsistence fishing is limited to two 36-hour fishing periods per week. Fishing time could be increased only if ADF&G were to project that the lower end of the sustainable escapement goal (SEG) range would be reached.

In 2007, the BOF changed the classification of Subdistrict 1 chum salmon from a “stock of management concern” to a “stock of yield concern.” Subdistricts 2 and 3 chum salmon stocks, as well as subdistricts 5 and 6 Chinook salmon stocks, continued as “stocks of yield concern” (Soong et al. 2008:34).

New state regulations governing customary trade of fish caught in the Norton Sound and Port Clarence areas became effective July 1, 2007. These regulations allowed cash sales, up to \$200, of subsistence-caught finfish per household per year. Persons who wanted to participate had to obtain a customary trade record keeping form from Nome ADF&G. Sales could not be made to a fishery business nor the fish resold by the buyer. Sales could also occur only within the Norton Sound–Port Clarence Area (Soong et al. 2008:34). Effective April 13, 2013, the Alaska Board of Fisheries increased the annual limit for selling subsistence-taken finfish as customary trade from \$200 per year to \$500 total per household in a calendar year (5 AAC 01.188).

Subsistence Salmon Harvest Data Collection Methods

Two methods were used to assess subsistence salmon harvests in the Norton Sound and Port Clarence districts in 2012: (1) fishing permits in Subdistrict 1 (Nome), the Cape Woolley Area, Subdistrict 2 (Golovin), Subdistrict 3 (Moses Point/Elim), and the Port Clarence District (Brevig Mission and Teller); and (2) postseason household surveys conducted by the Division of Commercial Fisheries in 5 communities: Koyuk in the Norton Bay area (Subdistrict 4), Shaktoolik (Subdistrict 5) and Unalakleet (Subdistrict 6) as well as St. Michael and Stebbins, which are not within any subdistrict’s boundaries. 2012 subsistence fisheries harvest information resulting from Division of Subsistence household surveys conducted in Golovin are also presented below, although this research is not part of the annual subsistence fisheries monitoring program implemented by the Division of Commercial Fisheries.

Norton Sound Subdistricts 1, 2, and 3: Fishing Permits

Permits have been required for subsistence salmon fishing in Norton Sound Subdistrict 1 (Nome) since 1974. Beginning in 1999, Tier II chum salmon fishing permits were also issued to a limited number of Nome households with the intent that these households would have first priority over other subsistence fishers if only a small number of chum salmon were available for harvest. This priority would allow these households to fish earlier in the season, when weather conditions were more suitable for drying salmon. Tier I fishing permits were available to all other households when run strength was determined to be adequate. In 2012, chum salmon run abundance was projected to achieve the biological escapement goal for the subdistrict and provide amounts reasonably necessary for subsistence. Because of this, a Tier II fishery was not implemented (Menard et al. 2013). The Nome ADF&G office issued 483 subsistence (Tier I) salmon permits, all of which were returned. This was slightly more than the 448 permits issued in 2011, but lower than the record 494 permits issued during the 2010 fishing season (Menard et al. 2012) (Table 3-1). A total of 340 households fished their permits, with the largest number of permits fished on the Nome River (242) and Snake River (61) (harvests largely came from those rivers, the Eldorado River, and marine waters) (Menard et al. 2013).

Since 1998, the Nome permit data have not been expanded to account for households whose permits were not returned. This contrasts with earlier years when permit data were expanded by drainage, with expansion factors based upon the fraction of unreturned permits for that drainage. ADF&G staff believed that expansion of the permit data led to an overestimation of the salmon harvest because the unreturned permits were most likely from households that did not fish.

Subsistence fishing permits were also issued for the Cape Woolley Area, a traditional camp and fishing area for King Island households, who, although they settled in Nome more than 40 years ago, maintain a distinct community identity. Located in the Norton Sound District west of Nome, this area lies outside Subdistrict 1 but within the boundaries of the area for which fishing permits are required (Rocky Point to Cape Douglas). In 2012, 20 permits were issued for the Cape Woolley Area; all were returned to ADF&G (Table 3-1). Only 4 households fished their permits (Menard et al. 2013).

Subsistence permits have been required for salmon fishing in Subdistrict 2 (Golovin) and Subdistrict 3 (Moses Point/Elim) since 2004. In 2012, 151 permits were issued for Subdistrict 2; more than the 145 permits issued in 2011, but fewer than in 2010 (159), 2005 (174) and 2004 (199) (Fall et al. 2012a:23; Menard et al. 2013). All 151 permits were returned (Table 3-1); 97 households reported fishing (Menard et al. 2013). The number of Subdistrict 2 permits issued to Nome residents has dropped since 2004, and fishery managers have attributed the decline to the easing of fishing restrictions in the Nome Subdistrict and rising fuel costs. The number of permits issued to residents of White Mountain and Golovin has held steady. In 2012, ADF&G issued 63 permits for Subdistrict 3, slightly more than the 60 permits issued in 2011 and slightly less than 64 permits issued in 2010. All permits were returned; 54 households reported fishing (Menard et al. 2012) (Table 3-1).

Port Clarence District: Salmon Lake and Pilgrim River Fishing Permits

Permits have been required to fish the Pilgrim River since 1974 (Magdanz 1992:27). This requirement was expanded to all Port Clarence waters in 2004. In 2012, 335 Port Clarence and Pilgrim River permits were issued, compared to 271 in 2011, 295 in 2010, and 328 in 2009 (tables 3-2 and 3-3). Of the permits issued in 2012, 188 were to fish the Pilgrim River only; no permits were issued for Salmon Lake because it was closed; and 147 were issued for other waters in the district (Menard et al. 2012:61). The number of permits for the Pilgrim River grew substantially between 2003 and 2008, likely in response to several consecutive years of record sockeye salmon runs. The decline in permits issued may, in turn, be a response to a poor run in 2008 followed by the crash in 2009, when only 953 salmon were counted passing through the weir. Poor runs continued in 2010, with a count of 1,654 salmon. There was improvement to the sockeye run in 2011, with a count of 8,449 sockeye salmon (Menard et al. 2012). In 2012, 7,085 sockeye salmon were counted passing through the weir, slightly less than 2011 (Menard et al. 2012). All Pilgrim River permits were returned, as well as all the permits issued for other waters of the Port Clarence District.

In 2007, the BOF adopted regulations that closed the southwestern half of Salmon Lake and allowed for fishing on the northeastern half by emergency order. No salmon fishing has been allowed in recent years in Salmon Lake due to the crash of the sockeye salmon run in 2009 and poor runs in 2010–11 (Menard et al. 2012). Although fishing was prohibited in Salmon Lake in 2012, this marked the second year in a row since 2008 that the sockeye salmon escapement goal has been met (Menard et al. 2013).

Household Surveys

In 2012, ADF&G Division of Commercial Fisheries conducted subsistence fisheries household surveys in Koyuk, Shaktoolik, Unalakleet, Saint Michael, and Stebbbins. Researchers attempted to contact all of the households in each of the surveyed communities. Actual sample rates varied: 200 of 223 Unalakleet households (90%) were contacted, as were 63 of 64 Shaktoolik households (98%), 82 of 83 Koyuk households (99%), 82 of 82 St Michael households (100%), and 106 of 117 Stebbbins households (97%). The salmon survey data were expanded by community to account for the households not contacted (Table 3-2).

The goals of the postseason household survey were to:

- collect harvest data that would result in a total harvest estimate for subsistence salmon by species and by community;

- compile information on harvest by gear types, participation rates, household size, use of salmon for dog food, and participation in customary barter and trade; and
- assess the quality of chum salmon fishing and what affected it.

Subsistence Salmon Harvests in 2012

Norton Sound District Subsistence Salmon Harvest

The estimated 2012 subsistence harvest of salmon by communities in the Norton Sound District was 76,524 fish (tables 3-1, 3-3). This was the second lowest total harvest for the district for an even numbered year on record since 1994, and 19% lower than the average even year harvest from 1994–2012 (Table 3-3). Pink salmon abundance commonly follows an even–odd year cycle. Their abundance in Norton Sound is usually significantly higher in even-numbered years (2004, 2006, 2008, etc.) with districtwide harvests usually reflecting this difference. Between 1994 and 2012, odd-year harvests of all salmon have ranged from a low of 43,883 in 2011, to a high of 113,612 in 1995, with an average of 72,810 salmon. Even-year harvests have ranged from the low in 2010 of 67,149 to a high of 134,050 in 1996, with an average of 93,432 salmon. Extensive June sea ice in 2012 made accurate assessments of early run strength for chum salmon impossible. Shorefast ice thickness reports ranged from 3 to 5 feet from St Michael Bay to Unalakleet as late as mid-May, and many residents of Norton Sound commented that the extent, thickness, and duration of sea ice in Norton Sound was unusual. Large pans of first year ice and shorefast ice melted in place in southern Norton Sound, which resulted in lower water temperatures nearshore. These factors likely delayed the migration timing of most Chinook, pink, and chum salmon stocks, and led to difficulties with setting subsistence and commercial fishing schedules. Anticipated weak runs resulted in no commercial fishing in Norton Sound’s Subdistricts 5 and 6 for Chinook salmon for the seventh consecutive year, and escapement was estimated to be weak for this species in many locations in 2012. The majority of Chinook salmon that returned was from the 2007 brood year; this brood year had unexpectedly poor productivity, which was even lower than the poor production in the 2005–2006 brood years. Pink salmon escapements in 2012 were all well below long-term, even-numbered year average escapements, but all escapement goals were easily met. Surpluses were adequate to allow for subsistence needs and commercial harvests. Record rainfall and flooding in August made projections of early run strength of coho salmon runs unattainable and reliable estimates of escapement could not be made at any project in 2012 (Menard et al. 2013).

Subdistrict 1 Harvest

For the seventh year in a row, Subdistrict 1 opened on June 15 for subsistence salmon fishing as per regulation. The chum salmon fishing schedule (72 hours in marine waters and two 48-hour fishing periods in fresh waters per week) was observed from mid-June to mid-July. Because chum runs were expected to exceed the ANS and because of the late spring, ADF&G also allowed the use of beach seines during freshwater gillnet periods beginning in mid-June in order to allow for more efficient harvest. By late June and early July, excellent marine subsistence catches of chum salmon were reported, but weir counts were lower than expected at the Snake River and Nome River weir projects. Aerial surveys were conducted in mid-July in the Flambeau, Elorado, and Bonanza rivers of the eastern Nome subdistrict and the Sinuk River in the western Nome subdistrict. Several thousand chum salmon were observed on these surveys, and the Eldorado River weir-based project salmon escapement goal was projected to be easily met by July 12. Management biologists were able to project that the subdistrict-wide biological escapement goal would be met, and chum salmon subsistence gillnet fishing proceeded at the standard freshwater and marine schedules for the remainder of the season. In addition, several beach seining opportunities were issued via emergency order to increase the efficiency of subsistence chum and pink salmon harvests during periods of good drying weather. Accurate assessment of coho salmon escapement in the Nome Subdistrict was not possible after mid-August because of flooding events, making all counting operations inoperable and causing poor aerial survey conditions (Menard et al. 2013). The

reported 2012 subsistence salmon harvest in the Nome Subdistrict was 11 Chinook, 2,521 chum salmon, 8,376 pink, 1,150 coho, and 171 sockeye salmon (Table 3-1).

Subdistrict 2 and 3 Harvest

No subsistence catch limits are in place in subdistricts 2 and 3. Most salmon harvested there are caught by residents of the communities of White Mountain, Golovin, and Elim. Pink and coho salmon made up the majority of the salmon harvest, followed by chum, Chinook, and a few sockeye salmon. In 2012, a total of 9,943 salmon were harvested in Subdistrict 2 (Golovin) (Table 3-1), the fifth lowest number in the 2000s and ranked last for even-numbered years in the same timeframe (Menard et al. 2013). Pink salmon composed 77% of the number of salmon harvested, with 11% chum, 11% coho, less than 1% Chinook, and less than 1% sockeye salmon making up the rest of the harvest. Early indicators of 2012 chum salmon abundance to the Golovin Subdistrict were limited to subsistence catch reports of fair harvests. Escapement of chum salmon in the subdistrict, however, was well below average. Directed commercial fishing of chum salmon was limited to one 36-hour period on July 26 when it was determined that the Niukluk River salmon escapement goal would be narrowly exceeded. There were no restrictions on subsistence fishing. Because of the poor weather, reliable estimates of coho salmon escapement could not be determined, but projections of escapement made after the historical quarter point at the Niukluk River suggested that the escapement goal would have been achieved. Limited commercial fishing was allowed, but there was little effort due to stormy weather and high surf conditions. No restrictions were placed on subsistence fishing. Subdistrict 2 harvests, as noted earlier, largely reflect those of communities within the subdistrict (Menard et al. 2013).

Based upon subsistence fishing permits, residents of Golovin harvested an estimated 3,519 salmon in 2012, of which more than half, 2,415 were pink salmon (Table 3-2). Coho salmon harvests (246) contributed 7% to the total salmon harvest, with chum salmon (775) composing 22%, Chinook salmon (39) contributed 1%, and sockeye harvests (44) contributed 1% to the total Golovin salmon harvest. White Mountain residents harvested an estimated 4,217 salmon, 3,662 (87%) of which were pink salmon. The remainder of the harvest was chum salmon (237) at 6%, coho salmon (300) at 7%, and Chinook salmon (18) at less than 1%. There was no harvest of sockeye salmon reported by residents of White Mountain in 2012.

In February 2013, residents of 33 Golovin households responded to a Division of Subsistence survey asking about their subsistence harvests of fish, wildlife, and plants during the calendar year 2012, representing a community sample of 56%. Golovin residents harvested an estimated 16,222 edible pounds of salmon in 2012, accounting for 58% of the total fish harvests. Pink salmon harvests (3,666 fish) were the greatest, accounting for 47% of the total salmon harvest in 2012 by weight, followed by 824 chum salmon, 670 coho salmon, 50 Chinook salmon, and 6 sockeye salmon. Forty-five percent of Golovin households reported using less salmon in 2012 than in previous years (Braem et al. in prep).¹⁰ Harvest estimates resulting from the Division of Subsistence household surveys differ from the results of subsistence permit returns due, in part, to the fact that harvests were expanded for uncontacted households in Golovin. As a result, most estimates are slightly higher than permit returns indicate, with the exception of sockeye salmon harvests, where permit returns suggested a greater number of sockeye salmon were harvested by residents of Golovin than the Division of Subsistence study estimated; this likely results from not contacting a number of households who had subsistence fishing permits who had harvested sockeye salmon, such that the comprehensive subsistence research underestimated sockeye salmon harvests.

Results from the Division of Subsistence household surveys indicated that nonsalmon fishes accounted for 42% of the total fish harvest by Golovin residents in 2012, and 21% of the estimated total subsistence

10. Braem, N., A. Godduhn, A. Brenner, B. Retherford, and M. Kostick. In prep. Chukchi Sea and Norton Sound Observation Network: Golovin, Noorvik, and Point Lay, 2012. Fairbanks: Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 403. Hereinafter cited as (Braem et al. *In prep*).

harvest, with an estimated total harvest of 11,552 edible pounds. The largest contributions to nonsalmon fish harvests were 2,123 Dolly Varden, 18,496 saffron cod (“tomcod”), 143 broad whitefish, and 107 Bering cisco. Other fish harvests by Golovin residents can be found in Braem et al. (*In prep*).

In Subdistrict 3 (Moses Point/Elim), early projections of the chum salmon escapement by the Kwiniuk River tower counts indicated a very weak run, and no directed chum salmon commercial fishing periods occurred. Commercial fishing for pink salmon was delayed until July 7 because of conservation concerns for chum salmon. The number of salmon reported harvested in the subdistrict by subsistence users was the third highest since 1994. Chinook salmon harvests were a record low, while the pink salmon harvest was a record high (Menard et al. 2013). Subsistence fishers harvested an estimated 13,686 salmon, 79% of which were pink salmon. The remainder were 11% chum salmon, 9% coho salmon, and less than 1% of the subsistence salmon harvest was of Chinook salmon. There was no harvest of sockeye salmon in Subdistrict 3 reported by Elim residents in 2012.

Subdistrict 4 Harvest

In 2012, the fifth consecutive annual subsistence salmon survey was conducted in Koyuk by the Division of Commercial Fisheries. Fishers in the subdistrict caught an estimated 5,758 salmon, the second fewest since 1994, with most of the harvest being made up of pink and chum salmon (46% and 47%, respectively). Of the remainder, 5% were coho salmon, and 2% were Chinook salmon. There was no reported harvest of sockeye salmon in the subdistrict in 2012 (Table 3-1). It is important to note, however, that commercial fishing occurred in the subdistrict, and the combined commercial and subsistence harvest of 68,548 salmon was the largest on record. (Menard et al. 2013). By comparison, in 2011, fishers in the subdistrict harvested an estimated 5,592 salmon, 21% of which were pink salmon (1,161) and 65% chum (3,621). Coho salmon made up 10% of that year’s subsistence salmon harvest, another 4% came from Chinook salmon, and less than 1% of the harvest was sockeye salmon (Fall et al. 2013).

Table 3-2 presents harvests at the community level. Because of additional harvests in other subdistricts, Koyuk households caught slightly more salmon than the total harvest for the Norton Bay subdistrict. Households harvested an estimated 6,045 salmon, the majority of which were pink salmon (47%) and chum salmon (45%). Households caught lesser amounts of coho (6%) and chinook (2%) salmon.

Subdistrict 5 and 6 Harvests

Preseason forecasts by ADF&G called for another very poor Chinook salmon run to subdistricts 5 and 6. Restrictions were put in place on subsistence fishing per the management plan (5 AAC 04.395). Fishery managers limited fishing time with set gillnets to two 48-hour periods per week in marine waters and two 36-hour periods per week in the Unalakleet River drainage. Beach seining was permitted beginning on June 27 in order to target abundant chum and pink salmon, but retention of any incidentally caught Chinook salmon was prohibited. In order to gauge the early run strength of the pink salmon run, both subdistricts were given one 12 hour commercial opening on July 3, but restricted mesh size of gillnets to 4.5” or less. Due to evidence that the chum salmon run strength was building, managers opened commercial fishing for a 24 hour window on July 5 (subdistrict 6) and July 6 (Subdistrict 5). Mesh size was restricted to 6” or less. Other commercial openings for both pink and chum salmon occurred in both subdistricts throughout the month of July. On July 11, subsistence Chinook salmon fisheries in both the Shaktoolik and Unalakleet rivers were closed and mesh size for set gillnets was restricted to a mesh size of no greater than 4 ½”. Subsistence salmon fishing was opened 7 days a week with mesh restrictions in place, and sport fishing for king salmon was closed. The near average coho salmon run in both subdistricts allowed for limited commercial fishing and no subsistence restrictions. The decrease in commercial coho salmon harvests in both subdistricts in 2012 can be attributed to poor August weather rather than a lack of abundance (Menard et al. 2013).

Fishers in subdistrict 5 (Shaktoolik) caught an estimated 6,576 salmon in 2012, the bulk of which (4,609 or 70%) were pink salmon. Coho salmon (1,110) composed 17% of the total harvest; chum salmon (634)

and Chinook salmon (214), each of which was 10% and 3% of the total, respectively. Less than 1% of the harvest consisted of sockeye salmon (Table 3-1).

In subdistrict 6 (Unalakleet), subsistence fishers caught an estimated 16,182 salmon, over half of which (55%) were pink salmon. Coho salmon (4,318) made up 27% of the annual harvest, followed by chum salmon (2,164 or 13%), and Chinook salmon (673 or 4%). One percent of the total harvest was sockeye salmon (Table 3-1).

Table 3-2 presents harvests at the community level. Because a portion of the respective subdistrict harvests were taken by residents of communities outside of Shaktoolik and Unalakleet, the community harvests are slightly less than the total harvest for the individual subdistricts. Households in Shaktoolik harvested an estimated 6,290 salmon, the majority of which were chum salmon (70%). Unalakleet households harvested an estimated 16,053 salmon, the majority of which were pink salmon (54%).

St. Michael and Stebbins

Prior to 2012, household surveys had not been conducted in Stebbins and St. Michael since 2009. Both of the communities lie outside of the boundaries of the Norton Sound commercial fishing subdistrict designations, but are within the Norton Sound-Port Clarence Fisheries Management Area. In 2012, St. Michael households caught an estimated 3,640 salmon, the bulk of which (2,172 or 60%) were chum salmon. Coho salmon (911) composed 25% of the total harvest; pink salmon (457) and Chinook salmon (80) were 13% and 2% of the total harvest, respectively. Less than 1% (20) of the harvest was made up of sockeye salmon. All harvests in the area came from households in the community of St. Michael. (tables 3-1 and 3-2).

In the Stebbins area, subsistence fishers caught an estimated 8,480 salmon. Pink salmon composed 43% (3,659 fish) of the total salmon harvest, while chum salmon contributed 40% (3,456 fish). Coho salmon (1,256) made up 15% of the harvest and Chinook salmon composed 1% of the catch. No sockeye salmon were harvested for subsistence in 2012 (Table 3-1).

On the community level, households in Stebbins caught slightly more salmon than the total harvest for their area. This is due to reports by Stebbins fishers of harvests in other subdistricts. The total estimated harvest for the community was 8,625 salmon, the majority of which were pink (44%) and chum (40%) salmon (Table 3-2).

Norton Sound Harvest Overall

Of the total 2012 subsistence salmon harvest in Norton Sound, 1% were sockeye salmon, 2% were Chinook salmon, 15% were coho salmon, 21% were chum salmon, and 61% were pink salmon (Figure 3-1). Total harvest estimates for the Norton Sound District for 1975–2011 are presented in Table 3-5. However, the methods used to determine harvests prior to 1994 are substantially different from those used since 1994. As a consequence, the data are not directly comparable. Methods changed again in 2004 when permits replaced surveys in Norton Sound Subdistrict 2 (Golovin and White Mountain) and Norton Sound Subdistrict 3 (Moses Point/Elim). Very little of the documented 2012 subsistence salmon harvest was taken by residents from outside the district (Table 3-2). Eleven subsistence permits were issued to residents of Anchorage, Fairbanks, and Palmer; their combined total salmon harvest was 270 salmon.

Port Clarence District Subsistence Salmon Harvest

The estimated 2012 subsistence harvest of salmon in the Port Clarence District was 15,172 fish (tables 3-3 and 3-4). This harvest was the highest since 2008 (15,957 fish) and higher than the 10-year average (2002–2011) of 14,218 fish. Of the total salmon harvest, less than 1% was Chinook salmon, 5% were coho salmon, 52% were chum salmon, 34% were pink salmon, and 9% were sockeye salmon (Figure 3-2). Total harvest estimates for the Port Clarence District for 1975–2011 are presented in Table 3-5.

ARCTIC-KOTZEBUE AREA SALMON

Introduction

In 2012, the fisheries management district for the North Slope, called the Northern Area, was renamed the Arctic Area. At the same time, management was separated from the Yukon Area and combined with the Kotzebue Area. The new fisheries management area is called the Arctic-Kotzebue Area. They will be addressed separately below. Previous annual reports have not addressed subsistence fisheries information from the Northern Area, as there have been no harvest monitoring programs conducted by ADF&G. Ongoing Division of Subsistence research will serve to expand available information on subsistence fisheries by residents of North Slope communities. Some research results from the North Slope Borough Department of Wildlife Management are also summarized below to better document the extent of subsistence fisheries on the North Slope.

Background

Arctic Area residents of the North Slope have relied on fish for cultural and nutritional sustenance for generations. The only systematic subsistence fisheries harvest monitoring program has been conducted by the North Slope Borough's Department of Wildlife Management (Bacon et al. 2011rev.). The most recent report by NSB described subsistence fish harvests in the region from 1994–2003; this includes harvest amounts, harvest timing, locations, gear and other qualitative information (Bacon et al. 2011rev.). Most residents in the region continue to participate in a mixed subsistence-cash economy, harvesting a wide variety of wild foods. The Arctic Area includes the subsistence fishing areas used by Anaktuvuk Pass, Atkasuk, Barrow, Kaktovik, Nuiqsut, Point Hope, Point Lay, and Wainwright. The role of salmon and nonsalmon in the wild food diet varies from community to community and is affected primarily by resource availability. Chum and pink salmon are present in the greatest abundance, although sockeye, coho, and Chinook salmon are occasionally caught. Residents often refer to ocean bright salmon as “silvers” leading to the misidentification of chum harvests as coho salmon in some cases. Nonsalmon species important to subsistence include Arctic grayling, Dolly Varden, lake trout, burbot, rainbow smelt, various whitefishes, Arctic cod, and saffron cod. ADF&G Division of Subsistence has an ongoing subsistence fisheries research project along the western coast of the North Slope (OSM 12-154), focusing on subsistence fishing harvest and use patterns by residents of Point Lay and Wainwright.

Kotzebue Sound residents have relied on fish for cultural and nutritional sustenance for thousands of years. Most residents in the region continue to participate in a mixed subsistence-cash economy, harvesting a wide variety of wild foods. The Kotzebue Area includes the subsistence fishing areas used by Point Hope, Kivalina, Noatak, Kotzebue, Kiana, Noorvik, Selawik, Ambler, Shungnak, Kobuk, Buckland, Deering, Shishmaref, and Wales. The role of salmon in the wild food diet varies from community to community, and is affected primarily by salmon abundance. Communities that harvest few salmon typically harvest large numbers of nonsalmon fish, such as sheefish *Stenodus leucichthys*, other whitefishes *Prosopium* and *Coregonus* spp., and Dolly Varden *Salvelinus malma*. Along the Noatak and Kobuk rivers, where runs of chum salmon are strong, many households' activities in mid- and late summer revolve around the harvesting, drying, and storing of salmon for use during the winter. Chum salmon predominate in the district, composing 90% of the subsistence salmon harvest. Small numbers of other salmon species are present in the district. ADF&G Division of Subsistence has one ongoing subsistence fisheries research project in the Kotzebue Sound region (OSM 12-153). This research resulted in subsistence fish harvest estimates for 2012, discussed below.

Regulations

In the Arctic-Kotzebue Area, subsistence salmon fishing has few restrictions, other than the general statewide provisions (e.g., 5 AAC 01.010) and specifications regarding lawful subsistence gear and gear specifications (5 AAC 01.120). Standard conditions include prohibition of fishing within 300 ft of a dam, fish ladder, weir, culvert, or other artificial obstruction. Salmon may be taken in the Arctic-Kotzebue

Area at any time with no harvest limits and no required permits. Salmon may be taken only by gillnets, beach seines, or by hook and line attached to a rod or pole, but only in the state waters of, and all flowing waters that drain into the Chukchi Sea or Kotzebue Sound from Cape Espenberg to Cape Prince of Wales (5 AAC 01.120(f)).

Fish other than salmon may be taken by set gillnet, drift gillnet, beach seine, fish wheel, pot, longline, fyke net, dip net, jigging gear, spear, and lead, or, as specified in 5 AAC 01.120(f), by hook and line attached to a rod or pole. In the Kotzebue District, gillnets used to take sheefish may not be more than 50 fathoms in aggregate length nor 12 meshes in depth, nor have a mesh size larger than seven inches (5 AAC 01.120(e)).

Other regulatory restrictions associated with subsistence fishing in the Arctic-Kotzebue Area include the provision that a gillnet may not obstruct more than one-half the width of any fish stream and any channel or side channel of a fish stream. Furthermore, a stationary fishing device may not obstruct more than one-half the width of any salmon stream and any channel or side channel of a salmon stream (5 AAC 01.120(c)). Except when fishing through the ice or when a subsistence fishing permit is required, use of a hook and line attached to a rod or pole between Cape Espenberg and Cape Prince of Wales requires a subsistence fisherman to follow the methods and means specified in sport fishing regulations 5 AAC 70.011 and 5 AAC 70.030, and the bag and possession limits, by species, specified in 5 AAC 70.011.

Subsistence Salmon Harvest Data Collection Methods

From 1994 through 2004, with funding from the Division of Commercial Fisheries, the Division of Subsistence conducted household surveys in selected Kotzebue Sound communities to collect subsistence salmon harvest data (Fall et al. 2007:23–38). Since funding for that effort has not been available since 2004, no annual surveys have been conducted; therefore, no subsistence salmon harvest estimate is available for 2012, with a couple of exceptions discussed below.¹¹ The average yearly subsistence harvest between 1994 and 2004 was 59,650 salmon, the majority of which were chum salmon (Table 3-3). This average may be low due to incomplete datasets resulting in low harvest totals for several years during that period. Harvest estimates for 1994, 2002, 2003, and 2004 do not include the regional center of Kotzebue. Because Kotzebue is the largest community in the region, it is believed that residents typically harvest as much salmon as residents from all other communities in the region combined (Menard and Kent 2007:1). No harvest information is available for Ambler, a Kobuk River village, for 2001. Data for 2002 include only harvest information from Noatak and Noorvik.

While no Arctic-Kotzebue Area annual subsistence salmon or subsistence fisheries harvest monitoring program is conducted by the Division of Commercial Fisheries, Division of Subsistence collected subsistence fishery harvest information from a number of Arctic-Kotzebue Area communities in 2012 as part of other subsistence research projects. These data are presented below.

ARCTIC-KOTZEBUE AREA SALMON, SHEEFISH, WHITEFISHES, AND ARCTIC CHAR/DOLLY VARDEN

In addition to salmon, major subsistence fisheries take place in the Arctic-Kotzebue Fisheries Management Area for sheefish, other whitefishes, and Dolly Varden (known locally as “trout”). Where salmon are not abundant, these nonsalmon fish often replace salmon in local diets.

Past household surveys to collect harvest information for subsistence salmon harvests in Kotzebue Sound communities also collected harvest data for sheefish, whitefishes, and Dolly Varden in some years (Magdanz et al. 2011:49–50) (tables 3-6 and 3-7).

11. Although the subsistence salmon harvest surveys were eliminated in 2004, information on subsistence fish harvests has been collected occasionally since then during comprehensive subsistence harvest surveys and other projects in some Northwest Alaska communities.

In 2004, the last year Kotzebue Area was surveyed, nonsalmon harvest information was collected in Ambler, Kiana, Kobuk, Noatak, Noorvik, and Shungnak. Those 6 communities harvested an estimated 10,835 sheefish, 50,501 whitefishes, and 11,697 Dolly Varden in that year (Fall et al. 2007:33). Kotzebue Area's total harvest of those species is probably higher, but subsistence fish surveys are not usually conducted in other villages.

The Division of Subsistence collected fish harvest data for 2008 in the villages of Noatak and Kivalina as part of comprehensive community harvest surveys associated with a supplemental environmental impact statement for the Red Dog Mine. Kivalina harvested more than 54,000 fish in 2007; just over 610 were salmon species. Of the estimated 79,000 edible pounds of fish and shellfish harvested, the majority (86%) were Dolly Varden. Saffron cod, known locally as "tomcod," composed just 2% of the total fish harvest; salmon made up only 1% of the total. No other fish species provided even 1% of the total community harvest (Magdanz et al. 2010:26–27). Noatak harvested nearly 23,500 fish in 2007, which provided an estimated 78,454 edible pounds of food to that community. Fish (both salmon and nonsalmon species) made up 41% of the total subsistence harvest by edible pounds. Three species were particularly important: Dolly Varden (33,771 lb), chum salmon (25,002 lb), and whitefishes (14,234 lb) (Magdanz et al. 2010:48).

In October 2011, the Division of Subsistence collected fish harvest data in Selawik for the preceding 12 months as part of a comprehensive subsistence harvest survey. Selawik harvested an estimated 316,653 lb of fish in the study period, the overwhelming majority of which (79%) came from various whitefish species. The greater part of the whitefishes harvest was broad whitefish (47,394 fish, 151,722 edible pounds), followed by sheefish (6,190 fish, 68,958 edible pounds), and humpback whitefish (12,647 fish, 23,705 edible pounds). Salmon are present in only low abundance in the vicinity of Selawik; the only species harvested in any quantity was chum salmon (879 fish, 5,273 edible pounds). Selawik fishers took 15,956 northern pike, totaling 52,653 pounds, the greatest amount for any species other than whitefishes. Lesser harvests of smelt, burbot, and Arctic grayling were also documented (Braem et al. 2013).

Division of Subsistence collected fish harvest data for 2012 in the upper Kobuk River communities of Ambler, Shungnak, and Kobuk as a part of a comprehensive subsistence harvest survey associated with proposed development activities in the Ambler Mining District. Much like the rest of northwest Alaska, weather in 2012 had an impact on upper Kobuk River communities. Key respondents and survey respondents in all 3 communities commented upon the unusual rainfall and the ways in which environmental conditions negatively impacted their harvests (Braem et al. in prep).

In 2012, Ambler households harvested an estimated 59,432 lb of fish, the majority of which (78%) came from various whitefish species. Broad whitefish made up the largest component of the fish harvest (9,150 fish, 36,601 edible pounds) representing 22% of the total estimated subsistence harvest, followed by chum salmon (1,621 fish, 9,214 edible pounds), which represented 6% of the total subsistence harvest. Ambler residents harvested 10,096 edible pounds of salmon in 2012, accounting for 17% of the total fish harvests. Chum salmon harvest were the greatest, accounting for 91% of the total salmon harvest, followed by 126 sockeye salmon, 12 coho salmon, 9 pink salmon, and 2 Chinook salmon. Forty-seven percent of Ambler households reported using less salmon in 2012 than in previous years. Nonsalmon fishes accounted for 83% of the total Ambler fish harvest in 2012, and 30% of the estimated total subsistence harvest, with an estimated harvest of 49,336 edible pounds. The largest contributions to nonsalmon fish harvests were broad whitefish, followed by 1,156 sheefish, 1,544 humpback whitefish, 568 northern pike, 948 Arctic grayling, and 146 burbot. Information on other fish harvests will be presented in the technical report (Braem et al. *in prep*).¹²

12. Braem, N.M., C.L. Brown, M. Kostick, E. Mikow, and S. Wilson. In Prep. "Wild food harvests in 3 Upper Kobuk River Communities: Ambler, Kobuk, and Shungnak, 2012-2013." Fairbanks: Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 402.

Shungnak households harvested an estimated 15,417 edible pounds of salmon in 2012, accounting for 50% of the total fish harvests. Chum salmon harvests (2,595 fish) were the greatest, accounting for 96% of the total salmon harvest by weight, followed by 90 sockeye salmon, 15 coho salmon, and 9 pink salmon. Fifty-two percent of Shungnak households reported using less salmon in 2012 than in previous years. Nonsalmon fishes accounted for 50% of the total fish harvest, and 17% of the estimated total subsistence harvest, with an estimated total nonsalmon harvest of 15,223 edible pounds. The largest contributions to nonsalmon fish harvests were 1,556 sheefish, 1,125 least cisco, 888 broad whitefish, 660 humpback whitefish, 399 Arctic grayling, 99 Dolly Varden, 75 Bering cisco, 50 burbot, and 38 northern pike (Braem et al. *in prep*).

Kobuk residents harvested 15,142 edible pounds of salmon in 2012, accounting for 65% of the total fish harvests. Chum salmon harvests (2,637 fish) were the greatest, accounting for 99% of the total salmon harvest, followed by 14 coho salmon, 4 pink salmon, and 4 Chinook salmon. Seventy percent of Kobuk households reported using less salmon in 2012 than in previous years. Nonsalmon fishes accounted for 35% of the total fish harvest, and 19% of the estimated total subsistence harvest, with an estimated total harvest of 8,329 edible pounds. The largest contributions to nonsalmon fish harvests were 1,062 sheefish, 286 broad whitefish, 256 Arctic grayling, 157 humpback whitefish, 96 Northern pike, 40 Dolly Varden, and 23 burbot.

In a separate study, the Division of Subsistence collected subsistence fishery harvest data from Noorvik for 2012. Noorvik residents harvested 60,326 edible pounds of salmon in 2012, accounting for 31% of the total fish harvests. Chum salmon harvests (9,584 fish) were the greatest, accounting for 95% of the total salmon harvest by weight, followed by 338 coho salmon, 275 pink salmon, 81 sockeye salmon, and 7 Chinook salmon. Forty-seven percent of Noorvik households reported using less salmon than in previous years. Nonsalmon fishes accounted for 69% of the total fish harvest, and 38% of the estimated total subsistence harvest, with an estimated total harvest of 135,274 edible pounds. The largest contributions to nonsalmon fish harvests were 6,032 sheefish, representing 67,197 edible pounds and the second largest species contribution to the total subsistence harvest following caribou. Noorvik residents also harvested 10,087 broad whitefish, 5,134 northern pike, 6,406 humpback whitefish, 876 burbot, 1,445 least cisco, and 99 Dolly Varden. Information about other fish harvests will be summarized in the technical report (Braem et al. *in prep*).¹³

The Division of Subsistence's Office of Subsistence Management (OSM) project, entitled "Northwest Alaska Key Subsistence Fisheries" documented subsistence fishery harvests in Kiana and Noatak for 2012. Kiana residents harvested 16,975 edible pounds of salmon in 2012, accounting for 30% of the total fish harvests. Chum salmon harvests (2,442 fish) were the greatest, accounting for 86% of the total salmon harvest by weight, followed by 320 pink salmon, 240 coho salmon, 63 sockeye salmon, 3 Chinook salmon, and 8 unknown salmon. Twenty percent of Kiana households reported using less salmon in 2012 than in previous years. Nonsalmon fishes accounted for 70% of the total fish harvest, with an estimated total harvest of 39,955 edible pounds. The largest contributions to nonsalmon fish harvests included 1,787 sheefish, 3,596 broad whitefish, 2,307 humpback whitefish, 464 burbot, 278 northern pike, and 250 Dolly Varden.

Noatak residents harvested 50,721 edible pounds of salmon in 2012, accounting for 62% of the total fish harvests. Chum salmon harvests (7,814 fish) were the greatest, accounting for 92% of the total salmon harvest by weight, followed by 612 coho salmon, 94 sockeye salmon, 80 pink salmon, and 2 Chinook salmon. Forty-seven percent of Noatak households reported using less salmon in 2012 than in previous years. Nonsalmon fishes accounted for 38% of the total fish harvest, with an estimated total harvest of 30,849 edible pounds. The largest contributions to nonsalmon fish harvests included 6,437 Dolly Varden

13. Braem, N.M., A. Godduhn, A. Brenner, B. Retherford, and M. Kostick. In Prep. "Chukchi Sea and Norton Sound Observation Network: Golovin, Noorvik, and Point Lay, 2012". Fairbanks: Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 403.

(called “trout” by Noatak residents), 1,826 broad whitefish, 1,205 humpback whitefish, 100 sheefish, 352 Arctic grayling, and 26 northern pike.

Division of Subsistence conducted comprehensive subsistence research in Point Lay on the western North Slope of Alaska. Point Lay residents harvested 8,479 edible pounds of salmon in 2012, accounting for 58% of the total fish harvests. Chum salmon harvests (659 fish) were the greatest, accounting for 47% of the total salmon harvest by weight, followed by 1,120 pink salmon, 372 coho salmon, 14 Chinook salmon, and 13 sockeye salmon. Twenty-six percent of Point Lay households reported using less salmon in 2012 than in previous years. Nonsalmon fishes accounted for 42% of the total fish harvest, and 5% of the estimated total subsistence harvest, with an estimated total harvest of 6,076 edible pounds. The largest contributions to nonsalmon fish harvests included 1,945 Arctic grayling, 493 Dolly Varden, 279 Bering cisco, 99 gallons unknown smelt, 279 round whitefish, and 37 sheefish. Information about other fish harvests will be presented in the technical report (Braem et al. *in prep*).

The Division of Subsistence’s OSM project, entitled “Emerging North Slope Salmon Fisheries” documented subsistence fishery harvests in Wainwright for 2012. Wainwright residents harvested 1,711 edible pounds of salmon in 2012, accounting for 5% of the total fish harvests. Coho salmon represented the greatest contribution to salmon harvest by weight, with 105 coho salmon harvested, followed by 66 sockeye salmon, 51 chum salmon, 136 pink salmon, and 20 Chinook salmon. Seventeen percent of Wainwright households reported using less salmon in 2012 than in previous years. Nonsalmon fishes accounted for 95% of the total fish harvest, with an estimated total of 31,801 edible pounds. The largest contributions to nonsalmon fish harvests included 3,489 gallons of smelt (20,935 edible pounds), 1,562 broad whitefish, 1,606 humpback whitefish, 648 Bering cisco, 7,513 Arctic grayling, 101 burbot, 624 least cisco, 20 lake trout, and 20 flounder.

Arctic-Kotzebue Area Salmon Harvest Overall

Of the 2012 harvest of salmon in the Kotzebue Area for which we have data, the 6 communities harvested an estimated 29,092 salmon. The vast majority of the harvest was chum salmon (92%), followed by coho salmon (4%), pink salmon (2%), sockeye salmon (2%), and chinook salmon (<1%) (Table 3-8; Figure 3-3).

Of the 2012 harvest of salmon in the Arctic Area for which we have data, the 2 communities harvested an estimated 2,556 salmon. The majority of the harvest was pink salmon (49%), followed by chum salmon (28%), coho salmon (19%), sockeye salmon (3%), and chinook salmon (1%) (Table 3-9; Figure 3-4).

Table 3-1.—Subsistence salmon harvests by Norton Sound subdistricts, Northwest Alaska, 2012.

Subdistrict	Households surveyed or permits returned	Estimated salmon harvest ^a					
		Chinook	Sockeye	Coho	Chum	Pink	Total
Cape Woolley	20	1	0	0	29	0	30
Elim	63	42	0	1,302	1,494	10,848	13,686
Golovin	151	57	52	1,143	1,056	7,635	9,943
Nome	483	11	171	1,150	2,521	8,376	12,229
Norton Bay	77	103	0	310	2,721	2,624	5,758
Shaktoolik	63	214	9	1,110	634	4,609	6,576
St Michael	82	80	20	911	2,172	457	3,640
Stebbins	106	109	0	1,256	3,456	3,659	8,480
Unalakleet	200	673	185	4,318	2,164	8,842	16,182
Total	1,245	1,290	437	11,500	16,247	47,050	76,524

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

a. Includes subsistence harvests and commercial harvests retained for home use.

Table 3-2.—Subsistence salmon harvests by community, Norton Sound-Port Clarence Area, Alaska, 2012.

Community ^b	Households or permits		Estimated salmon harvest ^a					
	Total	Surveyed or returned	Chinook	Sockeye	Coho	Chum	Pink	Total
Anchorage	5	5	0	4	38	61	138	241
Brevig Mission	43	43	11	376	597	3,321	3,093	7,398
Diomedes	1	1	0	0	0	0	0	0
Elim	54	54	41	0	1,281	1,465	10,379	13,166
Fairbanks	1	1	0	0	0	0	0	0
Gambell	3	3	0	0	0	0	0	0
Golovin	29	29	39	44	246	775	2,415	3,519
Koyuk	83	82	104	0	373	2,731	2,837	6,045
Nome	471	471	16	878	1,724	3,168	10,385	16,171
Palmer	5	5	4	1	0	11	13	29
Savoonga	3	3	0	0	0	0	19	19
Shaktoolik	64	63	213	9	1,043	624	4,401	6,290
St. Michael	82	82	80	20	911	2,172	457	3,640
Stebbins	117	106	121	3	1,266	3,476	3,759	8,625
Teller	45	45	26	342	100	3,864	1,951	6,283
Unalakleet	223	200	661	182	4,324	2,144	8,742	16,053
Wales	1	1	0	0	0	0	0	0
White Mountain	40	40	18	0	300	237	3,662	4,217
Total	1,270	1,234	1,335	1,859	12,203	24,049	52,250	91,696

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

- a. Includes subsistence harvests and commercial harvests retained for home use.
- b. Harvest information from residents of non-local communities (e.g. Anchorage) is available only for Norton Sound and Port Clarence permit areas. Non-local residents might subsistence fish in other northwest Alaska areas, but these harvests are not documented in the regional household surveys.

Table 3-3.—Historical subsistence salmon harvests by district, Northwest Alaska, 1994–2012.

Norton Sound District							
Year	Number of households	Chinook	Sockeye	Coho	Chum	Pink	Total
1994	839	7,212	1,161	22,108	24,776	70,821	126,077
1995	851	7,766	1,222	23,015	43,014	38,594	113,612
1996	858	7,255	1,182	26,304	34,585	64,724	134,050
1997 ^a	1,113	8,998	1,892	16,476	26,803	27,200	81,370
1998 ^a	1,184	8,295	1,214	19,007	20,032	51,933	100,480
1999	898	6,144	1,177	14,342	19,398	20,017	61,078
2000	860	4,149	682	17,062	17,283	38,308	77,485
2001	878	5,576	767	14,550	20,213	30,261	71,367
2002	935	5,469	763	15,086	17,817	64,354	103,490
2003	940	5,290	801	14,105	13,913	49,674	83,782
2004	1,003	3,169	363	8,225	3,200	61,813	76,770
2005	1,061	4,087	774	13,896	12,008	53,236	84,000
2006	1,066	3,298	901	19,476	10,306	48,764	82,745
2007	1,041	3,744	923	13,564	18,170	21,714	58,116
2008	1,151	3,087	399	18,889	11,505	56,096	89,976
2009	1,200	5,131	388	15,852	10,599	26,110	58,080
2010	1,030	2,074	554	11,517	14,295	38,710	67,149
2011	925	1,645	562	10,155	12,946	18,576	43,883
2012	1,245	1,290	437	11,500	16,247	47,050	76,524

Port Clarence District							
Year	Number of households	Chinook	Sockeye	Coho	Chum	Pink	Total
1994	151	203	2,220	1,892	2,294	4,309	10,918
1995	151	76	4,481	1,739	6,011	3,293	15,600
1996	132	194	2,634	1,258	4,707	2,236	11,029
1997	163	158	3,177	829	2,099	755	7,019
1998	157	289	1,696	1,759	2,621	7,815	14,179
1999	177	89	2,392	1,030	1,936	786	6,233
2000	163	72	2,851	935	1,275	1,387	6,521
2001	160	84	3,692	1,299	1,910	1,183	8,167
2002	176	133	3,732	2,194	2,699	3,394	12,152
2003	242	176	4,436	1,434	2,425	4,108	12,578
2004	371	278	8,688	1,131	2,505	5,918	18,520
2005	329	152	8,532	726	2,478	6,593	18,481
2006	345	133	9,862	1,057	3,967	4,925	19,944
2007	362	85	9,484	705	4,454	1,468	16,196
2008	399	125	5,144	562	2,499	7,627	15,957
2009	328	40	1,643	799	3,060	1,887	7,429
2010	295	57	824	596	5,232	5,202	11,911
2011	271	56	1,611	393	4,338	2,610	9,008
2012	335	44	1,422	703	7,802	5,201	15,172

-continued-

Table 3-3.–Page 2 of 2.

Year	Kotzebue District ^{b,i}						
	Number of households	Chinook	Sockeye	Coho	Chum	Pink	Total
1994 ^c	557	135	33	478	48,175	3,579	52,400
1995 ^d	1,327	228	935	2,560	102,880	2,059	108,662
1996	1,187	550	471	317	99,740	951	102,029
1997	1,122	464	528	848	57,906	1,181	60,925
1998	1,279	383	392	461	48,979	2,116	52,330
1999	1,277	9	478	1,334	94,342	841	97,004
2000	1,227	211	75	2,557	65,975	75	68,893
2001 ^e	1,149	11	14	768	49,014	36	49,844
2002 ^f	216	3	9	56	16,880	8	16,955
2003 ^g	488	40	53	1,042	19,201	583	20,918
2004 ^g	440	54	18	1,502	23,348	1,259	26,181
2005 ^h	ND	ND	ND	ND	ND	ND	ND
2006 ^h	ND	ND	ND	ND	ND	ND	ND
2007 ^h	ND	ND	ND	ND	ND	ND	ND
2008 ^h	ND	ND	ND	ND	ND	ND	ND
2009 ^h	ND	ND	ND	ND	ND	ND	ND
2010 ^h	ND	ND	ND	ND	ND	ND	ND
2011 ^h	ND	ND	ND	ND	ND	ND	ND
2012 ^g	360	16	455	1,230	26,694	697	29,092

Year	Arctic District ^k						
	Number of households	Chinook	Sockeye	Coho	Chum	Pink	Total
2012	120	34	79	477	710	1,256	2,556

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

- a. Includes Gambell and Savoonga.
- b. Normally includes Ambler, Kiana, Kobuk, Kotzebue, Noatak, Noorvik, and Shungnak.
- c. Includes Deering and Wales; does not include Kotzebue.
- d. Includes Shishmaref.
- e. Does not include Ambler.
- f. Includes only Noatak and Noorvik.
- g. Does not include Kotzebue.
- h. Due to lack of funding, no collection of subsistence salmon harvest data took place in Kotzebue area communities from 2005–2011. The average yearly subsistence harvest of salmon in the Kotzebue area between 1994 and 2004 was 59,650 fish.
- i. Formerly Kotzebue Area.
- j. Limited data exist in 2006, 2007 and 2011 for Kiana (2006), Kivalina (2007), Noatak (2007), and Selawik (2011). These are available online through the Community Subsistence Information System (CSIS) at <http://www.adfg.alaska.gov/sb/CSIS/>
- k. Includes Point Lay and Wainwright.

ND = no data.

Table 3-4.—Subsistence salmon harvests by district, Northwest Alaska, 2012.

District	Households surveyed or permits returned	Estimated salmon harvest ^a					
		Chinook	Sockeye	Coho	Chum	Pink	Total
Norton Sound District ^b	1,245	1,290	437	11,500	16,247	47,050	76,524
Port Clarence District ^c	335	44	1,422	703	7,802	5,201	15,172
Kotzebue District ^{d,f}	360	16	455	1,230	26,694	697	29,092
Arctic District ^g	120	34	79	477	710	1,256	2,556
Total^e	1,412	1,384	2,393	13,910	51,453	54,204	123,344

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

- a. Harvests reported during household surveys are expanded into estimates to account for uncontacted households. Harvests reported on permits are not expanded.
- b. Household surveys conducted in Unalakleet, Koyuk, and Shaktoolik. Permits issued for Cape Woolley, Nome Subdistrict (Tier I), Golovin Subdistrict, and Elim Subdistrict.
- c. Permits issued for Port Clarence District, Pilgrim River, and Salmon Lake.
- d. Due to lack of funding, no collection of subsistence salmon harvest data took place in Kotzebue Sound communities from 2005-2011. The average yearly subsistence harvest of salmon in the Kotzebue area between 1994 and 2004 was 59,650 fish.
- e. Households surveyed or permits returned column does not add up to the total shown above due to individual households fishing in multiple districts.
- f. Formerly Kotzebue Area.
- g. Includes Point Lay and Wainwright.

Table 3-5.—Historical subsistence salmon harvests, Northwest Alaska, 1975–2012.

Year	Households or permits		Estimated salmon harvest ^a					
	Total	Surveyed or returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1975	117	79	3	225	102	3,698	7,298	11,326
1976	138	104	6	0	275	1,856	5,472	7,609
1977	195	181	35	64	623	12,222	2,839	15,783
1978	168	126	31	0	242	4,035	10,697	15,005
1979	138	119	519	0	1,007	3,419	5,842	10,787
1980	232	161	135	0	2,075	5,839	21,728	29,777
1981	236	169	47	88	1,844	9,251	6,100	17,330
1982	230	182	33	6	2,093	5,719	20,480	28,331
1983	243	189	74	40	1,950	7,013	8,499	17,576
1984	240	189	85	0	1,890	4,945	18,067	24,987
1985	215	198	56	114	1,054	5,717	2,117	9,058
1986	279	240	157	127	788	8,494	9,011	18,577
1987	235	173	97	102	812	7,265	705	8,981
1988	192	166	67	171	1,089	6,379	2,543	10,249
1989	173	130	24	131	549	3,456	924	5,084
1990	188	165	60	234	542	4,525	2,413	7,774
1991	155	128	83	166	1,279	3,715	194	5,437
1992	163	132	152	163	1,720	2,030	7,746	11,811
1993	142	104	51	74	1,780	1,578	758	4,241
1994	1,547	1,169	7,713	3,414	24,494	75,489	78,954	190,063
1995 ^b	2,329	1,445	8,070	6,639	27,314	151,905	43,947	237,874
1996	2,177	1,454	7,999	4,287	27,879	139,032	67,911	247,108
1997 ^c	2,398	1,645	9,620	5,597	18,153	86,808	29,135	149,314
1998 ^c	2,620	1,730	8,967	3,301	21,226	71,632	61,863	166,989
1999	2,351	1,300	6,242	4,046	16,706	115,676	21,644	164,315
2000	2,247	1,336	4,399	3,612	20,654	84,196	40,499	153,360
2001 ^d	2,192	1,259	5,671	4,473	16,617	71,138	31,480	129,378
2002 ^e	1,327	1,204	5,624	4,504	17,838	37,396	67,756	133,119
2003 ^f	1,670	1,488	5,505	5,289	16,580	35,540	54,365	117,279
2004 ^g	1,915	1,814	3,534	9,159	11,585	31,386	70,841	126,506
2005 ^{g,h}	1,129	1,104	4,239	9,306	14,622	14,486	59,829	102,481
2006 ^{g,h}	1,125	1,099	3,431	10,763	20,533	14,273	53,689	102,689
2007 ^{g,h}	1,122	1,073	3,829	10,407	14,269	22,624	23,182	74,312
2008 ^h	1,247	1,172	3,212	5,543	19,451	14,004	63,723	105,933
2009 ^h	1,274	1,206	5,171	2,031	16,651	13,659	27,997	65,509
2010 ^h	1,106	1,032	2,131	1,378	12,113	19,527	43,912	79,060
2011 ^h	1,044	932	1,701	2,173	10,548	17,284	21,186	52,891
2012 ^{f,i}	2,034	1,714	1,384	2,393	13,910	51,453	54,204	123,344
5-year average (2007–2011)	1,159	1,083	3,209	4,306	14,606	17,420	36,000	75,541
10-year average (2002–2011)	1,296	1,212	3,838	6,055	15,419	22,018	48,648	95,978
Historical average (1975–2011)	932	713	2,670	2,639	9,431	30,195	26,901	71,835

-continued-

Table 3-5.–Page 2 of 2.

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Note Since 1994 ADF&G has conducted an annual subsistence salmon harvest assessment effort in Northwest Alaska that provides more extensive and reliable estimates. Harvest estimates prior to 1994 cannot be directly compared.

- a. Includes selected communities in the Norton Sound District, Port Clarence District, Arctic District, and Kotzebue District (formerly Kotzebue Area).
- b. Includes Shishmaref.
- c. Includes Gambell and Savoonga.
- d. Does not include Ambler.
- e. For the Kotzebue District, includes only Noatak and Noorvik.
- f. Does not include Kotzebue.
- g. Does not include Koyuk.
- h. Does not include Kotzebue District.
- i. For the Arctic District, includes only Point Lay and Wainwright.

Table 3-6.–Subsistence salmon harvests by Kotzebue District^a communities.

Year	Community	Households		Estimated salmon harvest					
		Total	Surveyed	Chinook	Sockeye	Coho	Chum	Pink	Total
2007	Kivalina	42	81	41	0	33	401	120	594
	Noatak	90	119	11	42	247	4,167	163	4,630
Total, 2007		132	200	51	42	280	4,568	283	5,224
2011	Selawik	169	61	0	167	7	879	0	1,053
2012	Ambler	76	53	1	126	11	1,621	9	1,769
	Kiana	103	65	3	63	240	2,442	320	3,068
	Kobuk	36	30	4	0	14	2,637	4	2,659
	Noatak	126	83	2	94	612	7,814	80	8,601
	Noorvik	135	83	7	81	338	9,584	275	10,285
	Shungnak	69	46	0	90	15	2,595	9	2,709
Total, 2012		545	360	16	455	1,230	26,694	697	29,092

Source ADF&G Division of Subsistence, household surveys, 2008, 2012, 2013.

- a. Formerly Kotzebue Area.

Table 3-7.—Subsistence nonsalmon harvests by Kotzebue District^a communities.

Year	Community	Households		Estimated number of fish								Total
		Total	Surveyed	Dolly Varden	Arctic grayling	Burbot	Broad whitefish	Humpback whitefish	Northern pike	Saffron cod	Sheefish	
2007	Kivalina ^b	42	81	20,527	786	15	ND	ND	0	25,824	0	47,152
	Noatak ^b	90	119	10,234	1,222	42	ND	ND	144	192	99	11,933
Total, 2007		132	200	30,761	2,008	58	0	0	144	26,015	99	59,086
2011	Selawik	169	61	19	815	1,081	47,394	12,647	15,956	0	6,190	84,102
2012	Ambler	76	53	85	948	146	9,150	1,544	568	0	1,156	13,597
	Kiana	103	65	249	ND	464	3,596	2,307	278	ND	1,787	8,682
	Kobuk	36	30	40	256	23	286	157	96	0	1,062	1,919
	Noatak	126	83	6,437	352	ND	1,826	1,205	26	ND	100	9,946
	Noorvik	135	83	99	28	876	10,087	6,406	5,134	0	6,032	28,662
	Shungnak	69	46	99	399	50	888	660	38	0	1,556	3,689
Total, 2012		545	360	7,008	1,983	1,559	25,833	12,280	6,139	0	11,694	66,496

Source ADF&G Division of Subsistence, household surveys, 2008, 2012, 2013.

a. Formerly Kotzebue Area.

b. Harvest information is available for whitefishes as a species category only. Kivalina harvested 338 whitefishes and Noatak harvested 6,778 in 2007.

ND = no data

Table 3-8.—Subsistence salmon harvests by Arctic District communities.

Community	Households		Estimated salmon harvest					
	Total	Surveyed	Chinook	Sockeye	Coho	Chum	Pink	Total
Point Lay	67	42	14	13	372	659	1,120	2,178
Wainwright	152	78	20	66	105	51	136	378
Total	219	120	34	79	477	710	1,256	2,556

Source ADF&G Division of Subsistence, household surveys, 2013.

Table 3-9.—Subsistence nonsalmon harvests by Arctic District communities.

Community	Households		Estimated number of fish					
	Total	Surveyed	Arctic char / Dolly Varden	Arctic grayling	Sheefish	Smelt	Whitefishes	Total
Point Lay	67	42	493	1,945	37	99	1,278	3,851
Wainwright	152	78	0	7,513	0	3,489	4,440	15,442
Total	219	120	493	9,458	37	3,588	5,718	19,293

Source ADF&G Division of Subsistence, household surveys, 2013.




Figure 3-1.—Species composition of estimated subsistence salmon harvests, Norton Sound District, 2012.




Figure 3-2.—Species composition of estimated subsistence salmon harvests, Port Clarence District, 2012.




Figure 3-3.—Species composition of estimated subsistence salmon harvests, Kotzebue District, 2012.




Figure 3-4.—Species composition of estimated subsistence salmon harvests, Arctic District, 2012.

CHAPTER 4: YUKON AREA

BACKGROUND

Residents of the Yukon River drainage have long relied on fish for human food and other subsistence uses. While nonsalmon fish species are an important component of the overall fish harvest (Andersen et al. 2004; Brown et al. 2005), large numbers of Chinook salmon, summer and fall chum salmon, and coho salmon compose the majority of all subsistence harvests of fish in the Yukon River drainage. Indeed, subsistence salmon harvests occur alongside robust commercial, sport, and personal use harvests across species.

Yukon Area fishers use drift gillnets, set gillnets, and fish wheels to harvest the majority of salmon. Set gillnets are used throughout the Yukon Area, often in the main rivers and coastal marine waters. Drift gillnets are used extensively in about half of the river. Under state regulations, drift gillnets are allowed from the mouth of the Yukon River to approximately 18 miles downriver of Galena, and under federal permit in subdistricts 4B and 4C. Fish wheels are a legal subsistence gear type throughout the Yukon drainage, although due to river conditions and the availability of wood for building materials, they are used almost exclusively on the middle and upper Yukon and Tanana rivers.

Depending on the area of the Yukon River drainage and each salmon species' run timing, subsistence fishing for salmon occurs from late May through mid October. Subsistence harvesters usually base their fishing activities either from fish camps or from their home communities. Extended family groups, typically representing several households, often participate in subsistence salmon fishing together. Households and related individuals typically cooperate to harvest, process, preserve, and store salmon for subsistence uses.¹⁴

The majority of the subsistence salmon harvest is preserved for later uses by freezing, drying, or smoking; the head, viscera, backbones, and other scraps are often fed to dogs. Chinook salmon are harvested and processed primarily for human consumption, although those fish deemed not suitable for human consumption due to the presence of the fungus *Ichthyophonus hoferi* or some other disease or abnormality are often fed to dogs. Small male Chinook salmon ("jacks") or spawned-out salmon may also be fed to dogs. In addition, while fishers harvest chum and coho salmon primarily for human consumption, dog mushers harvest and process relatively large numbers of these species as food for sled dogs. Fall chum salmon and coho salmon typically arrive in the upper portion of the drainage late in the season, coinciding with freezing weather, during which time some dog mushers "crib" salmon for use as dog food. This method involves storing whole salmon outdoors in large wooden boxes or log cribs in late fall, and allowing them to freeze (Andersen 1992). The practice of keeping sled dogs is much more common in communities along the upper Yukon River than in the lower river area.

REGULATIONS

Regulation and management of Yukon River drainage subsistence salmon fishing follows the Yukon River Drainage Subsistence Salmon Fishery Management Protocol, which provides a framework for coordinated subsistence fisheries management between ADF&G and the federal subsistence management programs in the Yukon River drainage. This protocol is applied through a Memorandum of Agreement between state and federal agencies which formalizes the working relationships between state and federal managers and fosters cooperation with federal regional advisory councils and fisheries interest groups. State managers are responsible for management of state subsistence, commercial, recreational, and personal use fisheries in all waters. Federal managers are responsible for management of subsistence fishing by qualified rural residents in applicable federal waters. The protocol also directs state and federal

14. For more detail on subsistence uses of Yukon River salmon, see (ADF&G 1987a–b, 1988).

managers to solicit input from the Yukon River Drainage Fisheries Association (YRDFA), the Yukon River Coordinating Fisheries Committee (YRCFC), and other stakeholders during the decision-making process.

The majority of the United States' portion of the Yukon Area is open to subsistence fishing. However, the Joint Board has defined a portion of the Tanana River in the Yukon River drainage as lying within the Fairbanks Nonsubsistence Area (5 AAC 99.015). Subsistence fisheries may not be authorized within nonsubsistence areas; the harvest of fish for home uses in these nonsubsistence areas occurs under personal use and sport fishing regulations.

In 2011 area managers implemented a 2010 Board of Fisheries decision to reduce the maximum stretched mesh net size to 7.5". Prior to this, Yukon Area fishers widely used 8"-8.5" mesh nets to target Chinook salmon. This change was considered a conservation tool that should allow more of the older and larger Chinook salmon, especially females, to escape to the spawning grounds. During the spring of 2011, in cooperation with the Pacific States Marine Fisheries Commission (PSMFC), the Tanana Chiefs Conference (TCC) administered a net exchange program to help fishermen comply with this new regulation. Eligible fishermen could send in their 8" or larger mesh nets in exchange for new, 7.5" mesh nets (Frothingham 2011). Kwik'pak Fisheries, LLC, a community-based economic development organization in the lower Yukon River, also facilitated a net exchange program in lower river communities.

Over the last 2 decades, several other regulatory changes have affected the subsistence salmon fishery in the Yukon River drainage. In 1993, the BOF adopted regulations which separated subsistence and commercial salmon fishing times in districts 1, 2, and 3 and in the lower portion of District 4 (Subdistrict 4A) (figures 4-1 and 4-2). In these areas, subsistence salmon fishing is allowed 7 days per week but may not occur 24 hours prior to and immediately following the commercial salmon fishing season. By regulation, once the commercial season is open, subsistence salmon fishing may not occur 18 hours immediately before, during, and 12 hours after each district 1, 2, or 3 summer season commercial fishing period unless altered by emergency order. During the fall season in districts 1, 2, and 3, subsistence fishing may not occur 12 hours immediately before, during, and 12 hours after each commercial fishing period. In areas including and upriver from subdistrict 4B, subsistence salmon fishing periods coincide with commercial salmon fishing periods. Since 1988, subsistence fishing in the lower Tanana River drainage in subdistricts 6A and 6B has been allowed for two 42-hour periods per week unless altered by emergency order.¹⁵ In the upper Tanana River drainage upstream of the Volkmar (north bank) and Johnson (south bank)¹⁶ rivers, subsistence fishing is allowed 7 days per week.

In 2005, the Federal Subsistence Board (FSB) established a subsistence drift gillnet fishery in subdistricts 4B and 4C, which includes the mainstem Yukon River villages of Galena and Ruby. Participation in this fishery was open to qualified rural residents under a federal subsistence permit, and limited to gillnets that were no longer than 150 ft and no deeper than 35 meshes. The mesh size was unrestricted so as to target Chinook salmon. In previous years, the regulation allowed drift gillnet fishing in the last 18 hours of each subsistence salmon fishing opening in the federal public waters of subdistricts 4B and 4C. However, in 2008, the FSB liberalized the regulation to align it with the regulatory openings, usually two 42-hour periods per week.

15. In the lower Tanana River drainage, the fishery to harvest salmon for home uses in Subdistrict 6C is a personal use fishery. Its fishing schedule matches those of the 6A and 6B subsistence salmon fisheries; namely, that personal use fishing is allowed for two 42-hour periods per week unless altered by emergency order. In that portion of Subdistrict 6B from the downstream side of the upper Tolovana River to 3 miles upstream of Totchaket Slough (the Old Minto area), subsistence fishing is allowed 5 days per week.

16. Salmon fishing is closed in that portion of the Tanana River drainage upstream of Subdistrict 6C, from the Salcha River upstream to the Volkmar River (north bank) and to the Johnson River (south bank). The area is closed to salmon fishing other than sport fishing and is included in the Fairbanks Nonsubsistence Area. Whitefishes and longnose suckers may be harvested upstream of the Salcha River under a personal use permit.

Since 1996, Yukon River salmon stocks have fluctuated in terms of abundance. Since the disastrous run in 2000, when restrictions were imposed late in the summer subsistence salmon season to protect Chinook salmon and summer chum salmon populations, Chinook salmon runs have been variable. Because of the inability to maintain expected yields and harvestable surpluses above escapement goals for several years, the BOF classified the Yukon River Chinook salmon stock as a stock of yield concern at its September 2000 work session (Lingnau and Salomone 2003). After a modest increase in Chinook salmon abundance from 2004 to 2007, more severe restrictions were imposed on the summer season to protect declining Chinook salmon runs beginning in 2008. Restrictions have been implemented through both period closures and limited gear use in some districts. During its January 2010 and 2013 meetings in Fairbanks, the BOF continued the stock of yield concern designation for Yukon River Chinook salmon.¹⁷

Fall chum salmon returns have also been variable over time. Restrictions on subsistence fall season salmon fishing occurred intermittently throughout the 1990s. There was a complete closure of the fall season in 2000 severely affecting the subsistence harvest of fall chum and coho salmon. In 2001 the BOF declared Yukon fall chum salmon a stock of concern. In 2007, after the returns of fall chum rebounded, the BOF lifted the stock of concern designation.

In 2001, as a result of the disastrous runs the year before, the BOF instituted a new subsistence schedule on the Yukon River based on historical subsistence fishing opportunities structured around commercial openings, commonly referred to as the “windows” schedule. The schedule was intended to fulfill several goals: 1) increase the quality of escapement, 2) distribute subsistence opportunity among users during years with no commercial fishing, and 3) reduce the impact of harvest on any one stock by spreading the harvest throughout the run, thereby providing windows of time that salmon may migrate upriver with reduced exploitation. The schedule, based on past fishing schedules, is initiated each year based on the historical average time of Chinook salmon entry into the Yukon River. Once initiated, the schedule is implemented chronologically upriver. The BOF determined that the schedule provides reasonable opportunity for subsistence users to achieve their harvest goals when salmon runs are below average. Subsistence fishing is allowed 7 days per week in all areas prior to the established schedule dates. In 2003, the BOF clarified the window schedule to allow ADF&G to relax the schedule if Chinook salmon run abundance allowed commercial fishing.

Preseason outlooks for 2012 projected a below average to poor Chinook salmon run, especially for Canadian-origin fish. While YRDFA had facilitated a series of regional teleconferences and meetings for managers, fishers, and other stakeholders to discuss options and develop a preseason plan, no such meeting took place prior to the 2012 season due to funding constraints. However, state and federal managers collaborated with fishermen, tribal council representatives and other stakeholders to develop a preseason strategy and distributed an informational flyer to approximately 2,900 Yukon River households. (JTC 2013:4). The 2012 subsistence fishing schedule for the Lower Yukon Area is presented in Table 4-1. Table 4-2 displays the 2012 subsistence fishing schedule for the Upper Yukon Area. The 2012 season marked the twelfth annual implementation of the windows schedule. Historically, the windows schedule began around May 28 in District 1. In 2012 the regulatory subsistence fishing schedule began on May 31. The summer chum salmon run was expected to be similar to the run in 2011 of approximately 2 million fish and to provide for escapements, a normal subsistence harvest, and a surplus for commercial harvest. A projected run of 986,000–1,242,000 fall chum salmon was expected to provide for escapement, subsistence harvests, and a projected commercial harvest of 500,000–700,000 fish. Coho salmon runs were projected to be of average strength based on escapements observed in 2008 (JTC (Joint Technical Committee of the Yukon River US/Canada Panel) 2012).

Throughout the season, emergency orders were issued to modify the subsistence fishing schedule to protect Chinook salmon. Beginning on June 6 through June 12, 2012 gillnet fishing gear was restricted to

17. Alaska Department of Fish and Game Division of Commercial Fisheries. “Yukon Area regulatory changes,” news release, January 25, 2013. Accessed August 2014. <http://www.adfg.alaska.gov/static/applications/dcfnewsrelease/245219141.pdf>

6 inch or smaller mesh size in the southern portion of the Coastal District. Because of a later than average run timing of the first pulse of Chinook salmon, gillnets were restricted to 6 inch or less in Districts 1 and 2 beginning on June 18 and 20, respectively. Based on inseason information, subsistence closures were initiated in District 1 on June 20 to protect the first pulse of Chinook salmon and implemented chronologically upriver as the pulse migrated. As the Chinook salmon migration progressed upriver, managers were increasingly concerned that the run would likely be near or below the lower end of the preseason projection. In order to try to meet escapement goals and to equitably distribute available subsistence harvest throughout the drainage, managers implemented additional conservation measures. Managers restricted the southern portion of the Coastal District to a 6 inch or smaller mesh net for the remainder of the season; in the northern portion of the Coastal District and in Districts 1 through 5, they implemented a second pulse closure immediately following the first. This second pulse closure was followed by a shortened window opportunity restricted to 6 inch or less mesh nets in Districts 1 through 4 in order to harvest some summer chum salmon while conserving Chinook salmon. Low passage numbers at Eagle sonar necessitated further restrictions in Subdistrict 5-D; as a result, subsistence closures were most pronounced in Subdistrict 5-D (JTC 2013).

During the preseason, it was considered unlikely that there would be a directed commercial Chinook salmon fishery given previous years' failures to meet minimum treaty escapement goals for Canadian Chinook salmon and because of the likely restrictions on the subsistence fishery. Ultimately, given the need for subsistence restrictions indicated by the preseason outlook and early inseason indicators, ADF&G did not authorize any commercial fishing periods targeting Chinook salmon on the mainstem Yukon River.

Preseason projections expected the 2012 summer chum run to be average, to provide for escapement and subsistence uses, and to have a surplus for commercial harvest, noted above. A harvestable surplus of summer chum has been available for the last 10 years (2003–2012). Because of the concurrent run timing of Chinook and summer chum salmon, managers expected that the conservative management strategies, designed to protect a poor Chinook salmon run, would affect and reduce the commercial harvest of summer chum salmon. To reduce the incidental catch of Chinook salmon in the summer chum commercial fishery, commercial fishing was delayed until the estimated midpoint of Canadian-origin Chinook salmon run had passed through the Lower Yukon Area. Additionally, commercial openings were instituted concurrently with subsistence openings to streamline openings with the hope of reducing the amount of time that Chinook salmon were susceptible to harvest. Despite these measures, a total of 2,421 Chinook salmon were incidentally caught and recorded but not sold during the summer season (JTC 2013:10). A new regulation adopted by the BOF in 2012 allowed ADF&G to open a commercial summer chum fishery in Subdistrict 4-A using fish wheels that were attended at all times in order to immediately release Chinook salmon back to the water alive. A total of 59 Chinook salmon were reported caught and released back to the water alive in the fish wheel fishery (JTC 2013:10). Kwik'pak Fisheries LLC donated 382 Chinook and 2,280 summer chum salmon to 8 surveyed communities (Beaver, Birch Creek, Fort Yukon, Hughes, Huslia, Stevens Village, Tanana); these fish were donations from fishermen, commercial harvests, or caught in the test fishery near Emmonak (Jallen et al. In prep:19). The prohibition on the sale of Chinook salmon continued through the fall season.

The preseason outlook for fall chum salmon estimated a return of greater than 800,000 fish, enough to meet the escapement goal and provide for subsistence harvests, and support a commercial harvest (JTC 2013:11). In 2012, fall chum salmon returned in numbers great enough to satisfy escapement and subsistence needs and provide an opportunity for commercial fishing. A directed commercial fall chum fishery resulted in a harvest of 289,692 fall chum and 74,789 coho salmon. Both harvests were above their respective most recent 5-year (2007–2011) and 10-year (2002–2011) averages and were the largest since 1995 and 1991 respectively (JTC 2013:11). A total of 469 permit holders participated in the fall season salmon commercial fishery; 457 in districts 1 and 2 and 12 in districts 4, 5, and 6 combined. Participation in all districts during the 2011 fishing season was above historical averages and the 2012

permit totals were approximately 97% above the 2002–2011 average of 238 permit holders (JTC 2013:11).

Similarly, preseason projections expected the 2012 coho salmon run to be average and able to provide for escapement and subsistence uses.

SUBSISTENCE HARVEST ASSESSMENT METHODS

For the majority of villages within the Yukon Area, there are no regulatory requirements to report subsistence salmon harvests. For these villages, ADF&G utilizes a voluntary survey program to estimate the total subsistence salmon harvest. Harvest information is collected using a combination of subsistence harvest calendars mailed prior to fishing activities and postseason surveys conducted in person or by phone or letter. In road-accessible portions of the Yukon Area—including the majority of the Tanana River drainage (subdistricts 6A and 6B and the Upper Tanana River drainage), the Yukon River drainage between Hess Creek and the Dall River (known as the Yukon River Bridge Area), the upper portion of Subdistrict 5D between the upstream mouth of Twenty-two Mile Slough and the U.S.–Canada border, and, as of 2004, the Rampart Area (western end of Garnet Island to the mouth of Hess Creek), and the Middle and South Fork Area of the Koyukuk River—subsistence fishers are required to obtain an annual household permit prior to fishing, document their subsistence salmon harvests on the household permit, and return it to ADF&G at the end of the season.

Prior to salmon fishing activities, subsistence harvest calendars are mailed to all identified fishing households within the survey communities. The Lower Yukon Area calendars contain the months of May through September and the Upper Yukon Area calendars contain the months of June through October. Additional calendars are mailed to those households for which fishing activities are unknown and are also made available to households upon request from ADF&G offices in Emmonak and Fairbanks. The calendars provide space for fishers to record their daily subsistence harvests of salmon by species. Calendars are return-postage-paid and are mailed to ADF&G or given to ADF&G research staff during postseason trips to the villages, especially during the postseason salmon survey. Posters sent to village post offices and announcements on area radio stations remind fishers to give their calendars to research staff. In 2012, Division of Commercial Fisheries staff distributed calendars to all households identified as participating in some level of fishing or with unknown fishing harvests; households identified as nonfishing households did not receive calendars. A total of 1,610 calendars were sent to Yukon River households. Approximately 18% of calendar recipients (288) returned harvest calendars either by mail or through research staff during their fall surveys. Calendars provide additional Yukon Area run and harvest timing information that is not obtained by other data collection methods (Jallen et al. In prep).

In addition to the harvest calendars, ADF&G Division of Commercial Fisheries staff conduct postseason in-person interviews with a stratified random sample of all households in surveyed communities within the Yukon River drainage. Survey questions focus on Chinook, summer chum, fall chum, and coho salmon, but households are also asked about other species as well, such as pink salmon (primarily taken by coastal communities), northern pike *Esox lucius*, whitefishes, and sheefish. Some households that are not contacted in person by the surveyors are contacted by telephone. Those households not contacted by telephone are mailed a survey questionnaire and a postage-paid return envelope.

A subsistence permit is required in the road-accessible portions of the Yukon River drainage. Subsistence fishers record their daily salmon harvests on a household permit and return the permit within 10 days of the expiration date on the permit. Subsistence permit applications are mailed to all who returned the prior year's permit, along with instructions on how to apply by mail. In addition, ADF&G staff travel to select villages so that applicants can be issued permits in person. Permits are also issued in several ADF&G offices or by mail throughout the season. Those who do not return permits are sent up to 2 reminder letters. Telephone contacts with households that do not respond to the reminder letters are attempted as a final measure.

Subsistence salmon permit holders in a portion of Subdistrict 6B (the Tanana River drainage above a point 3 miles upstream of Totchaket Slough to the boundary with Subdistrict 6C) and the personal use fishers in Subdistrict 6C are required to report their harvests weekly for inseason management purposes. To maximize the return of permits, ADF&G staff also sent reminder letters to these households. A total of 478 salmon fishing permits were issued to households in the Yukon Area in 2012, including 406 subsistence and 72 personal use permits (Table 4-3). Of these permits, 380 (94%) subsistence permits and 70 (97%) personal use permits were returned to ADF&G (Table 4-3) (Jallen et al. In prep). Unreturned permits were considered to be unfished, and subsistence fishing households are not eligible to receive a permit the following year until the previous year's permit is returned.

Department staff surveyed 985 households in the Yukon Area concerning their subsistence salmon harvests). Additional information for 103 households was collected by telephone, and 37 households mailed in their harvest numbers on a survey or a calendar (Jallen et al. In prep:15). Based on these various methods of collecting harvest data, it was estimated that 1,575 Yukon Area households (out of approximately 3,133 households) participated in subsistence and personal use fishing in 2012 (Table 4-4).

SUBSISTENCE SALMON HARVESTS IN 2012

In 2012, 1,125 surveyed households and 450 permit holders that returned permits (50% of the 3,133 total estimated households in districts 1–6) provided harvest data for the Yukon Area subsistence–personal use salmon fishery (Table 4-3; Table 4-5). The estimated subsistence–personal use salmon harvest for the entire Yukon Area included 30,486 Chinook salmon (11% of the estimated total salmon harvest), 127,313 summer chum salmon (45%), 99,719 fall chum salmon (35%), 21,633 coho salmon (8%), and 5,150 pink salmon (2%), for a total of 284,301 salmon (Table 4-5; Figure 4-2). Note that this is an estimated total based on household surveys and returned permits and calendars, and it includes subsistence harvests, personal use harvests, commercial harvests retained for subsistence, and fish distributed from ADF&G test fisheries.

Since the disastrous harvest levels in 2000 (152,300 total salmon), subsistence salmon harvests have fluctuated by species. Chinook salmon harvest levels have remained relatively stable except in years when conservative management actions were taken. As shown in Table 4-6, the 2012 Chinook salmon harvest estimates were below the most recent Yukon Area 5-year averages (2007–2011), likely reflecting the restrictions put in place to protect them. The estimated subsistence and personal use harvest of 30,486 Chinook salmon in 2012 was 31% below the most recent 5-year average of 44,065 fish, and 37% below the most recent 10-year average of 48,136 fish. Other explanations for decreases in Chinook harvest include voluntary reduction of harvest by Yukon River communities and individual households. Regardless, subsistence Chinook salmon harvests have not fallen within the amounts necessary for subsistence (ANS) range for the last 5 years. Households could also replace some of their Chinook harvest with other, more abundant, salmon species. Summer and fall chum salmon for example, both experienced substantially increased harvests in 2012 from 2011 and 2010, possibly demonstrating species replacement strategies. With the exception of Chinook salmon, the harvests of all other salmon species in 2012 were higher than their respective 5-year averages.

From 1990 to 1997, when the salmon roe market declined, subsistence harvests of summer chum salmon were regularly estimated between 115,000 and 142,000 fish. Fishers harvested summer chum salmon for roe and kept most of the carcasses primarily for dog food; these fish were counted in the subsistence harvest. Since then, summer chum harvests have been relatively stable. The estimated 2012 subsistence harvest of 127,313 summer chum salmon was 43% above the 5-year average of 89,145 fish and 41% above the 10-year average of 90,530 fish. Summer chum salmon may continue to play a larger role in subsistence salmon harvests if Chinook salmon harvests continue to decline as subsistence users attempt to adapt to changes in Chinook salmon availability.

Fall chum salmon harvests steadily declined in the late 1980s through the 1990s. Since then, harvest has fluctuated. Unlike summer chum, fluctuations in harvest are less connected to the commercial market

(Figure 4-3). Fall chum salmon are used as both human food and dog food, depending on quality and timing of harvests within the run. 2012 marks the first in 5 years that the harvest of fall chum salmon fell within the ANS range (Table 4-7); declines in the maintenance of dog teams along the river likely account for this change in harvest levels. Historically, due to run timing, the management of coho salmon has been tied to the management of fall chum salmon. As such, it is difficult to assess actual trends in the harvest of coho salmon and reasons for these trends.

Pink salmon are on the calendar and survey for the entire survey area, but harvest is typically only reported in lower river communities. Although pink salmon can be abundant in lower Yukon River and coastal Yukon River delta communities, fishers do not typically harvest large numbers of this species.

Figure 4-4 shows the number of dogs reported by surveyed households in each fishing district, as well as the percentage of total dogs in the Yukon Area reported in each district. Of the estimated 1,655 households in the Yukon Area that own dogs, about 17% (287 households) fed whole salmon to their dogs in 2012 (Jallen et al. In prep:15). Most households that own dogs feed fish scraps, but do not harvest salmon to feed to dogs. Of the 6,171 dogs owned by Yukon Area households in 2012, upper Yukon households in districts 4, 5, and 6 owned 4,381 dogs (71% of the total number of dogs owned in Yukon River districts) (Figure 4-4). In 2012, the Division of Commercial Fisheries collected information on the number of each of the 5 species of salmon that fishers retained for dog food from subsistence harvests in surveyed communities. In permit communities, only the total number of whole salmon, and not the numbers of each species, was documented. In the Coastal District and in districts 1 through 5, an estimated 28,054 summer chum salmon, 37,302 fall chum salmon, and 2,572 coho salmon were retained for dog food from subsistence salmon harvests. Additionally, permit holders fed 30,970 whole salmon to dogs, including in District 6, which includes Manley Hot Springs, Minto, Fairbanks, Healy and other Upper Tanana villages (Jallen et al. In prep:34).

Primary gear types used by Yukon Area fishing households in 2012 included set gillnet (50%), drift gillnet (45%), and fish wheel (5%) (Figure 4-5), largely the same as 2009, 2010, and 2011.

Since 1992, ADF&G has asked surveyed households whether they were able to meet their subsistence salmon needs for each survey year. The disastrous fishing year in 2000 resulted in restrictions and closures in subsistence salmon fishing schedules and made it extremely difficult for fishing families to meet their needs (64% of surveyed households reported not meeting their needs in 2000) (Borba and Hamner 2001:98). In 2003, ADF&G began asking households to describe whether they met their subsistence needs for each species of salmon, measuring responses by community and by species. Specifically, surveyed households were asked whether 100%, 75%, 50%, or less than 25% of their harvest needs were met for each species. Two checkboxes, “0%” and “no need,” were added to the 2005 survey in order to distinguish those who had a need but no success in harvesting a species from those who had no need and therefore did not harvest any fish. Not all surveyed households supplied information about whether or not they met their needs for each species. The number of households responding to the needs met question was greatest for Chinook salmon, and lowest for coho salmon (Chinook salmon, 723 households; summer chum salmon, 494 households; fall chum salmon, 275 households; coho salmon, 114 households). According to the data, there was a marked change in responses to the needs met question from 2011 and previous years. Only 21% of all households reported meeting greater than 75% of their needs for Chinook salmon (Jallen et al. In prep). This represents a continued decrease since 2005 in the percentage of households reporting that they met the majority of their needs for Chinook salmon. In 2012, more than half (52%) of surveyed households reported meeting greater than 75% of their needs for summer chum salmon; only 32% reported meeting greater than 75% of their needs for fall chum salmon; and only 30% of surveyed households reported meeting greater than 75% of their needs for coho salmon. Seventy-one percent of households reported meeting less than one-half (<50%) of their needs for Chinook salmon; 40%, 64%, and 69% of households reporting meeting less than one-half their needs for summer chum salmon, fall chum salmon, and coho salmon, respectively.

In 1993, the BOF made a positive C&T use finding for all salmon in the Yukon–Northern Area. The ANS determination was established at 348,000–503,000 salmon for all species combined (5 AAC 01.236). Under these guidelines, 1992 marked the last year when total subsistence salmon harvests fell within the combined ANS range. Since 1990, the overall total subsistence salmon harvest in the Yukon Area has declined by approximately 40% (Table 4-6). In 2001, the BOF made species-specific ANS determinations for each of 4 species of salmon harvested in the Yukon Area, including separate ANS determinations for summer chum salmon and fall chum salmon. The ANS range provides one index of the extent to which reasonable opportunity is provided in each subsistence fishery. Harvests below the lower bound of the ANS range may indicate, with other evidence, that there was not a reasonable opportunity for subsistence harvests during the previous season. Harvests consistently below the lower bound of the ANS are an indication to the BOF to consider whether additional management actions are necessary to provide reasonable subsistence opportunities or if harvest and use patterns for a species have changed over time such that harvests fall outside of ANS ranges. With the exception of Chinook salmon, harvests of all salmon species in 2012 fell within their respective ANS ranges (Table 4-7). This was the fifth consecutive year that Chinook salmon harvests remained below the minimum bound of the ANS range; however, 2012 was the first year since 2007 that fall chum and coho salmon harvests fell within their respective ANS ranges (Table 4-7). See Table 4-7 for a comparison of ANS ranges and subsistence salmon harvests from 1998–2012.

NONSALMON FISH HARVESTS

Although salmon harvests dominate most of the regulatory actions in the Yukon Area, nonsalmon fish harvests remain significant components of the seasonal subsistence round for Yukon Area fishers. Salmon are only available seasonally, but most nonsalmon species are available year-round. Nonsalmon fishes not only provide an important source of nutrition for residents of the Yukon Area, they also represent a significant cultural resource for subsistence fishers in the region. In 1987, and again in 1993, the BOF made a positive C&T use determination for freshwater fish species in the Yukon Area, including sheefish, whitefish species, Arctic lamprey, burbot, longnose sucker, Arctic grayling, northern pike, and Arctic char (5 AAC 01.236). Subsistence fishing for nonsalmon species is generally open by regulation 7 days per week, 24 hours per day, year-round. These state regulations also apply to subsistence fisheries in waters adjacent to federal lands in the project study area (unless superseded on federal public lands by federal subsistence regulations, applicable only to federally qualified subsistence users). Under ANILCA, rural Alaskan residents of the Yukon–Northern Area (except those living in ADF&G Game Management Unit 26B) and residents of the Yukon River drainage have a C&T use determination for nonsalmon fishes, and are qualified to participate in subsistence activities on federal public lands, even if other uses and/or users have been prohibited from subsistence fishing in federal waters due to conservation concerns or user conflicts.¹⁸

ADF&G Division of Commercial Fisheries collects nonsalmon harvest data on an annual basis as part of a postseason salmon survey; however, collection of nonsalmon harvest data is not the primary purpose of the postseason subsistence salmon harvest survey. Furthermore, the implementation of this survey immediately following the salmon fishing season may not be timed to produce the most accurate results for nonsalmon harvests, nor is the stratified sample of salmon fishing households necessarily the best design for collecting nonsalmon harvest information. However, while other single-year harvest data collection efforts suggest that the postseason survey may significantly underestimate harvests (Andersen et al. 2004; Brown et al. 2005), these data have value as the only annual estimate of nonsalmon fish harvests in the Yukon Area (Table 4-8).

Table 4-8 estimates harvests of whitefish, sheefish, and northern pike by community. In 2012 Yukon area fishers from districts 1–5 harvested a total of 106,030 of these nonsalmon fish. This represents an increase

18. USFWS. 2008. Subsistence management regulations for the harvest of fish and shellfish on federal public lands and waters in Alaska, effective April 1, 2008–March 31, 2009. U.S. Fish and Wildlife Service, Anchorage, Alaska.

since 2010 and 2011 when the total harvest of these species was 76,967 and 74,571 fish, respectively (Jallen et al. 2012:111). The “large whitefish” category includes broad and humpback whitefishes while the “small whitefish” category includes least and Bering cisco species and round whitefish. Fishers in District 1 harvested the most number of whitefishes (26,144), followed by District 4 (16,857). On a drainagewide level, large whitefish species were harvested in greater numbers than any other nonsalmon fish. Approximately 41,549 large whitefishes were harvested by Yukon River fishers from districts 1–5—a 54% increase in harvest over 2011; fishers in District 4 harvested the most number of large whitefishes (13,508), while fishers in District 1 harvested the largest number of small whitefishes (18,570). It is important to note that these totals do not include large whitefish harvests from District 6 along the Tanana River; data there are not reported by large and small categories. Fishers from districts 1–5 reported harvesting 18,450 northern pike and 17,094 sheefish in 2012. District 4 households harvested the most pike (5,887), followed by District 1 (3,459). District 1 households harvested more sheefish than in any other district (6,961). Permit fishers, primarily along the Tanana River and a few other locations along the Yukon River reported an additional harvest of 3,944 whitefish, 825 northern pike, and 147 sheefish (Jallen et al. In prep).

The Division of Subsistence has conducted numerous subsistence surveys along the Yukon River over time. In the past 4 years for example, comprehensive surveys that included questions on nonsalmon species have been administered in Galena, Nulato, Ruby, Marshall, Mountain Village¹⁹, Anvik, Grayling, Russian Mission (Ikuta et al. 2014), Shageluk, Pilot Station²⁰, Minto and Manley Hot Springs (Brown et al. 2014). Additionally, studies on the traditional ecological knowledge of nonsalmon have been conducted in the middle Yukon River communities of Tanana, Ruby, Galena, Nulato and Kaltag, and the Yukon Flat communities of Beaver, Birch Creek Village, Central, Circle, and Fort Yukon (Brown et al. 2010; Koskey and Mull 2011). A 2005 study explored the contemporary use of nonsalmon in the lower middle Yukon River communities of Grayling, Anvik, Shageluk and Holy Cross (Brown et al. 2005). Information on historical and contemporary harvest and use of nonsalmon in communities along the Yukon River, where data are available, can be accessed through the Community Subsistence Information System (CSIS) on the ADF&G website.

THE ROLE OF SALMON WITHIN ANNUAL SUBSISTENCE HARVESTS

In addition to post-season salmon surveys conducted by ADF&G, Division of Commercial Fisheries staff, Division of Subsistence staff conducted comprehensive subsistence surveys in the communities of Minto and Manley Hot Springs in the Central Tanana Valley. In 2012, fish was among the most widely used category of wild foods used by Minto and Manley Hot Springs households; 91% of Minto households used 20,726 edible pounds of fish and 98% of Manley Hot Springs households used 46,915 edible pounds of fish (Brown et al. 2014). Salmon accounted for 43% of Minto’s total subsistence harvest and 82% of the total fish harvest in 2012. Salmon accounted for 82% of Manley Hot Spring’s total subsistence harvest and 92% of the total fish harvest in 2012. However, a poor king run and conservative management strategies in 2012 resulted in lower harvests of king salmon throughout the drainage, and 59% of Minto households and 41% of Manley Hot Springs households reported using less salmon in 2012 than in previous years. After salmon species, moose was the next most heavily harvested resource in both communities (Brown et al. 2014).

Nonsalmon fish harvests accounted for 9% of Minto’s total subsistence harvest and 18% of the total fish harvest in 2012. Northern pike (1,528 lb, 4% of total harvest) and broad whitefish (684 lb, 2% of total harvest) were the 2 largest components of Minto’s nonsalmon harvest. Nonsalmon fish harvests accounted for 7% of Manley Hot Spring’s total subsistence harvest and 8% of the total fish harvest in

19. Brown, Caroline L. et al. In prep. “Subsistence harvests in 5 Yukon River communities, 2010: an index approach.” Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. NNN, Fairbanks.

20. Ikuta, Hiroko and Marylynne Kostick, editors. In prep. “Subsistence harvests in 6 communities in the Bering Sea, Kuskokwim River drainage, and Yukon River, 2013.” Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. NNN, Fairbanks.

2012. Northern pike (1,018 lb, 2% of total harvest) and humpback whitefish (718 lb, 1% of total harvest) were the 2 largest components of Manley Hot Springs' nonsalmon harvest (Brown et al. 2014).

Table 4-1.—Subsistence fishing schedule by district, Lower Yukon Area, 2012.

	Coastal District ^a				
Date	Southern ^b	Northern ^c	District 1	District 2	District 3
5/31	Start of regulatory schedule, implemented chronologically with the upriver migration of salmon.				
6/1	Open 7.5" mesh	Open	Open	Open	Open
6/2	Open	Open	Close 8am	Open	Open
6/3	Open	Open	Close	Open	Open
6/4	Open	Open	Open 8pm	Open	Open
6/5	Open	Open	Open	Close 8am	Open
6/6	12pm, 6"mesh	Open	Close 8am	Open 8pm	Open
6/7	6"mesh	Open	Open 8pm	Open	Open
6/8	6"mesh	Open	Open	Close 8am	Close 8am
6/9	6"mesh	Open	Close 8am	Closed	Closed
6/10	6"mesh	Open	Close	Open 8pm	Open 8pm
6/11	6"mesh	Open	Open 8pm	Open	Open
6/12	12pm, 7.5" mesh	Open	Open	Close 8am	Close 8am
6/13	Open 7.5" mesh	Open	Close 8am	Open 8pm	Open 8pm
6/14	Open	Open	Open 8pm	Open	Open
6/15	Open	Open	Open	Close 8am	Close 8am
6/16	Open	Open	Close 8am	Closed	Close
6/17	Open	Open	Close	Open 8pm	Open 8pm
6/18	Open	Open	Open 8pm 6"mesh	Open	Open
6/19	Open	Open	Open 6"mesh	Close 8am	Close 8am
6/20	Open	Close 8pm Closed Closed Closed Closed Closed	Close 8am	Open 8pm 6"mesh	Open 8pm
6/21	Open		Closed	Open 6"mesh	Open
6/22	Open		Closed	Close 8am	Close 8am
6/23	Open		Closed	Closed	Close
6/24	Open		Closed	Closed	Open 8pm 6"mesh
6/25	8pm 6"mesh		Closed	Closed	Close 2pm
6/26	Open 6"mesh	Closed	Closed	Closed	Closed
6/27	Open 6"mesh	Closed	Closed	Closed	Closed
6/28	Open 6"mesh	Closed	Closed	Closed	Closed
6/29	Open 6"mesh	8pm 6"mesh	Open 8pm 6"mesh ^{d,e}	Close	Close
6/30	Open 6"mesh	Close 8am	Close 8am	Close	Close
7/1	Open 6"mesh	8pm 6"mesh	Open 8pm 6"mesh ^{d,e}	Close	Close
7/2	Open 6"mesh	Close 8am	Close 8am ^e	Open 8pm 6"mesh ^{d,f}	Close
7/3	Open 6"mesh	Closed	Closed ^e	Close 8am	Close
7/4	Open 6"mesh	Closed	Closed	Open 8pm 6"mesh	Close
7/5	Open 6"mesh	8pm 6"mesh	Open 8pm 6"mesh ^{d,e}	Close 8am	Open 8pm 6"mesh
7/6	Open 6"mesh	Open 6"mesh	Close 2pm ^e	Close	Close 8am
7/7	Open 6"mesh	Open 6"mesh	Close	Close	Close
7/8	Open 6"mesh	Open 6"mesh	Close	Open 8pm 6"mesh ^{d,g}	Open 8pm 6"mesh
7/9	Open 6"mesh	Open 6"mesh	Open 8pm 6"mesh ^{d,e}	Close 2pm	Close 2pm
7/10	Open 6"mesh	Open 6"mesh	Close 2pm ^h	Close	Close
7/11	Open 6"mesh	Open 6"mesh	Close	Open 8pm 6"mesh ^{d,g}	Open 8pm 6"mesh
7/12	Open 6"mesh	Open 6"mesh	Open 8pm 6"mesh	Close 2pm ⁱ	Open 6"mesh
7/13	Open 6"mesh	Open 6"mesh	Close 2pm ^h	Close	Close 8am
7/14	Open 6"mesh	Open 6"mesh	Close	Open 6pm 6"mesh	Close

-continued-

Table 4-1.–Page 2 of 3

Date	Coastal District ^a		District 1	District 2	District 3
	Southern ^b	Northern ^c			
7/15	8pm 7.5"mesh	8pm 7.5"mesh	Open 8pm, Close 10pm	Close 6am ^{i, j}	Open 8pm 6"mesh
7/16	Open	Open	Closed ^{h, j, k}	Open 11am 7.5"mesh	Open 6"mesh
7/17	Open	Open	Open 10am 7.5"mesh	Close 12am ^{i, j}	Close 8am
7/18	Open	Open	Close 10pm ^{h, k, j}	Closed	Open 8pm 7.5" mesh
7/19	Open 7.5"mesh	Open 7.5"mesh	Closed	Open 9am 7.5" mesh	Open 7.5"mesh
7/20	Open	Open	Open 10am 7.5"mesh	Open	Open
7/21	Open	Open	Open	Open	Open
7/22	Open	Open	Open	Close 3am ^{i, j}	Open
7/23	Open	Open	Close 1am ^{h, j, k}	Open 9am 7.5"mesh	Open
7/24	Open	Open	Open 10am 7.5" mesh	Open	Open
7/25	Open	Open	Open	Close 3am ^{i, j}	Open
7/26	Open	Open	Close 1am ^{h, j, k}	Open 9am 7.5" mesh	Open
7/27	Open	Open	Open 10am 7.5" mesh	Open	Open
7/28	Open	Open	Open	Open	Open
7/29	Open	Open	Open	Open	Open
7/30	Open	Open	Close 1am ^{h, j, k}	Close 3am ^{i, j}	Open
7/31	Open	Open	Open 10am 7.5" mesh	Open 9am	Open
8/1	Open	Open	Open	Close 3am ^{i, j}	Open
8/2	Open	Open	Close 1am ^{h, j, k}	Open 9am	Open
8/3	Open	Open	Open 10am	Open	Open
8/4	Open	Open	Open	Close 3am ^{i, j}	Open
8/5	Open	Open	Close 5pm ^{h, j, k}	Open 9am	Open
8/6	Open	Open	Open 10am	Close midnight	Open
8/7	Open	Open	Open	Closed ^{i, j}	Open
8/8	Open	Open	Close 10pm ^{h, j, k}	Open 9am	Open
8/9	Open	Open	Closed	Open	Open
8/10	Open	Open	Open 10am	Open	Open
8/11	Open	Open	Open	Open	Open
8/12	Open	Open	Open	Open	Open
8/13	Open	Open	Open	Open	Open
8/14	Open	Open	Open	Open	Open
8/15	Open	Open	Open	Open	Open
8/16	Open	Open	Open	Open	Open
8/17	Open	Open	Close 10pm ^{h, j, k}	Open	Open
8/18	Open	Open	Closed	Open	Open
8/19	Open	Open	Open 7am, Close 10pm	Close 12am ^{i, j}	Open
8/20	Open	Open	Closed ^{h, j, k}	Open 9am	Open
8/21	Open	Open	Open 10 am	Open	Open
8/22	Open	Open	Close 9pm ^{h, j, k}	Close 3am ^{i, j}	Open
8/23	Open	Open	Closed	Open 9am	Open
8/24	Open	Open	Open 9am	Open	Open
8/25	Open	Open	Open	Close 9pm ^{i, j}	Open
8/26	Open	Open	Close 9pm ^{h, j, k}	Closed	Open
8/27	Open	Open	Closed	Open 6am	Open
8/28	Open	Open	Open 9am	Close 9pm ^{i, j}	Open

-continued-

Table 4-1.–Page 3 of 3

Date	Coastal District ^a		District 1	District 2	District 3
	Southern ^b	Northern ^c			
8/29	Open	Open	Close 9pm ^{h, j, k}	Closed	Open
8/30	Open	Open	Closed	Open 6am	Open
8/31	Open	Open	Open 9am	Close 2am ^{i, j}	Open
9/1	Open	Open	Open	Open 6am	Open
9/2	Open	Open	Open	Open	Open
9/3	Open	Open	Open	Open	Open
9/4	Open	Open	Open	Open	Open
9/5	Open	Open	Open	Open	Open
9/6	Open	Open	Open	Open	Open
9/7	Open	Open	Open	Open	Open
9/8	Open	Open	Open	Open	Open
9/9	Open	Open	Open	Open	Open
9/10	Open	Open	Open	Open	Open

Note Shaded areas indicate fishery closures; outlined shaded days were closed to protect the first and second pulses of Chinook salmon. Dates with dark shading were closed for subsistence fishing for 12 hours before, during and 12 hours after commercial fishing periods. Unless noted, mesh size was restricted to 7.5 inch or less in all districts and subdistricts in 2012. The Innoko River remained open 24 hours a day 7 days a week, but was restricted to 6 inch or smaller mesh from 8pm June 24 to 8pm July 18.

- a. The Coastal District was split for management purposes based on which mouths various salmon species were entering the delta.
- b. The portion of the Coastal District from the Naskonat Peninsula north to 62 degrees North latitude, including communities of Chevak, Hooper Bay, and Scammon Bay.
- c. The portion of the Coastal District from 62 degrees North latitude to Point Romanoff and 3 miles offshore.
- d. Commercial period open during all or part of a subsistence fishing period.
- e. Commercial fishing limited to the South Mouth of District 1 and restricted to 6 inch or smaller mesh. The South Mouth area included waters down river of the lower point of Head of Passes to Chris Point, Black River, Kwiguk Pass, and coastal waters from Chris Point to one mile north of Kwiguk Pass. Middle and North Mouth including Aproka Pass were closed to commercial fishing.
- f. Commercial fishing limited to the portion of District 2 downstream of the Andreafsky River Mouth and restricted to 6 inch or smaller mesh.
- g. Commercial fishing limited to the portion of District 2 downstream of Pilot Station Slough and restricted to 6 inch or smaller mesh.
- h. Commercial fishing open in the entire District 1 area during all or part of a subsistence fishery closure and restricted to 6 inch or smaller mesh.
- i. Commercial open in the entire District 2 area during all or part of a subsistence fishery closure and restricted to 6 inch or smaller mesh.
- j. Commercial fishing during a subsistence closure. Subsistence fishing closed for 12 hours before, during and 12 hours after a commercial period.
- k. Commercial fishing in District Y-1 open only in the Coastal setnet only area of District Y-1 for all or part of a commercial period and restricted to 6 inch or smaller mesh.

Table 4-2.--Subsistence fishing schedule by district, Upper Yukon Area, 2012.

Date	Subdistrict 4-A ^a		Sub 4-B /	5-A/5-B /	Subdistrict 5-D ^b			Koyukuk River	Tanana River		
	Lower	Upper			Lower ^c	Middle ^d	Upper ^e		6A	6B	6C
6/11	Open	Open	Open	Open	Open	Open	Open	Open	Open 6pm	Open 6pm	Open 6pm
6/12	Close 6pm	Close 6pm	Open	Open	Open	Open	Open	Open	Open	Open	Open
6/13	Open 6pm	Open 6pm	Open	Open	Open	Open	Open	Open	Close 12pm	Close 12pm	Close 12pm
6/14	Open	Open	Open	Open	Open	Open	Open	Open	Closed	Closed	Closed
6/15	Close 6pm	Close 6pm	Open	Open	Open	Open	Open	Open	Open 6pm	Open 6pm	Open 6pm
6/16	Close	Close	Open	Open	Open	Open	Open	Open	Open	Open	Open
6/17	Open 6pm	Open 6pm	Open	Open	Open	Open	Open	Open	Close 12pm	Close 12pm	Close 12pm
6/18	Open	Open	Open	Open	Open	Open	Open	Open	Open 6pm	Open 6pm	Open 6pm
6/19	Close 6pm	Close 6pm	Close 6pm	Open	Open	Open	Open	Open	Open	Open	Open
6/20	Open 6pm	Open 6pm	Open 6pm	Open	Open	Open	Open	Open	Close 12pm	Close 12pm	Close 12pm
6/21	Open	Open	Open	Open	Open	Open	Open	Open	Closed	Closed	Closed
6/22	Close 6pm	Close 6pm	Close 6pm	Open	Open	Open	Open	Open	Open 6pm	Open 6pm	Open 6pm
6/23	Closed	Closed	Closed	Open	Open	Open	Open	Open	Open	Open	Open
6/24	Open 6pm	Open 6pm	Open 6pm	Close 6pm	Open	Open	Open	Open	Close 12pm	Close 12pm	Close 12pm
6/25	Open	Open	Open	Close	Open	Open	Open	Open	Open 6pm	Open 6pm	Open 6pm
6/26	Close 6pm	Close 6pm	Close 6pm	Open 6pm	Open	Open	Open	Open	Open	Open	Open
6/27	Open 6pm 6" mesh	Open 6pm	Open 6pm	Open	Open	Open	Open	Open	Close 12pm	Close 12pm	Close 12pm
6/28	Closed 6pm	Open	Open	Close 6pm	Open	Open	Open	Open	Closed	Closed	Closed
6/29	Closed	Close 6pm	Close 6pm	Open 6pm	Open	Open	Open	Open	Open 6pm	Open 6pm	Open 6pm
6/30	Closed	Close	Close	Open	Open	Open	Open	Open	Open	Open	Open
7/1	Close ^f	Open 6pm gear ^{f, g}	Open 6pm	Close 6pm	Open	Open	Open	Open	Close 12pm	Close 12pm	Close 12pm
7/2	Close ^f	Close 6pm ^f	Open	Close	Open	Open	Open	Open	Open 6pm	Open 6pm	Open 6pm

-continued-

Table 4-2.—Page 2 of 6

Date	Subdistrict 4-A ^a		Sub 4-B /	5-A/5-B /	Subdistrict 5-D ^b			Koyukuk	Tanana River		
	Lower	Upper	4-C	5-C	Lower ^c	Middle ^d	Upper ^e	River	6A	6B	6C
7/3	Closed ^f	Closed ^f	Closed 6pm	Open 6pm	Open	Open	Open	6pm, 6" mesh	Open	Open	Open
7/4	Closed ^f	Closed ^f	Closed	Open	Open	Open	Open	Open 6" mesh	Close 12pm	Close 12pm	Close 12pm
7/5	Closed ^f	Closed ^f	Closed	Close 6pm	Open	Open	Open	Open 6" mesh	Closed	Closed	Closed
7/6	Closed ^f	Closed ^f	Closed	Closed	Open	Open	Open	Open 6" mesh	Open 6pm	Open 6pm	Open 6pm
7/7	Closed ^f	Closed ^f	Closed	Closed	Open	Open	Open	Open 6" mesh	Open	Open	Open
7/8	Closed ^f	Closed ^f	Closed	Closed	Open	Open	Open	Open 6" mesh	Close 12pm	Close 12pm	Close 12pm
7/9	Open 6pm - gear ^{f,g}	Closed ^f	Closed	Closed	Open	Open	Open	Open 6" mesh	Open 6pm	Open 6pm	Open 6pm
7/10	Close 6pm ^f	Closed ^f	Closed	Closed	Open	Open	Open	Open 6" mesh	Open	Open	Open
7/11	Closed ^f	Closed ^f	Closed	Closed	Open	Open	Open	Open 6" mesh	Close 12pm	Close 12pm	Close 12pm
7/12	Open 6pm - gear ^{f,g}	Open 6pm gear ^{f,g}	Closed	Closed	Close 7pm	Open	Open	Open 6" mesh	Closed	Closed	Closed
7/13	Close 6pm ^h	Close 6pm ^h	Closed	Closed	Closed	Open	Open	Open 6" mesh	Open 6pm	Open 6pm	Open 6pm
7/14	Closed ^h	Closed ^h	Closed	Closed	Closed	Open	Open	Open 6" mesh	Open	Open	Open
7/15	Open 6pm - gear ^{g,h}	Closed ^h	Open 6pm	Closed	Closed	Close 7pm	Open	Open 6" mesh	Close 12pm	Close 12pm	Close 12pm
7/16	Open - gear ^{g,h}	Open 6pm gear ^{g,h}	Close 12pm	Closed	Closed	Closed	Open	Open 6" mesh	Open 6pm	Open 6pm	Open 6pm
7/17	Close 6pm ^h	Close 6pm ^h	Closed	Closed	Closed	Closed	Close 7pm	Open 6" mesh	Open	Open	Open
7/18	Open 6pm ^{h,i}	Open 6pm gear ^{g,h}	Open 6pm	Open 6pm	Closed	Closed	Closed	Open 6" mesh	Close 12pm	Close 12pm	Close 12pm
7/19	Open ^h	Open ^{g,h}	Close 12pm	Close 12pm	Closed	Closed	Closed	Open 6" mesh	Closed	Closed	Closed
7/20	Close 6pm ^h	Close 6pm ^h	Closed	Closed	Closed	Closed	Closed	Open 6" mesh	Open 6pm ^{j,k}	Open 6pm ^{j,k,l}	Closed ^k
7/21	Closed ^h	Closed ^h	Closed	Closed	Closed	Closed	Closed	Open 6" mesh	Open ^{j,k}	Open ^{j,k,l}	Closed ^k
7/22	Open 6pm ^h	Open 6pm ^{h,i}	Open 6pm	Closed	Open 8am	Closed	Closed	7.5" mesh, 6pm	Close 12pm ^{k,l}	Close 12pm ^{j,k,l}	Closed ^k
7/23	Open ^h	Open ^h	Close 12pm	Closed	Close 8pm	Closed	Closed	Open	Open 6pm ^{j,k}	Open 6pm ^{j,k,l}	Closed ^k

-continued-

Table 4-2.—Page 3 of 6

Date	Subdistrict 4-A ^a		Sub 4-B /	5-A/5-B /	Subdistrict 5-D ^b			Koyukuk	Tanana River		
	Lower	Upper	4-C	5-C	Lower ^c	Middle ^d	Upper ^e	River	6A	6B	6C
7/24	Close 6pm ^h	Close 6pm ^h	Close 12pm	Closed	Closed	Closed	Closed	Open	Open ^{j, k}	Open ^{j, k, l}	Closed ^k
7/25	Open 6pm ^h	Open 6pm ^h	Open 6pm	Closed	Closed	Closed	Closed	Open	Close 12 pm ^k	Close 12pm ^{k, l}	Closed ^k
7/26	Open ^h	Open ^h	Open	Closed	Closed	Open 8am	Closed	Open	Closed ^k	Closed ^{k, l}	Closed ^k
7/27	Open ^h	Open ^h	Close 6pm	Open 6pm	Closed	Close 8pm	Closed	Open	Open 6pm ^{j, k, m}	Open 6pm ^{j, k, l}	Closed ^k
7/28	Open ^h	Open ^h	Closed	Open	Closed	Closed	Open 8am	Open	Open, gear ^k	Open, gear ^{j, k}	Closed ^k
7/29	Close 6pm ^h	Close 6pm ^h	Open 6pm	Close 6pm	Closed	Closed	Close 8pm	Open	Close 12pm ^k	Close 12pm ^k	Closed ^k
7/30	Closed ^h	Closed ^h	Open	Closed	Open 7pm	Closed	Closed	Open	Open 6pm ^k	Open 6pm ^k	Open 6pm ^k
7/31	Open 6pm	Open 6pm	Open	Open 6pm	Open	Closed	Closed	Open	Open ^k	Open ^k	Open ^k
8/1	Open	Open	Open	Open	Open	Closed	Closed	Open	Close 12pm ^k	Close 12pm ^k	Close 12pm ^k
8/2	Open	Open	Open	Open	Open	Closed	Closed	Open	Closed	Closed	Closed
8/3	Open	Open ⁿ	Close 6pm	Open	Open	Closed	Closed	Open	Open 6pm ^o	Open 6pm ^o	Open 6pm ^o
8/4	Open	Open	Closed	Open	Open	Closed	Closed	Open	Open ^o	Open ^o	Open ^o
8/5	Close 6pm	Close 6pm	Open 6pm	Close 6pm	Open	Closed	Closed	Open	Close 12pm	Close 12pm	Close 12pm
8/6	Closed	Closed	Open	Closed	Open	Closed	Closed	Open	Open 6pm ^o	Open 6pm ^o	Open 6pm ^o
8/7	Open 6pm	Open 6pm	Open	Open 6pm	Open	Closed	Closed	Open	Open ^o	Open ^o	Open ^o
8/8	Open	Open	Open	Open	Open	Closed	Closed	Open	Close 12pm	Close 12pm	Close 12pm
8/9	Open ^o	Open ^o	Open	Open	Open	Closed	Closed	Open	Closed	Closed	Closed
8/10	Open ^o	Open ^o	Close 6pm	Open	Open	Closed	Closed	Open	Open 6pm ^o	Open 6pm ^o	Open 6pm ^o
8/11	Open ^o	Open ^o	Closed	Open	Open	Closed	Closed	Open	Open ^o	Open ^o	Open ^o
8/12	Close 6pm	Close 6pm	Open 6pm	Close 6pm	Open	Closed	Closed	Open	Close 12pm	Close 12pm	Close 12pm
8/13	Closed	Closed	Open	Closed	Open	Closed	Closed	Open	Open 6pm ^o	Open 6pm ^o	Open 6pm ^o
8/14	Open 6pm ^o	Open 6pm ^o	Open	Open 6pm ^p	Open	Closed	Closed	Open	Open ^o	Open ^o	Open ^o
8/15	Open ^o	Open ^o	Open	Open ^p	Open	Open 6pm	Open 6pm	Open	Close 12pm	Close 12pm	Close 12pm
8/16	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Closed	Closed	Closed
8/17	Open ^o	Open ^o	Close 6pm	Open ^p	Open	Open	Open	Open	Open 6pm	Open 6pm	Open 6pm

-continued-

Table 4-2.—Page 4 of 6

Date	Subdistrict 4-A ^a		Sub 4-B / 4-C	5-A/5-B / 5-C	Subdistrict 5-D ^b			Koyukuk River	Tanana River		
	Lower	Upper			Lower ^c	Middle ^d	Upper ^e		6A	6B	6C
8/18	Open ^o	Open ^o	Closed	Open ^p	Open	Open	Open	Open	Open	Open	Open
8/19	Close 6pm	Close 6pm	Open 6pm	Close 6pm	Open	Open	Open	Open	Close 12pm	Close 12pm	Close 12pm
8/20	Closed	Closed	Open	Closed	Open	Open	Open	Open	Open 6pm	Open 6pm	Open 6pm
8/21	Open 6pm ^o	Open 6pm ^o	Open	Open 6pm ^p	Open	Open	Open	Open	Open	Open	Open
8/22	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Close 12pm	Close 12pm	Close 12pm
8/23	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Closed	Closed	Closed
8/24	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Open 6pm	Open 6pm	Open 6pm
8/25	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Open	Open	Open
8/26	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Close 12pm	Close 12pm	Close 12pm
8/27	Open ^o	Open ^o	Open	Open	Open	Open	Open	Open	Open 6pm	Open 6pm	Open 6pm
8/28	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Open	Open	Open
8/29	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Close 12pm	Close 12pm	Close 12pm
8/30	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Closed	Closed	Closed
8/31	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Open 6pm ^q	Open 6pm ^q	Open 6pm ^q
9/1	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Open ^q	Open ^q	Open ^q
9/2	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Close 12pm	Close 12pm	Close 12pm
9/3	Open ^o	Open ^o	Open	Open	Open	Open	Open	Open	Open 6pm ^q	Open 6pm ^q	Open 6pm ^q
9/4	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Open ^q	Open ^q	Open ^q
9/5	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Close 12pm	Close 12pm	Close 12pm
9/6	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Closed	Closed	Closed
9/7	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Open 6pm ^q	Open 6pm ^q	Open 6pm ^q
9/8	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Open ^q	Open ^q	Open ^q
9/9	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Close 12pm	Close 12pm	Close 12pm
9/10	Open	Open	Open	Open	Open	Open	Open	Open	Open 6pm ^q	Open 6pm ^q	Open 6pm ^q
9/11	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Open ^q	Open ^q	Open ^q
9/12	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Close 12pm	Close 12pm	Close 12pm
9/13	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Closed	Closed	Closed
9/14	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Open 6pm ^q	Open 6pm ^q	Open 6pm ^q
9/15	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Open ^q	Open ^q	Open ^q
9/16	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Close 12pm	Close 12pm	Close 12pm
9/17	Open	Open	Open	Open	Open	Open	Open	Open	Open 6pm ^q	Open 6pm ^q	Open 6pm ^q

-continued-

Table 4-2.—Page 5 of 6

Date	Subdistrict 4-A ^a		Sub 4-B /	5-A/5-B /	Subdistrict 5-D ^b			Koyukuk River	Tanana River		
	Lower	Upper			Lower ^c	Middle ^d	Upper ^e		6A	6B	6C
9/18	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Open ^q	Open ^q	Open ^q
9/19	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Close 12pm	Close 12pm	Close 12pm
9/20	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Closed	Closed	Closed
9/21	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Open 6pm ^q	Open 6pm ^q	Open 6pm ^q
9/22	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Open ^q	Open ^q	Open ^q
9/23	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Close 12pm	Close 12pm	Close 12pm
9/24	Open	Open	Open	Open	Open	Open	Open	Open	Open 6pm ^q	Open 6pm ^q	Open 6pm ^q
9/25	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Open ^q	Open ^q	Open ^q
9/26	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Close 12pm ^r	Close 12pm ^r	Close 12pm ^r
9/27	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Closed ^r	Closed ^r	Closed ^r
9/28	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Open 6pm ^q	Open 6pm ^q	Open 6pm ^q
9/29	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Open ^q	Open ^q	Open ^q
9/30	Open ^o	Open ^o	Open	Open ^p	Open	Open	Open	Open	Close 12pm	Close 12pm	Close 12pm

Note shaded areas indicate windowed fishery closures, outlined shaded days were closed to protect the first and second pulses of Chinook salmon. Unless noted, mesh size was restricted to 7.5 inch or less in all districts and subdistricts in 2012.

- a. Subdistrict 4-A was divided into two separate areas above and below Stink Creek to protect the first pulse of Chinook salmon as it passed through this long section of river.
- b. Subdistrict 5-D was divided into three separate areas to protect the first pulse of Chinook salmon as it passed through this long section of river.
- c. Subdistrict 5-D Lower: from the ADF&G marker 2 miles downstream of Waldron Creek upstream to the Hadweenzic River.
- d. Subdistrict 5-D Middle: from the Hadweenzic River upstream to 22 Mile Slough.
- e. Subdistrict 5-D Upper: from 22 Mile Slough to the US/Canada border.
- f. Commercial fishing period in Subdistrict 4A with fish wheels only. Fish wheels were required to be manned at all times and any Chinook salmon caught were to be immediately released alive. From July 1 to July 12, twelve commercial periods occurred that were each 12 hours in length.
- g. Gear restrictions to conserve Chinook salmon in place for all or part of a subsistence opening. Gillnets restricted to mesh sizes of 6 inch or less. Fishwheels must be closely attended, equipped with a chute, and all Chinook salmon released to the water alive.
- h. Commercial fishing period in Subdistrict 4A with fish wheels only. Fish wheels were required to be manned at all times and any Chinook salmon caught were to be immediately released alive. Starting at 8pm Friday 7/13, commercial fishing was open continuously and extended until 8pm Monday July 30.
- i. District returned to normal regulatory subsistence schedule with no additional gear restrictions.
- j. Gillnets not allowed. Subsistence fishing restricted to fish wheels equipped with a chute that must be closely attended and all Chinook salmon caught must be returned to the water alive.
- k. Commercial fishing restricted to continuously manned fish wheels and all Chinook salmon must be immediately returned to the water alive.

-continued-

Table 4-2.—Page 6 of 6

- l. The Old Minto Area of 6B remained on the regulatory schedule of one five day a week subsistence fishing period from 6pm Friday until 6pm Wednesday. Gear was restricted to only fishwheels with a chute and no retention of Chinook salmon from 6pm Friday 7/21 to 6pm Friday 7/27. For the remainder of the season fishermen could use gillnet with 7.5" or smaller mesh and fishwheels in the Old Minto Area.
- m. Gear restrictions discontinued at 6pm.
- n. Effective 12:01 am, drift gillnets can be used for subsistence fishing in the upper portion of Subdistrict 4A from Stink Creek to Cone Point.
- o. Commercial fishing opening concurrent with subsistence periods and restricted to fish wheels or gillnets with six-inch or smaller mesh.
- p. Commercial fishing periods open for 120 hours concurrent with subsistence periods in Subdistricts 5B and 5C and restricted to fish wheels or gillnets with six-inch or smaller mesh.
- q. Commercial fishing opening concurrent with subsistence periods. No additional gear restrictions.
- r. Commercial fishing period extension through a subsistence closure. No additional gear restrictions.

Table 4-3.–Household subsistence and personal use permits, listed by fishery and community of residence, Yukon Area, 2012.

Community	Permits		Percent returned	Number of permits returned that fished
	Issued	Returned		
Subsistence permits				
Central	4	4	100%	3
Circle	19	19	100%	7
Eagle	33	31	94%	17
Rampart	5	5	100%	5
Fairbanks (FNSB) ^a	157	154	98%	78
Healy	5	5	100%	4
Manley	17	14	82%	10
Minto	37	33	89%	10
Nenana	45	41	91%	26
Stevens Village	3	3	100%	1
Upper Tanana villages ^b	53	44	83%	22
Other subsistence ^c	28	27	96%	14
Subsistence permit subtotal	406	380	94%	197
Personal use permits				
Fairbanks (FNSB) ^a	66	65	98%	29
Other personal use ^d	6	5	83%	3
Personal use permit subtotal	72	70	97%	32
Total	478	450	94%	229

Source Jallen et al. (In prep).

- a. Fairbanks North Star Borough (FNSB) residents from the communities of Ester, Fairbanks, North Pole, Salcha, and Two Rivers.
- b. Upper Tanana River (UTV) residents from the communities of Delta Junction, Dot Lake, Northway, Tanacross, and Tok.
- c. Includes residents from Anchorage, Anderson, Copper Center, Denali Park, Eagle River, Palmer, Tanana, Wasilla, and Wiseman who were issued a subsistence fishing permit for the Yukon, Tanana, Tolovana, Kantishna, and Upper Koyukuk rivers.
- d. Includes residents of Nenana and Delta Junction that applied for a personal use permit.

Table 4-4.—Estimated number of subsistence fishing households in surveyed communities, with community and district totals, Yukon Area, 2012.

Community	Households		Estimated number of fishing households
	Total	Surveyed	
Hooper Bay	209	79	114
Scammon Bay	92	37	73
Coastal district subtotal	301	116	187
Alakanuk	133	54	85
Emmonak	166	92	88
Kotlik	104	40	99
Nunam Iqua	36	28	25
District 1 subtotal	439	214	297
Marshall	70	28	48
Mountain Village	163	66	103
Pilot Station	109	55	47
Pitkas Point	28	17	14
Saint Marys	121	48	89
District 2 subtotal	491	214	301
Holy Cross	56	29	32
Russian Mission	66	25	53
Shageluk	31	18	11
District 3 subtotal	153	72	96
Alatna	9	7	5
Allakaket	65	25	14
Anvik	28	24	25
Bettles	24	14	1
Galena	169	66	76
Grayling	48	18	47
Hughes	35	25	5
Huslia	82	28	41
Kaltag	61	17	35
Koyukuk	51	16	26
Nulato	76	24	50
Ruby	57	20	47
District 4 subtotal	705	284	372
Beaver	31	25	17
Birch Creek	16	7	0
Chalkyitsik	21	12	3
Fort Yukon	219	76	50
Stevens Village	15	11	8
Tanana	99	37	44
Venetie	78	28	14
District 5 subtotal	479	196	136
Total	2,568	1,096	1,389

Source Jallen et al. (In prep).

Table 4-5.—Estimated subsistence salmon harvests by community, Yukon Area, 2012.

Community	Households or permits		Estimated salmon harvest ^a					
	Total	Surveyed or returned	Chinook	Coho	Summer chum	Fall chum	Pink	Total
Hooper Bay	218	79	1,090	7	15,799	1	1,101	17,998
Scammon Bay	99	44	1,014	86	7,442	10	1,343	9,895
Coastal district subtotal	317	123	2,104	93	23,241	11	2,444	27,893
Alakanuk	158	54	1,081	252	9,012	449	174	10,968
Emmonak	180	92	1,864	2,660	15,829	5,890	199	26,442
Kotlik	110	37	1,173	420	8,552	1,073	195	11,413
Nunam Iqua (Sheldon Point)	42	34	195	18	1,977	210	1,051	3,451
District 1 subtotal	490	217	4,313	3,350	35,370	7,622	1,619	52,274
Marshall	69	26	1,409	567	5,903	184	5	8,068
Mountain Village	152	57	1,789	256	9,031	685	207	11,968
Pilot Station	118	57	1,078	329	5,716	1,031	23	8,177
Pitkas Point	27	23	261	53	1,153	9	2	1,478
Saint Marys	127	49	2,344	141	10,763	1,423	643	15,314
District 2 subtotal	493	212	6,881	1,346	32,566	3,332	880	45,005
Holy Cross	55	31	576	237	1,147	339	0	2,299
Russian Mission	72	26	1,711	319	2,508	282	76	4,896
Shageluk	29	11	75	0	5,035	16	24	5,150
District 3 subtotal	156	68	2,362	556	8,690	637	100	12,345
Alatna	10	4	0	0	100	18	0	118
Allakaket	63	21	5	38	3,850	508	0	4,401
Anvik	35	27	435	214	1,371	569	0	2,589
Bettles	22	17	3	0	7	0	0	10
Galena	169	51	742	276	718	2,947	3	4,686
Grayling	47	18	1,081	26	2,616	804	0	4,527
Hughes	31	25	0	0	428	2	0	430
Huslia	95	33	165	165	7,306	1,909	101	9,646
Kaltag	58	15	1,346	928	186	2,830	0	5,290
Koyukuk	49	22	614	62	828	1,331	0	2,835
Nulato	72	23	1,955	41	254	2,729	0	4,979
Ruby	66	21	1,316	1,806	3,891	4,408	0	11,421
District 4 subtotal	717	277	7,662	3,556	21,555	18,055	104	50,932
Beaver	31	24	71	2	27	174	0	274
Birch Creek	16	12	0	0	0	0	0	0
Central	4	4	66	0	0	0	0	66
Chalkyitsik	28	18	0	0	0	162	0	162
Circle	19	19	280	5	0	161	0	446
Eagle	33	31	167	0	0	18,731	0	18,898
Fairbanks	223	219	687	1,602	607	5,073	0	7,969
Fort Yukon	211	87	2,141	4	0	12,659	0	14,804
Rampart	5	5	190	0	71	190	0	451
Stevens Village	21	14	330	0	188	277	0	795
Tanana	103	52	2,100	3,060	4,333	20,465	3	29,961
Venetie	75	24	86	0	0	295	0	381
District 5 subtotal	769	509	6,118	4,673	5,226	58,187	3	74,207
Healy	5	5	0	760	0	595	0	1,355
Manley	17	14	174	1,374	58	2,164	0	3,770
Minto	37	33	99	0	64	2	0	165
Nenana	45	41	296	5,904	370	8,671	0	15,241

-continued-

Table 4-5.–Page 2 of 2

Community	Households or permits		Estimated salmon harvest ^a					Total
	Total	Surveyed or returned	Chinook	Coho	Summer chum	Fall chum	Pink	
District 6 subtotal	104	93	569	8,038	492	11,432	0	20,531
Other communities	87	76	477	21	173	443	0	1,114
Total	3,133	1,575	30,486	21,633	127,313	99,719	5,150	284,301

Source Jallen et al. (In prep).

- a. Includes subsistence harvests, personal use harvests, commercial harvests retained for home use, and fish distributed from ADF&G test fisheries.

Table 4-6.–Historical subsistence salmon harvests, Yukon Area, 1976–2012.

Year	Households or permits ^a		Estimated salmon harvest ^a					Total
	Total	Surveyed or returned	Chinook	Coho	Summer chum	Fall chum	Pink	
1976			17,530	12,737		1,375		31,642
1977			16,007	16,333		4,099		36,439
1978			30,785	7,965	213,953	95,532		348,235
1979			31,005	9,794	202,772	233,347		476,918
1980			42,724	20,158	274,883	172,657		510,422
1981			29,690	21,228	210,785	188,525		450,228
1982			28,158	35,894	260,969	132,897		457,918
1983			49,478	23,905	240,386	192,928		506,697
1984			42,428	49,020	230,747	174,823		497,018
1985			39,771	32,264	264,828	206,472		543,335
1986			45,238	34,468	290,825	164,043		534,574
1987			55,039	46,213	300,042	226,990		628,284
1988	2,700	1,865	45,495	69,679	229,838	157,075		502,087
1989	2,211	983	48,462	40,924	169,496	211,303		470,185
1990	2,666	1,121	48,587	43,460	115,609	167,900		375,556
1991	2,521	1,261	46,773	37,388	118,540	145,524		348,225
1992	2,751	1,281	47,077	51,980	142,192	107,808		349,057
1993	3,028	1,397	63,915	15,812	125,574	76,882		282,183
1994	2,922	1,386	53,902	41,775	124,807	123,565		344,049
1995	2,832	1,391	50,620	28,377	136,083	130,860		345,940
1996	2,869	1,293	45,671	30,404	124,738	129,258		330,071
1997	2,825	1,309	57,117	23,945	112,820	95,141		289,023
1998	2,986	1,337	54,124	18,121	87,366	62,901		222,512
1999	2,888	1,377	50,515	19,984	79,250	83,420		233,169
2000	3,209	1,341	36,844	16,650	77,813	19,402	1,591	152,300
2001	3,072	1,355	56,103	23,236	72,392	36,164	403	188,298
2002	2,775	1,254	44,384	16,551	87,599	20,140	8,425	177,100
2003	2,850	1,377	56,872	24,866	83,802	58,030	2,167	225,737
2004	2,721	1,228	57,549	25,286	79,411	64,562	9,697	236,506
2005	2,662	1,406	53,547	27,357	93,411	91,667	3,132	269,114
2006	2,833	1,473	48,682	19,985	115,355	84,320	4,854	273,196
2007	2,819	1,495	55,292	22,013	93,075	99,120	2,118	271,618

-continued-

Table 4-6.–Page 2 of 2

Year	Households or permits ^a		Estimated salmon harvest ^a					
	Total	Surveyed or returned	Chinook	Coho	Summer chum	Fall chum	Pink	Total
2008	3,030	1,664	45,312	16,905	86,652	89,538	9,529	247,936
2009	2,853	1,508	33,932	16,076	80,847	66,197	2,300	199,352
2010	3,066	1,659	44,721	14,107	88,692	71,854	4,199	223,573
2011	3,060	1,574	41,069	12,576	96,459	80,549	2,291	232,944
2012	3,046	1,546	30,486	21,633	127,313	99,719	5,150	284,301
5-year average (2007-2011)	2,966	1,580	44,065	16,335	89,145	81,452	4,087	235,085
10-year average (2002-2011)	2,867	1,464	48,136	19,572	90,530	72,598	4,871	235,708
Historical average (1976-2011)	2,840	1,389	44,845	26,873	150,353	112,969	4,226	328,096

Source Jallen et al. (In prep).

Note cells that do not contain data have no data available.

a. Estimates prior to 1988 are based on fish camp surveys and sampling information is unavailable.

Table 4-7.–Comparison of amounts necessary for subsistence (ANS) and estimated subsistence salmon harvests, Yukon Area, 1998–2012.

ANS range	Chinook	Coho	Summer chum	Fall chum
	45,500–66,704	20,500–51,980	83,500–142,192	89,500–167,900
Year	Estimated number of subsistence salmon harvested ^a			
1998 ^b	52,910	<u>16,606</u>	<u>81,858</u>	<u>59,603</u>
1999 ^b	50,711	<u>20,122</u>	<u>79,348</u>	<u>84,203</u>
2000 ^b	<u>33,896</u>	<u>11,853</u>	<u>72,807</u>	<u>15,152</u>
2001	53,462	21,977	<u>68,544</u>	<u>32,135</u>
2002	<u>42,117</u>	<u>15,619</u>	<u>79,066</u>	<u>17,908</u>
2003	55,221	22,838	<u>78,664</u>	<u>53,829</u>
2004	55,102	24,190	<u>74,532</u>	<u>61,895</u>
2005	53,409	27,250	93,259	91,534
2006	48,593	<u>19,706</u>	115,093	<u>83,987</u>
2007	55,156	21,878	92,891	98,947
2008	<u>45,186</u>	<u>16,855</u>	86,514	<u>89,357</u>
2009	<u>33,805</u>	<u>16,006</u>	<u>80,539</u>	<u>66,119</u>
2010	<u>44,559</u>	<u>13,045</u>	88,373	<u>68,645</u>
2011	<u>40,980</u>	<u>12,344</u>	96,020	<u>80,202</u>
2012	<u>30,415</u>	21,533	126,992	99,309

Source Jallen et al. (In prep).

a. Estimates for 1998–2004 do not include personal use harvests, ADF&G test fishery distributions, or salmon removed from commercial harvests. Estimates for 2005–2012 include test fishery distributions because the amounts necessary for subsistence (ANS) are based on harvests from 1990–1999 and included test fishery distribution. **Bold underlined** cells indicate harvest amounts are below the minimum ANS.

b. Species-specific ANS ranges do not apply before 2001.

Table 4-8.—Estimated subsistence harvest of nonsalmon fish by community, Yukon Area, 2012.

Community	Households		Estimated nonsalmon harvest				
	Total	Surveyed ^a	Large whitefish ^b	Small whitefish	Northern pike	Sheefish	Total
Hooper Bay	218	78	447	1,447	456	25	2,375
Scammon Bay	99	44	1,066	1,578	1,854	836	5,334
Coastal district subtotal	317	122	1,513	3,025	2,310	861	7,709
Nunam Iqua (Sheldon Point)	42	34	377	2,381	171	1,189	4,118
Alakanuk	158	54	3,006	7,406	1,732	2,355	14,499
Emmonak	180	89	3,497	6,122	1,181	2,295	13,095
Kotlik	110	37	694	2,661	375	1,122	4,852
District 1 subtotal	490	214	7,574	18,570	3,459	6,961	36,564
Mountain Village	152	56	1,302	612	1,579	426	3,919
Pitkas Point	27	22	502	59	78	48	687
Saint Marys	127	49	3,969	834	496	757	6,056
Pilot Station	118	57	931	127	66	416	1,540
Marshall	69	26	1,840	48	387	296	2,571
District 2 subtotal	493	210	8,544	1,680	2,606	1,943	14,773
Russian Mission	72	26	2,468	0	2,758	331	5,557
Holy Cross	55	31	291	73	53	44	461
Shageluk	29	11	1,027	0	118	85	1,230
District 3 subtotal	156	68	3,786	73	2,929	460	7,248
Anvik	35	27	658	0	62	97	817
Grayling	47	18	2,872	26	298	637	3,833
Kaltag	58	15	870	232	116	394	1,612
Nulato	72	23	354	25	65	465	909
Koyukuk	49	22	858	58	167	179	1,262
Galena	169	51	1,241	216	154	288	1,899
Ruby	66	21	21	0	104	149	274
Huslia	95	33	1,325	399	698	390	2,812
Hughes	31	25	1,086	973	0	103	2,162
Allakaket	63	21	3,812	420	4,200	2,394	10,826
Alatna	10	4	411	1,000	23	83	1,517
Bettles	22	17	0	0	0	3	3
District 4 subtotal	717	277	13,508	3,349	5,887	5,182	27,926
Tanana	103	52	4,695	1,473	257	1,194	7,619
Stevens Village	18	11	47	14	83	36	180
Birch Creek	16	12	454	0	60	5	519
Beaver	31	24	47	12	67	13	139
Fort Yukon	211	86	1,370	741	562	415	3,088
Venetie	75	24	0	0	131	7	138
Chalkyitsik	28	18	11	0	99	17	127
District 5 subtotal	482	227	6,624	2,240	1,259	1,687	11,810
Total	2,655	1,118	41,549	28,937	18,450	17,094	106,030

Source Jallen et al. (In prep).

- The number of households contacted per species may vary. The number of households indicated is the greatest number of households contacted for a given species.
- Whitefish that are greater than 4 lb in weight are considered large whitefish, and those that are less than 4 lb in weight are considered small whitefish.




Figure 4-5.—Map of the Alaska portion of the Yukon River drainage, showing communities and districts.




Figure 4-6.—Yukon Area estimated subsistence salmon harvests, 2012.




Figure 4-7.—Estimated subsistence salmon harvests by species, Yukon Area, 1988–2012.




Figure 4-8.—Estimated number of dogs by district, Yukon Area, 2012.




Figure 4-9.—Primary gear type utilized for subsistence salmon fishing, Yukon Area, 2012.

CHAPTER 5: KUSKOKWIM AREA

BACKGROUND

The subsistence salmon fisheries in the Kuskokwim Area are some of the largest in the state of Alaska, in terms of the number of residents who participate and the number of salmon harvested (Fall et al. 2013). Since 1994, when the Alaska Department of Fish and Game (ADF&G) began acquiring reasonably complete statewide coverage of subsistence harvest survey data, over 50% of king salmon harvested under subsistence regulations have been taken in the Kuskokwim Area, mostly in the Kuskokwim River drainage. Between 2010 and 2013 (study years 2009–2012), the Division of Subsistence conducted comprehensive subsistence harvest and use surveys in 18 Kuskokwim River communities. The results indicate that on average salmon contributes 42% of the total wild resource harvest (in edible pounds) in the Lower Kuskokwim communities, 65% in the Central Kuskokwim communities, and 25% in the Upper Kuskokwim communities (Brown et al. 2012, 2013; Ikuta et al. 2014).²¹ Residents of the Kuskokwim Area harvest 5 species of Pacific salmon for subsistence purposes: Chinook salmon *Oncorhynchus tshawytscha*, chum salmon *O. keta*, coho salmon *O. kisutch*, pink salmon *O. gorbuscha*, and sockeye salmon *O. nerka*. Drift gillnetting, set gillnetting, and hook and line fishing are the primary methods used when harvesting salmon, although additional gear types are allowed as specified in 5 AAC 01.270. Kuskokwim Area communities are heavily reliant upon the annual returns of salmon not only for basic nutrition, but also for maintenance of cultural identity and cultural values, in addition to economic opportunities for commercial sales (Andrews and Coffing 1986; Andrews 1989:154; Barker 1993; Brown et al. 2012, 2013; Coffing 1991; Fienup-Riordan 1990:184, 1995:120, 123; Himmelheber 1987:32; Ikuta et al. 2014, 2013; Oswalt 1963a–b, 1990; Pete 1993; Senecal-Albrecht 1998, 1990; Walker and Coffing 1993; Wolfe et al. 1984).

ADF&G has been estimating Kuskokwim Area subsistence salmon harvests annually by postseason subsistence harvest survey since 1960. Simon et al. (2007) discussed the history of annual harvest monitoring methods used by the Division of Commercial Fisheries from 1960–1987 as well as the different methods used from 1988–2007 by the Division of Subsistence (see also Walker and Coffing 1993). Beginning in 2008, the Division of Commercial Fisheries once again assumed responsibility for the annual post-season subsistence salmon harvest monitoring program using methods outlined in Carroll and Hamazaki (2012a). In the Kuskokwim Area, there are 38 communities, 28 of which are surveyed each year on a voluntary basis. As Table 5-1 shows, in 2012, there were approximately 4,294 households in 32 communities excluding the 6 Bering Sea communities.²² Bethel is the largest community in the region, consisting of approximately 2,128 households in 2012. The north Kuskokwim Bay communities of Kwigillingok, Kongiganak, and Kipnuk are not located on the Kuskokwim River, but many subsistence salmon fishing households from these communities have traveled to the Kuskokwim River to fish, but may have also harvested salmon from coastal areas and local tributaries (Himmelheber 1987:7; Stickney 1984:60–61; Walker and Coffing 1993:1). Except in 2000 and 2004, only the community of Kongiganak (Carroll and Hamazaki 2012a) has participated in the voluntary ADF&G harvest survey. The communities of Quinhagak, Goodnews Bay, and Platinum, located in south Kuskokwim Bay, comprise 7% of the total Kuskokwim Area households (Carroll and Hamazaki 2012b), and harvest salmon primarily from the drainages of the Kanektok, Arolik, and Goodnews rivers (Walker and Coffing 1993:1; Wolfe et al. 1984:321–322). Subsistence users from Bering Sea coastal communities have chosen to not participate in the ADF&G study for most years. These include the communities of Mekoryuk (on

21. See also Ikuta, Hiroko, David M. Runfola, and David S. Koster. In prep. Bethel Subsistence, 2012: Wild Resource Harvests and Uses, Land Use Patterns, and Subsistence Economy in the Hub Community of the Yukon–Kuskokwim Delta. Fairbanks: Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. NNN. Hereinafter referred to as (Ikuta et al. *In prep*).

22. Household number estimates are not available for the coastal communities. Subsistence users from these communities harvest salmon in coastal waters as well as in area rivers. Relatively little documentation exists of subsistence salmon harvests of Bering Sea coastal communities because the communities are not included in either the Kuskokwim or the Yukon postseason subsistence salmon harvest monitoring programs.

Nunivak Island), Newtok, Tununak, Toksook Bay, Nightmute, and Chefornak (Carroll and Hamazaki 2012a–b). While little information is available, residents of Bering Sea coastal communities harvest salmon from local rivers and coastal waters, which likely include coastal stocks as well as mixed stocks that were not bound for the Kuskokwim River (Fienup-Riordan 1983:112; Walker and Coffing 1993:1). In 2011, sponsored by the Arctic-Yukon-Kuskokwim Sustainable Salmon Initiative, the Association of Village Council Presidents (AVCP) collected subsistence salmon harvest data in 7 coastal communities: Chefornak, Kipnuk, Mekoryuk, Newtok, Nightmute, Tooksook Bay, and Tununak (Kwigillingok chose not to participate in the AVCP project) (Wolfe et al. 2012). This project provides the only reliable subsistence salmon harvest data in the recent years for this portion of the Kuskowkim Area (Table 5-2), and the data were used for the Amount Reasonably Necessary (ANS) for subsistence determination for the remainder of the Kuskokwim Area in by the Alaska Board of Fisheries in 2013.

In 2012, sharp declines in Chinook salmon abundance caused severe hardship for fishery-dependent communities in the Kuskokwim Area. Subsistence fishers were affected by the 12-day rolling closures of all subsistence salmon fishing in the Kuskokwim River and its tributaries. A poor king salmon run and 35 days of management restrictions resulted in low harvests of Chinook salmon that were approximately 70% below the recent 10-year average (Shelden et al. 2014). As a result, the U.S. Department of Commerce declared a resource disaster for the Kuskokwim River Chinook salmon fishery on September 13, 2012.

REGULATIONS

Statewide eligibility criteria require individuals to be Alaska residents for the preceding 12 months before harvesting salmon for subsistence uses. Most subsistence salmon fishers in the region are Kuskokwim Area residents; however, some subsistence fishers are domiciled in other parts of Alaska and return to fish alone, or to assist family or friends with the harvesting or processing of salmon (Simon, Krauthoefer, Koster, and Caylor 2007:5).

Prior to 1990, there were additional restrictions on participation in the subsistence fishery related to the state's rural priority for subsistence, which subsequently was determined by the Alaska Supreme Court to be unconstitutional. As a result of the passage of Alaska National Interest Lands Conservation Act (ANILCA) and in light of a 1989 Alaska Supreme Court decision, the federal government established the federal subsistence program, which provides subsistence opportunity for qualified rural residents on applicable federal public lands and in applicable federal public waters. Individuals must be Kuskokwim Area residents to participate in the Kuskokwim federal subsistence salmon fishery (50 CFR § 100.5). Federal subsistence schedules, openings, closings, and fishing methods are generally the same as those for state subsistence salmon fisheries, unless superseded by federal special action. Regulatory authority for Kuskokwim river salmon management is shared by the Federal Subsistence Board (FSB) and the State of Alaska Board of Fisheries (BOF). On the Kuskokwim, ADF&G is responsible for implementing regulations in accordance with the Kuskokwim River Salmon Management Plan (5 AAC 07.365) and also has inseason discretionary management authority of salmon in Alaska navigable waters. Waters of the lower Kuskokwim River are largely within or adjacent to federal public lands, namely the Yukon Delta National Wildlife Refuge. As such, the U.S. Fish and Wildlife Service (USFWS) shares inseason subsistence fishing management decision making with ADF&G. USFWS holds final decision-making authority over management of salmon in these waters only in the event that the federal subsistence program determines that all non-federally qualified subsistence uses must be eliminated in order to meet the federal subsistence priority. The Kuskokwim River Salmon Management Working Group (KRSMWG) is composed of knowledgeable stakeholders acting in a representative fashion for communities throughout the Kuskokwim River drainage, processors, sport fishery representatives, as well as an ADF&G management biologist. The working group advises state and federal managers through an established process and is currently the primary forum through which management decisions are made regarding Kuskokwim River subsistence, commercial, and sport salmon fisheries (Smith and Linderman

Jr. 2008:1). The highest priority in state and federal management of the Kuskokwim River's salmon populations is biological sustainability of the resources based on principles of sustained yield. In the event that returning salmon numbers are not sufficient to meet established escapement goals that will allow for the maintenance of future generations of salmon populations, consumptive uses of salmon may be restricted. Under conditions that there is a harvestable surplus beyond these minimum escapement levels, consumptive uses of salmon are prioritized for different user groups.

Alaska Statute 16.05.258, "Subsistence use and allocation of fish and game," establishes the subsistence use priority (above sport, commercial, and personal uses) when resources are not abundant enough to provide for all consumptive uses, while remaining in accordance with principles of sustained yield. Subsistence uses protected by the subsistence priority are those practices identified as customary and traditional practices, as determined by the BOF. In 1993, the BOF made positive findings for customary and traditional uses of all salmon species in the entire Kuskokwim Area.²³ As part of these findings, the BOF then determined the amount reasonably necessary for subsistence (ANS) in these respective areas as one means to provide reasonable opportunities for subsistence uses. Based on historical harvest information, an ANS of 192,000–242,000 for salmon of all species in the Kuskokwim Area was determined (5 AAC 01.286). In 2001, the BOF amended these ANS ranges for Kuskokwim River using subsistence harvest data from the years 1990 to 1999. After reviewing various options, the BOF made new customary and traditional use and ANS findings for the Kuskokwim area by species. Although not in effect during the study year of 2012, in January 2013, the BOF again modified ANS ranges by species for each river system²⁴ (Ikuta 2012).

Subsistence harvest of Pacific salmon species in the Kuskokwim River is allowed without a permit (5 AAC 01.280) and with no closed season (5 AAC 01.260), unless otherwise noted for conservation purposes. Alaska regulations allow a variety of gear types to be used in the Kuskokwim River for subsistence salmon fishing and include specifications regarding the use of gillnets (5 AAC 01.270) and hook and line gear (5 AAC 01.295). There are no federal or state bag or possession limits for subsistence salmon harvests in the Kuskokwim River, except from June 1 through August 31, when subsistence fishing with a hook and line attached to a rod or pole, in that portion of the Aniak River drainage upstream of Doestock Creek, the bag and possession limit is 2 Chinook salmon and rainbow trout, *O. mykiss*, may not be retained (5 AAC 01.295). Federal regulations of all subsistence fish harvests in Alaska federal public lands and waterways are administered under 50 CFR §100.27, including seasons, gear types, and bag and possession limits on all salmon and nonsalmon species.

By regulation, therefore, the subsistence salmon fishing season is open unless a subsistence fishing schedule closure is implemented. If closures to the fishery are necessary, they are implemented by emergency order prior to, during, and after commercial fishing periods, or closures to the fishery are implemented by emergency order for conservation purposes (see 5 AAC 01.260, and 5 AAC 07.365). On the Kuskokwim River, a subsistence fishing schedule with periodic fishing closures (openings between these closures were often referred to as "windows" or "openers") was implemented from 2001–2006 and has since been discontinued.

Subsistence Fishery

In 2012, preseason outlooks suggested a weak return of Chinook salmon to the Kuskokwim River.²⁵ With a concern for area tributary escapements and the outlook for overall returns, ADF&G and USFWS

23. The Kuskokwim Area includes the Kuskokwim River drainage, all waters of Alaska that flow into the Bering Sea between Cape Newenham and the Naskonat Peninsula, and Nunivak and St. Matthew islands. 38 communities are located within this area.

24. The current ANS ranges for salmon in the Kuskokwim River drainage, determined by the BOF in January 2013, are as follows: 67,200–109,800 Chinook salmon; 41,200–116,400 chum salmon; 32,200–58,700 sockeye salmon; 27,400–57,600 coho salmon; and 500–2,000 pink salmon; in districts 4 and 5 combined: 6,900–17,000 salmon; and in the remainder of the Kuskokwim area: 12,500–14,400 salmon (5 AAC 01.286).

25. Alaska Department of Fish and Game Division of Commercial Fisheries. "2102 Kuskokwim Area salmon outlook and management strategy," news release, March 8, 2012.

inseason managers suggested that harvestable surplus of Chinook salmon might not be enough to meet subsistence needs. Preseason tributary closures were followed by inseason mainstem closures. In the June 8 KRSMWG meeting, the managers recommended a 7-day rolling closure for all subsistence salmon fishing to begin in the lower section of Kuskokwim River Subdistrict 1-B effective June 10, 2012.^{26,27} The rolling closure was implemented in a stepwise progression up the Kuskokwim River consistent with salmon run timing. By June 15, data from the Bethel Test Fishery catch per unit effort (BTF CPUE) indicated that continued returns of king salmon were still not sufficient to meet the management objective. In the June 15 KRSMWG meeting, the managers recommended a 5-day extension to the original 7-day rolling closure in an effort to conserve Chinook salmon.

During the June 20 KWRMWG meeting, the inseason managers presented a recommendation for a 3-day subsistence fishing opening allowing 6-inch or smaller mesh gillnets after each section's 12-day rolling closure.²⁸ Subsistence salmon fishing opened for 6 consecutive days following the rolling closures. After the 6-day opening, ADF&G and USFWS inseason managers initiated a 2-day rolling closure, followed by the opening with the use of gillnets with 6-inch or less mesh permitted. Subsistence salmon fishing remained open with a 6-inch mesh restriction through July 15, when BTF CPUE indicated the end of the king salmon run in the lower river. On July 16, inseason managers ended restrictions on subsistence salmon fishing, allowing the unrestricted use of gillnets, hook and line gear, and fish wheels in the lower section of Subdistrict 1-B. The restrictions were lifted in the upriver sections with the rolling schedule. On July 23, subsistence salmon fishing with unrestricted gear resumed in Section 4 and 5.

In 2012, the Kuskokwim Area Chinook salmon subsistence harvest was the lowest on record and about 70% below the 10-year average. Furthermore, in 2011 and 2012, estimated Chinook salmon escapement on monitored tributaries was the lowest since 1990, and escapement goals were not met at Kwethluk, Tuluksak, and George River (Shelden et al. 2014). Kuskokwim Area harvests of Chinook salmon fell below the lower limit of the ANS range. Subsistence harvests of chum and coho salmon in the Kuskokwim River were within or exceeded the ANS ranges defined for the area. Sockeye salmon subsistence harvests were above the upper range of the ANS.

Throughout salmon fishing season, many subsistence fishers expressed that harvesting and storing salmon is critical to many families' survival each year. People explained that restrictions to salmon fishing and the resulting disruptions in the seasonal round cause serious limitations to food supplies and the threat of extreme hardship in months to come.

Commercial Fishery

Since 1987, ADF&G has not provided a directed commercial harvest opportunity for Chinook salmon in the Kuskokwim River drainage. In January 2004, the BOF granted ADF&G discretionary emergency order authority to close the subsistence salmon fishery around commercial salmon fishing periods in districts 1 and 2. Prior to this action, areas within commercial salmon fishing districts were closed to subsistence salmon net and fish wheel gear 16 hours before, during, and 6 hours after commercial fishing periods (Simon, Krauthoefer, Koster, and Caylor 2007:7). Since 2004, areas within commercial salmon fishing districts were closed to subsistence salmon net and fish wheel gear 6 hours before, during, and 3 hours after commercial fishing periods (5 AAC 01.260). The purpose of these closures was to discourage illegal fishing activities, such as the sale of subsistence-caught salmon in the commercial fishery (Simon, Krauthoefer, Koster, and Caylor 2007:7).

26. Fishers were prohibited from harvesting king salmon with hook and line gear and restricted to the use of gillnets with 4-inch or less mesh not exceeding 60 feet in length. Subsistence fishers were permitted to retain incidental catches of king salmon with the use of a legal gillnet. Fish wheels were permitted; however, they were required to be equipped with a livebox, which fishers were required to check at least every six hours and return all king salmon to the water alive.

27. Alaska Department of Fish and Game Division of Commercial Fisheries. "2012 Preliminary Kuskokwim Area salmon season summary," news release, October 3, 2012.

28. Alaska Department of Fish and Game Division of Commercial Fisheries. "2012 Preliminary Kuskokwim Area salmon season summary," news release, October 3, 2012.

In 2012, a total of 393,319 salmon were commercially harvested from the Kuskokwim Area. A total of 477 individual permit holders participated in area commercial fisheries with an estimated exvessel value of \$2,040,296; this was approximately 41% above the most recent 10-year average value.²⁹ The District 1 commercial fishing season was from July 13 to August 27. The initiation of the commercial fishery was delayed until the majority of the Chinook run had passed through the district to ensure ongoing Chinook salmon conservation. As a result, commercial fishing occurred after the peak of the sockeye and chum salmon runs had passed through the district. Subsistence salmon fishing was closed by emergency order 6 hours before, during, and 3 hours after commercial fishing periods. There were 23 commercial fishing periods. The District 4 (Quinhagak) and District 5 (Goodnews Bay) commercial fishing seasons were from June 27 to August 29 with 22 commercial fishing periods in Quinhagak and with 28 fishing periods in Goodnews Bay. Subsistence fishing was closed 8 hours before, during, and 6 hours after commercial fishing periods.

SUBSISTENCE SALMON HARVEST ASSESSMENT METHODS

ADF&G has been estimating Kuskokwim Area subsistence salmon harvests annually by postseason subsistence harvest survey since 1960: by the Division of Commercial Fisheries in 1960–1987, by the Division of Subsistence in 1988–2007, and by the Division of Commercial Fisheries since 2008 (Carroll and Hamazaki 2012a–b). The purpose of the survey is to collect data about the number and species of salmon harvested by area residents.

For data collection in 2012, under a cooperative program between ADF&G and the USFWS Office of Subsistence Management, subsistence salmon harvest data collection in Bethel was conducted by staff from the Orutsararmuit Native Council (ONC), which has been involved in subsistence salmon harvest monitoring in Bethel since 1999. Subsistence harvest data collection in Aniak was conducted under a similar agreement by staff from the Kuskokwim Native Association (KNA), which has been involved in subsistence salmon harvest monitoring in Aniak since 2002 (Simon, Krauthoefer, Koster, and Caylor 2007).

The data from the postseason subsistence harvest survey are analyzed to provide an estimate of the number of salmon harvested for subsistence purposes. This information has been used by ADF&G, the U.S. Fish and Wildlife Service (USFWS), the BOF, and the Federal Subsistence Board to manage customary and traditional uses of salmon and to provide reasonable opportunities for continued customary and traditional (C&T) uses of salmon throughout the area. More detailed descriptions of subsistence salmon harvest monitoring methods utilized in the Kuskokwim Area are found elsewhere (Hamazaki 2011; Simon, Krauthoefer, Koster, and Caylor 2007; Walker and Coffing 1993).

Household Harvest Surveys

Study Design

The postseason subsistence harvest survey for the majority of communities was designed based on a stratified random survey methodology (Scheaffer 1990). From 1989 to 2010, each household was classified into three strata based on the household's recent 2-year history of participation in the subsistence fishery. Beginning in 2005, surveys collected harvest data for pink salmon. In 2011, the above household classification was expanded into 5 strata based on a household's most recent 2 known years of participation within the past 5 years in the subsistence fishery.

An attempt was made to census the 2 highest harvest strata groups, and the strata group for which no harvest level could be determined. The remaining 2 strata groups of light and non-harvesters were sampled at 30% each. A full description of methods used for harvest level classification and sampling are available in Shelden et al. (2014).

29. Alaska Department of Fish and Game Division of Commercial Fisheries. "2012 Preliminary Kuskokwim Area salmon season summary," news release, October 3, 2012.

The data were entered into the subsistence harvest database maintained by the Division of Commercial Fisheries. The data were subsequently extracted and processed by Commercial Fisheries staff, as described in Shelden et al. (2014). The analyzed data were transmitted to Division of Subsistence analysts for final formatting and inclusion in the Alaska Subsistence Fishery Database, maintained by the Division of Subsistence. All subsistence harvest data were treated as confidential, such that individual harvest data are not shared and all analysis is aggregated and anonymous. The study was generally conducted in accordance with the *Alaska Federation of Natives Guidelines for Research*.³⁰

Estimating Bethel Salmon Harvests

In Bethel, the Division of Commercial Fisheries was responsible for designing and producing the survey instrument and selection of survey households, and ONC was responsible for conducting household surveys. Due to the impracticality of maintaining an accurate household list in order to stratify Bethel, a 50% random survey was conducted based on a simple random survey methodology where each dwelling (physical location instead of household) was the primary sampling unit. Before the harvest survey, ADF&G oriented ONC technicians to the project and instructed them in the proper implementation of the survey. ONC technicians conducted surveys in Bethel from October through November. Survey data were entered and analyzed by Division of Commercial Fisheries staff to generate subsistence salmon harvest estimates by species.

Estimating Aniak Salmon Harvests

Like Bethel, Aniak is too large to effectively maintain a reliable household list, but due to its smaller size than Bethel, an attempted census is required to obtain the level of reliability of the estimate desired for this project.

ADF&G Division of Commercial Fisheries was responsible for designing and producing the survey instrument and selection of survey households, and KNA was responsible for conducting household surveys in Aniak. Before the harvest survey, ADF&G oriented KNA technicians to the project and instructed them in the proper implementation of the survey. KNA technicians conducted surveys in Aniak from October through December 2012. Survey data were entered and analyzed by ADF&G Commercial Fisheries staff to generate subsistence salmon harvest estimates by species.

Estimating Kuskokwim Area Community Subsistence Salmon Harvests

For the remaining 36 communities in the Kuskokwim Area, the goal was to collect subsistence harvest data through harvest surveys conducted by Division of Commercial Fisheries staff beginning in the first week of October and continuing through November. The survey crew consulted with community officials before arriving in the community to update community household lists. Other resources were also useful in updating household lists, including telephone and utility records. Communities were prioritized based on transportation scheduling, staff time and community willingness to participate in the program. Participation in the surveys was voluntary, and some community leaders requested that the surveys not take place in their communities.

Harvest Calendars

In addition to systematic household harvest surveys, subsistence salmon harvest calendars were mailed in late April or early May so that they were available to fishers prior to the start of the salmon fishing season. The calendar data continue to be instrumental for examination of subsistence salmon harvest timing. The calendars are also used by some area fishers throughout the fishing season so that they can be referenced during postseason household harvest surveys.

30. Alaska Federation of Natives. 2013. "Alaska Federation of Natives Guidelines for Research." Alaska Native Knowledge Network. Accessed May 14, 2014. <http://www.ankn.uaf.edu/IKS/afnguide.html>.

Calendar mailings were based on the most up-to-date household lists used in the harvest monitoring program. Extra calendars were printed and maintained at the Bethel ADF&G office for distribution as needed or upon request. In an effort to increase the use and return rate of subsistence salmon harvest calendars, public service announcements were broadcast on local radio stations during the fishing season reminding fishers to keep logging their catches on their calendars and describing the importance of calendars for documenting subsistence salmon uses.

Most subsistence salmon harvest data obtained from the returned calendars were not used to directly calculate Kuskokwim Area subsistence salmon harvest estimates, but these data were used to corroborate household survey data. Calendars were occasionally used as the primary source of harvest data when contact was not made with a particular household. Calendars often include harvests from multiple households that fished together, so reported harvests may represent the efforts of multiple households. In such cases, every effort was made to contact the head of household to verify harvest information when using the calendar data. Calendars also provided data for assigning households to the 3-user strata, especially in cases where households were not contacted as part of the household surveys. Calendar data are not provided in this report.

Data Correction and Archiving

Division of Commercial Fisheries staff reviewed and edited all completed surveys and periodically sent reviewed surveys to staff in Bethel and Anchorage for further processing. The survey data were entered into an Arctic-Yukon-Kuskokwim Database Management System (AKDBMS) data server. The database is structured to ensure data are entered completely and accurately, and there were periodic back-ups to prevent data loss.

Data Analysis

Community estimates of subsistence salmon harvest for surveys collected in communities outside of Bethel and Aniak were generated using a stratified random sampling expansion technique. This approach applies means to unsurveyed households within each strata group, and sums total estimates of the 5 strata groups to give a community harvest estimate. Communities where harvest survey data were inadequate or unavailable, for 2011 and earlier, were estimated by employing a Bayesian hierarchical multiple imputation method. The details of these approaches are described in (2014).

2012 SAMPLING SUMMARY

From an estimated total of 4,294 households located in the Kuskokwim Area, excluding households in the coastal communities that declined to participate, surveys were conducted with 1,569 unique households within 25 Kuskokwim Area communities (Table 5-1). In 2012, chum salmon contributed 43% of the estimated subsistence salmon harvest, followed by sockeye salmon (27%, 50,616), coho salmon (16%, 30,221 fish), Chinook salmon (13%, 25,336 fish), and pink salmon (1%, 2,160 fish) (Figure 5-1). As noted above, a new method was developed in 2008 to estimate subsistence salmon harvests in communities in which no household surveys took place if adequate harvest data for previous years existed; however, there are Kuskokwim Area communities for which there are insufficient historical data to develop annual harvest estimates using a Bayesian hierarchical multiple imputation method. As a result, the Kuskokwim Management Area total should be viewed as a minimum estimate because data for some communities are not available (Simon, Krauthoefer, Koster, and Caylor 2007:20).

For lower Kuskokwim River communities, 1,020 (31%) of the 3,279 households were contacted. Based upon 2012 data, the region represents 76% of the estimated total number of households in the Kuskokwim Area.

In the south Kuskokwim Bay region (Quinhagak, Goodnews Bay, and Platinum), 130 (52%) of the 249 households were contacted. The Bering Sea coastal communities of Mekoryuk, Newtok, Nightmute, Toksook Bay, Tununak, and Chefornak had an estimated 453 total households as of 2009, but none were

surveyed in 2012, and data for previous years are incomplete. Currently, subsistence salmon harvest information collected by AVCP for 2011 is the only available and reliable data source for the region (Wolfe et al. 2012).

The 13 communities of the middle and upper Kuskokwim River regions are generally smaller than lower river communities, and together compose 16% of total households in the Kuskokwim Area. In the middle Kuskokwim River region, defined here as communities located on the Kuskokwim River from Lower Kalskag upriver to Chuathbaluk, 255 (71%) of 361 households were contacted in 2012. For upper Kuskokwim communities, defined here as communities located on the Kuskokwim River from Crooked Creek upriver to Telida (in addition to Lime Village located on the Stony River and Takotna located on the Takotna River), 164 (52%) of 315 households were contacted. Takotna and Telida were not surveyed in 2012. The communities of Georgetown and Napaimute are not currently included in the community sampling list due to limited permanent populations and primarily seasonal use patterns for these 2 communities; the large majority of Georgetown and Napaimute community members are surveyed during their residence in other Kuskokwim River communities.

2012 SUBSISTENCE SALMON HARVEST SUMMARY

A summary of the subsistence salmon harvest estimates by community and fishing area is presented in Table 5-1. In 2012, subsistence salmon harvest estimates for communities contacted in the Kuskokwim Area totaled 25,336 Chinook salmon (13%), 81,912 chum salmon (43%), 50,616 sockeye salmon (27%), 30,221 coho salmon (16%), and 2,160 pink salmon (1%), for a total estimate of 190,245 salmon (Figure 5-1). A poor Chinook salmon run and 35 days of management restrictions resulted in lowest harvest of Chinook salmon subsistence harvest, which was about 70% below the 10-year average (Table 5-3). The total chum salmon harvest was up sharply, 38% and 27% above the recent 5- and 10-year averages. The total harvest of sockeye salmon was above the recent 10-year average. It is possible that subsistence harvesters have been targeting more abundant species in years of lower Chinook salmon abundance, and they are tied to both voluntary and involuntary changes in gear usage. Chinook salmon abundance in the Kuskokwim River drainage has decreased since 2007, with some of the lowest total runs occurring in 2010–2012 (Bue et al. 2012; personal communication with Kevin Scharberg, Kuskokwim Area Research Biologist, September 12, 2013). Lower Kuskokwim River Area communities accounted for 73% of the 2012 estimated subsistence salmon harvests in the Kuskokwim Area and 74% of the entire estimated Chinook salmon subsistence harvest. Residents of Bethel accounted for 35% of the Kuskokwim Area subsistence salmon harvests and 29% of subsistence-caught Chinook salmon and 44% of the estimated total of subsistence-caught coho salmon.

As noted, several coastal communities within the Kuskokwim Area have chosen not to participate in the post-season subsistence harvest surveys conducted by ADF&G. However, 7 of these communities participated in a study conducted by AVCP to estimate subsistence salmon harvests for 2011 (Wolfe et al. 2012; Table 5-2). The total estimated subsistence harvest of salmon for these 7 communities in 2011 was 16,593 fish, including 7,226 chum (44%), 4,439 sockeye (27%), 2,864 coho (17%), 1,298 Chinook (8%), 746 pink (4%), and 20 salmon of unknown species (<1%).

Use of Salmon for Dog Food

Historically, salmon harvested for use as dog food were a large portion of the overall subsistence salmon harvest; specifically, chum and coho salmon. In recent years, the number of households harvesting salmon specifically for dog food has declined, likely due to decreased use of dog teams for transportation. In 2012, data show a reported harvest of 23,241 salmon for use as dog food (Table 5-4). The majority of the salmon harvested for dog food were chum salmon, at 14,361 fish, while coho salmon accounted for 6,230 fish. Sockeye salmon contributed 652 fish and pink salmon 1,870 fish to the harvest for dog food. Households do not target Chinook salmon for dog food; however, 128 Chinook salmon, likely unfit for human consumption, were reported to have been fed to dogs in an effort to avoid wasting the fish. It is

common for most households to feed scraps—backbones, entrails, and salmon unfit for human consumption—to their dogs.

Gear Types

Kuskokwim Area subsistence fishers deploy a variety of gear types to harvest salmon (e.g., set gillnet, drift gillnet, fish wheel, or rod and reel) (Table 5-5). Households that harvested salmon were asked to provide information on the primary gear type used by their household for harvesting salmon. In 2012, out of 878 contacted fishing households that responded to gear type questions, 695 (79%) reported drift gillnets as their primary subsistence salmon fishing gear type, 101 (12%) reported set gillnets, and 81 (9%) reported subsistence rod and reel gear. Preferred gear types vary between regions of the Kuskokwim Area, and fishers often select which gear type to use based on local environmental factors such as river morphology. In recent decades drift gillnets have been the most common gear type deployed by fishers in the lower and middle Kuskokwim River communities where river depth and width permit the efficient use of this type of net. In communities of the upper Kuskokwim River, a narrower and generally shallower river channel typically restricts fishers to the use of set gillnets and occasionally fish wheels. Also, subsistence fishers who reside near clear water streams often harvest salmon by rod and reel (e.g., Kwethluk, Takotna, and Nikolai). Perhaps equally important in determining selection of gear types are local and familial customs and traditions associated with subsistence salmon fishing.

Salmon Retained from Commercial Fishing for Subsistence Uses

Households involved in commercial salmon fishing occasionally keep a portion of their commercial harvest for subsistence uses; however, the number of salmon retained from commercial fishing activities for subsistence is usually low. In 2012, few households reported retaining commercially-caught salmon for subsistence uses. Data show a reported total of 901 salmon were retained from commercial catches, including 289 Chinook, 76 chum, 158 sockeye, 322 coho, and 56 pink salmon (Table 5-6).

OTHER FISH

Harvest data for nonsalmon fish species are also collected as part of the postseason salmon survey. In 2012, reported harvests of nonsalmon species in the Kuskokwim Area included 15,073 humpback whitefish; 22,706 broad whitefish; 15,344 cisco; 2,109 sheefish; 4,961 burbot; 15,403 northern pike; 148,179 blackfish; 1,683 grayling; 6 char; 2,658 herring; 67,417 smelt; and 378 rainbow trout (Table 5-7). Humpback and broad whitefish harvests were expanded to total harvest estimates for all communities surveyed in 2012. The estimated harvest of humpback whitefish was 36,144 fish ($\pm 10,047$ CI), and the estimated harvest of broad whitefish was 49,550 fish ($\pm 15,126$ CI).

The Division of Subsistence has recently conducted comprehensive subsistence harvest and use surveys in the following 20 Kuskokwim River drainage communities: in 2010 (study year 2009), Aniak, Chuathbaluk, Crooked Creek, Lower Kalskag, Red Devil, Sleetmute, Stony River, and Upper Kalskag (Brown et al. 2012); in 2011 (study year 2010), Akiak, Kwethluk, Oscarville, Tuluksak, Georgetown, and Napaimute (Brown et al. 2013); in 2012 (study year 2011), Napakiak, Napaskiak, McGrath, Nikolai, and Takotna (Ikuta et al. 2014); and in 2013 (study year 2012), Bethel (Ikuta et al. *In prep*). These comprehensive surveys included questions about salmon and nonsalmon harvests as well as harvests of wildlife and plants. In addition, the division conducted an ethnographic project to understand socioeconomic patterns and trends of subsistence Chinook salmon fishing in Tuntutuliak, Kwethluk, Kalskag, Sleetmute, and Nikolai in 2009 and in the Bethel area in 2012 (Ikuta et al. 2013). Studies focusing on the traditional ecological knowledge of nonsalmon fishes and nonsalmon harvest amounts have been conducted in Aniak and Chuathbaluk for 2001–2003 (Krauthoefer et al. 2007), Bethel for 2001–2003 (Simon, Krauthoefer, Koster, Coffing, et al. 2007) as well as Eek, Tuntutuliak, and Nunapitchuk for 2005–2009 (Ray et al. 2010). Information on historical and contemporary harvest and use of salmon and nonsalmon in communities along the Kuskokwim River, where data are available, can be accessed through the Community Subsistence Information System (CSIS) on the ADF&G website.

THE ROLE OF SALMON WITHIN ANNUAL SUBSISTENCE HARVESTS

In addition to post-season salmon surveys conducted by ADF&G, Division of Commercial Fisheries staff, Division of Subsistence staff conducted comprehensive subsistence surveys for the year 2012 in Bethel. While moose, as an individual resource, contributed the most to subsistence harvests by edible weight in 2012, fish were the most widely used category of subsistence food, by 90% of households, with an estimated harvest of 579,202 edible pounds, representing 61% of the total subsistence harvest in 2012. Salmon represented 67% of the total estimated fish harvest and 41% of the total estimated subsistence harvests. However, a poor Chinook salmon run and 35 days of restrictions in 2012 resulted in lower harvests of Chinook salmon in Bethel (7,846 Chinook salmon) and throughout the drainage, approximately 70% below the recent 10-year average. Forty-nine percent of Bethel households reported using less Chinook salmon than in previous years.

Nonsalmon fishes represented 20% of the estimated total subsistence harvest in Bethel, with an estimated total of harvest of 189,180 edible pounds of nonsalmon fishes. The largest contributions to nonsalmon fish harvests were northern pike (57,619 pounds), representing 6% of the total subsistence harvests, and smelt (31,694 pounds), humpback whitefish (31,280 pounds), and burbot (28,936 pounds), each of which contributed 3% of the total subsistence harvest by Bethel residents in 2012. Other fish harvest data are included in the technical report (Ikuta et al. *In prep*).

Table 5-1.—Subsistence salmon harvests by community, Kuskokwim Area, 2012.

Community	Households		Estimated salmon harvest					
	Total	Contacted	Chinook	Sockeye	Coho	Chum	Pink	Total
Kipnuk ^b	—	—	—	—	—	—	—	—
Kwigillingok ^b	—	—	—	—	—	—	—	—
Kongiganak ^a	90	0	571	1,211	458	1,901	0	4,141
North Kuskokwim Bay	90	0	571	1,211	458	1,901	0	4,141
Tuntutuliak	90	53	1,123	1,516	565	2,614	15	5,833
Eek	86	45	1,004	1,490	612	1,552	50	4,708
Kasigluk	104	51	552	1,451	303	3,261	0	5,567
Nunapitchuk	111	61	845	2,396	319	5,312	32	8,904
Atmautluak	61	35	234	1,623	383	2,701	22	4,963
Napakiak	99	46	457	1,141	402	1,711	0	3,711
Napaskiak	97	42	1,108	2,065	269	3,216	122	6,780
Oscarville ^c	14	14	51	323	38	599	0	1,011
Bethel ^d	2,128	447	7,321	18,282	13,280	26,872	305	66,060
Kwethluk	164	83	1,709	2,884	1,013	3,849	91	9,546
Akiachak	157	74	2,862	3,443	714	4,150	53	11,222
Akiak ^c	79	16	856	1,820	474	2,416	0	5,566
Tuluksak	89	53	651	1,380	341	2,585	8	4,965
Lower Kuskokwim	3,279	1,020	18,773	39,814	18,713	60,838	698	138,836
Lower Kalskag	79	41	459	891	1,107	3,284	25	5,766
Kalskag (Upper)	62	31	562	770	360	1,930	30	3,652
Aniak	187	155	993	1,375	3,365	5,667	940	12,340
Chuathbaluk	33	28	103	297	179	796	2	1,377
Middle Kuskokwim	361	255	2,117	3,333	5,011	11,677	997	23,135
Crooked Creek	37	31	124	234	149	610	2	1,119
Red Devil	13	10	225	511	238	516	42	1,532
Sleetmute	40	35	132	715	784	1,004	120	2,755
Stony River ^c	16	3	212	398	372	619	0	1,601
Lime Village	14	10	29	780	117	419	129	1,474
McGrath	136	45	68	233	2,257	885	14	3,457
Takotna ^a	23	0	0	2	22	0	0	24
Nikolai	34	30	276	0	214	1,044	0	1,534
Telida ^b	2	0	—	—	—	—	—	—
Upper Kuskokwim	315	164	1,066	2,873	4,153	5,097	307	13,496
Kuskokwim River	4,045	1,439	22,527	47,231	28,335	79,513	2,002	179,608
Quinhagak	162	77	2,396	2,015	1,380	2,001	70	7,862
Goodnews Bay	68	37	389	1,197	382	322	72	2,362
Platinum	19	16	24	173	124	76	16	413

-continued-

Table 5-1.–Page 2 of 2.

Community	Households		Estimated salmon harvest					
	Total		Chinook	Sockeye	Coho	Chum	Pink	Total
South Kuskokwim Bay	249	130	2,809	3,385	1,886	2,399	158	10,637
Mekoryuk ^b	–	–	–	–	–	–	–	–
Newtok ^b	–	–	–	–	–	–	–	–
Nightmute ^b	–	–	–	–	–	–	–	–
Toksook Bay ^b	–	–	–	–	–	–	–	–
Tununak ^b	–	–	–	–	–	–	–	–
Chefornak ^b	–	–	–	–	–	–	–	–
Bering Sea Coast	–	–	–	–	–	–	–	–
Total	4,294	1,569	25,336	50,616	30,221	81,912	2,160	190,245

Source Shelden et al. (2014).

Note includes harvests using rod and reel and the removal of salmon from commercial harvests as well as subsistence nets.

Note – indicates data not available.

- These communities were not contacted during the 2012 study period. Harvests were estimated using historical average household harvest expanded by the number of households.
- These communities were not contacted during the 2012 study period. Not enough data was available to estimate harvest.
- Communities were contacted, but numbers of selected households or total number of surveyed households were insufficient. Harvests were estimated using historical average household harvest expanded by the number of households.
- A total of 888 Bethel households were contacted. Of these, 447 were preselected, and these were used for determining harvest estimates for this community.

Table 5-2.—Subsistence salmon harvests in 7 coastal Kuskokwim communities, 2011.

Community	Households		Percent surveyed	Estimated salmon harvest						
	Total	Surveyed		Chinook	Sockeye	Coho	Chum	Pink	Other ^a	Total
Chefornak	83	69	83.1%	161	261	61	338	13	5	839
Kipnuk	131	49	37.4%	479	1,160	781	716	11	0	3,147
Mekoryuk	59	54	91.5%	0	2	201	3670	47	0	3,920
Newtok	63	58	92.1%	144	394	262	103	46	0	949
Nightmute	50	40	80.0%	98	289	64	475	13	3	942
Toksook Bay	104	94	90.4%	365	1834	1040	1637	433	4	5,313
Tununak	68	36	52.9%	51	499	455	287	183	8	1,483
Total	558	400	71.7%	1,298	4,439	2,864	7,226	746	20	16,593

Source Wolfe et al. (2012:17-18).

a. Unidentified species of salmon.

Table 5-3.—Historical subsistence salmon harvests, Kuskokwim Area, 1989–2012.

Year	Households		Estimated salmon harvest				
	Total	Surveyed	Chinook	Sockeye	Coho	Chum	Total
1989	3,422	2,135	85,322	37,088	57,786	145,106	325,287
1990	3,317	1,448	114,219	48,752	63,084	157,335	314,513
1991	3,340	2,033	79,445	50,383	44,222	89,008	298,561
1992	3,308	1,308	87,663	46,493	57,551	120,126	246,914
1993	3,269	1,786	91,973	53,631	31,971	64,551	240,103
1994	3,169	1,801	110,922	46,127	40,815	89,553	251,111
1995	3,638	1,907	105,787	31,736	39,582	71,789	236,885
1996	3,630	1,524	100,352	41,532	45,279	102,079	241,572
1997	3,501	1,919	83,022	39,827	31,324	38,073	198,466
1998	3,497	1,940	85,781	38,228	27,435	72,860	218,595
1999	4,165	2,512	79,752	50,988	30,184	51,200	202,413
2000	3,317	1,448	75,299	53,468	49,469	72,851	204,714
2001	4,469	2,215	82,106	55,290	33,474	57,060	212,338
2002	4,804	2,687	84,512	34,331	44,588	94,998	205,599
2003	4,513	2,292	70,579	33,821	36,953	46,666	194,474
2004	4,638	2,398	103,183	43,425	53,186	68,068	214,959
2005	4,603	1,593	89,538	44,637	35,793	59,220	186,762
2006	4,671	1,439	96,857	49,467	43,880	96,021	286,225
2007	4,620	1,279	101,554	50,092	37,481	76,187	265,314
2008	4,734	992	103,080	63,802	49,755	71,177	287,814
2009	4,810	1,699	81,853	37,779	31,613	45,101	196,346
2010	4,215	2,247	69,242	41,042	34,169	47,885	192,338
2011	4,241	1,822	65,852	46,296	33,943	55,995	202,086
2012	4,294	1,569	25,336	50,616	30,221	81,912	188,085
5-year average (2007–2011)	4,524	1,608	84,316	47,802	37,392	59,269	228,780
10-year average (2002–2011)	4,585	1,845	86,625	44,469	40,136	66,132	223,192
15-year average (1997–2011)	4,320	1,899	84,814	45,500	38,216	63,557	217,896
Historical average (1989–2011)	3,995	1,845	89,039	45,141	41,458	77,953	235,799

Sources ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014); Shelden et al. (2014).

Table 5-4.—Number of households that own dogs, fed salmon to dogs, and total number of salmon fed to dogs, Kuskokwim Area, 2012.

Community	Households		Households		Total number of dogs	Reported salmon fed to dogs					
	Total	Contacted	Own dogs	Fed salmon		Chinook	Sockeye	Coho	Chum	Pink	Total
Kipnuk ^a	—	—	—	—	—	—	—	—	—	—	—
Kwigillingok ^a	—	—	—	—	—	—	—	—	—	—	—
Kongiganak	90	0	—	—	—	—	—	—	—	—	—
North Kuskokwim Bay	90	0	0	0	0	0	0	0	0	0	0
Tuntutuliak	90	52	42	1	103	0	0	0	40	0	40
Eek	86	44	36	4	70	5	15	5	42	0	67
Kasigluk	104	50	42	1	93	12	45	0	30	0	87
Nunapitchuk	111	58	47	3	102	0	0	1	30	0	31
Atmautluak	61	33	27	6	113	0	27	65	89	21	202
Napakiak	99	41	24	2	31	0	4	0	14	0	18
Napaskiak	97	37	22	2	69	0	0	0	0	0	0
Oscarville	14	13	10	3	20	0	0	0	175	0	175
Bethel	2,128	384	182	5	276	0	0	300	111	15	426
Kwethluk	164	80	62	5	143	15	22	30	65	12	144
Akiachak	157	72	38	5	177	3	89	0	149	4	245
Akiak	79	13	8	2	44	2	20	0	30	12	64
Tuluksak	89	48	34	4	96	10	15	0	45	0	70
Lower Kuskokwim	3,279	925	574	43	1,337	47	237	401	820	64	1,569
Lower Kalskag	79	36	28	5	72	0	15	150	260	0	425
Kalskag (Upper)	62	28	20	2	73	0	0	0	550	30	580
Aniak	187	151	90	13	255	0	25	1,532	3,376	770	5,703
Chuathbaluk	33	27	19	1	38	0	0	0	30	0	30
Middle Kuskokwim	361	242	157	21	438	0	40	1,682	4,216	800	6,738
Crooked Creek	37	30	22	5	61	2	0	0	140	0	142
Red Devil	13	9	7	4	14	0	8	0	235	9	252
Sleetmute	40	35	19	5	36	0	18	337	368	60	783
Stony River	16	3	2	0	3	0	0	0	0	0	0
Lime Village ^a	14	10	2	0	2	0	0	0	0	0	0
McGrath	136	39	17	3	29	0	0	561	462	0	1,023
Takotna	23	0	—	—	—	—	—	—	—	—	—
Nikolai	34	28	24	2	56	0	0	115	900	0	1,015
Telida ^a	2	0	—	—	—	—	—	—	—	—	—
Upper Kuskokwim	315	154	93	19	201	2	26	1,013	2,105	69	3,215
Kuskokwim River	4,045	1,321	824	83	1,976	49	303	3,096	7,141	933	11,522
Quinhagak	162	75	48	10	90	0	0	0	1	0	1
Goodnews Bay	68	34	23	3	42	15	23	19	39	2	98
Platinum	19	15	12	1	28	64	326	3,115	7,180	935	11,620

-continued-

Table 5-4.—Page 2 of 2

Community	Households		Households		Total number of dogs	Reported salmon fed to dogs					
	Total	Contacted	Own dogs	Fed salmon		Chinook	Sockeye	Coho	Chum	Pink	Total
South											
Kuskokwim Bay	249	124	83	14	160	79	349	3,134	7,220	937	11,719
Mekoryuk ^a	—	—	—	—	—	—	—	—	—	—	—
Newtok ^a	—	—	—	—	—	—	—	—	—	—	—
Nightmute ^a	—	—	—	—	—	—	—	—	—	—	—
Toksook Bay ^a	—	—	—	—	—	—	—	—	—	—	—
Tununak ^a	—	—	—	—	—	—	—	—	—	—	—
Chefornak ^a	—	—	—	—	—	—	—	—	—	—	—
Bering Sea Coast											
Total	4,294	1,445	907	97	2,136	128	652	6,230	14,361	1,870	23,241

Source Shelden et al. (2014).

Note Includes harvests using rod and reel and the removal of salmon from commercial harvests as well as subsistence nets.

Note — indicates data not available.

a. These communities were not contacted during the 2012 study period.

Table 5-5.—Gear types used for subsistence fishing, Kuskokwim Area, 2012.

Community	Total households ^c	Gear types ^a			
		Setnet	Driftnet	Rod and reel	Fish wheel
Kipnuk ^b	—	—	—	—	—
Kwigillingok ^b	—	—	—	—	—
Kongiganak ^b	—	—	—	—	—
North Kuskokwim Bay	0	0	0	0	0
Tuntutuliak	43	—	43	—	—
Eek	31	5	26	—	—
Kasigluk	32	1	31	—	—
Nunapitchuk	41	—	41	—	—
Atmautluak	23	1	22	—	—
Napakiak	30	1	29	—	—
Napaskiak	28	4	24	—	—
Oscarville	8	—	8	—	—
Bethel	168	10	145	13	—
Kwethluk	50	5	43	2	—
Akiachak	55	10	45	—	—
Akiak	5	2	3	—	—
Tuluksak	41	6	33	2	—
Lower Kuskokwim	555	45	493	17	0
Lower Kalskag	24	—	24	—	—
Kalskag (Upper)	19	—	19	—	—
Aniak	89	6	60	23	—
Chuathbaluk	17	—	14	3	—
Middle Kuskokwim	149	6	117	26	0
Crooked Creek	16	1	13	2	—
Red Devil	6	3	2	1	—
Sleetmute	14	6	6	2	—
Stony River	1	1	—	—	—
Lime Village	6	5	—	1	—
McGrath	14	7	3	3	1
Takotna ^b	—	—	—	—	—
Nikolai	20	10	—	10	—
Telida ^b	—	—	—	—	—
Upper Kuskokwim	77	33	24	19	1
Kuskokwim River	781	84	634	62	1
Quinhagak	58	6	41	11	—
Goodnews Bay	29	8	18	3	—
Platinum	10	3	2	5	—
South Kuskokwim Bay	97	17	61	19	0
Mekoryuk ^b	—	—	—	—	—
Newtok ^b	—	—	—	—	—
Nightmute ^b	—	—	—	—	—
Toksook Bay ^b	—	—	—	—	—
Tununak ^b	—	—	—	—	—
Chefornak ^b	—	—	—	—	—

-continued-

Table 5-5.–Page 2 of 2

Community	Total households ^c	Gear types ^a			
		Setnet	Driftnet	Rod and reel	Fish wheel
Bering Sea Coast	–	–	–	–	–
Total	878	101	695	81	1

Source Sheldon et al. (2014).

Note – indicates data not available.

- Only data regarding the primary gear type from each household was collected.
- Community was not contacted during the 2012 study period.
- Number of households responding to the question about their primary gear type.

Table 5-6.–Reported number of salmon retained from commercial harvest for subsistence use, Kuskokwim Area, 2012.

Community	Households		Reported salmon					
	Total	Responding	Chinook	Sockeye	Coho	Chum	Pink	Total
Kipnuk ^a	–	–	–	–	–	–	–	–
Kwigillingok ^a	–	–	–	–	–	–	–	–
Kongiganak ^a	90	0	–	–	–	–	–	–
North Kuskokwim Bay	90	0	0	0	0	0	0	0
Tuntutuliak	90	22	39	7	40	0	6	92
Eek	86	20	16	0	40	0	0	56
Kasigluk	104	17	40	46	74	30	0	190
Nunapitchuk	111	20	19	8	51	5	3	86
Atmautluak	61	6	6	0	26	15	0	47
Napakiak	99	11	8	0	0	2	0	10
Napaskiak	97	4	5	0	0	0	0	5
Oscarville	14	3	4	15	9	1	0	29
Bethel	2,128	20	17	0	1	0	0	18
Kwethluk	164	18	19	5	28	0	6	58
Akiachak	157	30	76	35	5	17	4	137
Akiak	79	2	5	0	0	0	0	5
Tuluksak	89	7	0	0	0	0	0	0
Lower Kuskokwim	3,279	180	254	116	274	70	19	733
Lower Kalskag	79	1	0	0	0	0	0	0
Kalskag (Upper)	62	0	0	0	0	0	0	0
Aniak	187	0	0	0	0	0	0	0
Chuathbaluk	33	0	0	0	0	0	0	0
Middle Kuskokwim	361	1	0	0	0	0	0	0
Crooked Creek	37	0	0	0	0	0	0	0
Red Devil	13	0	0	0	0	0	0	0
Sleetmute	40	0	0	0	0	0	0	0
Stony River	16	0	0	0	0	0	0	0
Lime Village	14	0	0	0	0	0	0	0
McGrath	136	0	0	0	0	0	0	0

-continued-

Table 5-6.–Page 2 of 2

Community	Households		Reported salmon					Total
	Total	Responding	Chinook	Sockeye	Coho	Chum	Pink	
Takotna ^a	23	0	—	—	—	—	—	—
Nikolai	34	0	0	0	0	0	0	0
Telida ^a	2	0	—	—	—	—	—	—
Upper Kuskokwim	315	0	0	0	0	0	0	0
Kuskokwim River	4,045	181	254	116	274	70	19	733
Quinhagak	162	28	16	29	11	5	6	67
Goodnews Bay	68	11	14	8	10	0	26	58
Platinum	19	4	5	5	27	1	5	43
South Kuskokwim Bay	249	43	35	42	48	6	37	168
Mekoryuk ^a	—	—	—	—	—	—	—	—
Newtok ^a	—	—	—	—	—	—	—	—
Nightmute ^a	—	—	—	—	—	—	—	—
Toksook Bay ^a	—	—	—	—	—	—	—	—
Tununak ^a	—	—	—	—	—	—	—	—
Chefornak ^a	—	—	—	—	—	—	—	—
Bering Sea Coast	—	—	—	—	—	—	—	—
Total	4,294	224	289	158	322	76	56	901

Source Shelden et al. (2014).

Note includes harvests using rod and reel and the removal of salmon from commercial harvests as well as subsistence nets.

Note — indicates data not available.

a. These communities were not contacted during the 2012 study period.

Table 5-7.—Subsistence nonsalmon fish harvests by community, Kuskokwim Area, 2012.

	Households		Reported salmon harvest												
	Total	Contacted	Humpback whitefish	Broad whitefish	Cisco	Sheefish	Burbot	Blackfish	Smelt	Northern pike	Herring	Grayling	Char	Rainbow trout	Total
Kipnuk	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
Kwigillingok	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
Kongiganak	90	0	--	--	--	--	--	--	--	--	--	--	--	--	--
North Kuskokwim Bay	90	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Tuntutuliak	90	49	1,443	1,448	60	76	741	8,260	175	1,294	0	28	75	4	13,604
Eek	86	45	506	260	255	53	412	3,491	0	944	0	69	11	7	6,008
Kasigluk	104	51	961	2,487	0	21	134	5,250	1,325	1,563	0	0	0	0	11,741
Nunapitchuk	111	58	4,298	3,930	34	94	333	13,930	1,000	3,257	0	0	1	1	26,878
Atmautluak	61	34	1,359	1,705	0	174	186	3,450	1,150	922	0	0	0	3	8,949
Napakiak	99	45	619	315	2	11	255	4,690	2,310	1,509	0	0	1	0	9,712
Napaskiak	97	41	404	299	64	68	117	1,539	7,375	930	0	0	2	0	10,798
Oscarville	14	13	260	161	20	22	84	2,310	1,025	199	0	0	0	0	4,081
Bethel	2,128	437	1,042	2,715	198	197	379	76,296	18,136	1,663	190	109	12	23	100,960
Kwethluk	164	80	741	645	52	91	237	410	5,975	685	0	113	159	53	9,161
Akiachak	157	68	719	458	130	288	424	17,010	6,975	820	0	0	2	7	26,833
Akiak	79	11	30	28	0	16	85	2,100	50	74	0	0	0	5	2,388
Tuluksak	89	52	930	375	41	115	311	7,350	6,675	471		18	8	6	16,300
Lower Kuskokwim	3,279	984	13,312	14,826	856	1,226	3,698	146,086	52,171	14,331	190	337	271	109	247,413
Lower Kalskag	79	39	466	134	289	37	104	590	1,895	217	0	3	3	6	3,744
Kalskag (Upper)	62	29	274	139	25	75	15	420	2,125	43	0	2	4	3	3,125
Aniak	187	152	473	6,474	12,333	187	61	103	2,825	83	0	74	121	77	22,811
Chuathbaluk	33	27	167	27	0	35	1,017	0	400	8	0	156	12	2	1,824
Middle Kuskokwim	361	247	1,380	6,774	12,647	334	1,197	1,113	7,245	351	0	235	140	88	31,504
Crooked Creek	37	31	74	88	1	136	6	0	0	12	0	109	26	0	452
Red Devil	13	10	56	50	80	34	12	0	0	91	0	222	76	1	622
Sleetmute	40	34	70	164	60	233	12	0	0	68	0	378	7	3	995
Stony River	16	3	0	0	0	0	0	0	0	0	0	0	0	0	0
Lime Village	14	10	5	55	50	0	2	0	0	44	0	100	0	0	256

-continued-

Table 5-7.-Page 2 of 2.

Table 3-17 Page 2 of 12

Community	Households		Reported salmon harvest													Total
	Total	Contacted	Humpback whitefish	Broad whitefish	Cisco	Sheefish	Burbot	Blackfish	Smelt	Northern pike	Herring	Grayling	Char	Rainbow trout		
McGrath	136	41	25	32	5	96	1	350	0	83	0	65	1	1	659	
Takotna	23	0	--	--	--	--	--	--	--	--	--	--	--	--	--	
Nikolai	34	28	15	637	453	50	1	0	0	147	0	56	6	0	1,365	
Telida	2	0	--	--	--	--	--	--	--	--	--	--	--	--	--	
Upper Kuskokwim	315	157	245	1,026	649	549	34	350	0	445	0	930	116	5	4,349	
Kuskokwim River	4,045	1,388	14,937	22,626	14,152	2,109	4,929	147,549	59,416	15,127	190	1,502	527	202	58,753	
Quinhagak	162	76	136	75	801	0	32	630	6,610	276	498	155	3,346	150	12,709	
Goodnews Bay	68	35	0	0	214	0	0	0	885	0	1,370	7	696	22	3,194	
Platinum	19	15	0	5	177	0	0	0	506	0	600	19	1,725	4	3,036	
South Kuskokwim Bay	249	126	136	80	1,192	0	32	630	8,001	276	2,468	181	5,767	176	18,939	
Mekoryuk	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	
Newtok	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	
Nightmute	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	
Toksook Bay	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	
Tununak	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	
Chefornak	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	
Bering Sea Coast	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	
Total	4,294	1,514	15,073	22,706	15,344	2,109	4,961	148,179	67,417	15,403	2,658	1,683	6,294	378	60,193	

Source Sheldon et al. (2014).

Note -- indicates data not available.




Figure 5-1.—Kuskokwim subsistence salmon harvest composition, 2012.

CHAPTER 6: BRISTOL BAY AREA

BACKGROUND

In spite of numerous social, economic, environmental, and technological changes, Bristol Bay residents continue to depend on salmon and other fish species as an important source of food. Subsistence harvests still provide important nutritional, economic, and sociocultural benefits to most Bristol Bay households. The 5 species of salmon found in Alaska are utilized for subsistence purposes in Bristol Bay, but the most popular are sockeye, Chinook, and coho salmon. Many residents continue to preserve large quantities of fish through traditional methods, such as drying and smoking, and fish are also frozen, canned, salted, pickled, fermented, and eaten fresh.

REGULATIONS

Permits are required to harvest salmon for subsistence purposes in Bristol Bay. Standard permit conditions include prohibition of fishing within 300 ft of a dam, fish ladder, weir, culvert, or other artificial obstruction. Since 1990, under state regulations, all Alaska state residents have been eligible to participate in subsistence salmon fishing in all Bristol Bay drainages. From 1998 through 2006, with 2 exceptions, only gillnets were recognized as legal subsistence gear. The first exception occurred in the Togiak District, where spear fishing was also allowed. Second, in 1998 the BOF adopted new regulations for the taking of “redfish” (postspawn sockeye salmon) in portions of the Naknek District. Gillnets, spears, and dip nets may be used along a 100 yd length of the west shore of Naknek Lake near the outlet to the Naknek River from August 20 through September 30, at Johnny’s Lake from August 15 through September 25, and at the mouth of the Brooks River from October 1 through November 15. In the Bristol Bay Area in 2010, gillnet lengths were limited to 10 fathoms in the Naknek, Egegik, and Ugashik rivers; Dillingham beaches; and within the Nushagak commercial district during emergency openings. Up to 25 fathoms could be used in the remaining areas, except that nets were limited to 5 fathoms in the special “redfish” harvest areas in the Naknek District.

At its regulatory meeting in Dillingham in December 2006, the BOF adopted 3 changes to subsistence salmon fishing regulations that affected portions of the Bristol Bay Area. The first change allowed salmon to be taken with drift gillnets no more than 10 fathoms in length in the lower 2 miles of the Togiak River.³¹ The second change allowed spears to be used to take salmon in Lake Clark, and the third change allowed the use of beach seines and gillnets to take salmon in Iliamna Lake, Six Mile Lake, and Lake Clark.

Along the Dillingham beaches, and in the Naknek, Egegik, and Ugashik rivers, subsistence fishing was limited to several fishing periods per week during the peak of the sockeye salmon run. All commercial districts were open for subsistence fishing during commercial openings. In addition, all commercial districts were open for subsistence fishing in May and October, from Monday to Friday. In the late 1990s and early 2000s, declining Chinook salmon and coho salmon stocks resulted in longer commercial closures, and some residents had difficulty obtaining fish for home uses. Since 2004, there have been improvements in abundance of all species (Jones et al. 2009:20). Since 1988, the Nushagak commercial district has been open to subsistence fishing by emergency order during extended commercial closures.

In May 2001, the National Park Service (NPS) announced that it would begin enforcing the prohibition of subsistence fishing with nets in Lake Clark National Park and Preserve, including all of Lake Clark, except by federally qualified area rural residents. This was a new enforcement action of an existing NPS regulation and was applied to individuals who were not permanent residents of Iliamna, Lime Village,

31. In addition to the Togiak River as described above 5 AAC 01.320 allows for the harvest of salmon by drift and set gillnet within commercial fishing districts.

Newhalen, Nondalton, Pedro Bay, or Port Alsworth, or who did not have a Section 13.44 subsistence use permit issued by the park superintendent.

ADF&G has continued to issue Bristol Bay subsistence salmon permits to those Alaska residents who request them. However, ADF&G informs permit applicants that unless they live in one of the above-named communities or have a Section 13.44 permit, they need to take this NPS closure into account when they subsistence fish in waters of the park and preserve. ADF&G also informs permittees that waters outside of national park and preserve boundaries remain open for subsistence salmon fishing to all permit holders.

INSEASON MANAGEMENT IN 2012

From June 1 through September 30 in all waters of a commercial salmon district within the Bristol Bay region, subsistence salmon could be taken only during commercial fishing periods. For a list of 2012 commercial fishing emergency orders for Bristol Bay in commercial districts, see Table 6 in Jones et al. (2014:30). In the Nushagak District, subsistence salmon fishing was provided for by emergency order during periods of extended commercial fishing closures.

SALMON HARVEST ASSESSMENT PROGRAM

A permit program was gradually introduced throughout the Bristol Bay region in the late 1960s to document the harvest of salmon for subsistence uses. Much of the increase in the number of permits issued during these years reflects: 1) a greater compliance with the permitting and reporting requirements, 2) an increased level of effort expended by ADF&G in making permits available (including issuance by area vendors), 3) contacting individuals to remind them to return the harvest forms, and 4) a growing regional population. Most fishers are obtaining permits and reporting their harvests, and overall permit returns have averaged between 85% and 90%. However, fish removed for home uses from commercial catches are not included in most reported subsistence harvest totals. Also, fish caught later in the season, such as coho salmon and spawning sockeye salmon, are probably not documented as consistently as Chinook and prespawn sockeye salmon.

In 2012, a total of 1,107 permits were issued for the Bristol Bay Management Area, of those 932, or 84%, were returned (Table 6-1; Table 6-2). The largest number of permits were issued for the Nushagak (517 permits) and Naknek–Kvichak (483 permits) districts (Table 6-1). The number of permits issued in 2012 was only slightly above the 5-year (1,102 permits), and the 10-year (1,101 permits) averages, but above the historical (1,093 permits) average (Table 6-2).

SUBSISTENCE SALMON HARVESTS IN 2012

Estimated total Bristol Bay subsistence salmon harvests in 2012 were 122,582 fish (Table 6-1). The 2012 salmon harvest was below the 5-year (125,206 salmon), 10-year (124,453 salmon), and historical (1983–2011 of 146,948 salmon) averages (Table 6-2).

Chinook salmon harvests were estimated at 12,136 in 2012, a decrease from the previous year's harvest of 14,106, and lower than the 2003 record harvest of 21,231 fish. Estimated sockeye salmon harvests for 2012 were 100,728, which was above the recent 5-year average of 98,709 fish and the 10-year average of 95,785 fish, but below the historical average (1983–2011) of 115,072 fish. Because the return of pink salmon to Bristol Bay are higher in even-numbered years than odd-numbered years, the number of pink salmon reported harvested was significantly higher in 2012 (1,874 fish) than in 2011 (333 fish). The estimated harvest of chum salmon in 2011 (4,007 fish) was lower than both the recent 5-year (4,648 fish) and 10-year averages (5,233 fish) and below the historical average (1983–2011) of 6,477 fish. The coho harvest in 2012 was much smaller harvest than the previous year (3,837 fish) and also lower than the 5-year average at 6,521 fish, the 10-year average at 6,724 (Table 6-2) and the historical 1983–2011 average at 8,320 fish.

In 2012, the Bristol Bay subsistence salmon harvest was composed of 82% sockeye salmon, 10% Chinook salmon, 3% coho salmon, 3% chum salmon, and 2% pink salmon (Figure 6-1). Of the entire Bristol Bay Area subsistence salmon harvest in 2012, residents of Bristol Bay communities harvested 113,320 salmon (92%), and other Alaska residents harvested 9,262 salmon (8%) (Table 6-3).

In 2012, as over the last several decades, most of the Bristol Bay Area subsistence harvest was taken in the Naknek–Kvichak (61%) and the Nushagak (31%) districts (Figure 6-1). The Naknek–Kvichak total harvest of 74,578 salmon in 2012 (Table 6-1) was higher than in 2011 (68,675) and in 2010 (64,445 salmon). Kvichak River drainage residents within the Kvichak River–Iliamna Lake Subdistrict and other permit holders fishing in the Kvichak drainage portion of the Naknek–Kvichak District harvested an estimated 178 Chinook salmon and 52,370 sockeye salmon in 2012, while those fishing in the Naknek River Subdistrict harvested 607 Chinook salmon and 20,338 sockeye salmon (Table 6-1). The 2012 subsistence harvest of 52,370 sockeye salmon in the Kvichak drainage (Table 6-1) was higher than the 2011 harvest of 45,226 sockeye, and the 2010 harvest of 40,688 sockeye (Fall et al. 2009a:69) and above historical levels (the most recent 5-year average harvest from 2007 through 2012 was 67,995 sockeye salmon) (Jones et al. 2014:93).³²

Subsistence sockeye salmon harvests in the Kvichak District have declined since the early 1990s (Salomone et al. 2011:113). From 1998 to 2011, estimated harvests were below the range of 55,000 to 65,000 sockeye salmon established by the BOF as the amount reasonably necessary for subsistence uses (5 AAC 01.336 (b)(1)). Poor sockeye salmon returns, like those seen in 2000–2002, are likely one factor responsible for declining harvests, but socioeconomic and sociocultural factors may be partly responsible as well (Fall et al. 2001, 2003, 2006; Fall, Turek, et al. 2009; Stickman et al. 2003).

In the Nushagak District, the total estimated subsistence harvest in 2012 of 37,960 salmon (Table 6-1) was a decrease from the previous year (45,226 salmon). The next lowest estimated harvests were 40,373 salmon in 2006 and 43,154 salmon in 2004 (Jones et al. 2013:91). The estimated harvest in 2008 of 51,395 salmon was the highest since 55,076 salmon in 2003 (Jones et al. 2013:91). The 2008 estimated harvest more accurately recorded harvest numbers for the season due to the administration of comprehensive baseline household subsistence harvest surveys by the Division of Subsistence in Aleknagik and Manokotak. For a more detailed description of these data see Fall et al. (2012b:75). The Nushagak District Chinook salmon harvest in 2012 was 10,350 (Table 6-1), and was a decrease from the year before (12,461 fish), but higher than in 2010, which was the lowest recorded harvest for the 20-year period from 1991 to 2010 (9,150 fish). The next lowest estimated harvests were 9,470 salmon in 2000 and 9,971 salmon in 2006 (Jones et al. 2013:99). The harvests in 2009 and 2008 (12,737 and 12,960 fish, respectively) were down from the 2003 estimate of 18,686 fish (the highest estimate on record), and below the 5-year (2007–2012) average of 12,128 fish, (Jones et al. 2014:94). The 2012 Nushagak District sockeye salmon harvest of 20,587 fish (Table 6-1) was lower than the 2011 harvest of 28,006 fish, and the 2010 estimate of 22,326 fish, and also the previous 5-year average (2007–2012) of 25,842 fish (Jones et al. 2014:94).

The estimated total subsistence salmon harvest for the Togiak District in 2012, 7,339 fish (Table 6-1), was higher than the previous year's estimate of 5,212 fish and higher than the previous 5-year average (5,756 salmon) (Jones et al. 2014:95). Estimated harvests in 2002 and from 2004 through 2007 were below those for 2001 and 2003; this likely reflects at least in part the result of postseason household surveys in Togiak and Twin Hills for 2001 and 2003. Postseason household surveys included more harvesters in the estimate because fishers who did not turn in their harvest permits were contacted. Comprehensive baseline household subsistence harvest surveys conducted in Togiak for the 2008

32. Note that the Kvichak River drainage sockeye salmon harvest total for 2007 listed in the 2008 Bristol Bay Area Annual Management Report (Jones et al. 2009:106) was adjusted down to the total of 47,473 by the Division of Subsistence. The updated data were included in the 2011 Annual Management Report (Jones et al. 2012:100). Additional harvest reports collected through subsistence salmon household harvest surveys in 4 Kvichak drainage communities increased the accuracy of the reported harvests and reduced the expansion factor for the estimated subsistence salmon harvest in the Kvichak drainage in 2007.

calendar year also showed an increase in the participation in the 2008 harvest assessment program. The estimated subsistence salmon harvest in the Ugashik District in 2012 was 1,281 fish, which was up by almost twice from the previous year at 687 fish, the lowest count in recorded history (Table 6-1). The 2012 harvest was lower than the 10-year average (2002–2012) of 2,000 fish (Jones et al. 2014:94). In the Egegik District, the estimated subsistence salmon harvest of 1,425 fish (Table 6-1) was much lower than the 2011 estimate of 2,265 fish; however, the 2012 estimate was notably lower than the 4,711 fish estimated for 2004 (the second highest estimate since 1984), and was less than the previous 5-year average of 1,732 salmon (Jones et al. 2014:93).

OTHER SUBSISTENCE FISHERIES

In May 2003, new federal regulations authorizing subsistence fishing for Pacific halibut came into effect. A harvest assessment program for the subsistence halibut fishery was implemented in 2004. Subsistence halibut harvest estimates for 2012 appear in Fall and Koster (2014). Beginning in 2003 subsistence fishing for rainbow/steelhead trout *O. mykiss* and Arctic char/Dolly Varden in the Bristol Bay Area under federal subsistence regulations required a federal permit. No permits were issued (Michael Edwards, Fisheries Biologist, USFWS, King Salmon Field Office, personal communication, 2004). The permit requirement was dropped in 2005. The following overview derives primarily from Fall and Chythlook (1997) and Fall et al. (2009).

Subsistence Regulations

The BOF determined that all finfishes of the Bristol Bay Management Area support customary and traditional uses (5 AAC 01.336). In addition, the BOF determined that approximately 250,000 lb usable weight (about 41 lb per person) was the amount reasonably necessary to provide for these uses. This amount was based upon estimates of fish harvests derived from systematic household surveys conducted by the Division of Subsistence. Amounts for specific species or more specific stocks were not established.

For the most part, subsistence fishing for species other than salmon and rainbow/steelhead trout is open year-round in the Bristol Bay Area with gear listed in 5 AAC 01.010 (a). There are no seasonal limits established by regulation. The following regulations apply to subsistence fishing for species other than salmon in the area:³³

- Rainbow/steelhead trout taken incidentally in other subsistence net fisheries and through the ice are lawfully taken and may be retained for subsistence uses (5 AAC 01.310 (g)).
- Subsistence fishing with a line attached to a rod or pole is prohibited except when fishing through the ice (5 AAC 01.320 (l)).
- Subsistence fishing with nets is prohibited in 18 waters of the Kvichak–Iliamna Lake drainage and within one-quarter mile of the terminus of those waters from September 1 through June 14.

Subsistence Harvests and Uses

A detailed description of subsistence uses of freshwater fishes in the Bristol Bay Area appears in Fall et al. (1996) and Holen and Lemons (2012). Holen et al. (2012) and Wright and Chythlook (1985) describe the uses of herring spawn on kelp in the Togiak District. Harvests of fishes other than salmon contribute about 10% of the annual subsistence harvests of wild foods in the Bristol Bay region, about 41 lb per person (Fall, Krieg, et al. 2009; Holen and Lemons 2012).

Subsistence harvests of fishes other than salmon are not annually monitored by ADF&G. Harvest and use data are available for most communities through Division of Subsistence household harvest surveys (BBNA and ADF&G 1996; Coiley-Kenner et al. 2003; Evans et al. 2013; Fall et al. 2006; Holen et al.

33. In 2004, the BOF eliminated a permit requirement for subsistence fishing for rainbow/steelhead trout and Arctic char/Dolly Varden in the Bristol Bay Area. ADF&G had not developed a program for issuing these permits.

2011; Holen, Stariwat, et al. 2012; Krieg et al. 2005, 2009). As part of an OSM project (02-034, Subsistence Fisheries Assessment: Kvichak River Watershed Resident Species), the Division of Subsistence and the Bristol Bay Native Association collected subsistence harvest data in Kvichak River watershed communities from October 2002 to September 2003. The final report for that project (Krieg et al. 2005) includes detailed information about uses of nonsalmon fishes in 8 study communities. Some of the findings of ADF&G research regarding nonsalmon fishes are summarized in Table 6-4. The vast majority of households in the Bristol Bay Area use fishes other than salmon for subsistence purposes. Most households also participate in the harvest of these fishes. Harvests, as measured in pounds usable weight per person for available study years, vary from community to community and from year to year, but generally make important contributions to annual subsistence harvests. Fishes other than salmon generally rank third behind salmon and land mammals in their contribution to the total subsistence harvests in Bristol Bay communities.

Harvests and uses of the nonsalmon fishes listed in Table 6-5 have been documented in Bristol Bay communities through Division of Subsistence research. Uses of other species may occur: fish taken in the largest quantities in the area as a whole include smelt, whitefishes, Dolly Varden, Arctic grayling, and northern pike (see Fall et al. [1996] for a summary of harvest data).

In the Bristol Bay Area, harvests of nonsalmon finfishes occur throughout the year. Harvest effort for these fish is generally lower among Bristol Bay residents in the summer as attention is focused on salmon. Spring is important for herring, herring spawn on kelp, and rainbow smelt. Harvests of nonsalmon fishes occur through the ice in winter. “Smelting” is a popular activity in October and in late winter when these fish can be caught by jigging through the ice. Halibut are mostly taken in June and July (Wright et al. 1985:34).

Many gear types are used to harvest nonsalmon fishes for home uses in the Bristol Bay Area. Rod and reel is used for most fish; some, such as Arctic char/Dolly Varden and herring and other marine fishes, are removed from commercial catches. Other methods are used, including (but not necessarily limited to) the following:

- Traps (fyke nets): Alaska blackfish, burbot;
- Set hooks: burbot;
- Handline jigging through the ice: Arctic grayling, Arctic char/Dolly Varden, lake trout, rainbow smelt, rainbow/steelhead trout, whitefishes, northern pike;
- Set gillnets: Arctic grayling, Arctic char/Dolly Varden, lake trout, longnose suckers, rainbow/steelhead trout, herring, northern pike, burbot, whitefishes;
- Beach seining: Arctic char/Dolly Varden, lake trout, rainbow smelt, herring, whitefishes;
- Hand line in open water: Pacific halibut, rainbow/steelhead trout; and
- Dip nets: rainbow smelt, herring.

Herring spawn on kelp is usually picked by hand, although rakes, knives, and *uluqaqs* (women’s knives) are also used (Schichnes and Chythlook 1988:127).

Maps of areas used by Bristol Bay communities to harvest nonsalmon fishes appear in the *Alaska Habitat Management Guide Reference Atlas* series (Alaska Department of Fish and Game 1985), and in Wright et al. (1985). Updated maps of harvest locations for 8 communities in the Kvichak watershed appear in Krieg et al. (2005). Harvest activities occur throughout the region in most rivers and lakes as well as along shorelines. It is likely that most effort occurs near each community and near seasonal camps such as Kulukak. See Wright and Chythlook (1985) and Schichnes and Chythlook (1988) for maps of herring camps at Kulukak Bay. For frequency of uses of various areas for freshwater fishing by Nushagak River

communities, see Schichnes and Chythlook (1991) and by Togiak and Manokotak, see BBNA and ADF&G (BBNA and ADF&G 1996).

Bristol Bay residents use a wide variety of methods to process and preserve their harvests of fishes other than salmon. These vary by species and community. Some freezing of harvests of most species occurs. Some examples of other methods include the following:

- Arctic grayling: dried, half-dried, fresh frozen, aged frozen and eaten with seal oil (various species);
- Dolly Varden: dried, smoked, half dried (*egamaarrluk*);
- Northern pike: dried, half-dried, fresh frozen, aged frozen and eaten with seal oil;
- Rainbow/steelhead trout: dried, half dried, smoked; and
- Whitefishes: dried, fresh frozen, aged frozen and eaten with seal oil.

Dried fish product is eaten with seal oil. Fat from brown bears *Ursus arctos* mixed with dry fish is also consumed. Rainbow smelt are fried, boiled, dried, or eaten frozen with seal oil (Fall et al. 1986:100; Fall, Krieg, et al. 2009). Herring are salted, or split, dried, and smoked (Schichnes and Chythlook 1988:126). The heads and stomachs of northern pike are boiled and eaten (Schichnes and Chythlook 1991:139). Freshwater fishes that are usually eaten frozen with seal oil form a category called *kumlaneq*. This includes Arctic grayling, whitefishes, lake trout, and northern pike (Fall et al. 1986:102; Fall, Krieg, et al. 2009).

There is much traditional knowledge of the subsistence uses of nonsalmon fishes in the Bristol Bay Area. For example, a Yup'ik taxonomic classification system for freshwater fish species has 3 entries, and thus 3 taxons, for the fish that Western science classifies in only 1: Dolly Varden. The Yup'ik distinctions are made depending upon the condition of the flesh for aging, freezing, and/or drying; harvest locations; and harvest methods (Fall et al. 1996; Fall, Krieg, et al. 2009).

The Division of Subsistence has compiled a traditional ecological knowledge (TEK) database, "From *Neqa* to *Tepa*," about the fishes of Bristol Bay based on interviews with area residents in 2003 as part of OSM project 01-109.³⁴ An expanded version of the database incorporating findings from 8 Kvichak watershed communities was renamed "From *Neqa* to *Tepa*, *Luq'a* to *Chuqilin*" to reflect the addition of Dena'ina Athabascan TEK (BBNA and ADF&G 1996; Krieg et al. 2005).

34. Coiley-Kenner, P. 2003. From Neqa to Tepa: a database with traditional knowledge about the fish of Bristol Bay and the northern Alaska Peninsula. Version 2.0. Alaska Department of Fish and Game Division of Subsistence, Juneau.

Table 6-1.—Estimated subsistence salmon harvests by district and location fished, Bristol Bay Area, 2012.

Area and river system	Number of permits issued ^a	Estimated salmon harvest					
		Chinook	Sockeye	Coho	Chum	Pink	Total
Naknek-Kvichak District	483	785	72,708	485	127	474	74,578
Naknek River Subdistrict	280	607	20,338	396	104	384	21,828
Kvichak River/Iliamna Lake Subdistrict:	207	178	52,370	89	23	90	52,750
Igiugig	2	0	555	0	0	0	555
Iliamna Lake-General	37	0	6,655	0	0	0	6,655
Kokhanok	26	161	15,148	0	0	1	15,310
Kvichak River	21	0	3,774	0	0	0	3,774
Lake Clark	55	0	4,610	0	0	0	4,610
Levelock	3	17	845	89	23	89	1,063
Newhalen River	46	0	13,829	0	0	0	13,829
Pedro Bay	17	0	4,059	0	0	0	4,059
Six Mile Lake	13	0	2,895	0	0	0	2,895
Egegik District	38	37	1,172	190	19	7	1,425
Ugashik District	20	31	997	228	25	0	1,280
Nushagak District	517	10,350	20,587	2,642	3,072	1,309	37,960
Igushik/Snake River	12	143	937	105	20	7	1,212
Nushagak Bay Commercial	42	368	1,238	291	176	196	2,269
Nushagak Bay Noncommercial	204	2,685	7,387	1,011	796	410	12,289
Nushagak River	119	4,896	4,448	808	1,559	426	12,136
Site Unknown	1	0	80	0	0	0	80
Wood River	156	2,259	6,497	427	522	270	9,974
Togiak District	53	933	5,265	293	764	84	7,339
Total	1,107	12,136	100,728	3,837	4,007	1,874	122,582

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Note Harvests are extrapolated for all permits issued, based on those returned and on the area fished as recorded on the permit. Due to rounding, the sum of columns and rows may not equal the estimated total. Of 1,107 permits issued for the management area, 932 were returned (84.2%).

- a. Sum of sites may exceed district totals, and sum of districts may exceed area total, because permittees may use more than one site.

Table 6-2.—Estimated historical subsistence salmon harvests, Bristol Bay Area, 1983–2012.

Year	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1983	829	674	13,268	143,639	7,477	11,646	1,073	177,104
1984	882	698	11,537	168,803	16,035	13,009	8,228	217,612
1985	1,015	808	9,737	142,755	8,122	5,776	825	167,215
1986	930	723	14,893	129,487	11,005	11,268	7,458	174,112
1987	996	866	14,424	135,782	8,854	8,161	673	167,894
1988	938	835	11,848	125,556	7,333	9,575	7,341	161,652
1989	955	831	9,678	125,243	12,069	7,283	801	155,074
1990	1,042	870	13,462	128,343	8,389	9,224	4,455	163,874
1991	1,194	1,045	15,245	137,837	14,024	6,574	572	174,251
1992	1,203	1,028	16,425	133,605	10,722	10,661	5,325	176,739
1993	1,206	1,005	20,527	134,050	8,915	6,539	1,051	171,082
1994	1,193	1,019	18,873	120,782	9,279	6,144	2,708	157,787
1995	1,119	990	15,921	107,717	7,423	4,566	691	136,319
1996	1,110	928	18,072	107,737	7,519	5,813	2,434	141,575
1997	1,166	1,051	19,074	118,250	6,196	2,962	674	147,156
1998	1,234	1,155	15,621	113,289	8,126	3,869	2,424	143,330
1999	1,219	1,157	13,009	122,281	6,143	3,653	420	145,506
2000	1,219	1,109	11,547	92,050	7,991	4,637	2,599	118,824
2001	1,226	1,137	14,412	92,041	8,406	4,158	839	119,856
2002	1,093	994	12,936	81,088	6,565	6,658	2,341	109,587
2003	1,182	1,058	21,231	95,690	7,816	5,868	1,062	131,667
2004	1,100	940	18,012	93,819	6,667	5,141	3,225	126,865
2005	1,076	979	15,212	98,511	7,889	6,102	1,098	128,812
2006	1,050	904	12,617	95,201	5,697	5,321	2,726	121,564
2007	1,063	917	15,444	99,549	4,880	3,991	815	124,679
2008	1,178	1,083	15,153	103,583	7,627	5,710	2,851	134,924
2009	1,063	950	14,020	98,951	7,982	5,052	442	126,447
2010	1,082	979	10,852	90,444	4,623	4,692	2,627	113,238
2011	1,122	1,039	14,106	101,017	7,493	3,794	333	126,744
2012	1,107	932	12,136	100,728	3,837	4,007	1,874	122,582
5-year average (2007–2011)	1,102	994	13,915	98,709	6,521	4,648	1,414	125,206
10-year average (2002–2011)	1,101	984	14,958	95,785	6,724	5,233	1,752	124,453
Historical average (1983–2011)	1,093	958	14,730	115,072	8,320	6,477	2,349	146,948

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Table 6-3.—Estimated subsistence salmon harvests by community, Bristol Bay Area, 2012.

Community	Permits		Estimated salmon harvest					Total
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	
Aleknagik	29	21	696	1,548	108	86	19	2,457
Clarks Point	13	13	99	365	189	80	149	882
Dillingham	328	277	5,055	12,921	1,420	1,331	651	21,378
Egegik	9	6	0	66	104	0	0	170
Ekwok	15	13	681	167	59	234	112	1,253
Igiugig	13	9	0	2,711	0	0	0	2,711
Iliamna	29	23	3	8,194	0	0	0	8,197
King Cove	1	1	2	24	6	2	4	38
King Salmon	81	74	173	5,329	49	17	100	5,667
Kokhanok	27	20	161	16,593	0	0	1	16,755
Koliganek	15	13	852	835	361	579	207	2,834
Levelock	3	2	0	825	0	0	0	825
Manokotak	12	9	143	937	105	20	7	1,212
Naknek	106	84	273	10,318	227	49	207	11,074
New Stuyahok	39	26	2,439	1,778	345	677	137	5,375
Newhalen	14	11	0	5,064	0	0	0	5,064
Nondalton	31	30	0	9,327	0	0	0	9,327
Pedro Bay	15	14	0	4,028	0	0	0	4,028
Pilot Point	6	5	18	307	60	24	0	409
Port Alsworth	52	49	2	4,445	0	0	0	4,447
Portage Creek	1	1	31	2	0	2	0	35
South Naknek	18	15	20	778	79	11	54	942
Togiak	53	38	951	5,364	298	779	85	7,478
Twin Hills	1	0	0	0	0	0	0	0
Ugashik	9	8	7	588	168	1	0	764
Subtotal, Bristol Bay	920	762	11,604	92,514	3,577	3,891	1,733	113,320
Anchorage	92	80	248	3,618	54	15	68	4,002
Anderson	1	0	0	0	0	0	0	0
Aniak	1	1	0	9	0	0	0	9
Barrow	2	2	64	42	0	5	0	111
Bethel	1	1	0	7	0	0	0	7
Big Lake	1	1	0	32	0	0	0	32
Chugiak	5	5	14	163	0	3	0	180
Copper Center	1	1	0	0	0	0	0	0
Cordova	2	2	0	86	0	0	0	86
Eagle River	4	4	11	238	0	19	1	269
Fairbanks	12	11	8	298	97	4	20	427
Girdwood	3	3	0	15	19	2	8	44
Homer	13	12	23	924	17	17	14	995
Kasilof	1	1	7	175	0	14	0	196
Kenai	3	3	7	146	8	0	0	161
Kipnuk	1	1	5	25	0	0	0	30
Kodiak City	9	9	55	247	1	24	0	327

-continued-

Table 6-3.-2 of 2

Community	Permits		Estimated salmon harvest					Total
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	
Kotzebue	1	1	0	8	10	0	4	22
McCarthy	1	1	0	50	0	2	0	52
Nikiski	2	2	2	74	47	1	2	126
Palmer	9	8	7	734	0	0	0	740
Sitka	1	0	0	0	0	0	0	0
Soldotna	2	2	31	55	0	0	16	102
Talkeetna	2	2	17	29	0	2	0	48
Tok	1	1	0	16	0	3	0	19
Trapper Creek	1	1	1	71	0	0	0	72
Wasilla	14	14	32	1,153	7	5	9	1,206
Willow	1	1	0	0	0	0	0	0
Subtotal, other Alaska	187	170	531	8,214	260	116	142	9,262
Total	1,107	932	12,136	100,728	3,837	4,007	1,874	122,582

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Table 6-4.—Uses and harvests of fish other than salmon, Bristol Bay communities.

Community	Year ^a	Percentage of households ^a					Average pounds harvested	
		Use	Fish for	Harvest	Receive	Give	Per household	Per person
Aleknagik	2008	78	69	66	50	44	95	26
Clark's Point	2008	100	100	100	73	73	71	34
Dillingham	1984	75	56	55	40	20	52	17
Egegik	1984	64	60	60	24	40	37	16
Ekwok	1987	76	72	62	62	38	229	69
Igiugig	2005	100	83	83	92	58	188	59
Iliamna	2004	92	77	77	39	31	113	34
King Salmon	2007	57	55	49	16	12	15	5
Kokhanok	2005	74	66	66	51	57	137	36
Koliganek	2005	96	93	93	75	68	323	90
Levelock	2005	86	86	86	50	57	71	40
Manokotak	2008	93	80	80	84	56	173	44
Naknek	2007	76	68	65	48	32	47	18
New Stuyahok	2005	88	78	78	67	47	123	28
Newhalen	2004	88	88	88	56	52	128	32
Nondalton	2004	82	76	76	45	53	129	34
Pedro Bay	2004	89	61	61	83	39	50	15
Pilot Point	1987	94	94	94	35	59	56	16
Port Alsworth	2004	73	64	64	46	41	44	12
Port Heiden	1987	92	62	62	70	46	33	12
South Naknek	2007	86	52	52	67	43	16	8
Togiak	2008	94	85	84	81	73	264	62
Twin Hills	1999	92	92	92	75	92	303	101
Ugashik	1987	1	100	100	0	40	72	36

Sources CSIS; BBNA and ADF&G 1996; Coiley-Kenner (2003); Krieg et al. (2005); Fall et al. (2006); Krieg et al. (2009); Holen et al. (2011); Holen et al. (2012); and Fall et al. (2013)

a. Most recent year for which data are available.

Table 6-5.—Nonsalmon finfish used for subsistence purposes in the Bristol Bay Area.

Common English name	Scientific name	Yup'ik name	Dena'ina name
Arctic grayling	<i>Thymallus arcticus</i>	<i>Nakrullugpak</i> <i>Culugpak</i>	<i>Ch'dat'an</i>
Alaska blackfish	<i>Dallia pectoralis</i>	<i>Can'giiq</i>	<i>Huzhegh</i>
Burbot	<i>Lota lota</i>	<i>Manignaq^a</i> <i>Atgiaq^b</i>	<i>Ch'unya</i>
Dolly Varden ^c	<i>Salvelinus malma</i>	<i>Yugyaq^d</i> <i>Anerrluaq</i> <i>Anyuk</i>	<i>Qak'elay</i>
Lake trout	<i>Salvelinus namaycush</i>	<i>Cikignaq</i>	<i>Zhuk'udghuzha</i>
Longnose sucker	<i>Catostomus catostomus</i>	<i>Cungartak</i>	<i>Duch'ehdi</i>
Northern pike	<i>Esox lucius</i>	<i>Cuukvak</i>	<i>Ghelguts'i</i>
Rainbow smelt	<i>Osmerus mordax</i>	<i>Iqalluaq</i>	
Rainbow/steelhead trout	<i>Oncorhynchus mykiss</i>	<i>Talaariq</i>	<i>Tuni</i>
Broad whitefish ^e	<i>Coregonus nasus</i>	<i>Akakiik</i>	<i>Telay</i>
Humpback whitefish ^e	<i>Coregonus pidschian</i>	<i>Uraruq</i>	<i>Q'untuq'</i>
Round whitefish ^e	<i>Prosopium cylindraceum</i>	<i>Uraruq</i>	<i>Hesten</i>
Least cisco	<i>Coregonus sardinella</i>	<i>Cavirrutnaq</i>	<i>Ghelguts'i k'una</i>
Pacific herring	<i>Clupea harengus pallasi</i>	<i>Iqalluarpak</i>	
Pacific herring spawn on kelp		<i>Melucuaq</i>	
Starry flounder	<i>Platichthys stellatus</i>	<i>Naterna</i>	
Pacific halibut	<i>Hippoglossus stenolepis</i>	<i>Naternarpak</i>	
Pacific cod	<i>Gadus macrocephalus</i>	<i>Ceturtnaq</i>	
Sculpin	Various species	<i>Kayutaq</i>	
Capelin	<i>Mallotus villosus</i>	<i>Cikaaq</i>	
Yellowfin sole	<i>Limanda aspera</i>	<i>Sagiq</i>	

Source Fall et al. (1996).

- a. Nushagak River villages.
- b. Manokotak, Aleknagik, Twin Hills, Togiak.
- c. Also includes the closely related Arctic char.
- d. At Togiak, Manokotak, and Aleknagik, and perhaps elsewhere, there are three Yup'ik names for Arctic char/Dolly Varden. Yugyak probably refers to resident char/Dolly Varden. Anerrluaq, called "Togiak trout" in the local English dialect, probably refers to anadromous fish taken in fresh water. Finally, anyuk or "sea run dollies" are Dolly Varden or Arctic char taken in salt water. See Fall et al. (1996:16–20) for further discussion of these distinctions.
- e. Broad whitefish are rare to absent in the Bristol Bay region. Akakiik is the word used at Aleknagik and Manokotak to refer to whitefish they receive from Kuskokwim River communities, where broad whitefish are common. Humpback whitefish are caught in the Iliamna Lake subregion and called uraruq. Uraruq is also used for round whitefish in the Togiak and Nushagak drainages.




Figure 6-1.—Bristol Bay Area subsistence salmon harvest composition, 2012.




Figure 6-2.—Bristol Bay Area subsistence salmon harvests by district, 2012.

CHAPTER 7: CHIGNIK MANAGEMENT AREA

BACKGROUND

The Chignik Management Area (CMA) encompasses all coastal waters and inland drainages on the south side of the Alaska Peninsula from Kilokak Rocks at the southern entrance to Imuya Bay at 57 degrees 10.34' N. lat., 156 degrees 20.22' W. long., then due south to Kupreanof Point at 55 degrees 33.98' N. lat. 159 degrees 35.88' W. long. (5AAC 15.100).³⁵ There are 5 communities in Alaska Department of Fish and Game's (ADF&G) salmon Chignik Management Area (CMA): Chignik with a 2012 estimated population of 90, Chignik Lagoon (population 82), Chignik Lake (population 70), Perryville (population 112), and Ivanof Bay (population 7) (Figure 7-1).³⁶ All of these communities are within the Lake and Peninsula Borough, and virtually all area residents participate in harvesting salmon in the CMA. Published Division of Subsistence reports for the CMA include annual salmon permit harvest reports, sporadic household surveys, and subsistence salmon ethnography studies.

The Division of Subsistence household harvest surveys show that salmon compose approximately 45% of all resources harvested, by weight, for subsistence in these communities (Fall et al. 1995). Chignik subsistence salmon permits are issued annually by CMA vendors, with harvest reports due to the department by December 31. The 2012 estimated total subsistence salmon harvest was 8,242 salmon; 68% sockeye salmon, 18% coho salmon, 10% pink salmon, 3% chum salmon, and 1% Chinook salmon.

In 1993, the BOF made a positive determination that salmon in the CMA are customarily and traditionally taken or used for subsistence (a "positive C&T finding") and specified amounts of salmon are reasonably necessary for subsistence (ANS) in each CMA district. In 2002, the BOF modified the original finding for ANS (5 AAC 01.466 (a) and (b)) (ADF&G 1994). The current amounts necessary for subsistence for Chignik Bay, Central, and Eastern districts combined are 5,200–9,600 early-run sockeye salmon; 2,000–3,800 late-run sockeye salmon; 100–150 Chinook salmon; and 400–700 salmon other than sockeye or Chinook salmon. In the Perryville and Western Districts combined are 1,400–2,600 coho and 1,400–2,600 salmon, other than coho salmon. The BOF has also set an ANS for rainbow/steelhead trout *O. mykiss* at 200–300 pounds and for finfish other than those listed above at 15,200–22,800 pounds of usable weight.

REGULATIONS

Current (2013–2014) State of Alaska regulations governing subsistence salmon fishing in the Chignik Management Area require that to fish, an individual must obtain an annual subsistence salmon permit, and must be an Alaska resident (5 AAC 01.480).³⁷ Annually, permits are available locally at the Chignik ADF&G weir facility and from local CMA community vendors. The permit holder must record daily salmon harvests directly on the permit and return it to the Alaska Department of Fish and Game by December 31. Catch information obtained from subsistence permits is compiled annually and used to assess regional subsistence salmon fisheries. There is an annual limit of 250 salmon per permit. (5 AAC 01.480(b)(c)).

Under state regulations, all waters within the CMA are open year-around for subsistence fishing except for the following areas or conditions:

- Only Alaska residents are eligible to obtain a CMA subsistence salmon permit and may fish in the areas open to subsistence at any time. A commercial Chignik Area

35. Alaska Department of Fish and Game. 2011–2014 Chignik and Kodiak Areas, commercial salmon fishing regulations, 89. Alaska Department of Fish and Game, Juneau. <http://www.adfg.alaska.gov/static/regulations/fishregulations/pdfs/commercial/ChigKod-2011-14.pdf>

36. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. "Research and Analysis Homepage: Population Estimates." Accessed July 2014. <http://laborstats.alaska.gov/pop/popest.htm>

37. Alaska Department of Fish and Game. 2013-2014 Subsistence and personal use statewide fisheries regulations, 37. Alaska Department of Fish and Game, Juneau. <http://www.adfg.alaska.gov/static/regulations/fishregulations/pdfs/commercial/Subsistence-PU-2013-2014.pdf>

salmon fishing license holder (includes CFEC Permit and crewmember license) may subsistence fish during a commercial salmon fishing period, except for 12 hours before a commercial salmon fishing period and 12 hours after a commercial salmon fishing period (5 AAC 01.485).

- Legal gear includes seines and gillnets. Purse seines may not be used in Chignik Lake (5 AAC 01.470(a)). Additionally, any gillnet that is fixed, anchored, or otherwise held in place may not obstruct more than one half of the width of any stream open to subsistence fishing. All subsistence salmon fishing gear must be marked with a buoy listing the first initial and last name as well as the address of the person operating the gear (5 AAC 01.010(h)). Subsistence users must carry their subsistence fishing permit with them while fishing.
- Subsistence salmon fishing is permitted in the Chignik River; however, salmon may not be taken upstream from the ADF&G weir to the outlet of Chignik Lake from July 1-August 31 (5 AAC 01.475(1)), which is closed to protect the spawning Chinook salmon. The Chignik River, beginning 100 yards below the weir, is open to subsistence salmon fishing year round.
- Subsistence fishing is closed within 100 yards above or below the Chignik weir when it is operational (5 AAC 01.470).
- Subsistence fishing is closed year-round in Black Lake or any tributary to Black Lake or Chignik Lake, except the waters of Clark River and Home Creek from each of their confluences with Chignik Lake to a point 1 mile upstream (5 AAC 01.475(2)). The Alaska Board of Fisheries amended the subsistence regulations in 2008 to include these tributaries for the purposes of providing additional harvest opportunities for subsistence users.

RECENT REGULATORY HISTORY

Subsistence harvest patterns in the CMA are often influenced by Chignik commercial salmon fisheries since many of those who commercial fish are also subsistence harvesters. Regulations for subsistence salmon fishing are tied to Chignik commercial fishing operations. Prior to 2002, the CMA commercial fishery was managed by ADF&G as a competitive limited entry permit fishery. Pre-2002 regulations allowed subsistence fishing with seine and gillnet gear and required an individual permit with a seasonal limit of 250 salmon. Purse seines could be used to harvest subsistence salmon, except in Chignik Lake, which has been open by regulation to subsistence salmon fishing with gillnets or hand seines since 1985 (Morris 1987:185). Also prior to 2002, CMA commercial salmon harvesters could not subsistence fish between June 10 and September 30, although they were allowed to remove salmon caught during commercial openings for home use. Subsistence salmon fishing was not allowed in Chignik River upstream of the ADF&G weir site to Chignik Lake, in tributaries to Chignik Lake, or in Black Lake.

From 2002 to 2005, the CMA commercial salmon fishery was managed based on 2 management plans: the Chignik Area Management Plan ("competitive fishery") and the Chignik Area Cooperative Purse Seine Salmon Management Plan ("cooperative fishery"; Stichert 2007a). After development of the cooperative fishery, ADF&G management staff initiated subsistence permit conditions in 2003 that increased subsistence harvest opportunities for commercial fishing license holders.³⁸ By regulation, commercial fishing permit holders could not subsistence fish for salmon from 48 hours before the first commercial salmon fishing opening through September 30. Subsistence fishing permit conditions allowed commercial permit holders who were not engaged in commercial fishing during an opening for

38. Regulations providing for a cooperative commercial salmon fishery in CMA were invalidated by a decision of the Alaska Supreme Court and have not been operative since 2005.

cooperative or competitive fleets to subsistence fish during commercial openings, after registering with ADF&G.

In 2004, through emergency order, ADF&G allowed subsistence salmon fishing within the Chignik River, excluding the waters 100 yards upstream and downstream of the Chignik weir, through June 30. Regulations had closed the Chignik River to subsistence salmon fishing (5 AAC 01.475) until 2006. In addition to obtaining a subsistence permit, commercial harvesters wishing to subsistence fish after the first commercial opening were allowed to do so, with a requirement to register with ADF&G staff working at the weir. ADF&G established a subsistence fishing schedule for these commercial harvesters depending upon whether they fished for the cooperative fleet or independently (Bouwens 2004).

At its 2004 meeting, the BOF adopted regulations to increase subsistence fishing opportunities for commercial salmon fishing license holders by allowing them, with certain restrictions (5 AAC 01.485), to harvest subsistence salmon during the commercial salmon fishing season. In addition, the BOF directed ADF&G to manage for an increase in escapement of sockeye salmon during the August commercial fishery (from 50,000 to 75,000), in order to enhance late-season subsistence opportunities in Chignik Lake. Although the commercial fishery was limited in August, the sockeye escapement goal was not achieved in 2005 (Bouwens 2005). In 2005, the BOF opened the Chignik River drainage to subsistence fishing, except for waters within 300 feet of the weir, and except for a July 1 through August 31 closure upstream of the weir to protect spawning Chinook salmon (Stichert 2007b).

The cooperative fishery plan was repealed by the Alaska Supreme Court in March 2005, but the BOF reestablished the cooperative management plan by emergency regulation because there was not enough time between the ruling and the 2005 fishing season for co-op members to revert back to fishing as independent fishers. Since 2006, however, the CMA commercial fishery has been managed solely under the *Chignik Salmon Management Plan* as a competitive fishery (Stichert 2007b).

During its January 2008 meeting, the BOF adopted regulatory changes to subsistence fishing in the CMA that allowed subsistence salmon fishing in Clark River and Home Creek from their confluences with Chignik Lake upstream 1 mile. The use of gillnets for subsistence fishing in the CMA remained legal, but when they are fixed, anchored or otherwise held in place, they may not obstruct more than one-half of the stream that is open to subsistence salmon fishing (Jackson 2009).

The Village Council of Chignik Lake submitted a regulatory proposal at the CMA BOF meeting in January 2011. The proposal, if adopted, would have legalized subsistence fishing in the only areas in the CMA closed under state regulations—Chignik Lake, Black Lake, and all tributaries to both lakes—as well as legalized the use of hook and line gear for late-run spawned-out sockeye salmon in Clark River and Home Creek (Alaska Board of Fisheries 2011a). The BOF took no action on the proposal; however the Federal Subsistence Board (FSB) adopted a similar regulation for the CMA at its January 2011 meeting (Alaska Board of Fisheries 2011b; 76 FR 45:12566, 12578–12579 [March 8, 2011]; 36 CFR 242.27 (e) (8)).

At the FSB regulatory meeting January 24, 2013, the FSB adopted a proposal submitted by the Chignik Lake Traditional Council to allow the harvest of salmon in the Chignik River, with rod and reel, from a point 300 feet upstream of the AF&G weir to Chignik Lake from January 1 through August 9, with no daily harvest or possession limit under the authority of a federal subsistence fishing permit. In addition the FSB adopted a regulation allowing the taking of salmon by gillnet in Black Lake or any tributary to Black or Chignik lakes. The BOF closed this portion of the river in 2004 to protect spawning Chinook salmon, and it remains closed for subsistence fishing July 1–August 31, but open to sport fishing, under state regulation.^{39,40}

39. Federal Subsistence Board. Public Regulatory Meeting Proceedings, Volume III. January 24, 2013, Anchorage. Accessed July 2014. <http://www.doi.gov/subsistence/library/transcripts/upload/FSB-Mtg-24-Jan-13-2.pdf>

The State Board of Fisheries at their Chignik Finfish December 2013 meeting adopted a board-generated proposal to codify a management measure first established through board intent language adopted in 2004. The management measure is intended to ensure in river harvest opportunities above the Chignik River weir to satisfy late run subsistence harvests. To do so, the department shall manage for 50,000 sockeye salmon, in addition to late run escapement needs, which shall be comprised of 25,000 fish in August and 25,000 fish from September 1–15 (5 AAC 15.357(b)(3)).⁴¹

HARVEST ASSESSMENT PROGRAM

Estimates of annual CMA subsistence salmon harvests are based on annual permit returns combined with post permit household surveys in the CMA communities. The Division of Commercial Fisheries conducted its first subsistence salmon harvest assessment in the CMA in 1976. Subsistence harvest assessments for salmon have been conducted annually since then. Beginning in 1980, any fisher who chose to harvest their subsistence salmon from the Chignik Management Area had to obtain a permit. The Division of Subsistence assumed responsibility of the harvest assessment program from 1993–2011. In 1993, the Division of Subsistence obtained copies of all available subsistence permits for the CMA from the Division of Commercial Fisheries' archive in Kodiak. Permits issued prior to 1980 and for 1987 could not be located. Since 1993, harvest data from returned permits have been expanded by community of residence to estimate the harvest by all permit holders. Data from returned permits are tabulated by species and fishing area. Starting in 2012, the Division of Commercial Fisheries resumed the responsibility due to funding losses for the Division of Subsistence's harvest assessment program.

All permit data were entered into a database. The estimated harvests developed in this database and reported in subsequent Division of Commercial Fisheries reports differ slightly from those reported in earlier reports for several reasons. There are small discrepancies in some years in the number of permits issued or returned. Estimated harvests in earlier reports were based on a simple expansion from harvests reported on returned permits to the total number of permits issued.

The method of permit issuance in the communities varies by community and year, depending on the availability of vendors and other arrangements in place with area organizations. Permits are also issued upon request at the Chignik River fish weir by Division of Commercial Fisheries' seasonal staff, as well as from local community vendors. Chignik subsistence salmon permits must be returned by mail to the Alaska Department of Fish and Game by December 31 of the year issued. Permits include a harvest report that fishers are required to complete. The report asks for the dates fished, the specific locations fished, and the number of each species of salmon caught on each day. Nonresponses are sent reminder letters, and telephone calls are made if further follow-up is required. In addition, from 1993–2008, and 2011 (surveys were not conducted in 2009, 2010, and 2012 due to budgetary constraints), the Division of Subsistence administered face-to-face household subsistence salmon harvest in order to collect harvest information from households that subsistence fished but did not obtain a permit, or did obtain a permit, but had not returned their permit to the department at the end of the year. Survey technicians trained and hired by the Division of Subsistence from each community attempted to contact all households in the CMA. Surveys were generally conducted during January, February, and March. Respondents were asked questions similar to those included on the permit as well as additional questions regarding late season harvests and whether their subsistence needs were met.

Increases in permits issued and returned beginning in 1993–2007, and consequently higher harvest estimates, reflect the use of area vendors to issue permits as well as postseason surveys conducted by ADF&G staff and area research assistants. Comparisons of household survey data and permit data

40. Federal Subsistence Board. Subsistence Management Regulations for the Harvest of Fish and Shellfish on Federal Public Lands and Waters in Alaska, 2013–2015, 47–48. Federal Subsistence Board, Office of Subsistence Management, Anchorage.

41. Alaska Board of Fisheries. 2013. Chignik Finfish, Meeting, December 5–6, 2013, RC015, Anchorage. Accessed September 2014. http://www.adfg.alaska.gov/static/f/regulations/regprocess/fisheriesboard/pdfs/2013-2014/chignik/rcs/rc015_Chignik_Area_Salmon_Management_Plan.pdf

collected for 1984, and 1989 suggested that permit data underestimated subsistence harvests in the Chignik Area subsistence salmon fishery (Hutchinson-Scarborough and Fall 1996:27).⁴² With the assistance of area permit vendors, ADF&G Chignik weir staff, research assistants, and area governments, subsistence salmon harvest assessments for most recent years, with some exceptions, have resulted in more reliable estimates of the total harvest.

While subsistence salmon permits are issued to an individual, other members of a household can acquire additional permits if more fish are needed. Therefore the number of permits per household, and per community, can vary each year and may not necessarily represent a change in population or household size.

The number of permits issued as well as total salmon harvest estimates have fluctuated over time; but have declined overall in recent years. The reason for decline is not clearly understood; however, year-round populations in most CMA communities have declined as well, and there have been fewer participants in CMA commercial fisheries, which provides the main economic mainstay of these communities. Declining participation may be due to lower salmon prices, increased costs in maintaining boats and associated fishing expenses, a downturn in Alaska and U.S. economies, and a co-op commercial salmon fishery that occurred between 2002–2005 that resulted in numerous boats not fished that faced considerable repair and maintenance costs when the co-op fishery terminated and the limited entry only system resumed.

CMA SUBSISTENCE SALMON HARVESTS

In 2012, the number of subsistence permits issued for the Chignik Area totaled 106 permits, and 87 (82%) were returned to the department. The previous year, there were fewer permits issued (95) with 76 returned, a return rate 80%. Since 1977, the number of subsistence salmon permits issued for the Chignik Area has averaged 104 per year, with 71 permits (68%) returned. Over the last 10 years (2002–2011), the average has been 113 permits issued and 85 permits (75%) returned, and the recent 5-year average (2007–2011) was 106 permits issued and 80 (76%) returned. In years when the Division of Subsistence has conducted post season harvest surveys or has been responsible for collecting permits, the return rates have generally been the highest (Table 7-1).

Harvest reports printed on the back of subsistence salmon permits direct fishers to record each species of salmon they harvest. In 2012, the total estimated CMA salmon harvest was 8,242 fish, which was 40% (5,490) fewer salmon than in 2011 (13,732 fish). The 2012 total harvest was also well below the recent 5-year (11,166 fish) and 10-year (11,633 fish) and 1977–2011 (11,340 fish) historical averages. However, the 2012 estimate was similar to the 2008 and 2009 total harvests of 8,783 and 8,907 respectively. Therefore, even though the 2012 harvest numbers indicate a decline, they are still within the range of previous years (Table 7-1).

The 2012 total salmon harvest consisted of 68% sockeye (5,607), a significant (47%) decline from the 2011 sockeye harvest (10,578), which was confirmed by residents who said the Chignik River late sockeye run did not come in as strong as usual. Coho salmon made up 18% (1,488) of salmon harvested in 2012, very similar to the 2011 harvest of 1,458. Pink salmon harvests in 2012 totaled 810, a 37% decline from 2011 (1,289), and 220 chum salmon were harvested in 2012 (2% of the total), a 38% decline from 2011 (355). Chinook salmon composed 1% (116) of the 2012 salmon harvest, a 55% increase from 2011 but within the recent 5-year (2007–2011) average of 94, 10-year average (2002–2011) of 138 and historical average (1977–2011) of 83 (Table 7-1; Figure 7-2).

The 2012 composition of harvest was consistent with the recent 5-year, 10-year, and historical averages for the Chignik Management Area. The recent 5-year average (2007–2011) composition of the total

42. See also Hutchinson-Scarborough, Lisa, and Meredith Ann Marchioni. *In prep.* Chignik Bay, Chignik Lagoon, Chignik Lake, and Perryville: An Ethnographic Study of Traditional Subsistence Salmon Harvests and Uses. Alaska Department of Fish and Game Division of Subsistence Technical Paper No. NNN, Anchorage. Hereinafter cited as (Hutchinson-Scarborough and Marchioni 2014 *In prep.*).

salmon harvest (11,166) in the CMA comprised 77% sockeye (8,578), 13% coho (1,453), 8% pink (853), 2% chum (187), and <1% Chinook (94) (Table 7-1; Figure 7-2). The 10-year average (2002–2011) composition of total salmon harvested (11,633) comprised 75% sockeye (8,725), 14% coho (1,655), 8% pink (907), 2% chum (208) and 1% Chinook (138) (Table 7-1; Figure 7-3). The historical average (1977–2011), the Chignik Management Area’s total salmon harvest has comprised an estimated 78% (8,829) sockeye, 11% (1,290) coho, 8% (885) pink, 2% (355) chum, and 1% (52) Chinook (Table 7-1; Figure 7-4).

CMA Subsistence Salmon Harvests by Community

The majority of individuals who do their subsistence salmon fishing in the Chignik area are residents of Chignik Lake, Chignik Lagoon, Chignik Bay, Perryville, and Ivanof Bay. CMA residents have consistently held the majority of the CMA subsistence salmon permits and are responsible for the majority of the reported salmon harvest each year. In 2012, 75% of permits were issued to CMA residents, and they were responsible for 94% of the harvest (7,748 fish) while residents of other parts of Alaska harvested 6% (494 fish) of the total salmon harvest (Table 7-2; Table 7-3; Figure 7-3).

Perryville harvested more salmon than all the other communities harvesting salmon in the CMA, with a harvest of 3,578 total salmon representing 43% of the total CMA subsistence salmon harvests. Chignik Lagoon harvested the second highest amount of salmon (1,884) representing 23%, followed closely by Chignik Lake, with 1,442 salmon representing 17% of total salmon harvested in the CMA. Chignik Bay (532 salmon or 6%) and Ivanof Bay (312 salmon or 4%) harvested the least amount of salmon by community in the CMA. All other communities outside the CMA that participated in the CMA fishery in 2012 harvested a combined total of 806 salmon representing 10% of the total 2012 Chignik Management Area salmon harvest (Table 7-2; Table 7-3; Figure 7-6; Figure 7-7).

Community Salmon Harvests by Species

In 2012, the sockeye salmon harvest in the CMA was apportioned as follows: Chignik Lake 1,338 (24%), Perryville 1,607 (29%), Chignik Bay 385 (7%), Chignik Lagoon 1,771 (32%), Ivanof Bay 70 (1%), and 436 (8%) sockeye taken by residents of other Alaskan communities. Perryville’s 2012 harvest of 1,607 sockeye was a 42% decline from their 2011 sockeye harvest of 2,780. Chignik Lagoon residents harvested 1,771 sockeye in 2012, similar to the previous year’s harvest of 1559 sockeye. Chignik Lake’s 2012 estimated sockeye harvest (1,338) was less than half (52%) what it was in 2011 (2,807), and Chignik Bay’s estimated harvest in 2012 (385) was 83% less than in 2011 (2,221). Non-local residents harvested an estimated 436 sockeye in 2012 which marked a notable decline (36%) from the non-local residents’ 2011 harvest of 1,208 sockeye (Table 7-3; Figure 7-6; Figure 7-7). Researchers were told by residents of Chignik Lake and Chignik Lagoon that in 2011 the early run of sockeye was one of the largest in history, but that the late sockeye runs in 2012 and 2013 had poor returns explaining the low harvest rates those years (Hutchinson-Scarborough and Marchioni *In prep*).

Perryville harvested the most coho salmon (1,094) in 2012, which was 74% of the total CMA coho salmon harvest. Perryville residents also harvested the highest numbers of pink and chum salmon in 2012, responsible for 78% (172) of chum and 84% (679) pink salmon harvests in 2012 (Table 7-3; Figure 7-6; Figure 7-7). Chignik Lake, Chignik Lagoon, and Chignik Bay are all within close proximity to strong sockeye runs and therefore the majority of their salmon harvest is sockeye. Perryville is far from the other 3 communities and the Chignik River sockeye runs, but they do have local silver, pink, and chum salmon runs that they target for subsistence. There are fewer flights to Perryville each week than Chignik Lake, Chignik Lagoon, and Chignik Bay, and Perryville has no grocery store, making the residents of Perryville especially reliant on subsistence foods.

Location of Harvest

Subsistence salmon permits require people to record their harvest by species, date, quantity, and location. Table 7-4 shows the 2012 reported subsistence salmon harvests by species and by general locations

within the CMA identified by the Division of Subsistence as: Chignik Bay and Lagoon Subarea, Chignik Lake Subarea, and Perryville Subarea. The following section describes salmon harvest locations which are reported on salmon harvest permits. Not all individuals who return a permit record the location of their harvest, so the numbers do not reflect the total salmon harvest numbers for the Chignik Management Area.

For 2012, harvest location data were provided on 73 of the 87 returned permit returns for a total of 4,267 salmon. This included 1,915 salmon in the Chignik Bay/Lagoon subarea (Central, Eastern, and Chignik Bay commercial management districts [CMD], excluding areas above Mensis Point at the mouth of Chignik River at high tide in Chignik Lagoon), which represented 45% of the total reported harvest by location. Sockeye salmon (1,756 fish) made up the largest portion of the harvest in this subarea and represented 92% of the subarea harvest and 55% of the overall CMA sockeye harvest (Table 7-4).⁴³

The Chignik Lake subarea includes all waters of the Chignik River drainage above Mensis Point in Chignik Lagoon, including the Chignik River, Chignik Lake, and Chignik Lake tributaries. The reported subsistence harvests in the Chignik Lake Subarea totaled 868 salmon, which represented 20% of reported harvests by location. The majority of salmon harvested in this subarea were sockeye, totaling 824 (95%), and sockeye harvested from this area represented 26% of sockeye taken from the entire CMA. The reported salmon harvests in the Perryville subarea totaled 1,483 salmon which represented 35% of all salmon harvested. The Perryville Subarea was responsible for the majority of the CMA's harvest of coho, pink, and chum salmon with 601 coho salmon (89%), 156 pink salmon (68%), and 107 chum salmon (86%) (Table 7-4).

Table 7-5 shows reported CMA subsistence salmon harvests by species, fishing location, and date, in 2012. Harvest dates are divided into two periods of time, before and after July 5th because of the early and late sockeye run up the Chignik River. In 2012, 61% of the subsistence salmon harvest took place on or after July 5th. Residents said the numbers of sockeye salmon harvested on or after July 5th was low in 2012 because the late sockeye run was exceptionally low. Many people said that because the late run had a poor return they had to rely on fish harvested during the early run, which was not sufficient.

GEAR TYPE

Purse seines, hand seines, and gillnets are all allowable gear types for the harvesting of salmon for subsistence in the Chignik Management Area under state regulations. CMA subsistence salmon permits do not require that fishers record their gear type. Rod and reel or hook and line are sometimes used to harvest subsistence caught salmon under federal subsistence regulations (Hutchinson-Scarborough and Fall 1996; Hutchinson-Scarborough et al. 2010; Hutchinson-Scarborough and Marchioni *In prep*).

FEDERAL SUBSISTENCE FISHERY IN CMA

Federal subsistence fisheries are authorized in portions of the CMA for the permanent residents of the CMA communities.

Federal regulations in the CMA apply to waters within or adjacent to the Alaska Peninsula National Wildlife Refuge, Aniakchak National Monument and Preserve, and the Alaska Maritime National Wildlife Refuge. Federal and state subsistence regulations in the CMA generally parallel each other, however federal regulations authorize additional gear, harvest locations, and harvest seasons in portions of the CMA not authorized by the state. The federal program does not have a separate harvest assessment program from the state, however federal regulations within the Chignik watershed state that depending upon the area that one may fish, a Federal subsistence permit may be required. A Federal subsistence

43. Data in Tables 7-4 and 7-5 are based on extrapolating harvests recorded on returned permits. Not all permits recorded location of harvest; therefore these data are not expanded and differ from the estimated salmon harvest totals in tables 7-1 and 7-2. Reported harvests in this table are from 2012 permit returns only.

permit is required, in addition to a State subsistence fishing permit, to take salmon upstream of the Chignik River weir from January 1–August 9 using a rod and reel, with no daily harvest or possession limit, and to take salmon by gillnet in Black Lake or any tributary to Black or Chignik lakes. Additional information about the federal subsistence fishery is available by contacting the United States Fish and Wildlife Service, Office of Subsistence Management in Anchorage, AK.

SALMON REMOVAL FROM COMMERCIAL HARVESTS FOR HOME USE (“HOME PACK”)

Commercial fishers may also retain finfish from lawfully taken commercial catch for their home use, including use for bait under 5 AAC 39.010 (called “home pack” by area residents). These fish, if taken, are required to be reported on the commercial fish ticket and not on the subsistence salmon permit. Reported harvests are included in the ADF&G Division of Commercial Fisheries CMA annual finfish management reports. In 2012, Chignik commercial fishing boats reported removing 513 sockeye, 51 Chinook, 240 chum, and 22 pink salmon from their commercial harvest for home pack (Anderson et al. 2013; Table 7-6).

OTHER CHIGNIK AREA SUBSISTENCE FISHERIES

Estimates of subsistence halibut harvests for eligible communities and tribes, including those of the CMA, are available for 2012 (Fall and Koster 2014).

Although state regulations require a subsistence permit for the harvest of rainbow/steelhead trout and Arctic char/Dolly Varden, there are no annual harvest assessment programs for the other subsistence fisheries of the Chignik Area. The BOF, in an update of its C&T finding in January 2002, identified positive subsistence uses of all finfishes in the Chignik Area. Table 7-7 lists the finfishes other than salmon for which subsistence uses have been documented through systematic household interviews conducted by the Division of Subsistence.

For purposes of subsistence shellfish management, the Chignik Finfish Management Area is within the ADF&G Alaska Peninsula–Aleutian Islands Area. The BOF identified positive subsistence uses of all shellfish stocks in the Alaska Peninsula–Aleutian Islands Area. There are no subsistence harvest assessment programs for these shellfish stocks in the Chignik Area. Table 7-8 lists the shellfish for which subsistence uses have been documented through systematic household interviews.

In early 2004, the Division of Subsistence and the Bristol Bay Native Association, in a project funded by the *Exxon Valdez* Oil Spill Trustee Council, conducted comprehensive household surveys in Chignik Bay, Chignik Lagoon, Chignik Lake, and Perryville that, among other things, collected updated harvest data for nonsalmon fishes and marine invertebrates. A fifth community, Ivanof Bay, was not included in the study because it had no permanent year-round population at the time. A summary of these findings appears in Fall (2006a).

The reader should consult Morris (1987), Fall et al. (1995), Hutchinson-Scarborough and Fall (1996), and ADF&G (2002) for more background on these subsistence fisheries for nonsalmon finfishes and for shellfish. For harvest estimates based on systematic household interviews, see the CSIS. Limited nonsalmon subsistence resource use and harvest information was observed and documented during the Division of Subsistence 2010–2013 Chignik Management Area Subsistence Salmon Ethnography study (Hutchinson-Scarborough and Marchioni *In Prep*).

DISCUSSION

Prior to 2002, the years before the Chignik cooperative commercial fishery, many families processed most of their spring salmon for subsistence uses just prior to the first commercial opening in early June. Salmon were caught in early June either by purse seine or beach seine. Many families from Chignik Lake and Perryville would occupy fish camps across Chignik Lagoon. Chignik Lagoon and Chignik Bay

families would, for the most part, put up their early subsistence fish as a family effort prior to the start of the first commercial fishing opening. These early-run fish (sockeye salmon) are especially important to subsistence users because these fish are traditionally smoked, and it is necessary to cure these fish before flies hatch and deposit eggs on the fish, which typically occurs in mid- to late June. Traditionally, subsistence users could maximize their early season subsistence harvests because of large pulses of early-arriving fish. Area subsistence fishers have also reported that the early-run fish taste better and freeze or salt better if harvested early in the season. The second run (late run) of sockeye salmon was traditionally taken either in Chignik Lagoon, Chignik Lake, or near the mouth of the Clark River. Gillnets and beach seines were typically used to harvest late-run salmon in Chignik Lake (Hutchinson-Scarborough and Fall 1996:49).

During the years of the cooperative fishery (2002–2005), some changes occurred within area subsistence fishing patterns. Because many of the permit holders for Chignik were also area families who relied on subsistence fishing to feed their families, they would often use their commercial fishing boats to fish prior to the commercial fishing season. During these years, ADF&G opened the cooperative commercial fishery in early June, when many participants would have been processing their subsistence fish, and the nature of the management of the cooperative fishery resulted in fewer fish passing in the lagoon but at a more steady rate rather than experiencing pulses of fish that historically arrived when the competitive-only fishery was in operation. The management of the cooperative fishery resulted in a decrease in efficiency and an increase in effort for harvesting subsistence salmon in Chignik Lagoon.

According to verbal testimony by some Chignik families to Division of Subsistence personnel, many families had to wait until later in the summer to subsistence fish, and then the flies created spoilage. Other area residents reported both positive and negative changes occurring with their subsistence harvests and cash economy as a result of the creation of the cooperative commercial fishery. The cooperative fishing years resulted in a regulatory change that removed the restrictions on subsistence fishing for commercial fishers who could fish for subsistence throughout the summer as long as it was not done at the same time as a commercial opening and a permit was obtained. Fishers without a commercial permit and who wished to subsistence fish as before could fish for subsistence at any time as long as they had a subsistence permit. In addition, there was a reported increase in the amount of fish removed from the cooperative boats for home pack that were given to area residents to supplement their subsistence harvests. At that time, subsistence users informed ADF&G that despite the adjustments to the CMA subsistence fishery, which allowed for more opportunity for subsistence fishing, some were still having difficulty obtaining their subsistence salmon in 2004 and 2005.

In 2002 and 2004, the USFWS implanted radio transmitters in sockeye salmon in August and early September to determine when sockeye salmon targeted in the late season subsistence fishery passed the Chignik weir. The results of the 2002 studies are described in Anderson (2003). As stated in the regulations section of this chapter, in 2004 the BOF modified the commercial fisheries management plan for late-run sockeye salmon to allow more fish to pass into Chignik Lake in September, thus providing for subsistence harvests. Late-run sockeye salmon, which are dried, are harvested from Chignik Lake in the fall by many Chignik Area residents, including some Perryville families. In 2006, several residents, particularly from Chignik Lake, commented to ADF&G that despite the limits to the August commercial fishery, they still had difficulty acquiring their late-run salmon because they were not seeing as many fish as in prior years. They needed to fish more days to achieve harvest goals, or they harvested fewer late-run salmon. By 2006, after the cooperative commercial fishery was abolished, area subsistence patterns generally returned to the historical patterns used prior to the cooperative fishery, but on a reduced level. In 2008, there was a decrease in participation in the Chignik subsistence fishery with 89 permits issued; 39 fewer permits issued than in 2007, and 32 fewer than the previous 10-year (1996–2007) average of 121. In 2010, however, there was an increase in permits with 124 issued, which was an increase from 2009 (95). In 2012, 106 permits were issued, which was nearly the same as the historical average of 104 permits, and slightly less than the recent 10-year average of 113 permits.

Numerous fish or summer camps located on the north side of Chignik Lagoon were abundantly utilized in the 1990s but were mostly abandoned by 2006. However, in 2007 through 2012, a few families that reside in Perryville continued to stay at their camps during the commercial fishing season. A Perryville family member who has maintained and continues to utilize one of these multi-generational camps indicated that these camps are summer homes that provide housing for the families when commercial fishing; however, subsistence salmon are also harvested and processed while occupying the camps (Hutchinson-Scarbrough and Marchioni *In prep*).

In 2010 and 2011, gillnets, and purse seines were used to catch subsistence salmon, primarily sockeye from both runs in Chignik Lagoon and the lower mouth of Chignik River. In Chignik River, Chignik Lake and tributaries to Chignik Lake (primarily Clark River) salmon, primarily sockeye from both early and late runs were harvested using gillnets or beach seines. In addition, beach seines and handlines were used to harvest the late-run “red fish” (sockeye that have entered freshwater and starting to spawn) and “spawned-outs” (spawning sockeye), primarily at Hatchery Beach on Chignik Lake, the mouth of Clark River, or in Clark River and Home Creek. Also in 2011, sockeye were harvested by handline in Black Lake, but not in 2012. Late-run or spawning fish are typically preserved by drying in the wind on racks after harvest. This method is preferred because residents report they have less fat than early-run sockeye salmon. Chinook salmon were caught in Chignik Lagoon and the Chignik River using gillnets, seines and rod and reel or removed from commercial harvests and preserved by smoking or canning (Hutchinson-Scarbrough and Marchioni *In prep*).

Perryville subsistence patterns have not changed greatly from historical times, though fewer families are going to fish camps or summer homes located on the northern side of Chignik Lagoon. In 2011 and 2012, 6 of these camps were occupied by Perryville residents that utilize these camps during commercial fishing and to subsistence fish for sockeye. Fresh sockeye salmon are brought back to the village by commercial fishing families. Area streams and beaches are used extensively for the harvest of the local runs of coho, chum, and pink salmon, as well as the occasional sockeye salmon. Due to the fluctuations in river locations and stream flow, and fluctuations in salmon runs to these systems, Perryville subsistence fishers may have to use other streams and bays as far east of the village as Mitrofanina Bay and as far west as Ivanof Bay to harvest their fish. Fish are smoked, dried, canned, salted, and frozen by Perryville residents. Some Perryville families have relatives in Chignik Lake and travel to Chignik Lake in the fall to harvest late-run sockeye salmon for drying (Hutchinson-Scarbrough and Marchioni *In prep*).

Table 7-1.—Historical subsistence salmon harvests, Chignik Area, 1977–2012.

Year	Permits		Estimated salmon harvest					Total
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	
1977	NA	NA	50	9,700	2,400	600	1,800	14,550
1978	NA	NA	50	6,000	500	600	2,100	9,250
1979	NA	NA	14	7,750	34	0	262	8,060
1980	82	37	6	12,475	32	169	478	13,160
1981	29	7	0	2,049	0	0	0	2,049
1982	59	15	3	8,532	12	0	2	8,548
1983	32	21	0	3,078	1,319	850	1,250	6,497
1984	77	64	23	8,747	464	204	330	9,768
1985	59	48	1	7,177	50	25	26	7,279
1986	74	38	4	10,347	205	77	98	10,730
1987	NA	NA	10	7,021	278	204	261	7,774
1988	80	34	9	9,073	1,455	142	54	10,733
1989	68	23	24	7,551	384	147	81	8,187
1990	72	23	103	8,099	210	115	470	8,996
1991	95	58	42	11,483	13	81	275	11,893
1992	98	19	55	8,648	709	145	305	9,862
1993	201	141	122	14,710	3,765	642	1,265	20,503
1994	219	122	165	13,978	4,055	382	1,720	20,300
1995	111	95	98	9,563	1,191	150	723	11,726
1996	119	104	48	7,357	2,126	355	2,204	12,089
1997	126	103	28	13,442	2,678	840	2,035	19,024
1998	104	72	91	7,750	1,390	186	1,007	10,424
1999	106	88	243	9,040	1,679	136	1,191	12,290
2000	130	112	163	9,561	1,802	517	1,185	13,227
2001	135	122	171	8,633	1,859	213	2,787	13,663
2002	120	86	74	10,092	1,401	23	390	11,980
2003	146	127	267	10,989	2,256	286	1,597	15,394
2004	104	57	88	7,029	1,981	202	1,047	10,347
2005	119	100	224	8,171	2,112	353	730	11,590
2006	113	79	259	8,079	1,539	275	1,035	11,187
2007	128	83	84	10,191	1,936	165	996	13,372
2008	89	69	41	7,189	877	57	619	8,783
2009 ^a	95	82	104	6,785	1,174	137	707	8,907
2010 ^a	124	90	188	8,148	1,820	222	656	11,034
2011	95	76	52	10,578	1,458	355	1,289	13,732
2012 ^a	106	87	116	5,607	1,488	220	810	8,242
5-year average (2007–2011)	106	80	94	8,578	1,453	187	853	11,166
10-year average (2002–2011)	113	85	138	8,725	1,655	208	907	11,633
Historical average (1977–2011)	104	71	83	8,829	1,290	253	885	11,340

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014); Quimby and Owen (1994) for 1976–1979 and 1987.

-continued-

Table 7-1.–Page 2 of 2

Note NA indicates data not available. Information regarding the number of permits issued and returned was collected; however, the records containing this information no longer exist. Harvest data for these years are also recorded in ADF&G Division of Commercial Fisheries and Division of Sport Fish area management reports.

- a. From 1993–2008 and 2011, post-season household surveys were conducted to supplement harvest data collected through returned permits. Limited budgets prevented administering the surveys for 2009, 2010, and 2012 likely resulting in an underestimate of subsistence harvests since not all subsistence fishing households obtained a permit. To compensate for this underestimate, the average annual harvest for the period 1999–2008 and 2011 reported during post-season surveys was added to harvests from returned permits to estimate the total subsistence harvest for 2009, 2010, and 2012.

Table 7-2.—Participation in Chignik area subsistence salmon fishery and estimated total salmon harvest by place of residence, 1980–2012.

Year	Local area residents ^a				Other residents			
	Subsistence permits		Salmon harvests		Subsistence permits		Salmon harvests	
	Issued	Percent of total	Estimated amount	Percent of total	Issued	Percent of total	Estimated amount	Percent of total
1980	51	62%	9,013	68%	31	38%	4,148	32%
1981	24	83%	2,049	100%	5	17%	0	0%
1982	52	88%	8,059	94%	7	12%	489	6%
1983	21	66%	5,585	86%	11	34%	912	14%
1984	46	60%	6,751	69%	31	40%	3,018	31%
1985	43	73%	6,072	83%	16	27%	1,207	17%
1986	53	72%	8,977	84%	21	28%	1,753	16%
1987	<i>Data unavailable</i>							
1988	61	76%	8,768	82%	19	24%	1,965	18%
1989	42	62%	6,999	85%	26	38%	1,188	15%
1990	50	69%	7,258	81%	22	31%	1,738	19%
1991	69	73%	8,815	74%	26	27%	3,078	26%
1992	91	93%	9,612	97%	7	7%	250	3%
1993	176	88%	19,070	93%	25	12%	1,433	7%
1994	199	91%	18,760	92%	20	9%	1,540	8%
1995	101	91%	10,818	92%	10	9%	908	8%
1996	113	95%	11,809	98%	6	5%	280	2%
1997	115	91%	18,190	96%	11	9%	834	4%
1998	98	94%	10,164	98%	6	6%	260	2%
1999	98	92%	11,662	95%	8	8%	628	5%
2000	112	86%	11,777	89%	18	14%	1,450	11%
2001	108	80%	12,486	91%	27	20%	1,178	9%
2002	102	85%	8,308	92%	18	15%	726	8%
2003	127	87%	14,315	93%	19	13%	1,079	7%
2004	90	87%	10,026	98%	14	13%	213	2%
2005	110	92%	11,148	96%	9	8%	442	4%
2006	107	95%	11,071	99%	6	5%	116	1%
2007	106	83%	12,427	93%	22	17%	944	7%
2008	66	74%	7,633	87%	23	26%	1,151	13%
2009	71	75%	7,876	88%	24	25%	1,031	12%
2010	100	81%	9,583	87%	24	19%	1,451	13%
2011	80	84%	12,524	91%	15	16%	1,208	9%
2012	80	75%	7,748	94%	26	25%	494	6%

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

^a Local residents include individuals reporting the addresses of Chignik Bay, Chignik Lagoon, Chignik Lake, Ivanof Bay, and Perryville for permits issued for the Chignik subdistrict fishery.

Table 7-3.—Estimated subsistence salmon harvests by community, Chignik Area, 2012.

Community	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
Chignik Bay	7	7	11	385	105	12	19	532
Chignik Lagoon	19	17	61	1,771	12	2	37	1,884
Chignik Lake	16	13	16	1,338	45	3	41	1,442
Ivanof Bay	2	2	1	70	182	27	32	312
Perryville	35	27	26	1,607	1,094	172	679	3,578
Subtotal, Chignik Area residents	80	67	116	5,171	1,438	215	807	7,748
Anchorage	9	6	0	101	11	0	0	111
Haines	1	1	0	27	0	0	0	27
Homer	2	2	0	63	0	0	0	63
Juneau	1	0	0	0	0	0	0	0
Kodiak	8	8	0	141	10	4	1	156
Other USA	1	1	0	85	30	1	2	118
Ouzinkie	1	0	0	0	0	0	0	0
Palmer	1	1	0	0	0	0	0	0
Slana	1	1	0	19	0	0	0	19
Sutton	1	0	0	0	0	0	0	0
Subtotal, other Alaska residents	26	20	0	436	51	5	3	494
Total	106	87	116	5,607	1,488	220	810	8,242

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Table 7-4.—Subsistence salmon harvests by species and subarea of harvest, Chignik Area, 2012.

Subarea of harvest ^a	Estimated salmon harvest ^b					
	Chinook	Sockeye	Coho	Chum	Pink	Total
<i>Chignik Bay and Lagoon</i>	41	1,756	55	7	56	1,915
Chignik Bay	0	87	10	4	0	101
Chignik Lagoon	41	1,669	45	3	56	1,814
<i>Chignik Lake</i>	2	824	17	9	16	868
Chignik Lake	2	160	0	0	12	175
Chignik River	0	470	17	9	4	500
Clark River	0	26	0	0	0	26
Mouth of Clark River	0	168	0	0	0	168
<i>Perryville</i>	22	597	601	107	156	1,483
Ivanof Bay	0	200	469	28	6	703
Perryville Area	22	397	133	79	149	780
Total	66	3,176	673	124	228	4,267

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

- a. The Chignik Bay-Lagoon Subarea corresponds to the portion of the Central District and the Chignik Bay District, not including any of the Chignik River from the outlet of Chignik Lake (“FRI Point” to the River’s outlet at Mensis Point in Chignik Lagoon). The Chignik Lake Subarea includes subsistence harvests in the Chignik River from Mensis Point in Chignik Lagoon up to Black Lake. The Perryville Subarea corresponds to the Perryville and Western districts, including Ivanof Bay, Mitrofanina Bay, the Kametolook River and other streams near Perryville and Ivanof Bay. In recent years there have been no subsistence harvests reported for the Eastern District.
- b. Harvest estimates are based on 2012 permit returns only. Of 73 permits issued for the Chignik Area, 54 permits were returned (74%).

Table 7-5.—2012 Chignik area subsistence salmon harvests by species, fishing location, and date.

Subsistence area	Estimated salmon harvest					
	Chinook	Sockeye	Coho	Chum	Pink	Total
<i>Harvest before 7/5</i>						
Chignik Bay	0	60	10	0	0	70
Chignik Lagoon	11	623	0	0	0	634
Chignik Lake	0	73	0	0	0	73
Chignik River	0	299	0	0	0	299
Perryville	1	201	97	39	91	429
Ivanof Bay to Humpback Bay	0	162	0	3	0	165
Subtotal, early harvest	12	1,419	107	42	91	1,671
<i>Harvest on or after 7/5</i>						
Chignik Bay	0	30	0	4	0	34
Chignik Lagoon	30	837	38	3	39	947
Chignik Lake	2	87	0	0	12	101
Chignik River	0	171	17	9	4	201
Clark River	0	26	0	0	0	26
Mouth of Clark River	0	168	0	0	0	168
Perryville	21	195	35	40	58	350
Ivanof Bay to Humpback Bay	0	38	469	26	6	538
Unknown location	0	209	7	0	17	233
Subtotal, late harvest	53	1,759	566	82	136	2,597
Total	65	3,178	673	124	227	4,267

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Table 7-6.—Chignik area salmon removed from commercial catch for home use, 1994–2012.

Year	Reported Salmon Harvest					Total
	Chinook	Sockeye	Coho	Chum	Pink	
1994	ND	ND	ND	ND	ND	ND
1995	64	0	913	5	0	982
1996	40	40	20	21090	5262	26,452
1997	88	664	0	0	0	752
1998	108	267	27	155	0	557
1999	211	26	200	3	0	440
2000	20	0	0	0	0	20
2001	90	217	7	129	7	450
2002	77	1,371	164	0	0	1,612
2003	309	2,411	74	0	407	3,201
2004	158	1690	0	0	0	1,848
2005	271	1,364	5	115	234	1,989
2006	68	267	175	0	0	510
2007	16	205	56	1	0	278
2008	15	0	0	0	0	15
2009	75	93	0	1	0	169
2010	118	973	0	0	7	1,098
2011	142	323	16	0	0	481
2012	51	513	0	240	22	826
Total	1,921	10,424	1,657	21,739	5,939	41,680
Historical average (1995–2012)	107	579	92	1,208	330	2,316

Source ADF&G CFMD Reported Commercial Salmon Ticket Reports

ND = No data

Table 7-7.—Subsistence uses of nonsalmon finfishes by community, Chignik Area, 1989.

Common English name	Scientific name, if not previously given	Percentage of households using in				
		Chignik Bay	Chignik Lagoon	Chignik Lake	Ivanof Bay	Perryville
Pacific herring		23	47	29	29	15
Pacific herring spawn on kelp		14	0	5	0	4
Walleye pollock	<i>Theragra chalcogramma</i>	3	0	0	0	0
Rainbow smelt ^a		11	0	48	0	0
Pacific halibut		89	100	67	100	96
Rainbow trout		3	0	24	57	7
Dolly Varden		23	7	38	86	56
Eulachon (candlefish)	<i>Thaleichthys pacificus</i>	23	40	33	100	78
Pacific cod (gray cod)		29	60	48	86	63
Sculpin	<i>Hemilepidotus sp.</i>	11	0	5	0	30
Starry flounder		6	0	19	14	0
Kelp greenling	<i>Hexagrammos decagrammus</i>	11	0	10	0	30
Arctic grayling		0	0	0	14	0
Sablefish (black cod)	<i>Anoplopoma fimbria</i>	0	7	5	0	0
Steelhead trout		0	13	5	0	0
Black rockfish	<i>Sebastes melanops</i>	0	7	0	0	22
Red (yelloweye) rockfish	<i>Sebastes ruberrimus</i>	3	0	0	0	4
Any nonsalmon fish		89	100	86	100	96

Source CSIS; Hutchinson-Scarborough and Fall (1996).

- a. Most likely harvested outside the Chignik Management Area; Chignik area households receive gifts of rainbow smelt from relatives and friends in Pilot Point, Ugashik, and Naknek, among other communities.

Table 7-8.—Subsistence uses of marine invertebrates by community, Chignik Area, 1989.

Common English name	Scientific name, if not previously given	Percentage of households using in				
		Chignik Bay	Chignik Lagoon	Chignik Lake	Ivanof Bay	Perryville
Alaska razor clam	<i>Siliqua patula</i>	14	33	24	43	37
Butter clam	<i>Saxidomus giganteus</i>	71	67	52	71	41
Gaper clam	<i>Tresus capax</i>	11	0	0	0	4
Nuttall cockle	<i>Clinocardium nuttallii</i>	37	7	48	100	70
Pinkneck clam (redneck)	<i>Spicula polynuma</i>	0	0	0	71	4
Pacific littleneck (steamer) clam	<i>Protothaca staminea</i> ^a	11	0	0	29	11
Chiton, black (leather)	<i>Katharina tunicata</i>	49	27	57	100	93
Chiton, red (gumboot)	<i>Cryptochiton stelleri</i>	0	0	0	86	11
Mussel (blue)	<i>Mytilus trossulus</i>	9	7	0	14	15
Octopus	<i>Octopus</i> spp	43	20	48	71	52
Sea urchin	<i>Strongylocentrotus</i> spp	29	0	48	100	89
Sea cucumber	<i>Varius</i> spp	0	0	0	0	4
Shrimp	<i>Pandalus</i> spp	9	0	5	0	0
Giant Pacific scallop	<i>Pecten caurinus</i>	3	0	0	0	0
Red king crab	<i>Paralithodes camtschatica</i>	40	20	33	43	0
Dungeness crab	<i>Cancer magister</i>	37	40	48	100	52
Tanner crab	<i>Chionoecetes bairdi</i>	63	67	14	0	4
Snail	<i>Neptunea</i> spp	3	0	0	0	4
Limpet	<i>Acmaeidae</i> spp	3	0	0	0	4
Any marine invertebrates		89	87	81	100	96

Source CSIS; Hutchinson-Scarborough and Fall (1996).

a. May also include smaller-sized individuals of other species and softshell clams of the genus *Mya*.




Figure 7-1.—Location of Chignik Management Area (CMA) and communities within the CMA on Alaska Peninsula.




Figure 7-2.—Composition of Chignik Area subsistence salmon harvest by species, 2012.




Figure 7-3.—Species composition of Chignik Area subsistence salmon harvests, 2007–2011.




Figure 7-4.—Species composition of Chignik Area subsistence salmon harvests, 2002–2011.




Figure 7-5.—Species composition of Chignik Area subsistence salmon harvests, 1977–2011.




Figure 7-6.—Subsistence salmon harvests by community, Chignik Area, 2012.




Figure 7-7.—Subsistence salmon harvests by community, Chignik Area, 2012.

CHAPTER 8: ALASKA PENINSULA AREA

BACKGROUND

The Alaska Peninsula Area includes all Pacific Ocean waters of Alaska on the north side of the Alaska Peninsula southwest of a line from Cape Menshikof to Cape Newenham and east of the longitude of Cape Sarichef Light and on the south side of the Alaska Peninsula from a line extending from Scotch Cap through the easternmost tip of Ugamak Island to a line extending 135 degrees southeast from Kupreanof Point. The communities of the Alaska Peninsula Area are Port Heiden (estimated population 123 in 2012), Nelson Lagoon (population 46), False Pass (population 26), Cold Bay (population 98), King Cove (population 962), and Sand Point (population 982).⁴⁴ Port Heiden is in the Lake and Peninsula Borough; the other communities are in the Aleutians East Borough (which also includes Akutan in the Aleutian Islands Area).

REGULATIONS

A subsistence permit, which must be used to record daily harvests, is required for fishing in the Alaska Peninsula Area. There is an annual limit of 250 salmon per household. Legal gear includes seines and gillnets. In waters open to commercial fishing, set and drift gillnets may not exceed 50 fathoms in length. In most other areas, set gillnets may not exceed 100 fathoms and drift gillnets may not exceed 200 fathoms. Purse seines may not exceed 250 fathoms in length. Other standard permit conditions include prohibition of fishing within 300 ft of a dam, fish ladder, weir, culvert, or other artificial obstruction. The Alaska Administrative Code (5 AAC 01.423) includes special provisions regarding subsistence gear for other areas, including Mortensens Lagoon, the False Pass vicinity, and Bear and Sandy rivers. Salmon may be taken at any time, except in those districts and sections that are open to commercial salmon fishing, salmon may not be taken during the 24 hours before and 12 hours following a commercial salmon fishing period. A few small areas closed to subsistence salmon fishing are listed in 5 AAC 01.425.

Federal regulations governing subsistence salmon fishing in waters under the jurisdiction of the FSB are generally identical to the state regulations summarized above, with the exception that rod and reel, in addition to gillnet and seine, is legal subsistence gear under federal rules. There is no separate federal subsistence permit; a state permit is required for subsistence fishing under the federal regulations.

HARVEST ASSESSMENT PROGRAM

The Division of Commercial Fisheries has issued subsistence permits for the Alaska Peninsula Area since 1979. Except for residents of Sand Point and Cold Bay, permits are mailed each year to fishers who returned their permits at the end of the previous fishing season. Sand Point and Cold Bay residents are issued permits upon request at the ADF&G offices in Sand Point and Cold Bay. Permits are also issued upon request at other ADF&G offices and by mail to people who telephone to request them. Regulations require that permits be returned to ADF&G by October 31. Reminder letters are sent around November 1 to people who have not yet returned their permits. If a person does not return the permit, his or her name is removed from the mailing list. Data from returned permits are tabulated by species and fishing area. Harvest data from returned permits are expanded by community of residence to estimate the harvest by all permit holders.

SUBSISTENCE SALMON HARVESTS IN 2012

From 1985 through 2011, the number of subsistence salmon permits issued for the Alaska Peninsula Area averaged 192 per year (Table 8-1). The recent 5-year average (2007–2011) was 166 permits. In 2012, 172 subsistence salmon fishing permits were issued for the Alaska Peninsula Area, up from 163 issued in

44. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. "Research and Analysis Homepage." Accessed July 2014. <http://labor.alaska.gov/research/pop/popest.htm>

2011. This compares to the 248 permits issued for the commercial salmon fishery for the Alaska Peninsula Area in 2012 (Poetter et al. 2012). The response rate for subsistence permits was 80% in 2012 (138 of 172 permits were returned). Of all subsistence permits issued, 149 (87%) were issued to residents of Alaska Peninsula Area communities, and 23 (13%) were issued to residents of other Alaska communities (Table 8-2). Most nonlocal residents fish at Mortensens Lagoon on the Cold Bay road system.

The estimated subsistence salmon harvest in the Alaska Peninsula Area in 2012 was 14,231 fish. This is a decrease from the year before (14,466 salmon) but is more than the recent 5-year average (12,993) and less than the 10-year average (14,288) (Table 8-1). The 2012 subsistence harvest was made up of 66% sockeye salmon, 14% coho salmon, 7% pink salmon, 12% chum salmon, and 2% Chinook salmon (Figure 8-1). Of the total harvest, the residents of Cold Bay took 7%, False Pass residents 3%, Sand Point residents 42%, Port Heiden residents 2%, Port Moller residents <1%, Nelson Lagoon residents 2%, and King Cove residents 37%. Other Alaska residents harvested 4% (Table 8-2; Figure 8-2). Following historical peak harvest levels recorded in 1997, existing data indicate a general decline in the Alaska Peninsula Area subsistence salmon harvest (Table 8-1).

In interviews with Division of Subsistence staff, fishery managers expressed the view that the subsistence permit program did not completely document all subsistence salmon harvesting activities because some fishers did not obtain permits. A comparison of permit and household interview data for 1992 for King Cove found that about 31% of interviewed households that reported subsistence fishing did not have permits (Fall, Mason, et al. 1993:58–62). The estimated total subsistence salmon harvest for the community based on the interviews was 7,036 ($\pm 1,773$), compared to 5,856 based on permit returns (Fall, Mason, et al. 1993:58–62.). At Sand Point in the same year, 41% of interviewed households reported that they harvested salmon for subsistence but did not have permits. The estimated total subsistence salmon harvest for Sand Point based on the household interviews was 11,338 ($\pm 2,551$), compared to 7,833 based on estimates using permit return information (Fall, Andersen, et al. 1993:61).

The subsistence permit program for the Alaska Peninsula Area does not account for salmon withheld from commercial catches for home uses. Fishery managers believe that this number is substantial, especially in years when commercial salmon prices are low. For 1992, it was estimated that 51% of the salmon harvested for home uses at King Cove (Fall, Mason, et al. 1993), and 45% at Sand Point (Fall, Andersen, et al. 1993), were removed from commercial harvests.

In 2002 and 2003, the Division of Subsistence conducted the Subsistence Fisheries Harvest Assessment and Traditional Ecological Knowledge, Lower Alaska Peninsula and Aleutian Islands project, funded in part by OSM under project number 02-032. The goals of the project were to generate harvest data for salmon to supplement estimates produced through the subsistence permit program and to collect TEK about fisheries resources. Among other findings, the research documented that King Cove households removed 2,304 salmon from their commercial harvests for home uses in 2003, representing 24% of the total salmon harvest for home uses in the community (Davis 2005:116). Another product was a searchable TEK database called “The View from the Beach.” For detailed study findings, consult Davis (2005).

OTHER SUBSISTENCE FISHERIES

Subsistence Pacific halibut fishing harvest estimates for communities and tribes in the Alaska Peninsula Area are available for 2003–2012 (Fall and Koster 2014).

There are no other annual harvest assessment programs for the other finfish and shellfish subsistence fisheries of the Alaska Peninsula Area. The Division of Subsistence has conducted 1 round of systematic household harvest surveys in each of the area’s communities, except for Cold Bay. The findings of these surveys, including species used, percentage of households harvesting each species in the study year, and estimated harvest quantities for the study year, appear in the CSIS. Table 8-3 reports the percentage of households in the surveyed communities that used selected nonsalmon finfish species in the study year.

Generally, Pacific cod, halibut, and Arctic char/Dolly Varden were the most frequently used by households in these communities.

Table 8-1.—Historical subsistence salmon harvests, Alaska Peninsula Area, 1985–2012.

Year	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1985	161	95	74	4,037	7,504	1,566	574	13,755
1986	147	84	101	5,396	2,996	1,455	1,779	11,727
1987	191	144	193	5,777	4,259	1,943	1,547	13,719
1988	183	114	257	5,501	5,646	1,692	1,666	14,762
1989	188	139	88	10,404	3,505	2,104	1,213	17,314
1990	201	157	246	8,588	4,029	1,589	736	15,188
1991	249	185	458	11,345	5,551	3,551	1,878	22,783
1992	229	177	385	10,739	4,267	2,574	1,840	19,805
1993	262	215	615	12,478	5,753	1,997	1,189	22,032
1994	256	213	674	11,884	6,086	4,406	2,206	25,256
1995	260	198	492	12,716	5,021	3,369	2,653	24,251
1996	234	178	362	12,176	7,743	2,728	2,569	25,578
1997	217	172	420	15,224	4,612	2,885	2,955	26,096
1998	233	153	407	12,920	5,820	1,326	2,286	22,759
1999	185	148	391	15,119	4,961	2,235	2,136	24,843
2000	180	152	341	9,955	5,239	1,699	950	18,185
2001	185	155	570	12,259	3,940	1,963	1,181	19,912
2002	157	133	345	9,384	3,188	1,603	532	15,052
2003	166	128	312	10,103	4,266	2,353	1,194	18,228
2004	147	135	218	9,484	3,787	951	609	15,049
2005	160	139	192	11,260	4,089	716	1,054	17,310
2006	153	131	110	7,847	2,452	910	961	12,280
2007	150	124	100	6,872	2,648	498	693	10,811
2008	199	164	280	7,623	4,355	1,078	1,687	15,022
2009	134	118	350	5,629	2,545	434	749	9,707
2010	183	138	338	9,464	2,898	1,274	985	14,959
2011	163	117	337	9,871	2,353	835	1,070	14,466
2012	172	138	287	9,429	1,936	1,637	941	14,231
5-year average (2007–2011)	166	132	281	7,892	2,960	824	1,037	12,993
10-year average (2002–2011)	161	133	258	8,754	3,258	1,065	953	14,288
Historical average (1985–2011)	192	148	321	9,780	4,426	1,842	1,440	17,809

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Table 8-2.—Subsistence salmon harvest estimates by community, Alaska Peninsula Area, 2012.

Community	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
Cold Bay	23	20	2	838	21	5	0	866
False Pass	4	4	10	288	29	42	27	396
King Cove	56	45	50	3,619	1,094	436	24	5,222
Nelson Lagoon	9	4	9	788	97	11	2	907
Port Heiden	5	4	29	193	64	55	0	340
Sand Point	52	41	178	3,234	621	1,073	805	5,912
Subtotal, area residents	149	118	277	8,959	1,926	1,621	858	13,642
Anchorage	6	5	0	144	0	0	0	144
Auke Bay	1	1	0	50	0	0	0	50
Eagle River	1	1	0	30	0	0	0	30
Homer	4	4	0	25	8	10	80	123
Kasilof	2	1	0	110	0	0	0	110
Kodiak City	4	4	10	24	0	0	0	34
Kotzebue	1	0	0	0	0	0	0	0
Seward	2	2	0	12	2	6	3	23
Wasilla	1	1	0	0	0	0	0	0
Willow	1	1	0	75	0	0	0	75
Subtotal, other Alaska residents	23	20	10	470	10	16	83	589
Total	172	138	287	9,429	1,936	1,637	941	14,231

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Table 8-3.—Percentage of households using selected nonsalmon finfishes, Alaska Peninsula Area communities.

Resource ^b	Percentage of households using in that study year ^a				
	False Pass	King Cove	Nelson Lagoon	Port Heiden	Sand Point
Pacific cod	65	44	0	3	61
Sablefish	15	8	ND	ND	13
Kelp greenling	10	5	ND	ND	7
Flounder	20	4	8	11	4
Pacific halibut	95	73	0	22	89
Pacific herring	30	23	ND	3	14
Herring spawn on kelp	0	3	ND	3	1
Smelt	0	1	ND	49	5
Rockfishes	5	36	ND	ND	61
Sculpin	35	7	ND	ND	4
Walleye pollock	ND	3	ND	ND	2
Lake trout	ND	ND	ND	11	ND
Arctic char/ Dolly Varden	75	67	54	76	51
Rainbow trout/steelhead	5	4	ND	3	31

Source CSIS.

Note ND indicates no data for that resource.

- a. Study year = 1987–1988 for False Pass; 1986–1987 for Nelson Lagoon and Port Heiden; 1992 for King Cove and Sand Point.
- b. Most commonly used types in the study year; uses of other species occurred, or may occur in other years.




Figure 8-1.—Composition of Alaska Peninsula Area subsistence salmon harvest by species, 2012.




Figure 8-2.—Subsistence salmon harvests by community, Alaska Peninsula Area, 2012.

CHAPTER 9: ALEUTIAN ISLANDS AREA

INTRODUCTION

The Aleutian Islands Management Area includes all waters of Alaska in, and surrounding, the Aleutian Islands west of Cape Sarichef Light and west of a line extending from Scotch Cap through the easternmost tip of Ugamak Island, including the waters in and surrounding the Pribilof Islands (5 AAC 01.350). For subsistence purposes, the Aleutian Islands Area is divided into 6 management districts. From east to west, they are the Akutan District, Unalaska District, Umnak District, Pribilof Islands District, Atka–Amlia Islands District, and the Adak District (5 AAC 01.355). The major communities of the Aleutian Islands Area are Akutan, Unalaska–Dutch Harbor, Atka, Nikolski, and Adak. Akutan is part of the Aleutians East Borough; the other communities are part of the Aleutians West Census Area, but they are not within an organized borough.

According to the US Census, Akutan’s population in 2010 totaled 1,027, of which only 90 people lived in a total of 40 households in the Native Village of Akutan, and the remaining 937 lived in group quarters at the adjacent Trident Seafoods processing plant. In 2012, the total Akutan population was estimated at 1,106; however, most of the people (1,023) were estimated as residing in group housing, and 83 resided in households. In 2010, the population of Unalaska–Dutch Harbor was 4,376 with 2,277 residents residing in a total of 927 households and the remainder (2,099) in group quarters (primarily seafood industry workers housing). In 2012, the estimated population of Unalaska–Dutch Harbor was 4,766 with 2,419 residing in households and 2,347 in group quarters. In Nikolski, the population in 2010 was 18 residing in a total of 13 households; and in 2012, the estimated population was 16. Atka in 2010 had a population of 61 residing in a total of 24 households; and the estimated population in 2012 totaled 59. Adak’s 2010 census population totaled 326 people which 109 lived in a total of 44 households and 217 in group quarters; and in 2012, the estimated population was 331 total people, 182 of which were estimated to be in group quarters.⁴⁵

The communities in the Pribilof Island District; St. Paul in 2010 had a population of 479 with 455 residing in a total of 162 households and 24 residing in group quarters; and in 2012, the population was estimated at 453 with 441 residing in households and 12 people residing in group quarters. St. George’s 2010 population was 102, with 98 residing in a total of 42 households, and 4 residing in group quarters; and the 2012 population estimate was 84 people; 121 of which residing in group quarters..⁴⁶⁴⁷

The Alaska Board of Fisheries found that halibut and all other finfish in the Aleutian Islands Area and the waters surrounding the Pribilof Islands are customarily and traditionally taken or used for subsistence. The board found that (1) 13,500–23,000 salmon and (2) 200,000–330,000 usable pounds of finfish other than salmon are reasonably necessary for subsistence uses in the Aleutian Islands area (5AAC01.366).⁴⁸ Subsistence salmon harvests are monitored annually only in the Unalaska and the Adak districts, where a permit is required for harvest. A permit is not required for subsistence salmon fishing in the waters fished by the communities of Akutan, Atka, Nikolski, or Adak; therefore, subsistence salmon harvests are not systematically monitored in these communities. Harvest estimates for Akutan are based upon data in Davis (2005) and from household surveys conducted in 2009 for the 2008 harvest year. Estimates for Atka and Nikolski are based upon data in Davis (2005). There are no native populations of salmon in the

45. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. “Research and Analysis Homepage.” Accessed July 2014. <http://labor.alaska.gov/research/pop/popest.htm>

46. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. “Research and Analysis Homepage.” Accessed July 2014. <http://labor.alaska.gov/research/pop/popest.htm>

47. U.S. Census Bureau, Washington D.C. n.d. “American FactFinder.” U.S. Department of Commerce. Accessed May 2014. <http://factfinder2.census.gov/faces/nav/jsf/pages/index.xhtml>

48. Alaska Department of Fish and Game. 2012–2013 Subsistence and personal use statewide fisheries regulations. Alaska Department of Fish and Game, Juneau.

Pribilof Islands, and therefore there are no local subsistence salmon fisheries available for the communities of St. Paul and St. George.

SALMON HARVESTS IN THE UNALASKA DISTRICT

The Unalaska District includes all waters west of Akutan Pass up to, and including, Umnak Pass (5 AAC 12.200 (b)).

Salmon Harvest Regulations

A permit is required for subsistence salmon fishing in the Unalaska District. Fishers must record their daily harvests on the permit and return it to ADF&G by October 31. Permit holders may harvest up to 25 salmon per permit plus an additional 25 salmon for each member of the same household who is listed on the permit. A permit holder may obtain an additional permit from the department if more fish are needed. A record of subsistence-caught fish must be recorded on the reverse side of the permit and the permit must be returned to the department by October 31, even if no salmon were harvested (5 AAC 01.380).

Other standard permit conditions include prohibition of fishing within 300 ft of a dam, fish ladder, weir, culvert, or other artificial obstruction. Salmon may be taken from 6:00 AM until 9:00 PM beginning January 1 through December 31, except that from June 1 through September 15, a salmon seine vessel may not be used to take salmon for subsistence purposes 24 hours before, during, or 24 hours after an open commercial fishing period within a 50-mi radius of the area open to commercial fishing. Salmon may be taken by seine or gillnet, but from June 1 through September 15, a purse seine vessel may be used to take subsistence salmon only with a gillnet. In the Unalaska District, subsistence gillnets must be attended at all times while fishing. (5 AAC 01.360–5 AAC 01.370). Waters within the Unalaska District that are closed to subsistence fishing for salmon are defined in 5 AAC 01.375.

Salmon Harvest Assessment Program

The Division of Commercial Fisheries has issued subsistence salmon harvest permits for the Unalaska District since 1979. Permits are only issued in person at the ADF&G Dutch Harbor office. Unalaska District permits are required by regulation to be returned by October 31; they may be returned in person or mailed to the ADF&G Dutch Harbor office. Reminder letters are sent on approximately November 1 to all permit holders who have not returned their permits. Data from returned permits are tabulated by species and fishing area. Harvest estimates are calculated by expanding reported harvest numbers from successfully and unsuccessfully fished permits to represent fish taken by all permit holders, including those who did not return their permits (Wilburn and Nichols 2013). Federal subsistence fisheries are authorized for permanent residents residing in the Aleutian Islands Area; however they are managed consistently with the State fisheries in the region.⁴⁹

Subsistence Salmon Harvests in 2012

In 2012, 211 subsistence salmon permits were issued for the Unalaska District which was fewer than the previous year, 2011, when 230 were issued, but similar to the to the recent 5-year (2007–2011) average of 208 permits (and 10-year (2002–2011) average of 212 permits issued (Table 9-1). This number was also higher than the historical annual average (1985–2011) of 169 permits. Harvest numbers are recorded on the permit and returned at the end of the harvest season to ADF&G. In 2012, the return rate for the Unalaska District was 80%, with 169 permits returned out of 211 permits issued. Dutch Harbor and

49. Additional information about the federal subsistence fishery is available by contacting the United States Fish and Wildlife Service, Office of Subsistence Management in Anchorage, AK (<http://www.doi.gov/subsistence/index.cfm>).

Unalaska residents accounted for 205, or 97%, of all permits issued in the Unalaska District, and returned 163 permits out of 169 permits (96%) (Table 9-2).

The estimated subsistence harvest of salmon in the Unalaska District in 2012 was 5,790 fish, which was 454 salmon less than the previous year (6,244), but higher than the recent 5-year average (4,648 fish) and the 10-year average (4,944 fish) for the district (Table 9-1). The composition of the 2012 subsistence salmon harvest was sockeye (86%, similar to 2011 (88%), coho (7%, up from 5% in 2011), pink (6% similar to 5% in 2011),), chum (1%), and Chinook (<1%) salmon (Figure 9-1).

In 2012, the primary subsistence salmon fishing locations used in the Unalaska District occurred primarily in Reese Bay where sockeye salmon are targeted as they are migrating to McLees Lake. Subsistence salmon harvests harvested in Reese Bay in 2012 was estimated to be 4,347 fish (92% of the total sockeye salmon harvests for the Unalaska District (Wilburn and Nichols 2013).

In interviews with Division of Subsistence personnel, ADF&G fishery managers expressed the view that the permit program captured most subsistence salmon harvests occurring in the Unalaska District (Fall and Shanks 2000). In their view, most subsistence fishers likely obtained permits, perhaps due to the presence of Alaska Wildlife Troopers from the Alaska Department of Public Safety as well as a population that is self-enforcing (likely to report violators). Fishery managers in the Unalaska District believe that few commercially caught salmon are retained for subsistence purposes in the Aleutian Islands Area since most commercial fishing activities in the area target shellfish and groundfish rather than salmon. A 1994 survey of randomly selected Unalaska households conducted by the Division of Subsistence supports this view: it found that 4% of all salmon harvested for home uses were removed from commercial catches, 62% were harvested with noncommercial nets, and 34% with rod and reel (CSIS).

SALMON HARVESTS IN THE ADAK DISTRICT

The Adak District of the Aleutian Islands Area consists of waters west of Atka Pass at 175°23.00' west longitude to the terminus of the Aleutian Islands. Adak Island hosted a U.S. Navy base and military community (population of 4,633 in 1990) that was phased out between 1993 and 1996. With the Navy base closure complete, the population was estimated at 0 in 1997; however, for several years following the base closure, the Aleut Corporation worked with the Department of Interior and Department of the Navy to lease the base facilities and ultimately secure a land transfer to the corporation in 2004, with the intention of repurposing and privatizing the facility's infrastructure to support the local fisheries industry with processing, refueling, and housing services. During the lease period, the Aleut Corporation processed its first commercially caught fish in 1998 and has continued working with the Alaska seafood industry to further develop and maintain Adak as a commercial fish processing location. In 2000, the Alaska Boundary Commission approved Adak's application to become a second-class city. In 2012, Aleut Enterprises and the Aleut Corporation owned and operated a seafood processing facility, and a few Adak residents held commercial fishing permits. In addition, Adak provides a fueling port and crew transfer facility for foreign fishing fleets.^{50,51} Adak's estimated population was 316 in 2000⁵² and 331 in 2010, with 21 students attending the Adak school. The estimated population for Adak in 2012 was 321 of which 217 resided in group quarters.⁵³

50. Gen. Hansford T. Johnson, USAF. 2002. Statement of H.T. Johnson Assistant Secretary of the Navy (Installations and Environment) Before the Subcommittee on Public Lands and Forests of the Senate Committee on Energy and Natural Resources To Ratify an Agreement Between the Aleut Corporation And the United States of America. Washington, D.C. Accessed September 2014. http://www.navy.mil/navydata/people/assistsecnav/asn_ie/htjohnson020509.txt.

51. Adak Update: Adak Land Transfer Fact Sheet. 2004. Engineering Field Activity (EFA) Northwest, Naval Facilities Engineering Command. Accessed September 2014. <http://www.navfac.navy.mil/content/dam/navfac/PDFs/factsheets/adak-alaska.pdf>.

52. U.S. Census Bureau, Washington D.C. n.d. "American FactFinder." U.S. Department of Commerce. Accessed May 2014. <http://factfinder2.census.gov/faces/nav/jsf/pages/index.xhtml>

53. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. "Research and Analysis Homepage: 2010 census: demographic profiles." Accessed May 2014. <http://live.laborstats.alaska.gov/cen/dparea.cfm>

Salmon Harvest Regulations

Prior to 1988, the noncommercial salmon net fishery at Adak was classified as a subsistence fishery, then a personal use fishery in 1988, followed by a return to a subsistence classification in 1998.

Subsistence regulations in place since 2001 require that fishers obtain a permit from ADF&G. Fishers must record their daily harvests on the permit, and return it to ADF&G by October 31. Permit holders may harvest up to 25 salmon per permit, plus an additional 25 salmon for each household member listed on the permit. Other standard permit conditions include prohibition of fishing within 300 ft of a dam, fish ladder, weir, culvert, or other artificial obstruction. Also, as specified in 5 AAC 01.380 (b)(2), “a permit holder may obtain an additional permit from the department to harvest more salmon.” Salmon may be taken at any time. All salt waters within 100 yards of a stream terminus, as well as all fresh waters of and around Adak Island and Kagalaska Island, are closed to subsistence fishing for salmon (5 AAC 01.375 (6)).

Salmon Harvest Assessment Program

Subsistence salmon permits are issued by the ADF&G Cold Bay office and are faxed upon request to Adak residents. Permits must be returned by mail or fax to the ADF&G Cold Bay office by October 31, after which reminder letters are sent to those who have yet to report their harvests. ADF&G fishery managers believe that the program provides reliable data on subsistence salmon fishing effort and harvests at Adak.

Subsistence Salmon Harvests in 2012, Adak District

Only two subsistence salmon permits were issued for the Adak District in 2012. This was fewer than the 5-year (4) and 10-year (4) averages, and also fewer than the historical 1988–2011 average (17) (Table 9-3). In 2012, both permits issued were issued to residents of Adak (Table 9-4). The total estimated harvest in 2012 was 25 salmon, all sockeye (Table 9-3). The 2012 estimated harvest based on permits issued and returned was more than the previous year where 0 permits were issued, similar to 2009 and 2010, when 1–2 permits were issued with estimated harvests from 25–50, but much less than in 2008 where 10 permits were issued with an estimated harvest of 400 salmon. The recent 5-year average of total salmon harvests (2007–2011) was 175 salmon, most of which were sockeye (166), followed by pink salmon. For the period 1988–1993, during the Navy’s occupation of their base at Adak, an average of 49 personal use permits were issued and the average annual harvest was 611 salmon, with an average of 529 sockeye, 62 pink, 20 coho and 0 Chinook harvested (Table 9-3).

SALMON HARVESTS AT AKUTAN, NIKOLSKI, AND ATKA

Permits are not required for subsistence salmon harvests in the Akutan, Umnak, and Atka–Amlia islands districts, and there are no annual harvest assessment programs in place. In these districts, no more than 250 salmon may be taken annually for subsistence purposes (5 AAC 01.380).

The Division of Subsistence conducted postseason household interviews in Akutan (Akutan District) and Nikolski (Umnak District) pertaining to 1991 subsistence harvests (all resources); again in Akutan pertaining to 2008 subsistence harvests (all resources); and in Atka (Atka–Amlia Islands District) pertaining to harvests in 1992 (salmon only), and 1994 (all resources). Salmon harvest data were also collected for Akutan and Nikolski (2002 and 2003 harvests) and Atka (2003 harvests) as part of the project reported in Davis (2005). In most years, subsistence harvests of salmon in Akutan, Nikolski, and Atka are primarily composed of sockeye salmon, but coho and pink salmon also account for a relatively large proportion of yearly harvests (Table 9-5). Subsistence salmon harvests in Akutan totaled 3,268 fish in 1991, decreasing to 1,070 fish in 2002 and 1,675 fish in 2003. In 2008, Akutan harvests totaled nearly the same as in 1991 with a total of 3,363 salmon; with sockeye (1,489) and pink salmon (1,366) harvests being near equivalent. Yearly salmon harvests in Nikolski also presented an apparent decreasing pattern, with 1,902 fish caught in 1991 and 604 fish in 2003; further data collection and analysis is necessary to

confirm the trend. In Atka, the yearly salmon harvest varied between 1,454 and 2,387 in the 3 years for which information is available (Table 9-5).

OTHER SUBSISTENCE FISHERIES IN THE ALEUTIAN ISLANDS AREA

Finfishes

Harvest estimates of subsistence halibut for the Aleutian Islands Area are available for 2012 (Fall and Koster 2014).

There are no annual harvest assessment programs for other subsistence finfish fisheries of the Aleutian Islands Area. Permits are required for the taking of rainbow/steelhead trout and Arctic char/Dolly Varden, but no harvest reporting program is in place. Fish other than salmon may be taken by gear specified in 5 AAC 01.010, except that under state regulations, halibut may be taken only by a single handheld line with no more than 2 hooks attached, while federal rules allow up to 30 hooks using a longline (skate). The Division of Subsistence has conducted systematic household surveys pertaining to a single year's harvests in Akutan (1991; 2008), Atka (1994), Nikolski (1991), Saint George (1994), Saint Paul (1994), and Unalaska–Dutch Harbor (1994). Results, including harvest estimates for finfishes, are available in the CSIS.

Shellfish

Permits for the taking of shellfish for subsistence purposes are required only for king and Tanner crabs in that portion of the Alaska Peninsula–Aleutian Islands Area west of Scotch Cap Light and east of 168° west longitude. Estimates of subsistence harvests of all marine invertebrates for single study years, based on systematic household surveys, are available in the CSIS.

Table 9-1.—Historical subsistence salmon harvests, Unalaska District, 1985–2012.

Year	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1985	65	22	0	897	208	20	1,293	2,418
1986	121	28	0	3,449	847	375	2,468	7,139
1987	81	49	0	1,097	378	151	1,780	3,406
1991	77	45	3	966	390	83	2,627	4,069
1989	74	42	2	1,112	470	36	1,292	2,912
1990	94	37	4	2,357	681	100	1,428	4,570
1991	89	48	0	1,294	666	45	1,075	3,080
1992	144	102	7	2,739	587	11	1,723	5,067
1993	139	102	17	2,831	697	136	587	4,268
1994	150	120	1	2,759	774	48	1,053	4,635
1995	160	129	23	4,484	484	23	791	5,805
1996	189	123	5	1,107	1,033	49	492	2,686
1997	221	163	8	4,192	864	110	554	5,728
1998	206	161	4	3,317	731	26	729	4,807
1999	208	154	0	2,485	1,234	16	1,044	4,779
2000	212	167	10	3,935	603	26	580	5,154
2001	204	165	6	4,202	724	77	784	5,793
2002	231	180	3	5,678	707	65	385	6,837
2003	227	179	25	5,124	572	40	378	6,139
2004	208	170	7	4,713	955	26	437	6,139
2005	217	152	8	4,066	424	14	527	5,038
2006	199	159	15	2,007	422	74	675	3,193
2007	178	126	14	2,575	254	42	683	3,569
2008	204	161	2	1,676	828	90	660	3,257
2009	210	130	5	3,171	616	182	443	4,416
2010	216	170	1	3,883	319	71	336	4,611
2011	230	156	8	5,525	303	65	343	6,244
2012	211	169	20	4,960	429	43	338	5,790
5-year average (2007–2011)	208	149	6	3,366	464	90	493	4,648
10-year average (2002–2011)	212	158	9	3,842	540	67	487	4,944
Historical average (1985–2011)	169	120	7	3,024	621	74	932	4,658

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Table 9-2.—Estimated subsistence salmon harvests by community, Unalaska District, 2012.

Community	Permits		Estimated salmon harvest					Total
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	
Dutch Harbor	111	93	14	2,847	136	7	75	3,079
Fairbanks	1	1	0	0	0	0	0	0
Homer	1	1	0	0	0	0	0	0
Kodiak City	2	2	0	6	11	0	0	17
Soldotna	1	1	0	0	0	0	0	0
Unalaska	94	70	5	2,107	282	36	263	2,694
Wasilla	1	1	0	0	0	0	0	0
Total	211	169	20	4,960	429	43	338	5,790

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Table 9-3.—Historical subsistence and personal use salmon harvests, Adak District, 1988–2012.

Year ^a	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1988	43	29	0	503	23	0	150	676
1989	64	47	0	382	0	0	117	499
1990	61	29	0	800	47	0	41	888
1991	37	31	0	281	6	0	34	321
1992	52	41	0	572	30	0	4	606
1993	36	26	0	638	12	0	26	676
1994 ^b	0	0	0	0	0	0	0	0
1995	4	3	0	156	0	0	0	156
1996	6	6	0	91	0	0	0	91
1997 ^c	18	12	0	229	0	4	0	233
1998	13	10	0	399	0	0	25	424
1999	5	5	0	164	4	0	0	168
2000	13	13	0	270	4	0	75	349
2001	17	15	14	489	18	0	16	537
2002	3	3	0	150	0	0	0	150
2003	6	5	0	338	0	0	0	338
2004	6	4	0	336	0	0	0	336
2005	2	2	0	188	0	0	0	188
2006	1	1	0	74	0	0	1	75
2007	9	8	0	367	2	0	29	398
2008	10	8	0	386	0	0	14	400
2009	1	1	0	25	0	0	0	25
2010	2	1	0	50	0	0	0	50
2011	0	0	0	0	0	0	0	0
2012	2	2	0	25	0	0	0	25
5-year average (2007–2011)	4	4	0	166	0	0	9	175
10-year average (2002–2011)	4	3	0	191	0	0	4	196
Historical average (1988–2011)	17	13	1	287	6	0	22	316

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

- a. Personal use fishery 1988 to 1997; subsistence fishery 1998 to present.
- b. Navy presence at Adak was reduced beginning in 1994; no requests for permits that year.
- c. In 1997, a number of civilians were hired to work on a clean-up effort at Adak.

Table 9-4.—Estimated subsistence salmon harvests by community, Adak District, 2012.

Community	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
Adak	2	2	0	25	0	0	0	25
Total	2	2	0	25	0	0	0	25

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Table 9-5.—Estimated subsistence harvests of salmon by residents of Akutan, Atka, and Nikolski.

Community	Year	Estimated number of harvesting households	Estimated salmon harvest ^a						Total
			Chinook	Sockeye	Coho	Chum	Pink	Other— unknown	
Akutan	1991	24	10	1,872	429	36	915	6	3,268
Akutan	2002	NA	0	809	147	44	70	0	1,070
Akutan	2003	NA	3	1,270	127	0	275	0	1,675
Akutan	2008	21	2	1,489	452	54	1,366	0	3,363
Atka	1992	18	4	502	465	24	459	0	1,454
Atka	1994	23	10	394	583	133	1,267	0	2,387
Atka	2003	NA	8	1,187	333	0	264	0	1,792
Nikolski	1991	12	0	957	547	54	327	17	1,902
Nikolski	2002	NA	0	312	643	0	182	0	1,137
Nikolski	2003	NA	12	287	270	0	35	0	604

Sources ADF&G Division of Subsistence household surveys, (ADF&G 2009), Davis (2005).

Note NA indicates the estimated number of harvesting households cannot be calculated using available data.

- a. Includes harvests for home uses by all methods, including subsistence nets, rod and reel, and removal from commercial harvests.




Figure 9-1.—Composition of Unalaska District estimated subsistence salmon harvest by species, 2012.

CHAPTER 10: KODIAK AREA

INTRODUCTION

The Kodiak Area encompasses the waters of the Gulf of Alaska surrounding the Kodiak Archipelago and those waters along that portion of the Alaska Peninsula that drains into Shelikof Strait (Figure 10-1). The portion of the Kodiak Island Borough's population living along the island's road system is the largest rural community in Alaska (as defined by the Federal Subsistence Board [FSB]) and the largest community outside the nonsubsistence areas defined by the Joint Board (Figure 10-1). The population of the Kodiak Island Borough according to the State of Alaska Department of Labor and Workforce Development (14,030 in 2012) comprises all individuals residing on Kodiak Island; however this population is often distinguished by which communities have access to the road system. Communities along the Kodiak Island road system include Kodiak City (6,425), the U.S. Coast Guard Base (1,295), Womens Bay (762), Chiniak (44), and the remainder of the road-accessible Kodiak Island Borough (this includes all residents of Kodiak Island who are on the road system but are not identified within the population of a census designated place [CDP] or city) (4,661). Communities on Kodiak Island that are located outside the range of the road system include Akhiok (87), Aleneva CDP (37), Karluk (42), Larsen Bay (92), Old Harbor (206), Ouzinkie (178), and Port Lions (201).⁵⁴

SALMON HARVEST IN THE KODIAK AREA

Salmon Harvest Regulations

Permits have been required to harvest salmon for subsistence purposes in the Kodiak Area since 1962. Since 1990, all Alaska state residents have been eligible to participate in subsistence salmon fishing in the Kodiak Area under state regulations. In 2012, legal gear for subsistence salmon fishing under state regulations included gillnets (maximum length 50 fathoms) and seines. Fishers were required to physically attend their net while fishing and should always have a valid subsistence salmon permit with them while fishing for salmon and record the numbers of all fish harvested on the permit before concealing the fish from plain view or transporting them from the harvest area. Generally, fishing was open year-round from 6:00 AM to 9:00 PM daily. From June 1 through September 15, commercial purse seine vessels may be used for subsistence fishing only before June 1 and after September 15. Purse seines could not be used for subsistence salmon fishing 24 hours before, during, and 24 hours after any period open for commercial salmon fishing. Only gillnets could be operated for subsistence purposes from purse seine vessels between the dates of June 1 and September 15, and only when no other salmon fishing gear was on board. Permits allowed individual fishers to harvest 25 salmon for their own use plus 25 additional salmon for each member of the permit holder's household. An additional permit could be obtained if the fisher could demonstrate a need for more fish. Permit holders were required to keep a record of their harvests on the permit. A list of waters closed to subsistence fishing within the Kodiak Area appears in 5 AAC 01.525 and 5 AAC 01.530.

In 2012, federal regulations governing subsistence salmon fishing in waters under jurisdiction of the FSB were generally identical to the state regulations summarized above, except that rod and reel (in addition to gillnets and seines) was legal subsistence gear under federal rules. Another difference was that federal regulations allowed subsistence salmon fishing 24 hours per day, while state regulations limited subsistence fishing to the hours of 6:00 AM to 9:00 PM daily. There was no separate federal subsistence permit; a state permit was required for subsistence fishing in waters under federal jurisdiction.

54. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. "Research and Analysis Homepage." Accessed July 2014. <http://labor.alaska.gov/research/pop/popest.htm>

Salmon Harvest Assessment Program

Staff in the Division of Commercial Fisheries' Kodiak office manage the subsistence salmon harvest assessment program for the Kodiak Area. Permits are mailed each year to people who turned in their permits at the end of the previous fishing season. People may request subsistence permits by mail or in person at the Kodiak ADF&G office. In June 2001, staff from the Division of Commercial Fisheries and the Division of Subsistence visited 6 communities off the road system in the Kodiak Island Borough (Akhiok, Karluk, Larsen Bay, Old Harbor, Ouzinkie, and Port Lions) to implement an area permit vendor program. A resident in each community was trained to issue subsistence fishing permits. Outreach activities were conducted in each community to encourage subsistence fishers to obtain permits, record their harvests, and return the permits at the end of the season. Research conducted in 2013 and 2014 by Division of Subsistence researchers showed that outreach in regards to subsistence fishing regulations and permitting is again needed in the smaller communities on Kodiak Island. Researchers witnessed a great deal of confusion surrounding subsistence regulations and the permit system. Area managers were contacted so researchers could provide accurate answers to subsistence fishers' questions. Subsistence fishers mail permits with their harvest record to ADF&G at the end of the season or return them in person at the Kodiak ADF&G office. ADF&G sends reminder letters in February to permit holders who have not returned their permits.

Subsistence Salmon Harvests in 2012

In the Kodiak Area, ADF&G sends permits to every permit holder who returned a permit in the previous year. The U.S. Postal Service returns a number of permits to ADF&G marked "undeliverable." No record is maintained regarding the number of "undeliverable" permits—as a result, the actual number of permits issued remains unknown. For this reason, harvest reports have not been expanded for this area since 1999 (Table 10-1). Results of the harvest monitoring program therefore reflect only the reported harvests of subsistence fishers who returned permits.

In 2012, 1,866 subsistence permits with harvest information were returned to ADF&G (tables 10-1 and 10-2). Of these, 1,558 (83%) were returned by residents of Kodiak Island Borough, 302 (16%) were returned by residents of other Alaska communities, and 6 (<1%) had been issued to Alaska residents who were serving in the military outside of the state. Following a well-established trend, permit holders with addresses in Kodiak Island Borough accounted for the majority (83%) of all permits returned for 2012 (Table 10-2).

The total reported subsistence salmon harvest for the Kodiak Area in 2012 was 28,159 fish, which is lower than the recent 5-year (2007–2011) average of 30,339 salmon, and the 10-year (2002–2011) average of 34,188 salmon (Table 10-1). Of the total harvest, 26,987 salmon (96%) were harvested by residents of Kodiak Island Borough communities and 1,135 salmon (4%) were harvested by permit holders in other Alaska communities (Table 10-2). Of the 26,987 salmon harvested by Kodiak Island Borough residents, 22,048 fish (82%) were taken by residents living along the Kodiak Island road system (figures 10-1 and 10-2). The Kodiak Island road system includes Kodiak City, the U.S. Coast Guard Base, Womens Bay, Chiniak, and the remaining residents who live along the Kodiak Island road system but who are not identified within the population of a CDP or city. This is consistent with the pattern between 2000 and 2009 when 72% to 83% of all salmon harvested by Kodiak Island Borough residents for subsistence purposes was taken by residents of Kodiak City and areas along the road system. Comparatively, the 6 villages and other populated remote locations that do not have access to the road system surrounding Kodiak City harvested 4,939 salmon in 2012 (Table 10-2).

In 2012, the Kodiak Area subsistence salmon harvest was composed of 85% sockeye salmon, 10% coho salmon, 4% pink salmon, 1% chum salmon, and <1% Chinook salmon (Figure 10-3). The commercial harvest retained for home use was different in 2012 in terms of the composition of the harvest. As shown in Figure 10-4, in 2012, 7,734 salmon, including 1,971 coho (26%), 1,421 pink (18%), 4,116 sockeye (53%), 195 Chinook (3%), and 31 chum salmon (<1%), were retained from commercial harvests for

home use (Jackson et al. 2012:38). The total number of sockeye salmon retained from commercial harvests for personal use in 2012 was 3 times as large as it was in 2011. The greatest decrease overall was for the harvests of pink salmon retained from commercial boats (Jackson et al. 2012:38).

In 2001, interviews were conducted with Division of Subsistence staff and fishery managers within the Division of Commercial Fisheries. During interviews, fishery managers expressed uncertainty regarding the accuracy of subsistence salmon harvest data collected through the Kodiak Area permit program. ADF&G staff suspected that a substantial amount of subsistence harvests occurred without permits, especially in areas off the Kodiak Island road system. Subsistence salmon harvest estimates for the Kodiak Area based on household harvest surveys and reported in the CSIS were substantially higher than harvests reported in the FMRs. Delivery of permits to subsistence fishers living in communities outside of the road system, including Akhiok, Karluk, Larsen Bay, Old Harbor, Ouzinkie, and Port Lions, has proven problematic in the past. As mentioned above, an outreach effort and an area permit vendor program were implemented in 2001 to address this issue. These actions appeared to result in increased participation in the permit program in these 6 communities. A total of 100 permits were returned in 2000; from 2001 through 2006 between 189 and 143 permits were returned (Table 10-3). Accordingly, the yearly reported subsistence salmon harvest also fluctuated between 2000 and 2006 with the lowest number harvested being 6,299 fish in 2000 and the highest number being 10,172 fish in 2005. The most recent years of 2007–2010 have marked the lowest reported salmon harvests, the lowest being 5,138 in 2007 and the highest being 5,896 in 2010. In 2012, both the number of permits returned by the 6 villages (112 permits) and the number of harvested salmon reported (4,939 fish) were the closest to data for 2000, which was prior to the implementation of the local permit vendor program and the outreach effort (Table 10-3). Due to lack of funding, in 2012 a limited local vendor program only was in place in Ouzinkie, Larsen Bay, and possibly Port Lions (personal communication, Amanda Dorner, Division of Commercial Fisheries, Kodiak). Likewise, no outreach efforts occurred in the small communities on Kodiak Island in 2012. Additional research and outreach are needed to assess the most recent harvest data. In 2013, the Division of Subsistence conducted subsistence salmon harvest surveys with residents of Larsen Bay, Old Harbor, and Kodiak City and communities along surrounding road system to address this need for current data and community outreach. This data are currently in the analysis stage with the Division of Subsistence Information Management team, and a final report is expected to be published in 2015.

Household surveys (with results reported in the CSIS) have documented noncommercial salmon harvests with rod and reel gear, which is legal subsistence gear under federal subsistence regulations but not under state regulations. Household surveys also document the number of salmon removed from commercial harvests for personal use. Information on these 2 types of harvests not documented by the permit program is needed for a better understanding of the household salmon harvest in the Kodiak Area. This need was addressed by Division of Subsistence researchers when administering 2013 subsistence salmon harvest surveys.

In early 2004, the Division of Subsistence and the Kodiak Area Native Association (KANA) conducted comprehensive household surveys in Akhiok, Larsen Bay, Old Harbor, Ouzinkie, and Port Lions in the context of a project funded by the *Exxon Valdez* Oil Spill Trustee Council. Among other objectives, this project provided updated harvest data for salmon, nonsalmon finfishes, and marine invertebrates (Fall 2006a).

In spring 2013, the Division of Subsistence conducted comprehensive household harvest surveys in the communities of Kodiak City (along the road system), Old Harbor and Larsen Bay. The purpose of this project is to update the existing subsistence salmon harvest data for these communities. The project is currently in the data analysis phase.

OTHER SUBSISTENCE FISHERIES IN THE KODIAK AREA

Finfishes

Federal halibut subsistence harvest data are currently available for communities and tribes in the Kodiak Area. For the findings for 2012, see Fall and Koster (2014).

There are no annual harvest assessment programs for other subsistence finfish fisheries in the Kodiak Area. Harvest estimates based on comprehensive household surveys conducted by the Division of Subsistence are available in the CSIS for freshwater and marine species spanning multiple years for each Kodiak Island Borough community. Fish harvested in the largest quantities and used by the majority of households include Pacific cod, lingcod *Ophiodon elongatus*, various species of flounders, Pacific halibut, rockfishes, and Arctic char/Dolly Varden.

Shellfish

Subsistence permits are required for the harvest of king, Tanner, and Dungeness crabs in the Kodiak Area (5 AAC 02.410). Regulations establish sex, size, and bag and possession limits for these species of crabs. Only male crabs may be taken. Other marine invertebrates used for subsistence purposes in the Kodiak Area include clams, cockles, mussels, chitons, octopuses, sea urchins, and more.

Table 10-1.—Historical subsistence salmon harvests, Kodiak Area, 1986–2012.

Year	Permits		Reported salmon harvest ^a					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1986	1,244	1,002	90	14,391	6,998	605	2,371	24,455
1987	1,124	880	101	13,198	6,463	1,299	2,421	23,482
1988	1,098	699	108	10,081	4,291	377	1,320	16,177
1989	2,800	717	43	12,638	4,123	419	1,553	18,776
1990	2,900	1,167	131	17,959	8,627	655	1,605	28,977
1991	1,406	1,225	177	21,835	8,208	714	1,743	32,677
1992	1,561	1,195	318	20,684	8,643	643	1,646	31,934
1993	1,496	959	243	19,471	7,176	838	2,696	30,424
1994	2,550	1,464	205	17,962	7,491	440	1,758	27,856
1995	1,950	1,194	175	19,416	5,603	293	1,548	27,035
1996	1,567	1,390	253	28,287	5,117	381	1,125	35,163
1997	2,098	1,638	383	33,293	6,369	234	1,458	41,737
1998	1,841	1,126	350	20,459	5,348	214	1,412	27,783
1999	ND	1,438	397	26,497	4,932	388	1,266	33,480
2000	ND	1,376	273	24,873	5,399	341	742	31,628
2001	ND	2,153	273	33,833	5,920	427	1,158	41,611
2002	ND	2,271	593	32,977	6,057	350	1,665	41,642
2003	ND	2,275	500	32,104	6,096	384	1,484	40,568
2004	ND	2,240	379	30,217	5,819	261	1,395	38,071
2005	ND	1,900	431	27,002	7,447	592	2,343	37,815
2006	ND	1,906	280	22,905	6,640	441	1,827	32,093
2007	ND	2,118	207	24,556	4,630	240	1,532	31,165
2008	ND	1,637	151	20,809	4,336	168	1,128	26,592
2009	ND	1,737	159	21,852	4,570	186	1,180	27,947
2010	ND	1,890	158	22,170	4,200	273	1,266	28,067
2011	ND	1,996	122	34,037	2,367	198	1,199	37,923
2012	ND	1,866	54	23,865	2,920	166	1,154	28,159
5-year average (2007–2011)	ND	1,876	159	24,685	4,021	213	1,261	30,339
10-year average (2002–2011)	ND	1,997	298	26,863	5,216	309	1,502	34,188
Historical average (1986–2011)	ND	1,523	250	23,212	5,880	437	1,571	31,349

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Note ADF&G sends permits to every permit holder who returned a permit in the previous year. The U.S. Postal Service returns a number of permits to ADF&G marked “undeliverable.” No record is maintained regarding the number of “undeliverable” permits. As a result the actual number of permits issued remains unknown (ND). For this reason, harvest reports have not been expanded.

Table 10-2.—Reported subsistence salmon harvests by community and species, Kodiak Area, 2012.

Community	Permits returned	Reported salmon harvest ^a					Total
		Chinook	Sockeye	Coho	Chum	Pink	
Kodiak Island Borough							
Akhiok	3	2	128	21	0	7	158
Chiniak	28	0	213	57	1	10	281
Karluk	2	0	33	6	0	0	39
Kodiak (city)	1,418	47	19,223	1,887	81	529	21,767
Larsen Bay	17	1	387	27	0	16	431
Old Harbor	20	3	604	248	77	253	1,185
Ouzinkie	31	1	954	438	5	71	1,469
Port Lions	39	0	1,261	160	0	236	1,657
Subtotal, Kodiak Island Borough	1,558	54	22,803	2,844	164	1,122	26,987
Other Alaska							
Anchor Point	4	0	21	0	0	0	21
Anchorage	123	0	488	25	1	20	534
Bethel	2	0	21	0	0	5	26
Bettles	1	0	0	0	0	0	0
Big Lake	2	0	0	0	0	0	0
Central	1	0	0	0	0	0	0
Chignik Lagoon	1	0	0	0	0	0	0
Chugiak	4	0	0	0	0	0	0
Cold Bay	1	0	0	0	0	0	0
Cordova	1	0	0	0	0	0	0
Douglas	1	0	4	0	0	0	4
Eagle River	17	0	3	0	0	0	3
Fairbanks	18	0	294	0	1	1	296
Girdwood	3	0	24	0	0	0	24
Glennallen	1	0	0	0	0	0	0
Homer	20	0	34	13	0	2	49
Hope	1	0	0	0	0	0	0
Juneau	3	0	33	5	0	0	38
Kasilof	5	0	0	0	0	0	0
Kenai	5	0	10	0	0	0	10
Kotzebue	0	0	0	0	0	0	0
Nikiski	1	0	0	0	0	0	0
Ninilchik	2	0	0	0	0	0	0
Nome	1	0	4	0	0	1	5
North Pole	4	0	0	0	0	0	0
Palmer	14	0	0	6	0	2	8
Sand Point	1	0	0	0	0	0	0
Seldovia	2	0	0	0	0	0	0
Seward	8	0	0	0	0	0	0
Sitka	2	0	0	0	0	0	0
Soldotna	22	0	24	21	0	1	46
Sterling	3	0	0	0	0	0	0
Tok	1	0	0	0	0	0	0

-continued-

Table 10-2.—Page 2 of 2.

Community	Permits returned	Reported salmon harvest ^a					
		Chinook	Sockeye	Coho	Chum	Pink	Total
Valdez	1	0	0	0	0	0	0
Wasilla	25	0	71	0	0	0	71
Willow	1	0	0	0	0	0	0
Subtotal, other Alaska	302	0	1,031	70	2	32	1,135
Other USA ^b	6	0	31	6	0	0	37
Total	1,866	54	23,865	2,920	166	1,154	28,159

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

- a. ADF&G sends permits to every permit holder who returned a permit in the previous year. The U.S. Postal Service returns a number of permits to ADF&G marked “undeliverable.” No record is maintained regarding the number of “undeliverable” permits. As a result the actual number of permits issued remains unknown (ND). For this reason, harvest reports have not been expanded.
- b. These are Alaska residents serving in the military who had a mailing address outside the state.

Table 10-3.—Permits returned and salmon harvests reported by the villages of Akhiok, Karluk, Larsen Bay, Old Harbor, Ouzinkie, and Port Lions.

Year	Permits returned	Reported salmon harvest	Source
2000 ^a	100	6,299	(Fall et al. 2002:105)
2001	189	9,034	(Fall et al. 2003a:117)
2002	167	9,386	(Fall et al. 2003b:121)
2003	165	8,714	(Brown et al. 2005b:123)
2004	170	7,845	(Fall et al. 2007a:118)
2005	147	10,172	(Fall et al. 2007b:105)
2006	143	7,114	(Fall et al. 2009a:113)
2007	143	5,138	(Fall et al. 2009b:105)
2008	117	5,850	Fall et al. 2011:111
2009	118	5,824	Fall et al. 2012:119
2010	118	5,896	Table 10-2
2011	125	5,786	Table 10-2
2012	112	4,939	Table 10-2

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

- a. Local permit vendor program and outreach efforts implemented in 2000.




Figure 10-1.–Kodiak Area map, 2012.




Figure 10-2.—Subsistence salmon harvests by community, Kodiak Area, 2012.




Figure 10-3.—Composition of Kodiak Area subsistence salmon harvest by species, 2012.




Figure 10-4.—Salmon retained from commercial harvests for home use, Kodiak Area, 2012.

CHAPTER 11: COOK INLET AREA

INTRODUCTION

As shown in Figure 11-1, most of the waters of the Cook Inlet Management Area are within the Anchorage–Matsu–Kenai Nonsubsistence Area as established by the Joint Board [5 AAC 99.015 (3)]. Because subsistence fisheries are not permitted within nonsubsistence areas, noncommercial harvesting opportunities occur under sport, personal use, and educational fishing regulations. Commercial harvesters may retain finfish from lawfully taken commercial catch for home use (“home pack”). These fish are required to be reported on the commercial fish ticket, not on the subsistence salmon permit or personal use permit. In some parts of Alaska, in addition to gear authorized under subsistence fishing regulations, subsistence users report that substantial numbers of fish for home uses are taken with rod and reel (Fall, Turek, et al. 2009), which, in this area, is allowable gear under sport fishing regulations. Harvest summaries for the personal use, sport, educational, and commercial fisheries of the Upper Cook Inlet (UCI) Management Area can be found in annual management reports prepared by the ADF&G divisions of Sport Fish and Commercial Fisheries. A summary of the personal use salmon fisheries of the Cook Inlet Area follows the discussion of Cook Inlet subsistence fisheries.

Waters outside the nonsubsistence area include the Tyonek Subdistrict; the western portion of the Susitna River drainage; waters north of Point Bede that are west of a line from the easternmost point of Jakolof Bay and north of the westernmost point of Hesketh Island, including Jakolof Bay, and that are south of a line west of Hesketh Island; and those waters south of Point Bede which are west of the easternmost point of Rocky Bay, in Lower Cook Inlet.

Communities within the areas excluded from the nonsubsistence area include Skwentna (population 35 in 2012), Tyonek (population 171), Beluga (population 16), Seldovia (population 398 in the city and village CDP), Port Graham (population 169), and Nanwalek (formerly called English Bay, population 286). The population of the entire Cook Inlet area in 2012 was 449,103, including the Municipality of Anchorage (population 298,576), the Kenai Peninsula Borough (56,718), and the Matanuska-Susitna Borough (93,809). This represented 61% of the state’s total population in 2012.⁵⁵

PORT GRAHAM AND KOYUKTOLIK SUBDISTRICTS

History and Regulations

Subsistence regulations for this subsistence setnet fishery were first established by the BOF in 1980. The fishery is located along the southern shore of outer Kachemak Bay in the Port Graham and Koyuktolik subdistricts of the Southern District, and, beginning in 2002, the Port Chatham and Wind Bay subdistricts. Two predominately Alaska Native communities, Nanwalek and Port Graham, are located in the Port Graham Subdistrict. For detailed description of this subsistence fishery and other subsistence harvests and uses in Nanwalek and Port Graham, see Stanek (1985).

The fishery is open in the Port Graham and Koyuktolik subdistricts from April 1 through September 30 and in the Port Chatham and Windy Bay subdistricts from April 1 through August 1, from 10:00 PM Thursday to 10:00 AM Wednesday. The area open for the subsistence setnet fishery includes the entire shoreline of the subdistrict to a regulatory marker near the head of Port Graham Bay. There are no household bag or possession limits. The 3 primary species harvested include sockeye, pink, and coho salmon. The gear allowed includes set gillnets no longer than 35 fathoms, no deeper than 45 meshes, and no larger than a 6-in stretched mesh. Returns of sockeye salmon, which are a majority of the harvest in the subsistence fishery, have been poor for the past 20 years. However, in 2011 the return of sockeye

55. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. “Research and Analysis Homepage.” Accessed July 2014. <http://labor.alaska.gov/research/pop/popest.htm>

salmon counted at the English Bay weir continued to surpass the inriver goal (Hollowell et al. 2012:5). In 2012, the return and harvest rates dropped again, with 961 sockeye harvests reported by permit holders.

Harvest Assessment Methods

In the past, The Division of Subsistence issued household permits through cooperative agreements with the Port Graham and Nanwalek village councils. However, in 2012, responsibility for the distribution, collection, and summarizing of subsistence permits for the communities of Seldovia, Port Graham, and Nanwalek were transferred from the Division of Subsistence to the Division of Commercial Fisheries, Homer Office. When permits are issued, a separate monthly harvest calendar is also issued for recording daily household harvests. Home use salmon harvests by the 2 communities occur with the use of setnets and rod and reel gear. While the recording of harvests in the setnet fishery is mandatory, it is not in the rod and reel fishery. Therefore, fishers are asked to voluntarily record their rod and reel harvests on their harvest permits. In order to accommodate the recording of harvests in both fisheries, the recording device has 2 pages, 1 for each gear type, and is issued separately from the permit. Area assistants hired by each village council collect the calendars periodically throughout the season. Arctic char/Dolly Varden harvests are also recorded on the calendars.

Sockeye salmon returns to the English Bay lakes were severely depressed for much of the late 1980s and early 1990s, with runs failing to achieve minimum escapement goals for 9 consecutive years between 1985 and 1993. Returns in the late 1990s were enhanced as a result of a rehabilitation enhancement project initiated by ADF&G and subsequently run by the Nanwalek Salmon Enhancement Project in association with the Chugach Regional Resources Commission (CRRC) and the village of Nanwalek (Hammarstrom and Dickson 2006:62). Inseason escapement monitoring has taken place since 1994, with openings and closures in the subsistence and commercial fisheries controlled by emergency order. Inconsistent runs in recent years have been the result of disease outbreaks in the lake-rearing portion of the program and erratic adult behavior that caused difficulty in capturing broodstock (Hammarstrom and Dickson 2006:41). A newer hatchery at Port Graham, run by the Port Graham Hatchery Corporation, contributed 6% to the subsistence harvest in Lower Cook Inlet (Hammarstrom and Ford 2011:2) and in 2010 released more than 200,000 sockeye salmon fry into the English Bay Lakes system (Hammarstrom and Ford 2011:6).

Harvest Estimates for 2012

In 2012, responsibility for the distribution, collection, and summarizing of subsistence permits for the communities of Seldovia, Port Graham, and Nanwalek were transferred from the Division of Subsistence to the Division of Commercial Fisheries, Homer Office. The change in administration authority included some change in methodology for Port Graham and Nanwalek; the Division of Subsistence contracted local research assistants in the villages to ensure distribution and collection of permits. This approach was dropped by Commercial Fisheries and instead the permits were sent to the IRA Councils for distribution.

In 2012, estimated salmon harvests for home uses in the Port Graham and Koyuktoalik subdistricts totaled 1,912 salmon, including both subsistence setnet and reported rod and reel harvests (Table 11-1). The 2012 harvest was significantly lower than the historical average of 5,368 salmon.

In 2012, residents of Port Graham returned 7 permits and harvested 1,007 salmon (Table 11-2), a major decrease in harvest from the 2009 harvest of 2,265 salmon by 25 permit holders (Fall et al. 2012b). (Thirteen Port Graham households reported harvesting 1,117 salmon in 2011 [Fall et al. 2013:146]). Nanwalek residents returned 1 permit and harvested a total of 905 salmon in 2012, a major decrease from the 2011 harvest of 9,272 salmon by 40 permit holders. As shown in Table 11-2 and Figure 11-2, the combined harvest of the 2 communities of Nanwalek and Port Graham included 961 sockeye salmon, the species with the highest harvest (50% of the overall harvest), followed by coho salmon (414; 27%), pink salmon (482; 25%), chum salmon (31; 2%), and Chinook salmon (24; 1%). Sockeye salmon harvests decreased from 5,702 salmon in 2011 to 961 salmon in 2012 and are a major reason for the overall

decrease in salmon harvests. These extremely low participation rates likely account for the unusually low harvest estimates for 2012.

SELDOVIA SUBSISTENCE FISHERY

History and Regulations

The BOF established this subsistence set gillnet fishery in 1995. The fishery is located on the south side of Kachemak Bay, near Seldovia, which is in the Southern District of the Lower Cook Inlet Fisheries Management Area. The subsistence fishery operates in a split season. The spring fishery, open April 1–May 30, targets natural Chinook salmon migrating through Lower Cook Inlet. The fall fishery, open the first 2 weekends of August, targets coho salmon.

In the spring season, fishing is allowed during two 48-hour periods each week, while in the fall season, fishing is open continuously during the 2-day weekends. The BOF has set a guideline harvest level (GHL) of 200 Chinook salmon and an annual possession limit of 20 Chinook salmon per household. There are no seasonal limits for other salmon species.

The area open to subsistence set gillnetting includes those waters along the eastern shore of Seldovia Bay as well as a short stretch outside Seldovia Bay to the west of Point Naskowhak. Seasons and bag limits were designed in 1995 to reduce potential interceptions of enhanced Chinook salmon bound for the stocking site in the Seldovia small boat harbor (Hollowell et al. 2012:14). The gear allowed includes set gillnets no longer than 35 fathoms, no deeper than 45 meshes, and no larger than a 6-in stretched mesh. Other standard permit conditions include prohibition of fishing within 300 ft of a dam, fish ladder, weir, culvert, or other artificial obstruction.

Harvest Assessment Methods

Household permits are issued by ADF&G prior to fishing, and harvests are recorded on the permits. Permits are also available from the harbormaster in Seldovia. Fishers are required to telephone daily harvest numbers to ADF&G or the harbormaster as well as to return their permits after each of the 2 fisheries. ADF&G sends reminder letters to permit holders if harvest records have not been returned in a timely manner, and telephone calls are also made to enhance permit returns.

The 2012 Season

There were 20 permits issued for the Seldovia subsistence fishery in 2012; 7 were returned (Table 11-3). The estimated harvest was 79 sockeye salmon (56%), 54 pink salmon (39%), and 8 Chinook salmon (6%), and no reported coho, or chum salmon (Figure 11-3). Out of the 20 permits issued, 19 permits were issued to residents of Seldovia and one was issued in Anchor Point (Table 11-3).

Total salmon harvests in 1998 through 2005 were higher than the first 2 years of the fishery, the result of a longer season that began in 1998 when the BOF lengthened the season by 10 days in May. The additional fishing time resulted in increased harvests of both Chinook and sockeye salmon from 1998 through 2003 (Table 11-4). However, Chinook salmon harvests have declined since 2004, with 53 harvested in 2005, 23 harvested in 2006, 24 in 2007, 4 harvested in 2008, 15 harvested in 2009, 3 harvested in 2010, and no harvests in 2011; 2012 marked an increase with 8 Chinook salmon harvests. Since the extension of fishing time in 1998, the 2006 season resulted in the lowest harvest estimate on record for total salmon harvested. The 2012 harvest was less than the 5-year (2007–2011) average of 217 salmon, 10-year (2000–2011) average of 257 salmon, and the historical average of 248 salmon (Table 11-4).

TYONEK SUBDISTRICT

History and Regulations

Subsistence salmon fishing regulations for the Tyonek Subdistrict were established by court order in 1980 and subsequently permanently established by the BOF. This setnet fishery is located in the Tyonek Subdistrict of the Northern District of Upper Cook Inlet. The subdistrict includes the area from 1 mile south of the mouth of the Chuitna River south to the easternmost part of Granite Point and from the mean point of high tide to the mean point of lower low tide. The area is unique in that all the lands within the subdistrict are owned by the Tyonek Native Corporation. This feature often raises issues of trespass for those individuals living outside the Tyonek Area who do not seek prior permission to land their boats or set their nets on the privately-owned uplands. For a detailed discussion of this fishery and other subsistence uses at Tyonek, see Fall et al. (1984) and Holen and Fall (2011).

In 2011 the Alaska Board of Fisheries modified the Northern District King Salmon Management Plan (5 AAC 21.366). This modification was in response to reduced abundance of Chinook salmon in the Northern District. The sport fishery on the Chuitna River, which is at the northern edge of the Tyonek Subdistrict, was closed, and commercial fishing was closed from a point just south of the community to the Susitna River in 2011 (Shields and Dupuis 2012:10).

The season in this subsistence fishery also operates in 2 parts. The first part, which focuses on Chinook salmon, is open on Tuesdays, Thursdays, and Fridays from May 15–June 15. The second part is open Saturdays from June 16–October 15. In 2011 the Alaska Board of Fisheries specified the amounts reasonably necessary for subsistence of Chinook salmon and other salmon in the Tyonek subdistrict as 700–2,700 Chinook salmon and 150–500 other salmon. A permit is required and 5 AAC 01.595 (a)(3) specifies that each permit holder may harvest 70 Chinook salmon in the Tyonek Subdistrict and 25 other salmon for the head of household and an additional 10 salmon for each dependent of the permit holder.

Allowable gear for the Tyonek Subdistrict subsistence fishery includes set gillnets 10 fathoms in length, no deeper than 45 meshes, and a stretched mesh sized no larger than 6 in. When fishing, permit holders are required to be present at the net site. Other standard permit conditions include prohibition of fishing within 300 ft of a dam, fish ladder, weir, culvert, or other artificial obstruction.

Harvest Assessment Methods

Household permits are issued by ADF&G prior to fishing, and harvests are recorded on the permits. Two separate permits are required, 1 for the early season and 1 for the late season. A Division of Subsistence staff person travels to Tyonek each May and issues approximately 40–50 permits within several hours. Permits are also available in the Anchorage ADF&G office or in the Tyonek village office. Reported harvests are not expanded in this fishery. Because of the high compliance with the permit requirement and the strong support of the Tyonek village government, ADF&G views the harvest estimates for this fishery as very reliable.

The 2012 Season

In 2012, 89 permits were issued for the Tyonek Subdistrict subsistence salmon fishery, including 63 permits issued to Tyonek residents (71%) and 26 permits issued to other Alaska residents, including 19 to residents of Anchorage (21%; Table 11-5). Residents of Tyonek accounted for 81% of the reported harvest total (942 salmon), including 89% of the reported Chinook salmon harvest (745 Chinook salmon).

The 2012 reported harvest of 1,160 salmon was lower than the historical average of 1,512 salmon. The 2012 harvest was notably higher than the 2011 harvest of 789 salmon which was the lowest reported harvest since 1981, although the number of returned permits was higher than the historical average of 58 permits (Table 11-6). Of the total reported subsistence salmon harvest in 2012, 840 were Chinook salmon

(73%), 176 were sockeye salmon (15%), 138 were coho salmon (12%), 2 were chum salmon (<1%) , and 4 were pink salmon (<1%) (Figure 11-4).

UPPER YENTNA RIVER FISH WHEEL FISHERY

History and Regulations

This subsistence fish wheel fishery began in 1996 as a personal use fishery and was reclassified as a subsistence fishery by the BOF in 1998. It is located in the mainstem of the Yentna River from its confluence with Martin Creek upstream to its confluence with the Skwentna River. The fishery occurs from July 15 through July 31. Fishing periods are from 4:00 AM to 8:00 PM Mondays, Wednesdays, and Fridays. For a more detailed discussion of this fishery see Holen and Fall (2011).

Legal gear includes a fish wheel equipped with a live box. Permit holders must be present at the fish wheel while the wheel is fishing. A season limit of 2,500 salmon was established for the fishery. Chinook salmon and rainbow/steelhead trout must be returned alive to the water. Seasonal limits for households are 25 salmon for a household of 1 plus 10 salmon for each additional household member. Other standard permit conditions include prohibition of fishing within 300 ft of a dam, fish ladder, weir, culvert, or other artificial obstruction.

Harvest Assessment Methods

A permit issued by ADF&G is required prior to fishing. Permits are available through the Division of Sport Fish offices in Palmer and Anchorage. Permit holders must record their harvests on the permit and return it to ADF&G. Participants must also report their daily harvest of salmon to the Palmer ADF&G office by noon of the day following an open period. In the view of ADF&G, compliance with the permit requirement is high and harvest estimates for this fishery are very reliable.

Harvests in 2012

In 2012, 21 subsistence permits were issued for the Yentna River subsistence fish wheel fishery and all were returned (tables 11-7 and 11-8). In 2012, 9 of the 21 permit holders resided in the Skwentna area (43%), with the remaining 12 permits held by residents of other Cook Inlet area communities (Figure 11-5). Permit holders living in the community of Skwentna in 2012 harvested 196 of the reported 343 salmon, or 57% of the harvest (Table 11-7).

Of the total harvest of 343 salmon reported in 2012 were 279 sockeye salmon (81%), 24 coho salmon (7%), 21 pink salmon (6%), and 19 chum salmon (6%) (Figure 11-6). There were no reported harvests of Chinook salmon nor is it legal to retain the harvest. The 2012 harvest of 343 salmon was the lowest recorded harvest since 2005, almost half of the the 5-year average of 637 salmon, less than the 10-year average of 605 salmon, and also less than the historical average of 592 salmon (Table 11-8).

FEDERAL SUBSISTENCE SALMON FISHERIES IN COOK INLET

Since 2007 federal regulations allow for the harvest of salmon, trout, and Dolly Varden by residents of Cooper Landing, Hope, and Ninilchik in the Kenai National Wildlife Refuge and Chugach National Forest. This includes the harvest of salmon by dip net in the Kenai River. In 2012, the total harvest in the federal fishery on the Kenai and Kasilof rivers was 1,438 salmon, all of which were sockeye salmon (Table 11-9). There were a total of 133 permits issued to residents of these 3 communities, with 76 permits issued to residents of Cooper Landing, 29 to residents of Hope, and 28 to residents of Ninilchik (Table 11-9).

Table 11-10 shows the harvest over time, but only includes the years 2007–2012 because this is a new fishery. In all 6 years, sockeye salmon are a majority of the harvest, with 2008 being the highest harvest, at 1,716 sockeye salmon harvested by residents of the 3 Kenai Peninsula communities.

COOK INLET PERSONAL USE SALMON FISHERIES

Background

The BOF first established personal use salmon fisheries in the Cook Inlet Area in 1981 (Nelson et al. 1999:146). Since Alaska statehood in 1959, opportunities had been provided to harvest salmon for home uses with noncommercial set gillnets along various Cook Inlet beaches under subsistence regulations (Braund 1982rev.). In 1978, the new Alaska subsistence statute defined, for the first time, subsistence fishing as fishing for “customary and traditional” uses (AS 16.05.940(31, 33)). In 1980, the BOF determined that only the noncommercial net fisheries in the Tyonek and Port Graham subdistricts met the criteria to qualify as customary and traditional subsistence fisheries. Therefore, the BOF created the “personal use” category of fishing regulations to continue providing opportunities for Alaskans to harvest salmon for home use with nets in areas of Cook Inlet that are generally accessible along the road system. In 1992, the Joint Board classified most of the Cook Inlet Area as a “nonsubsistence area,” where subsistence fishing may not be permitted. Thus, in these areas, personal use fisheries are the primary means by which Alaska residents may obtain salmon for home uses using setnets or dip nets.

Due primarily to court decisions and legislation, personal use fishing regulations for Cook Inlet changed frequently in the 1980s and early 1990s. In 1981, the BOF created personal use dip net fisheries targeting sockeye salmon in the Kasilof and Kenai rivers. Until 1996, these fisheries opened only after achievement of escapement goals was projected. Since then, they have taken place within a fixed season. In 1986, the BOF created a personal use dip net fishery at the mouth of Fish Creek (Knik Arm) focusing on sockeye salmon. A fourth Upper Cook Inlet dip net fishery began in 2008 in the lower portion of the Beluga River on the western shore of Cook Inlet; this fishery is open only to Alaska residents 60 years of age or older. In most years since 1981, personal use set gillnet fisheries in the Cook Inlet Area have been limited to Kachemak Bay and an area at the mouth of the Kasilof River. For more detail on the history of subsistence and personal use salmon fisheries in the Cook Inlet Area, see Braund (1982rev.), Fall and Stanek (1990), Brannian and Fox (1996), Nelson (1994), Nelson (Nelson 1995), Nelson et al. (1999), and Dunker (2010). Table 11-11 summarizes harvest data for selected Cook Inlet personal use and subsistence fisheries that are no longer authorized by state regulations.

Upper Cook Inlet Personal Use Salmon Fisheries

Presently, personal use salmon fisheries in the Upper Cook Inlet Area are governed by the provisions of the Upper Cook Inlet Personal Use Salmon Fishery Management Plan (5 AAC 77.540). Participants must possess an Alaska resident sport fishing license and obtain an Upper Cook Inlet Personal Use Fishing Permit for their household. Permit holders and household members may participate in any of the upper inlet personal use salmon fisheries (except, as noted, the Beluga River fishery is only open to Alaska residents 60 years of age or older). For all the fisheries combined, the annual limit is 25 salmon for the permit holder and 10 salmon for each additional household member. Permits must be returned to ADF&G at the end of the season with a record of the harvest.

In 2012, 34,315 permits were issued for Upper Cook Inlet personal use fisheries, excluding the Beluga River dip net fishery. For the 4 fisheries combined (including unknown fishing locations), the estimated harvest was 640,757 salmon, including 629,597 sockeye (98%), and there were lesser totals for the other 4 species (Table 11-12). The estimated harvest in 2012 was the second highest in the history of these fisheries. For 1996 through 2011, the average annual harvest was 311,061 salmon, although participation and harvest grew steadily (Table 11-13).

Table 11-14 reports the number of permits issued for these 4 Upper Cook Inlet personal use fisheries and the estimated harvest by place of residence of the permit holder. Residents of the Municipality of Anchorage (including Anchorage, Chugiak, Eagle River, JBER [Joint Base Elmendorf/Richardson], and Girdwood) held the most permits (57%) and accounted for 57% of the harvest, followed by Kenai Peninsula Borough residents (20% of permits; 18% of harvests), Matanuska–Susitna Borough residents

(16% of permits; 17% of harvest), residents of other Alaska communities (3% of permits; 4% of harvest), and permit holders for whom a community of residence could not be established (3% of permits; 3% of harvests).

Kasilof River Personal Use Setnet Fishery

This fishery takes place at the mouth of the Kasilof River between regulatory markers approximately 1 mile on either side of the river. Legal gear is a set gillnet no more than 10 fathoms in length, 6 inches in mesh size, and 45 meshes in depth. The fishery is open daily from 6:00 AM to 11:00 PM from June 15 through June 24. In 2012, the total estimated harvest in the fishery was 15,970 salmon, of which 15,638 (98%) were sockeye salmon. (Note that the harvests for this set net fishery plus the dip net fisheries in the Kasilof River, the Kenai River, and Fish Creek are reported through a single permit system, the combined estimated totals are reported above.) The average annual harvest from 1996 through 2011 was 20,111 salmon (Table 11-15).

Kasilof River Dip Net Fishery

This dip net fishery takes place in the lower mile of the Kasilof River 24 hours per day from June 25 through August 7. Retention of Chinook salmon in this fishery is prohibited. The estimated harvest in 2012 was 75,648 salmon, of which 97% was sockeye salmon. From 1996 through 2011, the average annual harvest in this fishery was 44,963 salmon (Table 11-16).

Kenai River Dip Net Fishery

This dip net fishery takes place in the lower Kenai River downriver of the Warren Ames Bridge. Fishing is open from July 10 through July 31, 7 days per week from 6:00 AM to 11:00 PM; when the abundance of sockeye salmon is greater than 2 million fish, the fishery may be open by emergency order 24 hours a day. No more than 1 Chinook salmon per permit may be retained in this fishery. Estimated harvests totaled 535,235 salmon in 2012, including 526,992 sockeye salmon (98%). The average annual harvest from 1996 through 2011 was 231,864 salmon, with harvest—along with participation—rising markedly over that period. (Table 11-17).

Fish Creek Dip Net Fishery

This dip net fishery opens by emergency order if the department projects an escapement into Fish Creek (Knik Arm) of more than 50,000 sockeye salmon. The season is July 10 through July 31. Open waters extend from the terminus of Fish Creek upstream to one-quarter of a mile above the Knik–Goose Bay Road. No Chinook salmon may be retained in this fishery. The fishery did not open in 2012. The most recent harvest numbers are from 2011 when the estimated harvest totaled 6,370 salmon, 82% of which was sockeye salmon. This was substantially lower than the record harvest of 29,304 salmon estimated for 2010. The fishery did not open from 2002 through 2008. The average annual harvest for those years with an open fishery from 1996 through 2010 was 9,664 salmon (Table 11-18).

Unknown Upper Cook Inlet Personal Use Dip Net Fishery

Because not all participants in the Upper Cook Inlet personal use dip net fisheries indicate the location of their fishing activities when they return their permits, an estimate of harvests in an “unknown” Upper Cook Inlet dip net fishery is produced annually. Harvests that could not be attributed to one of the 4 Upper Cook Inlet personal use fisheries (3 dip net fisheries and 1 set net fishery) (excluding the Beluga River fishery, which is discussed below) were estimated at 13,904 salmon in 2012, 97% of which was sockeye salmon (Table 11-19).

Beluga River Personal Use Salmon Fishery

Participation in this dip net fishery, which first took place in 2008, is limited to Alaska residents 60 years of age or older. The fishery is open 24 hours per day from July 10 to August 31 within the Beluga River, western Cook Inlet, from about one-quarter mile upstream of the Beluga River bridge to about 1 mile

below the bridge. The fishery operates under the single seasonal limit for Cook Inlet Area personal use salmon fisheries (25 salmon for the permit holder and 10 additional salmon for each dependent), except only 1 Chinook salmon may be retained. Participants must report their harvest weekly to ADF&G, and the fishery closes when 500 salmon have been harvested (5 AAC 77.540(g)). Harvests totaled 16 salmon in 2012, compared to 159 salmon in 2011, 53 salmon in 2010, 225 salmon in 2009, and 66 salmon in 2008 (Table 11-20). Harvest data by place of residence are presently not available for this fishery, and totals for this fishery are not included with other Upper Cook Inlet personal use fisheries summarized in Table 11-14.

Lower Cook Inlet Personal Use Salmon Fisheries

Kachemak Bay Setnet Fishery

This setnet fishery along Kachemak Bay in the Lower Cook Inlet Management Area was a subsistence fishery before being reclassified as a personal use fishery in the early 1980s. By regulation, the fishery is open from August 16 through September 15, from 6:00 AM Monday until 6:00 AM Wednesday and from 6:00 AM Thursday until 6:00 AM Saturday. The fishery closes when a guideline harvest range of 1,000–2,000 coho salmon has been achieved. Participants must obtain a permit from the Homer ADF&G office—this is separate from the permit program for the Upper Cook Inlet personal use fisheries. Seasonal limits are 25 salmon for the permit holder and 10 salmon for each additional household member (5 AAC 77.549). Fishers must phone the Homer ADF&G office to report their daily harvests.

In 2012, the reported harvest, based on 95 returned permits (97% of the 98 permits issued), was 1,894 salmon, of which 1,471 (78%) were coho. The recent 10-year average harvest for this fishery (2002–2011) was 1,619 salmon (Table 11-21). Harvest data by place of residence are presently not available for this fishery. Table 11-21 also provides historical harvests for this fishery for 1969 through 2012.

China Poot Dip Net Fishery

This personal use dip net fishery first opened in 1980. It takes place in China Poot Bay, approximately 4 miles southeast of the Homer Spit, on the south side of Kachemak Bay. This area is not accessible by road. The fishery targets enhanced sockeye salmon (stocked by the Cook Inlet Aquaculture Association) that have escaped the commercial fishery. Personal use fishers must have a valid Alaska resident sport fishing license, but a permit is not required. The season is July 1 through August 7. Only sockeye salmon may be retained in this fishery, with a bag and possession limit of 6 fish (5 AAC 77.545). Since 1996, ADF&G has not estimated harvests in this fishery. Table 11-22 summarizes historical harvest data for this fishery for 1980–1995. During those years, sockeye salmon harvests ranged between 794 (in 1985) and 8,605 (in 1995) and averaged 3,373 sockeye salmon. The annual average participation in this fishery was 1,215 fishers.

OTHER SUBSISTENCE FISHERIES IN COOK INLET

Federal halibut subsistence harvest data are currently available for communities and tribes in the Cook Inlet area. Residents of Port Graham, Nanwalek, and Seldovia participate in this program. For the findings for 2012, see Fall and Koster (2014).

There are no annual harvest assessment programs for other subsistence finfish fisheries in Cook Inlet. Harvest estimates based on comprehensive household surveys conducted by the Division of Subsistence are available in the CSIS for freshwater and marine species spanning multiple years for selected Cook Inlet communities. Of note in Lower Cook Inlet are rockfish (*Sebastes*) documented in Turek et al. (2009). Information on other fish species used in Upper Cook Inlet by Tyonek and Beluga residents can be found in Stanek et al. (2007).

Table 11-1.–Historical subsistence salmon harvests, Port Graham and Koyuktolik subdistricts, 1981–2012.

Year	Permits		Reported salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1981	ND	57	138	2,670	825	177	874	4,684
1982	ND	61	124	2,354	1,493	220	2,932	7,123
1983	ND	46	67	2,480	471	95	187	3,300
1984	ND	24	45	3,262	510	6	673	4,496
1985	ND	24	146	1,177	621	26	345	2,315
1986	ND	44	125	647	481	14	1,062	2,329
1987	ND	55	21	901	914	114	714	2,664
1988	ND	48	104	1,021	844	110	1,756	3,835
1989	ND	44	51	157	1,155	74	1,495	2,932
1990	ND	60	265	1,162	1,417	151	2,960	5,955
1991	ND	63	163	688	2,053	221	4,587	7,712
1992	ND	71	200	535	1,150	236	1,421	3,542
1993	ND	56	277	1,148	913	257	2,663	5,258
1994	ND	70	300	830	1,370	504	1,979	4,983
1995	ND	87	585	1,795	538	376	1,273	4,567
1996	ND	75	310	1,744	939	276	749	4,018
1997	ND	26	202	325	203	153	511	1,394
1998	ND	19	169	289	243	240	459	1,400
1999	ND	74	485	3,157	1,747	1,104	2,023	8,516
2000	ND	67	259	4,664	1,831	953	1,606	9,313
2001	ND	49	133	1,085	1,295	228	1,454	4,195
2002	ND	79	346	10,620	1,057	488	1,831	14,342
2003	ND	52	465	5,534	1,006	532	1,572	9,109
2004	ND	80	312	3,525	1,303	213	1,600	6,953
2005	ND	68	292	2,126	1,193	180	1,608	5,399
2006	ND	53	275	2,559	1,200	296	2,131	6,461
2007 ^a	ND	24	92	532	0	63	74	761
2008	ND	48	124	4,352	1,448	269	2,682	8,875
2009	ND	44	44	3,497	528	140	914	5,123
2010 ^a	ND	35	30	1,630	1,448	308	1,054	4,470
2011	ND	53	53	5,702	1,491	511	2,632	10,389
2012	ND	8	24	961	414	31	482	1,912
5-year average (2007–2011)	-	41	69	3,143	983	258	1,471	5,924
10-year average (2002–2011)	-	54	203	4,008	1,067	300	1,610	7,188
Historical average (1981– 2011)	-	53	200	2,328	1,022	275	1,543	5,368

Source Hollowell et al. (2013). ADF&G Division of Subsistence, 1981–2011.

Note There are no records indicating the numbers of permits issued for any year. Only the numbers of permits returned are recorded. For this reason, averages of the number of permits issued cannot be calculated (indicated with "-").

a. Harvest reports are incomplete.

Table 11-2.—Subsistence salmon harvests by community, Port Graham and Koyuktolik subdistricts, 2012.

Community	Permits		Reported salmon harvest					Total
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	
Nanwalek	ND	1	0	300	400	5	200	905
Port Graham	ND	7	24	661	14	26	282	1,007
Total	-	8	24	961	414	31	482	1,912

Source Hollowell et al. (2013).

Note There are no records indicating the numbers of permits issued for any year. Only the numbers of permits returned are recorded. For this reason, averages of the number of permits issued cannot be calculated (indicated with "-").

Table 11-3.—Subsistence salmon harvests by community, Seldovia, 2012.

Community	Permits		Estimated salmon harvest					Total
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	
Anchor Point	1	0	0	0	0	0	0	0
Seldovia	19	7	8	79	0	0	54	141
Total	20	7	8	79	0	0	54	141

Source Hollowell et al. (2013).

Table 11-4.—Historical subsistence salmon harvests, Seldovia, 1996–2012.

Year	Permits		Estimated salmon harvest					Total
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	
1996	43	42	51	9	0	0	0	60
1997	20	17	52	22	0	0	0	74
1998	22	20	143	65	0	8	0	216
1999	16	16	136	130	0	38	0	304
2000	22	22	179	252	0	16	0	447
2001	19	16	149	142	0	0	0	290
2002	20	20	124	234	13	11	31	413
2003	18	15	117	290	2	66	22	496
2004	14	12	102	69	5	18	65	258
2005	18	16	53	74	14	11	100	251
2006	17	11	23	12	0	0	31	66
2007	19	15	24	66	12	35	103	239
2008	11	9	4	38	50	6	79	177
2009	18	17	15	115	22	13	77	242
2010	16	12	3	133	41	47	88	312
2011	7	4	0	96	0	0	18	114

-continued-

Table 11-4.–Page 2 of 2.

Year	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
2012	20	7	8	79	0	0	54	141
5-year average (2007–2011)	14	11	9	90	25	20	73	217
10-year average (2002–2011)	16	13	46	113	16	21	61	257
Historical average (1997–2011)	19	17	73	109	10	17	38	248

Source Hollowell et al. (2013); ADF&G Division of Subsistence, 1996–2011.

Table 11-5.–Subsistence salmon harvests by community, Tyonek Subdistrict, 2012.

Community	Permits		Reported salmon harvests					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
Anchorage	19	15	45	54	37	2	4	142
Eagle River	2	2	11	1	0	0	0	12
Kenai	1	1	18	0	0	0	0	18
Palmer	2	2	10	0	0	0	0	10
Tyonek	63	47	745	121	76	0	0	942
Wasilla	2	2	11	0	25	0	0	36
Total	89	69	840	176	138	2	4	1,160

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Table 11-6.–Historical subsistence salmon harvests, Tyonek Subdistrict, 1981–2012.

Year	Permits		Reported salmon harvests					Total
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	
1981	70	NA	2,002	269	64	32	15	2,382
1982	69	NA	1,590	310	113	4	14	2,031
1983	75	NA	2,665	187	59	6	0	2,917
1984	75	NA	2,200	266	79	23	3	2,571
1985	76	NA	1,472	164	91	10	0	1,737
1986	65	NA	1,676	203	223	46	50	2,198
1987	64	61	1,610	166	149	24	10	1,959
1988	47	42	1,587	91	253	12	8	1,951
1989	49	47	1,250	85	115	1	0	1,451
1990	42	37	781	66	352	12	20	1,231
1991	57	54	902	20	58	0	0	980

-continued-

Table 11-6.–Page 2 of 2.

Year	Permits		Reported salmon harvests					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1992	57	44	907	75	234	19	7	1,242
1993	62	54	1,370	57	77	17	19	1,540
1994	58	49	770	85	101	22	0	978
1995	70	55	1,317	45	153	15	0	1,530
1996	73	49	1,039	68	137	7	21	1,272
1997	70	42	639	101	137	8	0	885
1998	74	49	1,027	163	64	2	1	1,257
1999	77	54	1,230	144	94	11	32	1,511
2000	60	59	1,157	63	87	0	6	1,313
2001	84	58	976	172	49	6	4	1,207
2002	101	71	1,080	209	115	4	9	1,417
2003	87	74	1,183	111	44	10	7	1,355
2004	97	75	1,345	93	130	0	0	1,568
2005	78	66	982	61	139	2	0	1,184
2006	82	55	943	20	14	1	0	978
2007	84	67	1,281	200	123	2	3	1,609
2008	94	77	1,178	121	194	9	13	1,515
2009	89	69	636	184	258	2	1	1,081
2010	105	77	843	212	167	2	2	1,226
2011	114	63	595	154	26	7	7	789
2012	89	69	840	176	138	2	4	1,160
5-year average (2007–2011)	97	71	907	174	154	4	5	1,244
10-year average (2002–2011)	93	69	1,007	137	121	4	4	1,272
Historical average (1981– 2011)	74	58	1,233	134	126	10	8	1,512

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Note NA = Information regarding the number of permits returned in 1981–1986 does exist; however, it was not available.

Table 11-7.–Subsistence salmon harvests by community, Upper Yentna River, 2012.

Community	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook ^a	Sockeye	Coho	Chum	Pink	Total
Anchorage	2	2	0	14	0	0	2	16
Big Lake	2	2	0	11	1	0	0	12
Eagle River	2	2	0	31	0	5	3	39
Skwentna	9	9	0	148	18	14	16	196
Wasilla	5	5	0	75	5	0	0	80
Willow	1	1	0	0	0	0	0	0
Total	21	21	0	279	24	19	21	343

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

a. Regulations prohibit the retention of Chinook salmon in this fishery (5 AAC 01.593).

Table 11-8.—Historical subsistence and personal use salmon harvests, Upper Yentna River, 1996–2012.

Year	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook ^b	Sockeye	Coho	Chum	Pink	Total
1996 ^a	17	17	0	242	46	51	115	454
1997 ^a	24	21	0	549	83	10	30	672
1998	21	18	0	495	113	15	30	653
1999	18	16	0	516	48	13	18	595
2000	19	19	0	379	92	7	4	482
2001	16	15	0	545	50	4	10	608
2002	25	22	0	454	133	31	14	632
2003	19	15	0	553	67	8	2	630
2004	21	19	0	441	146	3	36	625
2005	18	17	0	177	42	25	24	268
2006	22	22	0	368	175	26	14	583
2007	22	22	0	367	66	18	17	468
2008	16	16	0	310	57	7	23	397
2009	17	17	0	253	14	6	0	273
2010	32	32	0	642	50	18	38	748
2011	25	25	0	598	90	21	337	1,046
2012	21	21	0	279	24	19	21	343
5-year average (2007–2011)	21	19	0	480	58	14	84	637
10-year average (2002–2011)	21	19	0	448	89	17	51	605
Historical average (1996–2011)	20	19	0	448	83	17	45	592

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

- a. This fishery was classified as personal use in 1996 and 1997; it has been a subsistence fishery since 1998.
- b. Regulations prohibit the retention of Chinook salmon in this fishery (5 AAC 01.593).

Table 11-9.—Federal subsistence salmon harvests by community, Kenai and Kasilof rivers, 2012.

Community	Permits		Reported salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
Cooper Landing	76	70	0	1,052	0	0	0	1,052
Hope	29	25	0	287	0	0	0	287
Ninilchik	28	26	0	99	0	0	0	99
Total	133	121	0	1,438	0	0	0	1,438

Source Jeffry Anderson, USFWS, Kenai Fish & Wildlife Field Office, personal communication.

Table 11-10.—Historical federal subsistence salmon harvests, Kenai and Kasilof rivers, 2007–2012.

Year	Permits		Reported salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
2007	136	131	0	742	5	0	0	747
2008	160	151	2	1,716	12	0	0	1,730
2009	160	138	0	1,104	9	0	0	1,113
2010	169	151	0	943	0	0	0	943
2011	131	123	0	1,090	0	0	0	1,090
2012	133	121	0	1,438	0	0	0	1,438

Source Jeffry Anderson, USFWS, Kenai Fish & Wildlife Field Office, personal communication.

Table 11-11.—Miscellaneous Upper Cook Inlet personal use and subsistence salmon harvests, 1981–1995.

Year ^a	Permits		Reported salmon harvest					
	Total	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
<i>Noncommercial gillnet fishery</i>								
1981	1,108	NA	68	466	12,713	305	149	13,701
<i>Fall coho personal use/subsistence</i>								
1983	295	NA	0	0	712	0	0	712
1984	309	NA	1	2	2,261	7	10	2,281
1985	998	NA	50	805	11,265	53	108	12,281
1986	892	NA	0	0	2,422	0	0	2,422
1987	486	NA	8	9	2,213	37	2	2,269
1988	449	NA	2	19	2,662	10	38	2,731
1989	365	NA	0	0	2,376	0	0	2,376
1990	420	NA	0	0	2,290	0	0	2,290
1991 ^b	360	NA	0	0	2,703	8	0	2,711
1993	535	NA	0	0	1,168	0	23	1,191
<i>Northern/Central districts subsistence/personal use setnet</i>								
1985 ^d	638	NA	117	2,218	1,427	121	90	3,973
1991	7,065 ^e	NA	496	20,855	3,372	1,596	517	26,836
1992	9,200 ^e	NA	957	28,949	8,821	1,753	1,217	41,697
1994	10,127 ^e	NA	1,260	36,701	9,509	1,601	1,653	50,724
1995	9,300 ^e	NA	1,294	45,259	9,678	1,665	1,236	59,132
<i>Knik Arm subsistence</i>								
1985	405	NA	4	1,649	2,055	212	48	3,968

Source Ruesch and Fox (1996); Brannian and Fox (1996).

a. Years listed are only the years in which the fishery was open.

b. In 1991, the fall coho fishery operated as a personal use fishery separate from subsistence setnet fisheries (Ruesch and Fox 1992).

-continued-

Table 11-11.—Page 2 of 2.

- c. Summary data reported in Ruesch and Fox (1996) and in Brannian and Fox (1996) include dip net and setnet harvests. Here, only setnet harvests are included. See separate tables for the Kasilof River dip net fishery and the Kenai River dip net fishery for harvest data for those fisheries.
- d. In 1985, this subsistence fishery was open in areas generally open to commercial fishing, except for the Upper Subdistrict, which had a separate season and permit (called the "fall coho fishery" in this table). The Knik Arm subsistence gillnet fishery was also administered separately in 1985 (Ruesch 1987).
- e. For 1991, 1992, 1994, and 1995, the number of permits issued includes all Upper Cook Inlet dip net and setnet fisheries except the Tyonek subdistrict.

NA = Data not available.

Table 11-12.—Cook Inlet personal use salmon fisheries, 2012.

Year ^a	Permits		Estimated salmon harvest ^b					
	Total	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
<i>Lower Cook Inlet</i>								
Kachemak Bay setnet	98	95	5	137	1,471	6	275	1,894
China Poot Bay dip net ^a								
Subtotal, Lower Cook Inlet	98	95	5	137	1,471	6	275	1,894
<i>Upper Cook Inlet</i>								
Kasilof River setnet ^c			103	15,638	161	15	53	15,970
Kasilof River dip net ^c			16	73,419	1,170	147	896	75,648
Kenai River dip net ^c			40	526,992	4,008	425	3,770	535,235
Fish Creek dip net ^c			--	--	--	--	--	--
Unknown Upper Cook Inlet ^c			8	13,548	173	40	135	13,904
Subtotal, common permit fisheries^c	34,315	27,080	167	629,597	5,512	627	4,854	640,757
Beluga River dip net	7	7	0	9	7	0	0	16
Subtotal, Upper Cook Inlet	34,322	27,087	167	629,606	5,519	627	4,854	640,773
Cook Inlet Total	34,420	27,182	172	629,743	6,990	633	5,129	642,667

Source ADF&G Division of Sport Fish

- a. Permits are not issued for this fishery and harvest estimates are not produced.
- b. Estimated harvests for all fisheries except Kachemak Bay setnet. Only reported harvests are available.
- c. A single permit is issued for the Kasilof setnet, Kasilof dip net, Kenai dip net, and Fish Creek dip net fisheries. In some cases, returned permits did not indicate the area fished.

Table 11-13.—Estimated personal use salmon harvests, Upper Cook Inlet personal use fishery total, 1996–2012.

Year	Permits		Estimated salmon harvest					
	Total	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1996	14,576	13,452	452	145,545	4,811	350	2,973	154,131
1997	14,919	13,756	464	148,940	777	88	844	151,113
1998	15,535	13,190	549	176,581	2,685	220	1,933	181,968
1999	17,197	14,216	1,108	208,589	1,413	168	2,078	213,356
2000	16,107	13,582	1,102	149,267	3,638	290	2,482	156,779
2001	16,915	14,398	1,138	218,688	2,637	276	1,821	224,560
2002	17,568	14,284	1,070	259,623	3,271	757	8,470	273,191
2003	19,110	15,726	1,711	298,831	2,250	371	2,082	305,245
2004	21,910	17,748	1,098	350,091	3,754	502	2,715	358,160
2005	21,905	19,081	1,132	369,776	3,415	428	2,520	377,271
2006	18,563	16,532	1,405	216,047	3,759	746	12,434	234,391
2007	23,046	20,312	1,924	356,717	2,727	614	2,352	364,334
2008	23,722	20,259	1,601	318,594	3,249	727	11,869	336,040
2009	29,619	25,029	1,384	457,539	4,204	559	6,969	470,655
2010	31,590	25,222	1,059	514,255	8,405	1,090	6,482	531,291
2011	34,515	27,193	1,453	630,242	6,754	1,169	4,879	644,497
2012	34,315	27,080	167	629,757	5,512	627	4,854	640,757
5-year average (2007–2011)	28,498	23,603	1,484	455,469	5,068	832	6,510	469,363
10-year average (2002–2011)	24,155	20,139	1,384	377,172	4,179	696	6,077	389,508
Historical average (1996–2011)	21,050	17,749	1,166	301,208	3,609	522	4,556	311,061

Source ADF&G Division of Sport Fish

Note Does not include the Beluga River dip net fishery.

Table 11-14.—Personal use salmon harvest estimates by community, Upper Cook Inlet, 2012.

Community	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
Anchor Point	305	260	1	4,876	45	1	22	4,944
Clam Gulch	54	47	0	902	12	10	7	932
Cooper Landing	31	30	0	283	0	0	6	290
Fritz Creek	72	62	0	1,326	2	1	4	1,334
Homer	906	774	2	15,863	82	8	104	16,059
Hope	16	13	0	277	1	0	2	280
Kasilof	460	386	12	7,626	59	2	47	7,746
Kenai	1,747	1,413	11	29,965	153	17	140	30,286
Moose Pass	28	25	0	481	1	0	1	483
Nanwalek	2	1	0	38	0	0	0	39
Nikiski	264	211	1	4,230	15	7	20	4,274
Nikolaevsk	16	14	0	395	0	0	2	398
Ninilchik	194	163	0	2,968	7	3	6	2,983
Seldovia	11	9	2	223	4	11	0	241
Seward	258	210	0	4,492	18	1	11	4,523
Soldotna	2,068	1,766	9	34,913	113	12	119	35,167
Sterling	478	390	5	7,983	35	8	29	8,060
Subtotal, Kenai Peninsula Borough	6,910	5,774	44	116,844	548	82	521	118,039
Anchorage	16,244	12,627	75	297,009	3,017	409	2,877	303,388
Chugiak	715	618	4	14,617	116	2	69	14,808
Eagle River	1,976	1,689	9	37,700	261	27	277	38,275
Girdwood	257	210	0	4,313	74	1	30	4,418
Joint Base Elmendorf Richardson	297	216	0	5,190	30	3	44	5,267
Subtotal, Anchorage Municipality	19,489	15,360	89	358,830	3,499	443	3,297	366,157
Big Lake	195	159	0	3,406	8	1	14	3,429
Chickaloon	13	12	0	228	0	0	4	233
Houston	37	28	0	525	2	0	5	532
Palmer	1,635	1,362	9	31,405	225	30	184	31,853
Sutton	59	47	0	1,151	2	0	12	1,166
Talkeetna	87	68	0	1,631	4	0	12	1,647
Trapper Creek	26	19	0	576	1	0	1	579
Wasilla	3,353	2,697	9	67,979	765	45	500	69,298
Willow	163	130	0	3,197	31	1	10	3,240
Subtotal, Matanuska-Susitna Borough	5,568	4,522	18	110,100	1,039	77	742	111,975
Akiachak	1	1	0	25	0	0	0	25
Anderson	3	3	0	81	0	0	2	83
Atkasuk	2	0	0	28	0	0	0	29
Barrow	53	25	0	1,221	6	2	4	1,233

-continued-

Table 11-4.-Page 2 of 4.

Community	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
Bethel	14	11	0	236	1	0	0	237
Bettles	1	0	0	14	0	0	0	15
Brevig Mission	1	1	0	12	0	0	2	14
Cantwell	12	11	0	195	0	0	2	198
Chefornak	1	0	0	14	0	0	0	15
Chenega Bay	2	2	0	0	0	0	0	0
Chevak	4	2	0	28	2	0	0	31
Chitina	1	0	0	14	0	0	0	15
Clear	7	5	0	154	0	0	0	155
Copper Center	2	1	1	36	0	0	0	38
Cordova	6	5	0	70	0	0	0	71
Craig	1	1	0	6	0	0	0	6
Deering	1	1	0	63	0	0	0	63
Delta Junction	35	33	0	1,329	3	0	3	1,336
Denali Park	24	23	0	542	3	0	6	552
Dillingham	4	2	0	38	0	0	0	39
Eagle River	1	1	0	0	0	0	0	0
Eielson AFB	16	13	0	341	1	0	0	342
Elim	1	1	0	45	0	0	0	45
Ester	9	8	0	117	0	0	0	118
Fairbanks	598	485	4	12,218	154	4	54	12,435
Fort Greely	3	2	0	82	0	0	2	85
Fort Wainwright	17	14	0	297	1	0	7	305
Gakona	2	2	0	50	0	0	0	50
Galena	4	4	0	96	0	1	0	97
Gambell	1	0	0	14	0	0	0	15
Glennallen	8	7	0	71	0	0	0	72
Goodnews	1	1	0	20	0	0	0	20
Haines	4	3	0	79	0	0	0	80
Healy	47	40	0	858	19	0	1	879
Hoonah	2	1	0	39	0	0	0	40
Hooper Bay	2	1	0	14	0	0	0	15
Huslia	2	2	0	96	0	0	0	96
Iliamna	2	1	0	14	0	0	0	15
Juneau	34	28	0	715	1	0	3	719
Ketchikan	11	7	0	175	1	0	1	176
Kiana	1	0	0	14	0	0	0	15
Klawock	1	1	0	0	0	0	0	0
Kodiak	17	13	2	293	1	0	1	296
Kotlik	1	1	0	0	0	0	0	0
Kotzebue	10	9	0	252	0	0	0	253
Kwethluk	1	1	0	2	0	0	0	2
Levelok	1	1	0	16	0	0	0	16
Manley Hot Springs	1	1	0	25	0	0	0	25
Manokotak	1	0	0	14	0	0	0	15
McGrath	6	5	0	81	12	0	0	94

-continued-

Table 11-4.-Page 3 of 4.

Community	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
Mekoryuk	1	1	0	25	0	0	0	25
Minto	2	1	0	26	0	0	0	27
Naknek	1	1	0	25	0	0	0	25
Napakiaik	1	0	0	14	0	0	0	15
Nenana	9	8	0	233	0	0	0	234
Newhalen	1	1	0	3	0	0	0	3
Newtok	1	0	0	14	0	0	0	15
Nome	17	17	0	427	0	0	1	428
Noorvik	2	0	0	28	0	0	0	29
North Pole	173	137	2	3,982	37	1	15	4,037
Nuiqsut	3	1	0	28	0	0	0	29
Nulato	1	1	0	2	0	0	0	2
Point Hope	4	2	0	46	0	0	0	47
Port Lions	1	1	0	16	0	0	0	16
Quinhagak	1	0	0	14	0	0	0	15
Russian Mission	1	1	0	45	0	0	0	45
Saint Marys	1	1	0	0	0	0	0	0
Saint Paul Island	3	3	0	134	0	0	0	134
Salcha	9	8	0	168	0	0	0	169
Sand Point	1	0	0	14	0	0	0	15
Shishmaref	2	2	0	2	0	0	0	2
Shungnak	1	1	0	0	0	0	0	0
Sitka	9	9	0	163	0	0	0	163
Skagway	3	2	0	65	0	0	0	66
Skwentna	1	1	0	0	0	0	0	0
Stebbins	1	0	0	14	0	0	0	15
Togiak	1	0	0	14	0	0	0	15
Tok	10	7	0	233	1	0	1	235
Toksook Bay	2	2	0	32	0	0	3	35
Twin Hills	1	1	0	25	0	0	0	25
Two Rivers	4	2	0	70	0	0	0	71
Unalakleet	5	4	0	145	0	0	1	147
Unalaska	10	8	0	159	0	0	0	160
Valdez	26	20	0	587	4	0	1	592
Venetie	1	1	0	55	0	0	0	55
Wainwright	1	0	0	14	0	0	0	15
Wales	1	1	0	5	0	0	0	5
White Mountain	2	1	0	44	0	0	0	45
Whittier	10	6	0	155	1	0	1	156
Wrangell	3	2	0	68	0	0	0	69
Yakutat	1	1	0	6	0	0	0	6
Subtotal, other Alaska	1,308	1,035	10	27,253	257	9	120	27,649

-continued-

Table 11-4.–Page 4 of 4.

Community	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
Unknown Communities	1,040	389	6	16,570	169	17	175	16,937
Total	34,315	27,080	167	629,597	5,512	627	4,854	640,757

Source ADF&G Division of Sport Fish

Note Includes Kasilof River setnet fishery, Kasilof River dip net fishery, Kenai River dip net fishery, Fish Creek (Knik Arm) dip net fishery and unknown fishery.

Table 11-15.–Estimated personal use salmon harvests, Kasilof River setnet fishery, 1982–2012.

Year ^a	Permits		Estimated salmon harvest					
	Total	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1982	649	NA	372	7,543	24	NA	17	7,956
1983	684	NA	307	8,846	NA	NA	NA	9,153
1984	698	NA	165	12,926	NA	NA	NA	13,091
1985	692	NA	203	10,746	NA	NA	NA	10,949
1986	NA	NA	168	9,609	NA	NA	NA	9,777
1987	NA	NA	184	9,375	NA	NA	NA	9,559
1988	NA	NA	118	9,803	NA	NA	NA	9,921
1989	NA	NA	186	9,928	NA	NA	NA	10,114
1990	NA	NA	133	7,123	NA	NA	NA	7,256
1991 ^b	NA	NA	34	8,380	NA	NA	NA	8,414
1992	--	--	--	--	--	--	--	--
1993	NA	NA	47	7,942	NA	NA	NA	7,989
1994	--	--	--	--	--	--	--	--
1995	--	--	--	--	--	--	--	--
1996 ^c	NA	NA	46	9,506	0	1	8	9,561
1997	NA	NA	65	17,997	1	3	102	18,168
1998	NA	NA	126	15,975	0	12	15	16,128
1999	NA	NA	442	12,832	25	10	10	13,319
2000	NA	NA	514	14,774	9	10	17	15,324
2001	NA	NA	174	17,201	6	7	11	17,399
2002	NA	NA	192	17,980	12	13	30	18,227
2003	NA	NA	400	15,706	107	4	9	16,226
2004	NA	NA	163	25,417	58	0	6	25,644
2005	NA	NA	87	26,609	326	1	16	27,039
2006	NA	NA	287	28,867	420	6	11	29,591
2007	NA	NA	343	14,943	68	0	2	15,356
2008	NA	NA	151	23,432	65	23	35	23,706
2009	NA	NA	127	26,646	165	11	14	26,963
2010	NA	NA	136	21,924	23	1	23	22,107
2011	NA	NA	167	26,780	47	3	23	27,020
2012	NA	NA	103	15,638	161	15	53	15,970

-continued-

Table 11-15.–Page 2 of 2

Year ^a	Permits		Estimated salmon harvest					
	Total	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
5-year average (2007-2011)	NA	NA	185	22,745	74	8	19	23,030
10-year average (2002-2011)	NA	NA	205	22,830	129	6	17	23,188
Historical average (1996-2011) ^d	NA	NA	214	19,787	83	7	21	20,111

Source Ruesch and Fox (1996) for 1982–1995; Division of Sport Fish for 1996–2012.

- The fishery was closed 1992, 1994, and 1995.
- This fishery was administered separately from the subsistence setnet fisheries that operated in 1991 (Ruesch and Fox 1992).
- Current regulations in place since 1996. Permits since 1996 issued for 4 Upper Cook Inlet personal use salmon fisheries.
- Historical average based on years since 1996 when current regulations were adopted.

Table 11-16.—Estimated personal use salmon harvests, Kasilof River dip net fishery, 1981–2012.

Year ^b	Permits		Estimated salmon harvest ^a					
	Total	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1981	NA	NA	NA	10,300	NA	NA	NA	10,300
1982	NA	NA	NA	1,800	NA	NA	NA	1,800
1983	NA	NA	NA	11,124	NA	NA	NA	11,124
1984	NA	NA	NA	12,771	NA	NA	NA	12,771
1985	NA	NA	NA	16,284	NA	NA	NA	16,284
1986	NA	NA	NA	38,674	NA	NA	NA	38,674
1987	NA	NA	NA	18,454	NA	NA	NA	18,454
1988	NA	NA	NA	3,547	NA	NA	NA	3,547
1989	--	--	--	--	--	--	--	--
1990	--	--	--	--	--	--	--	--
1991 ^{cd}	7,065	5,480	10	907	2	0	3	922
1992	9,500	4,104	24	1,230	24	0	3	1,281
1993	--	--	--	--	--	--	--	--
1994 ^e	10,127	4,823	54	6,414	137	14	59	6,678
1995	NA	NA	NA	4,160	NA	NA	NA	4,160
1996 ^f	NA	NA	50	11,197	334	17	103	11,701
1997	NA	NA	35	9,737	90	19	19	9,900
1998	NA	NA	134	45,161	731	74	610	46,710
1999	NA	NA	127	37,176	286	52	264	37,905
2000	NA	NA	134	23,877	1,004	34	841	25,890
2001	NA	NA	138	37,612	766	23	307	38,846

-continued-

Table 11-16.–Page 2 of 2.

Year ^b	Permits		Estimated salmon harvest ^a					
	Total	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
2002	NA	NA	106	46,769	1197	139	1862	50,073
2003	NA	NA	57	43,870	592	30	286	44,835
2004	NA	NA	44	48,315	668	90	396	49,513
2005	NA	NA	16	43,151	538	102	658	44,465
2006	NA	NA	55	56,144	1,057	105	992	58,353
2007	NA	NA	35	43,293	487	136	383	44,334
2008	NA	NA	46	54,051	509	143	787	55,536
2009	NA	NA	34	73,035	1,441	173	1,274	75,957
2010	NA	NA	31	70,774	1,768	279	974	73,826
2011	NA	NA	24	49,766	977	144	652	51,563
2012	NA	NA	16	73,419	1170	147	896	75,648
5-year average (2007–2011)	NA	NA	34	58,184	1,036	175	814	60,243
10-year average (2002–2011)	NA	NA	45	52,917	923	134	826	54,846
Historical average (1996–2011) ^g	NA	NA	67	43,371	778	98	651	44,963

Source Nelson et al. (1999) for 1981–1990 and 1993–1995; Brannian and Fox (1996) for 1991, 1992, and 1994; Division of Sport Fish for 1996–2012.

- a. Personal use harvests are estimated based on the annual sport harvest survey conducted by the Division of Sport Fish prior to 1996, and are estimated based on permit returns since 1996. Only sockeye salmon harvests reported, 1981–1990.
- b. Fishery closed 1989–1990, and 1993. Classified as a subsistence fishery in 1991 and 1992.
- c. In 1991, 1992, and 1994, a single permit issued for all Upper Cook Inlet subsistence fisheries except Tyonek (central dip net, central setnet, northern setnet) (Brannian and Fox 1996). Permit return rate for 1992 was approximately 43.2% (Ruesch and Fox 1993).
- d. Harvests for 1991 and 1992, and subsistence harvests for 1994, are reported, not estimated.
- e. In 1994 both a subsistence and a personal use dip net fishery took place in the Kasilof River (Nelson 1999). Sockeye harvests included 3,679 salmon in the personal use fishery and 2,735 salmon in the subsistence fishery. Harvest data for other species in the personal use fishery are not available.
- f. Current regulations have been in place since 1996. Permits have been required since 1996 and are issued for 4 Upper Cook Inlet personal use fisheries.
- g. Historical average based on years since 1996 when current regulations were adopted.

NA = Data not available.

Table 11-17.—Estimated personal use salmon harvests, Kenai River dip net fishery, 1981–2012.

Year ^b	Permits		Estimated salmon harvest ^a					
	Total	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1981	—	—	—	—	—	—	—	—
1982 ^c	NA	NA	NA	NA	NA	NA	NA	0
1983	NA	NA	NA	7,562	NA	NA	NA	7,562
1984	—	—	—	—	—	—	—	—
1985	—	—	—	—	—	—	—	—
1986	—	—	—	—	—	—	—	—
1987	NA	NA	NA	24,086	NA	NA	NA	24,086
1988	NA	NA	NA	16,880	NA	NA	NA	16,880
1989	NA	NA	NA	48,976	NA	NA	NA	48,976
1990	—	—	—	—	—	—	—	—
1991 ^{de}	7,065	5,480	44	10,468	146	2	17	10,677
1992 ^f	9,500	4,104	158	28,429	1,475	74	598	30,734
1993	NA	NA	NA	33,467	NA	NA	NA	33,467
1994	10,127	4,823	187	13,897	2,535	114	1,263	17,996
1995	NA	NA	NA	14,352	NA	NA	NA	14,352
1996 ^g	NA	NA	295	102,821	1,932	175	2,404	107,627
1997	NA	NA	364	114,619	559	58	619	116,219
1998	NA	NA	254	103,847	1,011	85	1,032	106,229
1999	NA	NA	488	149,504	1,009	102	1,666	152,769
2000	NA	NA	410	98,262	1,449	193	1,457	101,771
2001	NA	NA	638	150,766	1,555	155	1,326	154,440
2002	NA	NA	606	180,028	1,721	551	5,662	188,568
2003	NA	NA	1,016	223,580	1,332	249	1,647	227,824
2004	NA	NA	792	262,831	2,661	387	2,103	268,774
2005	NA	NA	997	295,496	2,512	321	1,806	301,132
2006	NA	NA	1,034	127,630	2,235	551	11,127	142,577
2007	NA	NA	1,509	291,270	2,111	472	1,939	297,301
2008	NA	NA	1,362	234,109	2,609	504	10,631	249,215
2009	NA	NA	1,189	339,993	2,401	285	5,482	349,350
2010	NA	NA	865	389,552	2,870	508	3,655	397,450
2011	NA	NA	1,243	537,765	4,745	915	3,914	548,582
2012	NA	NA	40	526,992	4,008	425	3,770	535,235
5-year average (2007–2011)	NA	NA	1,234	358,538	2,947	537	5,124	368,380
10-year average (2002–2011)	NA	NA	1,061	288,225	2,520	474	4,797	297,077
Historical average (1996– 2011) ^h	NA	NA	816	225,130	2,045	344	3,529	231,864

Source Nelson et al. (1999) for 1981–1990 and 1993–1995; Brannian and Fox (1996) for 1991, 1992, and 1994; Division of Sport Fish for 1996–2012.

- a. Personal use harvests are estimated based on the annual sport harvest survey conducted by the Division of Sport Fish prior to 1996, and are estimated based on permit returns since 1996. Only sockeye salmon harvests reported, 1981–1990.
- b. Fishery closed 1981, 1984–1986, and 1990. Classified as a subsistence fishery in 1991, a portion of 1992 and 1994.
- c. The 1982 harvest is reported as "unknown" but "insignificant" (Nelson 1999; Brannian and Fox 1996).

-continued-

Table 11-17.–Page 2 of 2.

- d. Subsistence harvests for 1991, 1992, and 1994 are reported, not estimated.
- e. 1991, 1992, and 1994 permits: single permit issued for all Upper Cook Inlet subsistence fisheries except Tyonek.
- f. Harvests for 1992 include 16,240 sockeye salmon in the subsistence fishery and 12,189 sockeye in the personal use fishery. Harvests for other species are for the subsistence fishery only. Personal use harvests are not available for the other species.
- g. Current regulations have been in place since 1996. Permits have been required since 1996 and are issued for 4 Upper Cook Inlet personal use fisheries.
- h. Historical average based on years since 1996 when current regulations were adopted.
- NA = Data not available.

Table 11-18.–Estimated personal use salmon harvests, Fish Creek dip net fishery, 1987–2012.

Year ^b	Permits		Estimated salmon harvest ^a					
	Total	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1987	NA	NA	0	2,200	0	0	2,200	4,400
1988	NA	NA	0	3,000	0	0	3,000	6,000
1989	NA	NA	0	5,000	0	0	5,000	10,000
1990	NA	NA	0	6,500	0	0	6,500	13,000
1991	NA	NA	0	14,369	0	549	567	15,485
1992	NA	NA	0	19,002	0	607	678	20,287
1993	NA	NA	0	37,224	973	503	2,068	40,768
1994	NA	NA	0	16,012	1,336	248	632	18,228
1995	NA	NA	0	9,102	2,640	99	290	12,131
1996	NA	NA	37	17,260	2,414	153	331	20,195
1997	NA	NA	0	3,277	63	4	53	3,397
1998	NA	NA	1	4,036	649	29	80	4,795
1999	NA	NA	0	1,083	17	0	12	1,112
2000	NA	NA	0	6,925	958	29	83	7,995
2001	NA	NA	0	436	18	1	2	457
2002	–	–	–	–	–	–	–	–
2003	–	–	–	–	–	–	–	–
2004	–	–	–	–	–	–	–	–
2005	–	–	–	–	–	–	–	–
2006	–	–	–	–	–	–	–	–
2007	–	–	–	–	–	–	–	–
2008	–	–	–	–	–	–	–	–
2009	NA	NA	10	9,898	53	33	66	10,060
2010	NA	NA	12	23,705	3,576	290	1,721	29,304
2011	NA	NA	2	5,236	905	72	155	6,370
2012	--	--	--	--	--	--	--	--
Historical average (1996–2011)	NA	NA	7	7,984	961	68	278	9,298

Source Brannian and Fox (1996) for 1987-1994; Howe et al. (1996) for 1995; Division of Sport Fish for 1996–2011.

- a. Estimates derived from statewide sport harvest survey prior to 1996. Permits required since 1996.
- b. Fishery closed 2002 through 2008 and 2012.

Table 11-19.—Estimated personal use salmon harvests, unknown fishery, 1996–2012

Year	Permits		Estimated salmon harvest					
	Total	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1996	NA	NA	24	4,761	131	4	127	5,047
1997	NA	NA	0	3,310	64	4	51	3,429
1998	NA	NA	34	7,562	294	20	196	8,106
1999	NA	NA	51	7,994	76	4	126	8,251
2000	NA	NA	44	5,429	218	24	84	5,799
2001	NA	NA	188	12,673	292	90	175	13,418
2002	NA	NA	166	14,846	341	54	916	16,323
2003	NA	NA	238	15,675	219	88	140	16,360
2004	NA	NA	99	13,527	366	25	210	14,227
2005	NA	NA	32	4,520	39	4	40	4,635
2006	NA	NA	29	3,406	47	84	304	3,870
2007	NA	NA	37	6,729	61	6	28	6,861
2008	NA	NA	41	6,890	66	58	412	7,467
2009	NA	NA	25	7,968	144	57	133	8,327
2010	NA	NA	15	8,300	168	12	109	8,604
2011	NA	NA	17	10,695	80	35	135	10,962
2012	NA	NA	8	13,548	173	40	135	13,904
5-year average (2007–2011)	NA	NA	27	8,116	104	34	163	8,444
10-year average (2002–2011)	NA	NA	70	9,256	153	42	243	9,764
Historical average (1996–2011)	NA	NA	65	8,393	163	36	199	8,855

Source ADF&G Division of Sport Fish.

Table 11-20.—Beluga River senior personal use dip net fishery summary, 2008–2012.

Year	Permits		Reported salmon harvest					
	Total	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
2008	20	20	0	31	35	0	0	66
2009	11	11	0	140	78	0	7	225
2010	14	14	0	47	1	5	0	53
2011	13	12	0	137	17	5	0	159
2012	7	7	0	9	7	0	0	16
Historical average (2008– 2011)	15	14	0	89	33	3	2	126

Source ADF&G Division of Sport Fish.

Table 11-21.—Personal use/subsistence salmon harvests, Kachemak Bay setnet fishery (excluding the Port Graham/Nanwalek subsistence fishery and the Seldovia subsistence fishery), Lower Cook Inlet, 1969–2012.

Year	Households or permits		Reported salmon harvest					
	Total	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1969	47	44	0	9	752	0	38	799
1970	78	73	0	12	1,179	13	143	1,347
1971	112	95	2	16	1,549	7	44	1,618
1972	135	105	1	11	975	69	48	1,104
1973	143	128	0	18	1,304	40	84	1,446
1974	148	118	0	16	376	77	43	512
1975	292	276	4	47	1,960	61	632	2,704
1976	242	221	16	46	1,962	56	1,513	3,593
1977	197	179	12	46	2,216	119	639	3,032
1978	311	264	4	35	2,482	34	595	3,150
1979	437	401	6	37	2,118	41	2,251	4,453
1980	533	494	43	32	3,491	25	1,021	4,612
1981	403	383	15	73	4,370	68	718	5,244
1982	395	372	41	49	7,398	154	956	8,598
1983	344	328	5	17	2,701	44	305	3,072
1984	368	346	3	25	3,639	105	804	4,576
1985	328	302	5	49	3,317	34	138	3,543
1986	349	310	7	68	3,831	56	3,132	7,094
1987	363	339	5	50	3,979	61	279	4,374
1988	439	417	14	73	5,007	75	1,445	6,614
1989	477	453	41	156	7,219	53	883	8,352
1990	578	543	12	200	8,323	69	1,846	10,450
1991	472	459	8	47	4,931	23	366	5,375
1992	365	350	5	63	2,277	21	643	3,009
1993	326	317	6	44	1,992	18	463	2,523
1994	286	284	66	80	4,097	18	1,178	5,439
1995	235	232	118	108	2,916	7	343	3,492
1996	299	293	302	102	3,347	24	1,022	4,797
1997	276	264	384	191	1,817	12	257	2,661
1998	227	214	135	20	1,461	5	167	1,788
1999	146	141	276	119	1,803	3	168	2,369
2000	213	206	104	28	2,064	4	304	2,504
2001	154	148	86	27	1,579	16	150	1,858
2002	122	113	61	33	1,521	12	251	1,878
2003	104	96	17	57	1,071	9	170	1,324
2004	91	83	7	56	1,554	16	172	1,805
2005	108	96	8	57	833	13	296	1,207
2006	89	82	15	41	1,295	5	221	1,577
2007	141	133	10	113	1,431	34	641	2,229
2008	146	142	2	92	1,844	14	687	2,639
2009	145	142	9	273	646	4	101	1,033
2010	128	122	14	149	875	17	251	1,306
2011	119	112	15	223	806	5	145	1,194

-continued-

Table 11-21.–Page 2 of 2.

Year	Households or permits		Reported salmon harvest					Total
	Total	Returned	Chinook	Sockeye	Coho	Chum	Pink	
2012	98	95	5	137	1,471	6	275	1,894
5-year average (2007–2011)	136	130	10	170	1,120	15	365	1,680
10-year average (2002–2011)	119	112	16	109	1,188	13	294	1,619
Historical average (1969–2011)	254	238	44	70	2,565	36	594	3,309

Source Hallowell et al. (2013).

Table 11-22.–Estimated personal use salmon harvests, China Poot dip net fishery, 1980–1995.

Year	Fishers	Estimated salmon harvest					Total
		Chinook	Sockeye	Coho	Chum	Pink	
1980	NA	0	1,000	0	0	0	1,000
1981 ^a	—	—	—	—	—	—	—
1982	NA	0	1,320	0	0	0	1,320
1983	1,956	0	5,910	0	0	0	5,910
1984	1,237	0	1,794	0	0	0	1,794
1985	398	0	794	0	0	12	806
1986	993	0	1,815	0	0	673	2,488
1987	1,016	0	1,231	0	0	0	1,231
1988	1,361	0	1,910	0	127	36	2,073
1989	1,428	0	5,416	0	0	239	5,655
1990	1,537	0	5,835	0	178	68	6,081
1991	395	0	1,528	0	0	33	1,561
1992	810	0	3,468	0	76	183	3,727
1993	1,036	0	4,260	0	0	45	4,305
1994	1,372	0	5,715	0	0	34	5,749
1995 ^b	2,261	0	8,605	0	0	77	8,682
Historical average (1980–1995)	1,215	0	3,373	0	25	93	3,492

Source Fall and Stanek (1990), for 1980 to 1989, based on annual reports of the sport fish harvest survey. 1990 through 1995: annual sport fish angler survey report. Harvest data as reported in annual sport fish angler survey reports differ from data reported in Nelson (1995:222), which reports "sport and personal use harvests combined."

a. Fishery was closed in 1981.

b. Harvest data not collected after 1995.

NA = Data not available.




Figure 11-1.—Anchorage Nonsubsistence Area map.




Figure 11-2.—Subsistence salmon harvests in the Port Graham and Koyuktolik subdistricts, 2012.




Figure 11-3.—Subsistence salmon harvests in Seldovia, 2012.




Figure 11-4.—Subsistence salmon harvests in the Tyonek Subdistrict, 2012.




Figure 11-5.—Permits issued, by place of residence, for the Upper Yentna River fishery, 2012.




Figure 11-6.—Subsistence salmon harvests in the Upper Yentna River, 2012.

CHAPTER 12: PRINCE WILLIAM SOUND AREA

INTRODUCTION

The Prince William Sound Management Area includes all waters of Alaska between the longitude of Cape Fairfield and the longitude of Cape Suckling. Subsistence fisheries are not permitted in the Valdez Nonsubsistence Area (5 AAC 99.015(a)(5)). In 2012, there were 9 subsistence or personal use salmon fisheries with annual harvest assessment programs in the Prince William Sound Management Area:

- Upper Copper River, Glennallen Subdistrict: state subsistence permit program,
- Upper Copper River, Glennallen Subdistrict: federal subsistence permit program,
- Upper Copper River, Chitina Subdistrict: state personal use permit program,
- Upper Copper River, Chitina Subdistrict: federal subsistence permit program,
- Batzulnetas: a federal subsistence permit program,
- Copper River Flats–Prince William Sound: state subsistence permit program,
- Prince William Sound, Eastern District–Tatitlek: state subsistence permit program,
- Prince William Sound, Southwestern District–Chenega Bay: state subsistence permit program, and
- Prince William Sound, general area: state subsistence permit program.

The Glennallen and Chitina subdistricts have had separate state and federal permit programs since 2000. The year 2012 was the eleventh in which there were separate state and federal permit programs for the Glennallen and Chitina subdistricts. It should also be noted that the dip net fishery that takes place in the Chitina Subdistrict of the Upper Copper River District under state regulations was classified as a personal use fishery through 1999. The BOF reclassified this fishery as subsistence in 2000, and again as personal use in 2003 (with no other regulatory changes). Historical data for this fishery, including years when it was classified as subsistence or personal use, are included in statewide summaries as personal use.

For both state and federal management purposes, the Upper Copper River District of the Prince William Sound Management Area consists of all waters of the mainstem Copper River from the mouth of the Slana River downstream to an east–west line crossing the Copper River approximately 200 yd upstream of Haley Creek as designated by ADF&G regulatory markers. There are 2 subdistricts: the Chitina Subdistrict consists of all waters of the Upper Copper River District downstream of the downstream edge of the Chitina–McCarthy Road Bridge, and the Glennallen Subdistrict consists of all remaining waters of the Upper Copper River District.

The state established the Glennallen and Chitina subdistricts in 1977. Prior to that time, the Upper Copper River was treated as one unit for management purposes. For a detailed discussion of the history of these fisheries, see Simeone and Fall (1996).

Under the provisions of 5 AAC 01.630 (h), a village council or other similarly qualified organization may obtain a permit to operate a fish wheel on behalf of its members upon approval of a harvest assessment plan submitted to ADF&G. These organizations may also issue household permits and register fish wheels. Table 12-1 summarizes data for the permits issued for village fish wheels by ADF&G from 1997 through 2011; in 2012, no fish wheel permits were issued.

UPPER COPPER RIVER STATE AND FEDERAL SUBSISTENCE FISHERIES: GLENNALLEN SUBDISTRICT

Regulations

In the Glennallen Subdistrict, permits are required to participate in the state and federal subsistence salmon fisheries. ADF&G issues state permits upon request at ADF&G offices under the authority of 5 AAC 01.630. In 2002, the FSB created a federal permit requirement for qualified rural residents (primarily residents of Copper River Basin and Upper Tanana communities), which is administered by the NPS. While the state subsistence permits limit fishers to either fish wheels or dip nets, federal permit holders may use fish wheels, dip nets, and rod and reel. In the state fishery, fishers may participate in either the Chitina Subdistrict personal use fishery or the Glennallen Subdistrict subsistence fishery in any given year, but not both. Federally-qualified rural resident households may hold permits for both the federal and state Glennallen Subdistrict fisheries, or for the Glennallen federal fishery and the Chitina state personal use fishery, although state and federal harvest limits are not additive. Other standard permit conditions include prohibition of fishing within 300 ft of a dam, fish ladder, weir, culvert, or other artificial obstruction.

In the Glennallen Subdistrict under state regulations, fishers may use either fish wheels or dip nets, but they may not use both types of gear. Federal subsistence permit holders may use rod and reel in addition to fish wheels and dip nets, and they may use all 3 types of gear, but not at the same time. The state season is June 1–September 30; the federal season is May 15–September 30. Annual limits are the same under state and federal regulations: 30 salmon for a household with 1 person or 60 salmon for a household of 2 persons, of which no more than 5 may be Chinook salmon if taken with a dip net. For a household of more than 2, 10 salmon for each additional person may be added to the annual limit. Upon request, permits can be issued for additional salmon, with limits of 200 salmon for 1-person households and 500 for households of 2 or more persons. The number of Chinook salmon (5) taken by dip net does not increase under state regulations; federal permit holders may take up to 5 additional Chinook salmon with rod and reel.

Harvest Assessment Program

Annual subsistence salmon harvest assessments have been conducted for Upper Copper River since 1960, originally by the Division of Commercial Fisheries, but currently by the Division of Sport Fish. Permits include harvest reports, and fishers are required to record the dates they fished and the number of each species harvested each day. Total harvest estimates for the fishery are made based on reported harvests expanded to all permit holders. Beginning in 2002, the NPS, on behalf of the FSB, has compiled the data from federal permit returns in a program separate from that administered by ADF&G.

The creation of a dual permit program for subsistence fishing in the Upper Copper River creates challenges for the compilation of a single subsistence harvest estimate for this subsistence fishery, which is the goal of this annual report. Issues include the following:

- As noted above, federal permits allow fishing with multiple gear types, including rod and reel, but state permits allow fishing with only 2 gear types—dip nets or fish wheels. Thus while past years' annual report summaries for the Glennallen Subdistrict showed the number of permits issued by gear type, this is not possible for the combined state and federal data summaries reported here.
- Some households obtain both the state and federal permit for the Glennallen Subdistrict. Of these “dual-permitted” households, some report harvest and effort only on their state permits (not returning the federal permit), some report harvest and effort only on their federal permits (not returning the state permit), some report identical harvests and efforts on both permits, and some return neither permit. Editing the data to compensate for double-reporting of salmon harvest and effort requires 2 assumptions: 1) permittees returning only 1 permit did not report harvest or

effort on the other, and 2) permittees reporting identical harvests and efforts on both permits reported identical harvests twice rather than split their harvests between permits. These assumptions were employed in the analysis only after discussing the dual-permitted households with the program administrators in the Division of Sport Fish and the NPS. All households obtaining both state and federal permits were counted as receiving only 1 permit in the summary tables for the Glennallen Subdistrict included here.

- Until 2006, state permits collected only the permit holder's city in terms of their mailing address, but federal permits collected this and the "community of primary residence." Since the Copper River area has a number of smaller communities without their own post offices, state permits issued to residents of these communities prior to 2006 did not provide adequate information to assure analysis results accurately reflect the true community residency of harvesters. But because of the precision of the federal permit regarding place of residence, the federal permit place of residence data were used to compile the harvest tables, in combination with the mailing address data from state permits. Since there were several dual-permitted households in the Glennallen Subdistrict fishery, the federal residence community was used as the default where this information differed.

Subsistence Salmon Harvests in 2012

As shown in Table 12-2, ADF&G and NPS issued a total of 1,805 subsistence salmon permits for the Glennallen Subdistrict for 2012. This total is higher than both the recent 5-year average (1,490 permits) and 10-year average (1,367 permits), and represents a steady increase in issued permits since 1989. Of all Glennallen Subdistrict permits issued, both federal and state, residents of Copper Basin communities held 397 permits (approximately 22%) and other Alaska residents held 1,408 permits (78%) (Table 12-3).

As reported in Table 12-2, the estimated total subsistence salmon harvest in the Glennallen Subdistrict in 2012 was 98,110 salmon, including 94,991 sockeye salmon (approximately 98% of the year's salmon harvest), 2,649 Chinook salmon, and 470 coho salmon. Pink and chum salmon are not generally available in the Upper Copper River, although a few chum salmon are sometimes reported. This total includes fish wheel and dip net harvests in the state-administered fishery, and fish wheel, dip net, and rod and reel harvests in the federally-administered fishery. The 2012 harvest was higher than the recent 5-year average (81,546 salmon), 10-year average (80,856 salmon), and the historical average (1989–2009; 68,889 salmon). Table 12-3 reports subsistence salmon harvests in the Glennallen Subdistrict by place of residence of permit holders in 2012. Copper Basin residents caught 32% of the harvest (31,466 salmon) and other Alaska residents harvested 68% (64,782 salmon).

UPPER COPPER RIVER STATE PERSONAL USE FISHERY: CHITINA SUBDISTRICT

Background and History

The Chitina Subdistrict consists of all waters of the Upper Copper River District downstream of the downstream edge of the Chitina–McCarthy Road Bridge to an east–west line crossing the Copper River approximately 200 yd upstream of Haley Creek. In 1984, and from 1986 through 1999, the Chitina Subdistrict was closed to subsistence fishing, and the dip net fishery was operated as a personal use fishery. At its December 1999 meeting, the BOF reversed an earlier decision, determined that the Chitina Subdistrict supported C&T uses of salmon, and returned the classification to subsistence. In February 2003, the BOF reconsidered the subsistence classification of the Chitina dip net fishery, reversed its decision of 1999, made a negative C&T finding, and returned the classification to personal use. No other regulatory changes were made. For a detailed discussion of the history of these fisheries, see Simeone and Fall (1996) and ADF&G (2003).

Regulations

There are state and federal permit programs for the Chitina Subdistrict. Under state regulations, a household permit and an Alaska state resident sport fishing license, both issued by ADF&G, are required for personal use fishing in the Chitina Subdistrict. Households may not possess both the Chitina state personal use permit and the Glennallen state subsistence permit in the same year. Under state regulations, dip nets are the only legal gear in the Chitina Subdistrict. Annual limits are 15 salmon for a 1-person household and 30 salmon for households of 2 or more. Only 1 Chinook salmon may be harvested annually. If ADF&G authorizes a supplemental harvest period by emergency order, permit holders who have already filled their original limit may take 10 additional sockeye salmon during each announced supplementary period. Rainbow/steelhead trout taken by dip net under the state fishery must be released immediately and returned to the water unharmed. Other standard permit conditions include prohibition of fishing within 300 ft of a dam, fish ladder, weir, culvert, or other artificial obstruction.

Harvest Assessment Program

Annual subsistence–personal use salmon harvest assessments have been conducted by ADF&G, currently by the Division of Sport Fish, in the Upper Copper River Area since 1960. Chitina Subdistrict permits include harvest reports, and fishers are required to record the dates they fish, the number of each species harvested each day, whether they fished from a boat or from shore, and if they fished during a supplemental harvest period. Total harvest estimates for the state personal use fishery are made based on reported harvests expanded to all permit holders.

Personal Use Salmon Harvests in 2012

As reported in Table 12-4, the estimated total salmon harvest in the state-administered Chitina Subdistrict personal use fishery in 2012 was 138,465 fish, including 136,441 sockeye salmon (98%), 613 Chinook salmon, and 1,411 coho salmon, by 10,016 permit holders. The 2012 total estimated harvest for the Chitina Subdistrict was well above the recent 5-year (120,621 salmon) and 10-year (117,065 salmon) averages, as well as the historical average (1989–2011; 113,261salmon). It should be noted however that the 2012 Chinook salmon harvest was down significantly from the 5-year, 10-year, and historical averages, as was the 2012 coho salmon harvest.

Table 12-5 reports estimated salmon harvests in the Chitina Subdistrict personal use fishery by city of mailing address of state permit holders in 2012; most participants in this fishery lived in Fairbanks, Anchorage, or the Matanuska–Susitna Borough. Only 34 Copper Basin residents (<1%) obtained state personal use salmon permits for the Chitina Subdistrict in 2012. Non-area residents harvested all but 266 of the salmon harvested in this fishery in 2012 (>99%).

UPPER COPPER RIVER FEDERAL SUBSISTENCE FISHERY: CHITINA SUBDISTRICT

Regulations

In 2012, qualified Alaska rural residents could obtain federal subsistence permits for the Chitina Subdistrict from the NPS. Legal gear included fish wheels, dip nets, and rod and reel. Federally-qualified rural resident households may hold permits for both the federal and state Chitina Subdistrict fisheries, or for the Chitina federal fishery and the Glennallen state subsistence fishery, although state and federal harvest limits are not additive. Federal seasonal limits for the Chitina Subdistrict were the same as for the Glennallen Subdistrict, but were also not additive. Under federal regulations, rainbow/steelhead trout incidentally taken from fish wheels could be retained. Other standard permit conditions include prohibition of fishing within 300 ft of a dam, fish ladder, weir, culvert, or other artificial obstruction.

Federal Subsistence Harvests in 2012

As reported in Table 12-6, an estimated 996 salmon were harvested in the federal Chitina Subdistrict subsistence fishery in 2012, the lowest harvest year on record. The recent 5-year average was 2,478 salmon and the historical average (1989–2011) was 2,089 salmon.

The total harvest included 981 sockeye salmon (98%), 9 coho salmon (1%), and 5 Chinook salmon. A total of 90 permits were issued, which is greater than the 5-year average of 84. Table 12-7 reports harvest and permit numbers according to each permittee's community of residence in 2012 for the Chitina Subdistrict.

Native Village of Batzulnetas Subsistence Fishery

The state created the Batzulnetas fishery in 1987 through an emergency regulation to settle the federal district court case of *John vs. Alaska*. There is also a federal permit program for a federal fishery in this area. Participants in this fishery are largely from the community of Mentasta. Legal gear includes fish wheels and dip nets in the Copper River and dip nets and spears in Tanada Creek. For both state and federal fisheries, the open area is all waters of the Copper River from regulatory markers near the mouth of Tanada Creek and approximately one-half mile downstream from that mouth; and all waters of Tanada Creek between regulatory markers. The state fishing season is open June 1–September 1 or until the season is closed by emergency order; fishing periods are established by emergency order and are 2 days per week during June and 3½ days per week for the remainder of the season. The federal fishing season is May 15–September 30 or until the season is closed by special action.

Since 1987, subsistence permits have been issued in 14 of the 25 years (Table 12-8). One permit was issued and returned every year from 1998 through 2004. No permits were issued for the years 2005 through 2009. Three permits were issued and returned in 2012. The total 2012 harvest included 136 sockeye salmon and 1 Chinook salmon. The historical average (1987–2011) harvest for this fishery is 101 sockeye salmon, with the highest harvest occurring in 1994 with a take of 997 sockeye salmon.

COPPER RIVER DISTRICT SUBSISTENCE FISHERY

Background and Regulations

This fishery takes place in the Copper River District at the mouth of the Copper River (Copper River Flats) near the community of Cordova. Permits are required to participate in subsistence fishing for salmon and freshwater fish species under the authority of 5 AAC 01.630. Permits are issued upon request either in person or by telephone to the Cordova ADF&G office. Fishers must declare their intent to fish in the Copper River Flats Area or in Prince William Sound, since the permit is valid for only one or the other location. Legal gear is set or drift gillnet no longer than 50 fathoms; open season is May 15–September 30, with additional restrictions during times of commercial fishing activity. Annual limits for salmon are 15 salmon for a household of one; 30 salmon for a household of 2 or more; and 10 salmon for each additional person in the household. There is a limit of 5 Chinook salmon per permit. Other standard permit conditions include prohibition of fishing within 300 ft of a dam, fish ladder, weir, culvert, or other artificial obstruction.

Harvest Assessment Program

A permit program with annual subsistence salmon harvest assessments has been in place for Prince William Sound at least since 1960. Permits are returned to the Cordova ADF&G office either in person or by mail at the end of the fishing season. Permits include a harvest report, and fishers are required to record the dates fished and the number of each species of fish harvested each day.

Subsistence Salmon Harvests in 2012

As reported in Table 12-9, 378 permits were issued for this fishery in 2012, and 359 (94%) were returned. Participation in this fishery as represented by issued permits has fluctuated over time, generally increasing from the historical average (167) and remaining somewhat consistent for the 5-year (379) and 10-year (381) averages. The estimated harvest in 2012 of 4,767 salmon was a significant increase from the previous year (2,096). The 2012 harvest was composed mainly of 4,499 sockeye salmon (94%) and included 248 Chinook salmon (5%), and 19 chum salmon (<1%). Most permit holders lived in Cordova (294; 78%) and took 79% of the total harvest (Table 12-10).

EASTERN DISTRICT SUBSISTENCE SALMON FISHERY

Although the Eastern District is defined as those waters of the eastern mainland shore from the radio tower at Whitshed Village to Point Freemantle, including Bligh Island, Goose Island, and other adjacent islands (5 AAC 24.200 (c)), under regulations in place since 1988, salmon may be taken for subsistence purposes only in those waters north of a line from Porcupine Point near Goose Island to Granite Point near Glacier Island, and south of a line from Point Lowe to Tongue Point in Valdez Arm (5 AAC 01.648 (b)). The primary participants in this fishery are residents of Tatitlek. Prior to 1992, permits were issued only in Tatitlek, but since 1992, they have been issued at the Cordova ADF&G office as well. Permits may be returned in person to the Cordova ADF&G office or the Tatitlek Village Council office, or mailed at the end of the fishing season. Permits include a harvest report, and fishers are required to fill in the dates fished and the number of each species of salmon caught each day.

Legal gear for this fishery includes seines up to 50 fathoms in length and 100 meshes deep with a maximum mesh size of 4 in, or gillnets up to 150 fathoms in length with a maximum size of 6¼ in. Pink salmon may be taken in fresh water with dip nets only. The open season is May 15–October 31, under the following fishing periods: 7 days per week from May 15 until 2 days before the commercial opening of the Eastern District; during the commercial fishing season, but only during commercial openers; and 7 days per week from 2 days after the closure of the commercial season through October 31. There are no bag or possession limits for this fishery.

In 2012, there were 16 permits issued for this fishery, up 6 permits from the year before (Table 12-11). The permittees reported a total harvest of 1,052 salmon, down slightly from the year before but representing a general increase over the 5-year, 10-year, and historical averages. The 2012 harvest numbers align with survey efforts from 2003 (Fall 2006), indicating that until just recently, the harvest assessment program for this fishery has consistently and substantially underestimated harvests. As shown in Table 12-12, household surveys in Tatitlek provided an estimate of 1,075 salmon taken with subsistence methods in 2003, compared to the 298 salmon (Table 12-11) based on returned permits for that year. In Tatitlek, salmon for home use have also been acquired via rod and reel and removal from commercial harvests. However, all salmon that were reported harvested in the 2003 surveys were taken with subsistence nets or seines (Fall 2006b).

SOUTHWESTERN DISTRICT SUBSISTENCE SALMON FISHERY

The Southwestern District is described as the mainland waters from the outer point of the north shore of Granite Bay to Cape Fairfield, as well as the waters surrounding Knight, Chenega, Bainbridge, Evans, Elrington, and Latouche islands and their adjacent islands (5 AAC 24.200 (i)). Under regulations in place since 1988, salmon may be taken in the Southwestern District as well as in waters along the northwestern shore of Green Island from the westernmost tip of the island to the northernmost tip (5 AAC 01.648 (a)). The primary participants in this fishery are residents of Chenega Bay. Prior to 1992, permits were issued only in Chenega Bay, but since 1992, they have also been issued at the Cordova ADF&G office. Permits may be returned in person to the Cordova ADF&G office or the Chenega Village Council office, or mailed at the end of the fishing season. Permits include a harvest report, and fishers are required to fill in the dates fished and the number of each species of salmon caught each day.

Legal gear for this fishery includes seines up to 50 fathoms in length and 100 meshes deep with a maximum mesh size of 4 in, and gillnets up to 150 fathoms in length with a maximum size of 6¼ in. The open season is May 15–October 31 under the following fishing periods: 7 days per week from May 15 until 2 days before the commercial opening of the Eastern District; during the commercial fishing season at the time of commercial openers; and 7 days per week from 2 days after the closure of the commercial season through October 31. There are no bag or possession limits for this fishery.

In 2012, 23 permits were issued for this fishery and 14 were returned; the highest number of recorded permits returned since 1988. Because permit return rates for this fishery have been low in the past, data in Table 12-13 reflect reported harvests only. The reported harvest for 2012 was 700 salmon, consisting of 603 sockeye salmon, 77 chum salmon, and 20 coho salmon. The 2012 harvest represented an increase of the reported harvest in comparison with the recent 5-year (344 salmon), 10-year (439 salmon), and historical averages (556 salmon). However, it is likely that the harvest assessment program for this fishery continues to underestimate harvests. As shown in Table 12-14, household surveys in Chenega Bay in 2003 (Fall 2006b) provided an estimate of 1,690 salmon taken with subsistence methods compared to 677 (Table 12-13) based on returned permits for that same year.

PRINCE WILLIAM SOUND: GENERAL DISTRICTS

Subsistence fishing for salmon in the other districts of the Prince William Sound Area (other than the Upper Copper River, Copper River, Eastern, and Southwestern districts; also, the Valdez Nonsubsistence Area is closed to subsistence fishing) is open in conformance with regulations regarding gear, open areas and open periods. Permits are required and may be obtained from the Cordova ADF&G office. Annual limits are 15 salmon for a household of one, 30 salmon for a household of 2, and 10 salmon for each additional person in the household.

Since the creation of separate regulations for the waters fished by Tatitlek and Chenega Bay residents in 1988, it appears that participation in the general Prince William Sound fishery has been limited. However, further collection and analysis of data is necessary to support this hypothesis. In the last 22 years issued permits have been typically low with a 5-year average of 4 and a 10-year average of 8 (Table 12-15). In 2011, 14 permits were issued, and 12 were returned; the second highest return on record. The reported harvest for 2012 was 99 salmon, consisting of 67 sockeye salmon and 32 chum salmon. The 2012 harvest for the Prince William Sound general area is significantly higher than the 5 and 10-year average; 41 and 33 respectively (tables 12-15 and 12-16).

OTHER SUBSISTENCE FISHERIES IN THE PRINCE WILLIAM SOUND AREA

Subsistence halibut harvest estimates for eligible communities and tribes in the Prince William Sound Area communities of Cordova, Chenega Bay, and Tatitlek, are available for 2012 (Fall and Koster 2014).

In 2012, on-going harvest assessment programs did not exist for other subsistence finfish fisheries in the Prince William Sound Area. However, there is a subsistence permit available for the harvest of freshwater finfish species, which is issued out of the Glennallen office. Also, in the Upper Copper River watershed, resident species such as Arctic grayling, burbot, and whitefishes, among other species, are harvested for home use. Harvest estimates based on household surveys are available in the CSIS.

The Division of Subsistence, in collaboration with the Copper River Native Association, the Cheesh'Na Tribal Council, the Mentasta Tribal Council, and the Chitina Tribal Council, conducted a household survey to collect nonsalmon fish harvests and use information in Copper Basin communities for a 12-month period from October 2000–September 2001. In total, 472 households were interviewed, 42% of the estimated 1,193 households living in Copper Basin communities. The study produced estimated harvests by study community and gear type for burbot, Arctic char/Dolly Varden, lake trout, Arctic grayling, northern pike, longnose suckers, rainbow/steelhead trout, and whitefishes. Detailed summaries of study methods and findings appear in Simeone and Kari (2005).

Residents of Cordova, Chenega Bay, Tatitlek, Valdez, and Whittier take a variety of shellfish and marine finfishes for subsistence uses. Harvest estimates are available in the CSIS based upon systematic household surveys. Subsistence fishing for shrimp is open April 15–September 15, with no more than 5 pots per person and 5 pots per vessel, and no bag or possession limits. The year 2006 was the first year in which a permit was not required. In March 2009, the BOF adopted a Prince William Sound Pot Shrimp Management Plan that allocated 40% of the harvestable surplus of shrimp to commercial users and 60% to noncommercial users. Harvestable surplus is estimated annually prior to the start of the fishing season (April 15) with a surplus production model that requires more timely and precise estimates of noncommercial harvest than are provided by the statewide harvest survey (SWHS). This made it necessary to reinstate the noncommercial shrimp permit prior to the start of the 2009 shrimp pot fishery season. The Prince William Sound noncommercial shrimp permit requires all noncommercial users to report the date, location, duration, number of pots, and harvest of shrimp (gallons) for each set of pot gear made throughout the fishing season (April 15–September 15). Detailed summaries of harvest estimates and data from returned permits appear in Hochhalter and Hansen (2011) for 2010. Data for 2012 is available from the ADF&G Division of Sport Fish statewide harvest survey.⁵⁶ Subsistence fishing for Dungeness, Tanner, and king crabs in the Prince William Sound Management Area was closed, either by regulation or by emergency order, due to low stock status.

56. “Alaska Sport Fishing Survey, Southcentral Alaska Region.” 2014. Alaska Department of Fish and Game, Alaska Sport Fishing Survey. Accessed September 2014. <http://www.adfg.alaska.gov/sf/sportfishingsurvey/index.cfm?ADFG=region.target&targetRegion=2>.

Table 12-1.—Subsistence harvests by village fish wheel permits, Glennallen Subdistrict, 1997–2012.

Year ^a	Village	Reported subsistence harvest					Total
		Chinook	Sockeye	Coho	Steelhead	Other	
1997	Chistochina	105	342	139	88	1	675
1997	Gakona	8	1,242	0	0	0	1,250
1997	Kluti-Kaah	12	61	0	0	0	73
1999	Chickaloon	1	5	0	0	0	6
1999	Gakona ^b	0	0	0	0	0	0
1999	Kluti-Kaah	46	85	0	0	0	131
2000	Chickaloon	73	200	0	0	0	273
2000	Chistochina	1	880	0	0	0	881
2000	Kluti-Kaah	20	110	0	0	0	130
2001	Chickaloon	20	120	0	0	0	140
2001	Chistochina	4	1,203	0	0	0	1,207
2001	Kluti-Kaah	3	259	114	0	0	376
2002	Chickaloon	0	91	0	0	0	91
2002	Chitina ^c	0	0	0	0	0	0
2003	Chickaloon	8	105	0	0	0	113
2004	Chickaloon	5	178	0	0	0	183
2004	Chistochina	17	1,563	0	0	0	1,580
2005	Chistochina	4	545	0	0	0	549
2005	Chickaloon	20	533	0	0	1	554
2005	Gakona	9	442	0	0	0	451
2006	Chistochina	8	559	0	0	0	567
2006	Chickaloon ^c	0	0	0	0	0	0
2006	Chitina ^c	0	497	0	0	0	497
2007	Chitina ^c	0	0	0	0	0	0
2008	Chickaloon ^c	0	0	0	0	0	0
2008	Gakona	1	241	15	0	0	257
2009	Chickaloon ^c	0	0	0	0	0	0
2009	Kluti-Kaah	0	30	0	0	0	30
2010	Chickaloon	2	237	0	0	0	239
2010	Gakona ^a	0	0	0	0	0	0
2010	Kluti-Kaah ^c	0	0	0	0	0	0
2011	Gulkana	2	50	0	0	0	52
2011	Gakona	5	37	0	0	0	42

Source Mark Somerville, ADF&G Division of Sport Fish, Glennallen personal communication.

a. No fish wheel permits were issued in 2012.

b. Did not fish

c. Did not return permit.

Table 12-2.—Historical subsistence salmon harvests, Glennallen Subdistrict, 1989–2012.

Year	Permits		Estimated salmon harvest ^a					Total
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	
1989	386	360	796	28,724	67	0	0	29,587
1990	406	384	639	32,219	91	0	0	32,949
1991	712	645	1,314	39,364	241	0	0	40,919
1992	655	619	1,440	45,115	345	0	0	46,900
1993	773	696	1,443	54,003	76	0	0	55,523
1994	970	776	1,979	69,143	71	0	0	71,193
1995	858	726	1,968	54,336	975	0	0	57,280
1996	850	788	1,483	52,269	552	0	0	54,305
1997	1,136	1,058	2,608	83,692	183	0	0	86,483
1998	1,010	951	1,846	64,876	553	0	0	67,275
1999	1,102	1,040	3,234	76,456	1,145	0	0	80,835
2000	1,251	1,197	4,937	60,551	539	5	0	66,032
2001	1,239	1,176	3,480	81,960	1,142	20	0	86,601
2002	1,308	1,162	4,446	63,028	686	1	0	68,161
2003	1,227	1,101	3,344	64,618	650	0	0	68,612
2004	1,212	1,032	4,503	82,174	880	0	0	87,557
2005	1,234	1,070	2,785	91,715	252	0	0	94,752
2006	1,239	1,100	3,233	78,244	266	0	0	81,743
2007	1,458	1,277	4,125	86,678	308	0	0	91,110
2008	1,455	1,269	3,417	59,293	694	0	0	63,404
2009	1,364	1,138	3,341	67,887	287	0	0	71,515
2010	1,587	1,331	2,653	92,632	422	0	0	95,706
2011	1,586	1,328	3,649	81,216	1,131	0	0	85,996
2012	1,805	1,557	2,649	94,991	470	0	0	98,110
5-year average (2007–2011)	1,490	1,269	3,437	77,541	568	0	0	81,546
10-year average (2002–2011)	1,367	1,181	3,550	76,748	558	0	0	80,856
Historical average (1989–2011)	1,088	966	2,725	65,661	502	1	0	68,889

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

- a. Starting in 2002, estimates include salmon harvested under federal as well as state subsistence fishing regulations and permits.

Table 12-3.—Subsistence salmon harvests by community, Glennallen Subdistrict, 2012.

Community	Permits		Estimated salmon harvest ^a					Total
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	
Chistochina	5	5	23	711	0	0	0	734
Chitina	26	25	85	2,257	10	0	0	2,352
Copper Center	112	93	384	10,453	82	0	0	10,919
Copperville	4	4	12	333	0	0	0	345
Gakona	35	33	41	2,738	59	0	0	2,839
Glennallen	81	75	202	5,532	51	0	0	5,784
Gulkana	2	1	32	600	0	0	0	632
Kenny Lake	33	33	69	2,138	42	0	0	2,249
Lake Louise	1	1	0	44	0	0	0	44
McCarthy	17	12	0	54	0	0	0	54
Mendeltna	3	3	2	174	0	0	0	176
Nelchina	5	5	0	358	0	0	0	358
Paxson	1	1	1	114	0	0	0	115
Slana	20	18	1	688	0	0	0	689
Tazlina	45	41	128	3,672	26	0	0	3,827
Tolsona	7	7	5	342	0	0	0	347
Subtotal, Copper Basin	397	357	986	30,208	271	0	0	31,466
Anchor Point	1	1	5	61	0	0	0	66
Anchorage	380	304	453	15,810	11	0	0	16,274
Anderson	2	2	0	35	0	0	0	35
Barrow	2	1	20	114	0	0	0	134
Big Lake	5	4	9	328	0	0	0	336
Chickaloon	4	2	0	144	0	0	0	144
Chugiak	12	11	21	439	0	0	0	459
Clear	1	1	0	0	0	0	0	0
Cooper Landing	3	3	0	166	0	0	0	166
Delta Junction	52	47	50	1,610	0	0	0	1,660
Denali Park	1	1	0	0	0	0	0	0
Dot Lake	2	2	0	0	0	0	0	0
Eagle River	67	64	119	2,859	0	0	0	2,978
Ester	5	5	13	393	0	0	0	406
Fairbanks	268	224	266	10,818	48	0	0	11,132
Fort Greely	3	3	0	32	0	0	0	32
Fox	3	2	0	0	0	0	0	0
Funny River	1	1	0	0	0	0	0	0
Girdwood	1	1	0	16	0	0	0	16
Glacier View	2	2	6	84	0	0	0	90
Haines	1	1	0	2	0	0	0	2
Healy	2	1	0	138	18	0	0	156
Homer	2	2	4	188	0	0	0	192
Houston	1	1	6	145	0	0	0	151

-continued-

Table 12-3.--Page 2 of 2.

Community	Permits		Estimated salmon harvest ^a					Total
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	
Indian	1	1	0	2	0	0	0	2
Kenai	2	2	1	23	0	0	0	24
Kennicott	1	1	0	0	0	0	0	0
Klawock	1	1	0	0	0	0	0	0
Meadow Lakes	1	1	0	20	0	0	0	20
Mentasta Lake	4	4	1	163	0	0	0	164
Nabesna	3	3	0	91	0	0	0	91
Nenana	2	2	2	246	9	0	0	257
North Pole	112	92	149	4,888	0	0	0	5,036
Northway	8	6	1	203	0	0	0	204
Palmer	101	91	168	5,897	3	0	0	6,068
Peters Creek	2	2	4	33	0	0	0	37
Salcha	7	6	6	204	0	0	0	210
Shishmaref	1	1	0	3	0	0	0	3
Soldotna	3	3	14	547	63	0	0	624
Sutton	9	7	1	117	0	0	0	118
Talkeetna	2	1	0	28	0	0	0	28
Tanacross	1	1	0	51	0	0	0	51
Tok	45	42	15	1,586	0	0	0	1,601
Tonsina	6	6	3	74	0	0	0	77
Trapper Creek	1	0	0	0	0	0	0	0
Two Rivers	1	1	0	29	0	0	0	29
Unknown Community	1	1	0	16	0	0	0	16
Valdez	51	44	54	2,647	0	0	0	2,702
Wasilla	218	192	271	14,528	47	0	0	14,846
Willow	3	3	2	6	0	0	0	8
Subtotal, other communities	1,408	1,200	1,663	64,782	199	0	0	66,644
Total	1,805	1,557	2,649	94,991	470	0	0	98,110

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

a. Includes salmon harvested under federal as well as state subsistence fishing regulations and permits.

Table 12-4.—Historical subsistence and personal use salmon harvests, state Chitina Subdistrict permits, 1989–2012.

Year	Permits		Estimated salmon harvest					Total
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	
1989	4,584	4,353	2,269	56,547	865	0	0	59,681
1990	5,689	5,475	2,711	66,435	1,516	0	0	70,662
1991	6,222	5,990	4,092	78,412	3,378	0	0	85,882
1992	6,387	6,229	3,422	87,090	1,524	0	0	92,036
1993	7,914	7,914	2,729	89,629	1,358	0	0	93,716
1994	7,060	5,939	4,198	106,163	2,204	0	0	112,566
1995	6,762	5,442	5,617	94,494	5,861	0	0	105,972
1996	7,196	6,962	3,607	95,645	3,404	0	0	102,656
1997	9,086	8,919	5,470	149,020	160	0	0	154,650
1998	10,002	9,751	6,746	137,530	2,156	0	0	146,431
1999	9,941	9,607	5,964	142,682	2,199	0	0	150,845
2000	8,145	7,676	3,219	109,370	3,758	0	0	116,347
2001	9,458	8,356	3,171	137,047	2,687	0	0	142,905
2002	6,804	5,736	2,093	90,655	2,034	0	0	94,782
2003	6,440	5,438	1,962	84,790	2,579	0	0	89,332
2004	8,153	6,855	2,521	111,203	2,751	0	0	116,476
2005	8,232	6,768	2,155	129,506	1,885	0	0	133,546
2006	8,497	6,762	2,598	128,469	2,343	0	0	133,410
2007	8,378	7,187	2,782	131,460	1,747	0	0	135,990
2008	8,041	6,861	1,991	82,961	2,747	0	0	87,699
2009	7,958	6,908	229	93,766	1,667	0	0	95,662
2010	9,308	7,757	700	140,089	1,892	0	0	142,680
2011	9,167	7,566	1,118	138,089	1,866	0	0	141,073
2012	10,016	8,030	613	136,441	1,411	0	0	138,465
5-year average (2007–2011)	8,570	7,256	1,364	117,273	1,984	0	0	120,621
10-year average (2002–2011)	8,098	6,784	1,815	113,099	2,151	0	0	117,065
Historical average (1989–2011)	7,801	6,976	3,103	107,872	2,286	0	0	113,261

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Note Under state regulations, this fishery was classified as personal use from 1986 through 1999; in 2000, 2001, and 2002, it was classified as a subsistence fishery, in 2003, it was reclassified as personal use.

Table 12-5.—Personal use salmon harvests by community, state Chitina Subdistrict permits, 2012.

Community	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
Copper Center	20	15	1	69	0	0	0	71
Glennallen	14	13	1	187	6	0	0	195
Subtotal, Copper Basin	34	28	2	257	6	0	0	266
Allakaket	2	1	0	0	0	0	0	0
Anaktuvuk Pass	1	1	0	36	0	0	0	36
Anchor Point	5	4	0	0	0	0	0	0
Anchorage	2,653	2,095	190	30,202	211	0	0	30,604
Anderson	2	1	0	0	0	0	0	0
Aniak	1	1	1	20	0	0	0	21
Barrow	13	8	2	128	0	0	0	130
Bethel	2	1	0	50	0	0	0	50
Big Lake	40	35	2	440	24	0	0	466
Brevig Mission	1	1	0	0	0	0	0	0
Cantwell	5	4	0	105	0	0	0	105
Central	3	2	0	3	0	0	0	3
Chenega Bay	1	1	0	1	0	0	0	1
Chevak	1	1	0	0	0	0	0	0
Chickaloon	13	11	1	253	0	0	0	254
Chuathbaluk	1	0	0	0	0	0	0	0
Chugiak	150	123	11	1,584	11	0	0	1,606
Clear	4	4	0	66	0	0	0	66
Cooper Landing	1	1	0	0	0	0	0	0
Cordova	1	1	0	3	0	0	0	3
Delta Junction	406	371	30	6,209	45	0	0	6,284
Denali Park	27	22	5	233	1	0	0	239
Dot Lake	1	0	0	0	0	0	0	0
Eagle	1	1	0	0	0	0	0	0
Eagle River	315	267	33	3,769	47	0	0	3,850
Eielson AFB	89	55	3	1,206	3	0	0	1,212
Elmendorf AFB	31	25	2	213	0	0	0	216
Ester	91	73	7	1,561	15	0	0	1,583
Fairbanks	2,944	2,355	129	46,611	590	0	0	47,330
Fort Greely	21	15	1	293	0	0	0	294
Fort Richardson	9	4	0	0	0	0	0	0
Fort Wainwright	96	64	5	1,202	15	0	0	1,221
Fort Yukon	3	2	0	23	0	0	0	23
Gakona	1	1	0	30	0	0	0	30
Girdwood	38	29	5	616	5	0	0	626
Grayling	1	0	0	0	0	0	0	0
Haines	1	1	0	0	0	0	0	0
Healy	26	22	4	364	6	0	0	373
Homer	11	11	1	187	0	0	0	188
Hooper Bay	2	1	0	0	0	0	0	0
Houston	6	6	2	107	0	0	0	109

-continued-

Table 12-5.—Page 2 of 2

Community	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
Indian	6	6	0	116	0	0	0	116
Juneau	12	11	0	109	0	0	0	109
Kaktovik	2	1	0	0	0	0	0	0
Kasilof	1	0	0	0	0	0	0	0
Kenai	2	1	0	20	0	0	0	20
Ketchikan	1	1	0	0	0	0	0	0
Kobuk	1	1	0	40	0	0	0	40
Kodiak	3	0	0	0	0	0	0	0
Kotzebue	5	3	0	77	0	0	0	77
Lake Minchumina	1	1	0	30	0	0	0	30
McCarthy	1	0	0	0	0	0	0	0
McGrath	1	0	0	0	0	0	0	0
Meadow Lake	1	0	0	0	0	0	0	0
Minto	1	1	0	0	0	0	0	0
Moose Creek	1	1	0	0	0	0	0	0
Nenana	28	25	2	440	0	0	0	442
Nikiski	1	1	0	30	0	0	0	30
Nikolaevsk	2	2	0	30	0	0	0	30
Ninilchik	3	2	0	56	0	0	0	56
Nome	4	3	0	3	0	0	0	3
North Pole	850	641	40	13,040	93	0	0	13,173
Northway	1	0	0	0	0	0	0	0
Nulato	1	1	0	0	0	0	0	0
Palmer	512	441	43	6,430	95	0	0	6,568
Petersburg	2	1	2	38	0	0	0	40
Prudhoe Bay	1	0	0	0	0	0	0	0
Rampart	2	1	0	0	0	0	0	0
Saint Marys	1	0	0	0	0	0	0	0
Salcha	58	40	1	784	44	0	0	829
Seldovia	2	1	0	0	0	0	0	0
Seward	5	5	0	120	0	0	0	120
Sitka	1	1	0	30	0	0	0	30
Skagway	2	2	2	78	0	0	0	80
Soldotna	4	3	0	59	0	0	0	59
Stebbins	1	0	0	0	0	0	0	0
Sutton	60	51	4	918	0	0	0	921
Talkeetna	15	12	1	205	0	0	0	206
Tanacross	1	1	0	15	0	0	0	15
Tanana	1	0	0	0	0	0	0	0
Thorne Bay	1	1	1	10	0	0	0	11
Tok	16	14	2	305	0	0	0	307
Trapper Creek	5	4	1	88	0	0	0	89
Two Rivers	20	17	1	346	0	0	0	347
Valdez	192	152	3	2,459	23	0	0	2,485
Wainwright	1	0	0	0	0	0	0	0
Ward Cove	1	1	0	17	0	0	0	17
Wasilla	976	789	62	12,457	171	0	0	12,689
Willow	50	39	6	532	5	0	0	544

-continued-

Table 12-5.—Page 2 of 2

Community	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
Wiseman	1	1	0	0	0	0	0	0
Unknown Community	102	102	5	1,789	0	0	0	1,794
Subtotal, other communities	9,982	8,002	611	136,185	1,404	0	0	138,200
Total	10,016	8,030	613	136,441	1,411	0	0	138,465

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Table 12-6.—Historical subsistence salmon harvests, federal Chitina Subdistrict permits, 2003–2012.

Year	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
2003	99	71	33	1,316	152	0	0	1,500
2004	109	83	9	1,631	28	0	0	1,668
2005	77	64	27	1,498	0	0	0	1,526
2006	76	62	16	1,681	26	0	0	1,723
2007	97	86	29	1,095	41	0	0	1,165
2008	81	65	26	939	97	0	0	1,062
2009	68	34	15	1,522	22	0	0	1,560
2010	92	38	36	5,352	88	0	0	5,476
2011	84	42	21	3,090	14	0	0	3,125
2012	90	80	5	981	9	0	0	996
5-year average (2007–2011)	84	53	25	2,400	52	0	0	2,478
Historical average (1989– 2011)	87	61	24	2,014	52	0	0	2,089

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Table 12-7.—Subsistence salmon harvests by community, federal Chitina Subdistrict permits, 2012.

Community	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
Chitina	9	9	0	65	0	0	0	65
Copper Center	16	13	0	81	0	0	0	81
Coppsville	1	1	0	0	0	0	0	0
Dot Lake	1	1	0	32	0	0	0	32
Gakona	2	2	0	0	0	0	0	0
Glennallen	10	7	0	19	0	0	0	19
Kennicott	2	1	0	118	0	0	0	118
Kenny Lake	14	13	0	349	0	0	0	349
McCarthy	15	13	0	88	9	0	0	97
Northway	1	1	0	1	0	0	0	1
Slana	1	1	0	0	0	0	0	0
Tazlina	8	8	1	14	0	0	0	15
Tok	6	6	0	164	0	0	0	164
Tonsina	4	4	4	51	0	0	0	55
Total	90	80	5	981	9	0	0	996

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Table 12-8.—Historical subsistence salmon harvests, Batzulnetas fishery, 1987–2012.

Year	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1987	8	8	0	22	0	0	0	22
1988	0	0	0	0	0	0	0	0
1989	0	0	0	0	0	0	0	0
1990	0	0	0	0	0	0	0	0
1991	0	0	0	0	0	0	0	0
1992	0	0	0	0	0	0	0	0
1993	1	1	0	160	0	0	0	160
1994	4	4	0	997	0	0	0	997
1995	4	2	0	32	0	0	0	32
1996	0	0	0	0	0	0	0	0
1997	0	0	0	0	0	0	0	0
1998	1	1	0	382	0	0	0	382
1999	1	1	0	55	0	0	0	55
2000	1	1	0	55	0	0	0	55
2001	1	1	1	61	0	0	0	62
2002	1	1	0	208	0	0	0	208
2003	1	1	0	164	0	0	0	164
2004	1	1	0	182	0	0	0	182
2005	0	0	0	0	0	0	0	0
2006	0	0	0	0	0	0	0	0
2007	0	0	0	0	0	0	0	0
2008	0	0	0	0	0	0	0	0
2009	0	0	0	0	0	0	0	0
2010	3	3	0	106	0	0	0	106
2011	3	3	0	101	0	0	0	101
2012	3	3	1	136	0	0	0	137
5-year average (2007–2011)	1	1	0	41	0	0	0	41
10-year average (2002–2011)	1	1	0	76	0	0	0	76
Historical average (1987– 2011)	1	1	0	101	0	0	0	101

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Table 12-9.—Historical subsistence salmon harvests, Copper River District (Copper River Flats), 1965–2012.

Year	Permits		Estimated salmon harvest					Total
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	
1965	31	20	19	711	132	0	0	862
1966	45	31	68	254	0	0	0	322
1967	61	56	90	167	0	0	0	257
1968	17	15	12	41	0	0	0	53
1969	49	33	24	94	126	0	0	244
1970	32	27	78	212	0	0	0	290
1971	29	26	11	36	4	0	0	51
1972	104	79	196	749	70	0	0	1,015
1973	94	89	162	344	190	0	0	696
1974	9	5	9	7	4	0	0	20
1975	2	2	0	5	0	0	0	5
1976	27	14	2	19	0	0	0	21
1977	23	22	10	74	0	0	0	85
1978	34	28	45	22	15	0	0	81
1979	49	41	54	31	20	0	0	105
1980	39	35	21	30	19	0	0	70
1981	72	51	68	205	147	0	0	419
1982	108	90	72	761	127	0	0	960
1983	87	73	94	128	68	0	0	290
1984	118	104	77	368	153	0	0	598
1985	94	94	88	261	83	0	0	432
1986	88	85	89	360	49	0	0	498
1987	95	89	52	383	15	0	0	450
1988	114	97	69	266	49	0	0	384
1989	75	64	66	397	60	0	0	523
1990	88	76	69	543	95	0	0	707
1991	129	115	153	931	43	0	0	1,126
1992	126	113	158	875	47	0	0	1,080
1993	111	93	143	511	35	0	0	689
1994	101	97	171	494	70	0	0	734
1995	126	112	173	779	35	0	0	987
1996	176	157	309	1,086	53	0	0	1,448
1997	269	243	223	1,144	1,967	0	0	3,333
1998	245	230	314	905	724	0	0	1,944
1999	294	275	377	1,422	729	0	0	2,528
2000	416	400	717	4,534	46	18	3	5,318
2001	468	439	881	3,275	75	2	0	4,232
2002	355	331	589	3,289	30	2	0	3,910
2003	384	367	730	1,655	37	0	16	2,439
2004	511	487	1,163	1,910	48	5	3	3,129
2005	237	224	260	830	15	0	1	1,106
2006	421	399	779	4,355	1	0	0	5,135
2007	469	445	1,211	6,458	16	2	6	7,694
2008	506	482	495	4,161	55	0	21	4,732
2009	323	293	232	1,916	23	1	0	2,173
2010	326	320	281	2,034	27	22	0	2,365
2011	273	263	220	1,839	35	2	0	2,096

-continued-

Table 12-9.–Page 2 of 2.

Year	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
2012	378	359	248	4,499	0	19	0	4,767
5-year average (2007–2011)	379	361	488	3,282	31	5	5	3,812
10-year average (2002–2011)	381	361	596	2,845	29	3	5	3,478
Historical average (1965–2011)	167	154	237	1,082	118	1	1	1,439

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Table 12-10.–Subsistence salmon harvests by community, Copper River District (Copper River Flats), 2012.

Community	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
Anchorage	38	35	10	261	0	0	0	270
Big Lake	1	0						
Chenega Bay	1	1	0	0	0	0	0	0
Chugiak	2	2	0	30	0	0	0	30
Cordova	294	282	207	3,569	0	0	0	3,776
Delta Junction	2	2	0	0	0	0	0	0
Eagle River	1	1	0	30	0	0	0	30
Fairbanks	1	1	0	0	0	0	0	0
Girdwood	2	2	0	19	0	0	0	19
Homer	13	11	7	278	0	7	0	292
Hope	1	1	5	10	0	0	0	15
Juneau	1	1	1	20	0	0	0	21
Kodiak	1	1	0	0	0	0	0	0
Northway	1	1	0	0	0	0	0	0
Palmer	3	3	0	0	0	6	0	6
Seward	2	2	2	34	0	0	0	36
Soldotna	1	1	0	0	0	0	0	0
Sutton	1	1	0	1	0	6	0	7
Tatitlek	2	2	3	2	0	0	0	5
Valdez	2	2	4	83	0	0	0	87
Wasilla	8	7	9	163	0	0	0	173
Total	378	359	248	4,499	0	19	0	4,767

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Table 12-11.—Historical subsistence salmon harvests, Prince William Sound, Eastern District, 1988–2012.

Year	Permits		Reported salmon harvest					Total
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	
1988	17	NA	2	210	249	297	143	901
1989	14	NA	1	107	653	43	28	832
1990	13	NA	0	5	241	4	10	260
1991	19	NA	0	107	984	28	320	1,439
1992	15	NA	2	441	369	49	30	891
1993	18	NA	2	512	305	74	144	1,037
1994	14	NA	0	50	143	70	50	313
1995	15	0						
1996	6	NA	0	0	38	0	0	38
1997	6	NA	0	107	45	54	0	206
1998	11	NA	0	2	71	28	4	105
1999	17	NA	0	344	541	31	31	947
2000	12	3	0	140	468	40	40	688
2001	14	9	0	114	230	12	60	416
2002	19	8	6	437	278	66	71	858
2003	15	8	0	81	185	12	20	298
2004	18	12	2	358	505	28	105	998
2005	16	3	0	98	286	16	200	600
2006	11	1	0	3	18	25	35	81
2007	14	0						
2008	1	1	0	60	0	0	0	60
2009	12	4	0	170	131	0	0	301
2010	8	5	0	165	142	10	50	367
2011	10	4	0	922	536	22	0	1,480
2012	16	8	15	954	75	8	0	1,052
5-year average (2007–2011)	9	3	0	329	202	8	13	552
10-year average (2002–2011)	13	5	1	241	231	19	54	546
Historical average (1988–2011)	13	4	1	202	292	41	61	596

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

NA = Data not available.

Table 12-12.–Estimated harvests of salmon for home use, Tatitlek, 2003.

Species	Estimated salmon harvest			
	Subsistence methods	Rod and reel	Removed from commercial harvests	All methods
Chinook	27	0	0	27
Sockeye	306	0	0	306
Coho	651	0	0	651
Chum	13	0	0	13
Pink	77	0	0	77
All salmon	1,075	0	0	1,075
Estimated number of households harvesting ^a	13 households	0 households	0 households	13 households (any method)

Source Fall (2006).

a. Number of households in the community = 27; 15 (93%) were interviewed.

Table 12-13.–Historical subsistence salmon harvests, Prince William Sound, Southwestern District, 1988–2012.

Year	Permits		Reported salmon harvest					Total
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	
1988	10	NA	1	50	8	294	251	604
1989	8	NA	0	322	0	180	554	1,056
1990	7	NA	1	36	5	2	20	64
1991	12	NA	3	345	42	53	195	638
1992	14	NA	1	526	23	99	313	962
1993	22	NA	2	835	50	124	232	1,243
1994	16	NA	5	192	77	161	402	837
1995	10	NA	2	152	67	41	67	329
1996	7	NA	0	107	7	46	105	265
1997	5	NA	44	193	30	272	110	649
1998	4	NA	13	114	20	119	65	331
1999	14	NA	57	499	62	101	168	887
2000	12	8	24	39	229	143	211	646
2001	16	9	2	119	92	146	95	454
2002	10	5	10	142	123	60	83	418
2003	13	7	6	219	156	147	149	677
2004	8	5	3	535	44	84	56	722
2005	13	8	10	515	84	174	124	907
2006	7	6	0	159	1	111	28	299
2007	4	3	2	293	27	55	4	381
2008	15	3	4	97	75	30	70	276
2009	5	4	2	168	26	84	5	285
2010	9	5	0	55	0	87	6	148
2011	17	6	2	134	26	60	50	272

-continued-

Table 12-13.—Page 2 of 2.

Year	Permits		Reported salmon harvest					Total
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	
2012	23	14	0	603	20	77	0	700
5-year average (2007–2011)	10	4	2	149	31	63	27	344
10-year average (2002–2011)	10	5	4	232	56	89	58	439
Historical average (1988– 2011)	11	6	8	244	53	111	140	556

Source ADF&G Division of Subsistence, ASFDB 2012 (ADF&G 2013).

NA = Data not available.

Table 12-14.—Estimated harvests of salmon for home use, Chenega Bay, 2003.

Species	Estimated salmon harvest			
	Subsistence methods	Rod and reel	Removed from commercial harvests	All methods
Chinook	79	36	19	134
Sockeye	829	100	0	929
Coho	331	263	0	594
Pink	201	131	0	333
Chum	250	81	0	331
Other/unknown	0	56	0	56
All salmon	1,690	668	19	2,376
Estimated number of households harvesting ^a	8 households	10 households	1 household	14 households (any method)

Source Fall (2006).

a. Number of households in the community = 20; 16 (80%) were interviewed.

Table 12-15.—Historical subsistence salmon harvests, Prince William Sound general, 1960–2012.

Year	Permits		Estimated salmon harvest					Total
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	
1960	50	NA	1	139	505	27	1,292	1,964
1961	12	NA	3	41	123	3	732	902
1962	9	NA	0	0	119	142	214	475
1963	9	NA	0	0	406	24	298	728
1964	15	NA	0	11	0	0	900	911
1965	22	16	0	0	0	34	246	281
1966	3	3	0	3	19	50	20	92
1967	4	3	0	0	5	0	5	11
1968	4	3	0	0	27	0	208	235
1969	7	3	0	0	37	0	0	37
1970	1	1	0	0	0	0	0	0
1971	3	2	0	0	0	0	69	69
1972	0	0	0	0	0	0	0	0
1973	19	16	0	0	343	0	0	343
1974	3	1	0	0	0	0	0	0
1975	2	0						
1976	0	0	0	0	0	0	0	0
1977	4	4	0	0	0	0	0	0
1978	3	2	0	0	0	0	0	0
1979	15	2	0	0	0	0	0	0
1980	26	15	0	12	10	0	0	23
1981	12	8	0	5	44	3	0	51
1982	35	27	0	109	5	31	40	185
1983	26	21	0	27	45	98	11	181
1984	8	8	0	10	0	2	11	23
1985	22	16	1	37	22	36	19	116
1986	25	14	0	9	27	0	0	36
1987	18	17	5	33	6	17	0	61
1988	7	7	2	51	7	9	10	79
1989	11	7	0	0	0	5	0	5
1990	8	8	0	0	7	0	4	11
1991	9	5	0	4	0	0	0	4
1992	10	6	0	33	0	0	0	33
1993	6	6	1	104	10	0	0	115
1994	5	4	0	0	0	0	0	0
1995	4	2	0	0	0	0	0	0
1996	10	7	0	0	0	0	0	0
1997	4	3	0	4	0	0	0	4
1998	4	3	0	0	0	0	0	0
1999	3	3	0	0	0	0	0	0
2000	3	3	0	0	0	0	0	0
2001	5	5	0	0	0	0	0	0
2002	11	9	0	38	0	9	11	57
2003	11	11	0	48	0	3	0	51
2004	8	7	0	12	0	5	0	17
2005	14	13	0	4	0	0	0	4
2006	11	9	0	20	30	0	0	50

-continued-

Table 12-15.–Page 2 of 2.

Year	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
2007	3	3	0	30	0	0	0	30
2008	11	10	1	33	0	0	0	34
2009	1	1	0	0	0	0	0	0
2010	1	1	0	0	0	0	0	0
2011	4	4	29	40	1	10	5	85
2012	14	12	0	67	0	32	0	99
5-year average (2007–2011)	4	4	6	21	0	2	1	41
10-year average (2002–2011)	8	7	3	22	3	3	2	33
Historical average (1960–2011)	9	7	1	14	26	10	56	107

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

NA = Data not available.

Table 12-16.–Subsistence salmon harvests by community, Prince William Sound general, 2012.

Community	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
Anchorage	5	4	0	5	0	20	0	25
Cooper Landing	1	1	0	0	0	0	0	0
Cordova	1	1	0	10	0	0	0	10
Manley Hot Springs	1	1	0	0	0	0	0	0
Palmer	1	1	0	0	0	0	0	0
Seward	1	1	0	0	0	0	0	0
Talkeetna	1	1	0	0	0	0	0	0
Wasilla	2	1	0	52	0	12	0	64
Willow	1	1	0	0	0	0	0	0
Total	14	12	0	67	0	32	0	99

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

CHAPTER 13: THE SOUTHEAST REGION

INTRODUCTION

The Southeast region is composed of 2 areas: the Southeastern Alaska Area, which includes all waters between a line projecting southwest from the westernmost tip of Cape Fairweather and Dixon Entrance, and the Yakutat Area, which includes all waters of Alaska between the longitude of Cape Suckling and the longitude of Cape Fairweather. Positive customary and traditional use (C&T) findings, which allow for subsistence fishing opportunity, have been made for most of the waters in the Southeast region. In areas where no C&T finding exists, personal use fisheries may be authorized. In addition, the Joint Board identified 2 nonsubsistence areas in the Southeast region: the Juneau Nonsubsistence Area and the Ketchikan Nonsubsistence Area (figures 13-1 and 13-2) (5 AAC 99.015). By regulation, no subsistence fisheries may be authorized in nonsubsistence areas.

The Southeast region is divided into 6 areas for management purposes:

- Yakutat Management Area,
- Haines Management Area,
- Juneau Management Area,
- Sitka Management Area,
- Petersburg Management Area, and
- Ketchikan Management Area.

HARVEST ASSESSMENT PROGRAMS

Since 1990, any Alaska resident may harvest salmon under state subsistence regulations. In the Southeast region permits are required for both subsistence and personal use salmon fisheries. In the Haines and Yakutat management areas, the permit is only for subsistence fisheries; no personal use fisheries are authorized in these areas. Permits in the other 4 management areas apply to area subsistence fisheries and area personal use fisheries. The Division of Commercial Fisheries is responsible for administering the subsistence and personal use salmon permit programs in the Southeast region. Permits are available at area offices. Department personnel or authorized designees also travel to Angoon, Hoonah, and Kake in the spring of each year to issue permits. Since 1985, annual subsistence salmon harvest assessments, based on the permit reporting program, have occurred in the Southeast Alaska area. Annual harvest assessments did not begin in the Yakutat area until 1989.

Subsistence and personal use permits are a mixture of personal information about the permittee, rules and regulations governing harvesting methods and means, and a harvest calendar that details all the fish caught under each permit. Permits are issued for each management area. Area management biologists have discretionary authority to change permit conditions, either before the season begins or inseason, through issuing emergency orders. While permit conditions vary based on the management area the permit applies to, some conditions are based on regionwide regulations and are therefore listed on all permits issued in the Southeast region. These conditions include: to be issued a permit a person must be an Alaska resident, only 1 permit per household will be issued, fishermen must record their harvests on a daily basis prior to leaving the fishing location, and the permit must be with the permittee, other authorized members of the household, or authorized proxy while taking or transporting subsistence salmon. Other standard permit conditions include removal of dorsal fins of subsistence salmon and both tips of the tail fin of personal use salmon and a prohibition on fishing within 300 ft of a dam, fish ladder, weir, culvert, or other artificial obstruction. Sport-caught and subsistence-caught salmon cannot be possessed on the same day. Salmon may not be harvested for subsistence use by a line attached to a rod or

pole, except in the Redoubt Bay and Lake subsistence salmon fishery. In addition, regulations specify that a permit will not be issued to anyone who has failed to return a permit issued for the previous year. Generally, however, area management offices will accept a harvest record for the previous year at the time a person applies for a current year's permit.

The information on the harvest calendars, such as locations and amount of harvest, species harvested, and date of harvest, is entered into *Alexander: the Integrated Fisheries Database for Southeast Alaska and Yakutat*. The database also includes the names and addresses of all who held subsistence or personal use permits, along with their harvest record. All of the harvest information collected each year composes the basis of the harvest assessment program in the Southeast region.

SUBSISTENCE SALMON HARVESTS IN 2012

In 2012, the total estimated subsistence and personal use salmon harvest for the Southeast region, based on returned permits, was 58,392 fish (Table 13-1). This is above the total estimated harvest for 2011 (52,350 salmon) as well as the recent 5-year average (54,751 salmon), but below the 10-year average (60,484 salmon) (Table 13-2). Sockeye salmon usually make up the largest proportion of subsistence salmon catches in Southeast Alaska, in contrast to the commercial fishery which has been dominated by pink salmon harvests since the early 1900s (Tingley and Davidson 2011). As expected, in 2012, sockeye salmon contributed the greatest amount to the overall harvest at 50,427 fish (86%), followed by 3,210 coho salmon (6%), 2,806 pink salmon (5%), 1,186 chum salmon (2%), and 763 Chinook salmon (1%) (Table 13-1; Figure 13-3). While the numbers of each species of salmon harvested differed from the 2011 harvest, the overall contribution of each species to the total harvest did not change significantly; the 2012 harvest was stronger in sockeye salmon and slightly weaker in coho salmon. Harvests of sockeye, coho, and chum salmon increased over 2011 estimates, while harvests of pink and Chinook salmon decreased. For a comparison, in the commercial fisheries in 2012, sockeye, pink, and Chinook salmon were below their 10-year and historic averages, coho salmon was below the ten-year average but on par with the historical average, and chum salmon harvests were above average (Conrad and Davidson 2013). Pink salmon have exhibited a strong odd-year, weak even-year return to the commercial fisheries since 2006, and this pattern appears visible in the subsistence harvests of 2012 as well. The estimated subsistence salmon harvests by management area were as follows: Sitka 15,650 (27%), Ketchikan 11,510 (20%), Haines 10,386 (18%), Juneau 9,530 (16%), Yakutat 8,036 (14%), and Petersburg 3,280 (5%) (Table 13-3, Figure 13-4). Compared to 2011, harvests in Sitka increased the most dramatically with 5,000 more salmon harvested than in 2011; small overall increases were also seen in the Juneau and Ketchikan Management Areas. Harvests in Petersburg and Haines management areas decreased since 2011.

The number of permits issued per year, on average, for the 10-year time period of 2002–2011, has been 3,260 (Table 13-2). In both 2011 and 2012, an average number of permits were issued. In 2012, a total of 3,267 permits were issued and 2,853 permits were returned. This corresponds to a region-wide response rate of 87%, approximately the same as the 2011 response rate, and higher than the 10 year average of 82%. The numbers reported on the returned permits are expanded to account for the unreturned permits. Prior to 1996, only permits returned with harvest data were included in the database and reported harvests were not expanded to account for permits not returned.

YAKUTAT MANAGEMENT AREA

Yakutat Area Subsistence Fisheries

Background and History

The Yakutat Management Area stretches from Cape Fairweather to Cape Suckling and encompasses Yakutat Area subsistence fisheries. Fishing areas used by Yakutat residents are under the management responsibility of the Division of Commercial Fisheries' Yakutat Area offices. C&T findings by the BOF for salmon identify the freshwaters upstream from the terminus of streams and rivers from the Doame

River to the Tsiu River, the waters of Yakutat Bay and Russell Fjord, and the waters of Icy Bay (5 AAC 01.666 (a)(3)). Unlike the other management areas, in the Yakutat Management Area, subsistence salmon fishing locations are not restricted to specific streams, nor are there daily or annual limits on the number of fish harvested.

Yakutat is the only community within the Yakutat Management Area. In 2012, the population of the Yakutat City and Borough was 621, a decrease of 26 residents over the 2011 estimate.⁵⁷

Regulations

There were no significant changes made to the 2012 subsistence permit. It remained valid only in the areas with positive C&T findings. No daily or annual limits or allowable subsistence gear was specified. In addition to the common permit conditions listed above, the weekly subsistence fishing period during the commercial salmon net season was from 6:00 AM Friday to 6:00 PM Saturday. On the Situk River, subsistence fishers were required to attend their nets when they were being used to harvest salmon. In Yakutat Bay, each permit holder needed to attend their net at least once a day. The permit was valid through December 31. An emergency order was released on May 17, 2012 which closed subsistence fishing for Chinook salmon in the Situk-Ahrnklin subsistence fishery at 6:00 am May 20. The order was issued because the preseason forecast of the Situk River Chinook salmon return suggested that the return would be low, and conservation actions were required to limit harvest on the stock.⁵⁸

Harvest Assessment Program

The estimated total subsistence salmon harvest for the Yakutat Management Area in 2012, based on returned permits, was 8,036 salmon, including 6,142 sockeye salmon (76%), 1,286 coho salmon (16%), 384 Chinook salmon (5%), 204 pink salmon (3%), and 20 chum salmon (<1%)(Table 13-3). An estimated 116 permits were fished in the Yakutat Area (Table 13-3). This reflects a slight increase in the number of permits fished compared to 2011 and an increased harvest, particularly of sockeye salmon.

Residents of Yakutat were issued 142 subsistence permits, with 112 returned (79%). The estimated total subsistence salmon harvest for the community of Yakutat in 2012 was 7,034 fish, including 5,336 sockeye salmon (76%), 1,116 coho salmon (16%), 369 Chinook salmon (5%), 202 pink salmon (3%), and 11 chum salmon (<1%) (Table 13-4).

HAINES MANAGEMENT AREA

Haines Area Subsistence Fisheries

Background and History

The Haines Management Area, encompassing the Haines Area subsistence fisheries, stretches from Little Island in Lynn Canal north to Chilkat Inlet, and includes the waters of the Chilkat River, as well as the waters in the Chilkoot Inlet to Skagway. Subsistence salmon fisheries in the waters traditionally used by the residents of the Haines area are under the management responsibility of the Division of Commercial Fisheries' Haines Area office. Positive C&T findings for salmon identify all the waters of the Chilkat River and Chilkat Inlet north of the latitude of Glacier Point, and in the Chilkoot River, Lutak Inlet, and Chilkoot Inlet north of the latitude of Battery Point, excluding waters of Taiya Inlet north of the latitude of the tip of Taiya Point (5 AAC 01.716 (a)(2)).

There are several communities in the Haines Management Area: the city of Haines and surrounding borough, which includes the settlements of Covenant Life, Lutak, Mosquito Lake, Mud Bay, and Excursion Inlet, as well as Klukwan on the Chilkat River and Skagway at the head of Chilkoot Inlet. In

57. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. "Research and Analysis Homepage." Accessed July 2014. <http://labor.alaska.gov/research/pop/popest.htm>

58. Alaska Department of Fish and Game Division of Commercial Fisheries, "Yakutat Subsistence Announcement," news release, May 17, 2012. Accessed July 23, 2014. <http://www.adfg.alaska.gov/static/applications/DCFnewsrelease/129954947.pdf>

2012, the combined population of these communities was 3,668, which is approximately the same estimate as in 2011.⁵⁹ The populations of Haines and Skagway are predominantly non-Native, while Klukwan continues to have a predominantly Alaska Native population.

Regulations

Permit conditions did not change from the 2011 permit. In 2012, the subsistence permit for the Haines Management Area was valid for the waters of Section 15A in all waters of the Chilkat River and Chilkat Inlet north of the latitude of Glacier Point and in the Lutak Inlet and Chilkoot Inlet north of the latitude of Battery Point. The permit provided for an open season of June 1–September 30 for sockeye, coho, pink, and chum salmon in the Chilkat River, Chilkat Inlet, and Lutak Inlet. In regulation, the subsistence net fisheries in Chilkat Inlet north of a line extending from approximately 1 mile south of Anchorage Point to just north of the Letnikof boat ramp are closed through July 15. The subsistence net fisheries in the Chilkat River, excluding that portion of the river from Haines Highway mile 19 upstream to Wells Bridge, are closed from the third week of June through the fourth week of July. Inseason, these closures were extended through 11:59 p.m. July 31 due to assessments indicating a weak and late inriver Chinook salmon run. Limits for the season for sockeye salmon were 25 in possession and 50 annually; for coho salmon, 20 in possession and 40 annually; and for pink and chum salmon, 75 in possession and 100 annually. Chinook salmon, rainbow/steelhead trout, and Arctic char/Dolly Varden could be retained only if taken incidentally by gear operated under the terms of the permit.

In the salt waters of Lynn Canal (District 15), including Chilkat, Chilkoot, and Lutak inlets, subsistence salmon could not be taken during closed periods of the commercial salmon net fishery, except salmon could be taken in the salt waters of Chilkoot Inlet north of the latitude of Battery Point and in Chilkat Inlet north of Glacier Point on the Saturday before any open period of the Section 15A commercial salmon net fishery.

Allowable gear types in the Haines Management Area subsistence fishery are set and drift gillnets. This gear could be used to take salmon in the mainstem and side channels, but not in the tributaries, of the Chilkat River from Mile 4 of the Haines Highway to 1 mile upstream of Wells Bridge. Drift and set gillnets could not exceed 50 ft in length when fishing in the Chilkat River, and drift gillnets fished in marine waters could not exceed 50 fathoms in length. In the Chilkat River, the permit holder was required to be physically present at the net while it was in use.

Harvest Assessment Program

The estimated subsistence salmon harvest in the Haines Management Area in 2012 was 10,386 salmon, including 8,663 sockeye salmon (83%), 786 pink salmon (8%), 456 chum salmon (4%), 257 Chinook salmon (2%), and 224 coho salmon (2%) (Table 13-3). The overall salmon harvest was more than 1,000 fish greater than the 2011 harvest. The majority of this increased harvest came from sockeye salmon. Pink salmon harvests decreased significantly, from composing 26% of the overall harvest in 2011 to just 8% in 2012. An estimated 384 permits were fished in the Haines Management Area in 2012.

In Haines Borough, 426 permits were issued and 406 were returned (95%). Permits issued to residents of the city of Haines, Mud Bay, Mosquito Lake, Covenant Life, or Lutak would be included in the Haines totals. In Klukwan, 10 permits were issued and 9 were returned. Sixteen residents of Skagway were issued permits and all were returned. In Excursion Inlet, 3 permits were issued and all were returned. The estimated salmon harvest by Haines, Klukwan, Skagway, and Excursion Inlet residents combined (9,339 salmon total) included 7,721 sockeye salmon (82%), 722 pink salmon (8%), 422 chum salmon (5%), 241 Chinook salmon (3%), and 234 coho salmon (3%) (Table 13-4).

59. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. "Research and Analysis Homepage." Accessed July 2014. <http://labor.alaska.gov/research/pop/popest.htm>

JUNEAU MANAGEMENT AREA

The Juneau Management Area encompasses subsistence fisheries in the Angoon Subsistence Area and the Hoonah Subsistence Area, as well as personal use fisheries in the Juneau area. Subsistence and personal use harvests by residents of Elfin Cove, Tenakee Springs, Gustavus, and Pelican occur primarily, but not exclusively, in the Juneau Management Area. Management responsibility for the area rests with both the Juneau and Sitka area offices. Overall, in 2012 there were an estimated 474 permits fished in the Juneau Management Area with an estimated harvest of 9,530 (Table 13-3). Sockeye salmon harvests constituted 89% of the total harvest.

Angoon Subsistence Area

Background and History

Subsistence salmon fisheries in the waters traditionally used by the residents of Angoon are under the management responsibility of the Division of Commercial Fisheries' Juneau and Sitka area offices. In 1989, the BOF adopted a positive C&T finding for salmon in the waters of District 12 south of a line from Fishery Point to South Passage Point and north of the latitude of Point Caution, and in waters of Section 13C east of the longitude of Point Elizabeth (5 AAC 01.716 (a)(6)).

The residents of Angoon are the principal subsistence fishers in this area. In 2012, Angoon had a population of 455, a slight decrease over the 2010 estimate.⁶⁰ Angoon Tlingit have traditionally used most of the west coast of Admiralty Island, from Hawk Inlet to the southern tip of Admiralty Island, and lands and waters of the east coasts of Chichagof and Baranof islands. Over the years, the waters of Kootznahoo Inlet, Favorite Bay, and Hood Bay to the south; Mitchell Bay, Salt Lake, and Kanalku bays further east; and Chatham Strait to the west have provided the people of Angoon with salmon and other marine resources.

Regulations

Subsistence permit conditions in 2012 did not differ from 2011, but some changes were made to subsistence salmon fishing limits and seasons. The 2012 subsistence permit was valid for waters with a positive C&T finding. The open season for sockeye salmon in Kanalku Bay and Basket Bay (Kook Lake outlet) was from June 1–July 31, with a limit of 20 fish in possession and annually at Kanalku Bay and 15 fish in possession with an annual limit of 30 fish in Basket Bay; in Sitkoh Bay from June 1–August 31, with a possession and annual limit of 50 fish; and in Hasselborg River–Salt Lake from July 1–August 15, with a limit of 25 fish in possession and annually. The open period for subsistence coho salmon fishing on Hasselborg River–Salt Lake was from July 1–October 31 with a possession and annual limit of 20 fish. Coho salmon could also be taken in other streams in the Angoon subsistence areas from June 1–October 31, with limits of 20 in possession and 40 annually from all combined streams. Pink salmon could be harvested in all streams in the area from June 1–September 30, with a possession and annual limit of 150 fish. The season for chum salmon in all streams of the area was from June 1–October 31, and the possession and annual limit was 50 fish. Chinook salmon, rainbow/steelhead trout, and Arctic char/Dolly Varden could be retained only if taken incidentally by gear operated under the terms of the permit.

Salmon taken under the subsistence or personal use regulations could not be subsequently used as bait for commercial fishing purposes. Gaffs, spears, beach seines, dip nets, drift gillnets, and cast nets were the allowable subsistence gear types. Drift gillnets could not exceed 50 fathoms in length; set gillnets could not be used. Snagging or fishing with a rod or reel was prohibited.

60. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. "Research and Analysis Homepage." Accessed July 2014. <http://labor.alaska.gov/research/pop/popest.htm>

Harvest Assessment Program

The estimated salmon harvest in the Angoon Area subsistence fisheries in 2012 was 1,385 salmon, including 1,161 sockeye salmon (84%), 165 pink salmon (12%), and 59 coho salmon (4%) (Table 13-3). The 2012 salmon harvest was approximately 700 salmon less than the 2011. The harvests of all species of salmon, except pink salmon, decreased. An estimated 54 permits were fished in the area, compared to 81 permits fished in 2011.

The estimated salmon harvest for the community of Angoon in 2012, based on 98 permits issued and 77 returned (79%), totaled 1,410 salmon, including 1,308 sockeye salmon (93%), 51 coho salmon (4%), and 51 pink salmon (4%)(Table 13-4). The number of permits issued in Angoon in 2012 was similar to the number issued in 2011, as was the overall salmon harvest. The composition of the harvest changed, however, with a higher percentage of sockeye salmon being caught, many fewer coho salmon being harvested, and no chum or Chinook salmon harvest being reported.

Hoonah Subsistence Area

Background and History

Subsistence salmon fisheries in the waters traditionally used by the residents of Hoonah are under the management responsibility of the Division of Commercial Fisheries' Juneau and Sitka area offices. In 1989, the BOF adopted a positive C&T finding for the salmon in those waters of District 12 that were in Basket Bay inside a line from lat. 57°30.83' N, long. 134°53.20' W, to lat. 57°39.28' N, long. 134°53.88' W; in those waters of District 13 that were along the western shore of Yakobi Island east of a line from Cape Spencer light to Surge Bay light; and in the waters of sections 14B and 14C (5 AAC 01.716 (a)(4)).

The residents of Hoonah are the principal users of waters in the Hoonah Subsistence Area. In 2012, Hoonah had a population of 776, continuing a slightly increasing trend in population estimates since 2010.⁶¹

Regulations

No changes were made to the 2012 subsistence salmon permit for the Hoonah area. The 2012 permit was valid in the waters identified above with a positive C&T finding and provided open seasons and limits for sockeye salmon at the following locations: Surge Bay, Hanus Bay (Lake Eva), and Neva Creek from June 1–August 15, Hoktaheen Cove from June 1–July 20, and Berg Bay from June 1–July 31. Limits at these locations varied: 50 sockeye salmon annually and in possession were allowed at Surge and Hanus bays and at Hoktaheen Cove; a limit of 40 fish in possession and annually was in effect at Neva Creek; and Berg Bay had a limit of 25 fish annually and in possession. Pink salmon could be harvested under a subsistence permit in all streams in the Hoonah Subsistence Area from June 1–September 30, with a possession and annual limit of 150 fish. Chum salmon could be harvested in the same waters from June 1–October 31, with a possession and annual limit of 50 fish. Coho salmon could be taken in streams in the subsistence area from June 1–October 31, with limits of 20 in possession and 40 annually. Chinook salmon, rainbow/steelhead trout and Arctic char/Dolly Varden could be retained only if taken incidentally by gear operated under the terms of the permit.

Salmon taken under subsistence regulations could not be subsequently used as bait for commercial fishing purposes. Gaffs, spears, beach seines, dip nets, drift gillnets, and cast nets were the types of subsistence gear allowed in the Hoonah Subsistence Area. Drift gillnets could not exceed 50 fathoms in length; set gillnets could not be used.

61. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. "Research and Analysis Homepage." Accessed July 2014. <http://labor.alaska.gov/research/pop/popest.htm>

Harvest Assessment Program

The estimated salmon harvest in the Hoonah Subsistence Area in 2012 was 1,270 salmon, including 969 sockeye salmon (77%), 142 pink salmon (11%), 97 chum salmon (8%), and 62 coho salmon (5%) (Table 13-3). The 2012 harvest was about 700 salmon fewer than the 2011 harvest. Decreases in harvests were seen most notably in sockeye salmon, which was estimated at 1,739 salmon in 2011 and coho salmon, the 2011 harvest of which was 156 fish. Pink and chum salmon harvests both increased. An estimated 54 permits were fished in the Hoonah Subsistence Area in 2012 in comparison to 75 permits fished in 2011.

For the community of Hoonah, in 2012, 122 permits were issued and 88 were returned (72%) with a total estimated harvest of 2,016. The harvest consisted of 1,543 sockeye salmon (77%), 326 pink salmon (16%), 103 chum salmon (5%), and 44 coho salmon (2%). No Chinook salmon were harvested (Table 13-4). More permits were issued to Hoonah residents as compared to 2011, but the return rate of those permits declined. The overall harvest was higher in 2012, with increases in all harvests except coho salmon. The greatest increase was seen in sockeye salmon harvests, which were estimated at 996 in 2011.

Elfin Cove, Gustavus, Pelican, and Tenakee Springs Subsistence and Personal Use Salmon Fishing

Background

Subsistence and personal use salmon fisheries in the waters traditionally used by the residents of Elfin Cove, Gustavus, Pelican, and Tenakee Springs are under the management responsibility of the Division of Commercial Fisheries' Juneau and Sitka area offices. Fishers from these communities fish primarily in districts 11, 12, 13, and 14; harvests are included in the Angoon Subsistence Area, Hoonah Subsistence Area, and Juneau Personal Use Area in Table 13-3. Elfin Cove fishers harvest salmon from Hoktaheen Cove in District 13. Gustavus fishers harvest salmon primarily from Surge Bay and Hoktaheen Cove in District 13, but also from the Taku River in District 11, the Berg River in District 14, and the Chilkat River in District 15. Residents of Pelican and Tenakee Springs harvest salmon at Kook Creek and Kook Lake Outlet in Basket Bay, and Takanis Bay and Hoktaheen Cove in District 13. Most of these areas have positive C&T findings as described in other sections of this report.

In 2012, Elfin Cove had a population of 20; Gustavus—489 residents; Pelican—82 residents; and Tenakee Springs—151 residents.⁶² The population of each community, with the exception of Pelican, increased slightly over 2011 estimates.

Regulations

Permit regulations applying to fishers in this area can be found under the Hoonah, Angoon, Haines, and Juneau subsections.

Harvest Assessment Program

In 2012, the number of salmon reported on permits from Elfin Cove, Gustavus, Pelican, and Tenakee Springs was modest. One permit was issued but not returned in Elfin Cove. In Gustavus, 22 permits were issued and 21 were returned. The estimated harvest for Gustavus was 88 total salmon, a significant decrease from 2011, when 481 fish were reported. The harvest consisted of 84 sockeye salmon (96%), 2 pink salmon (2%), and 1 chum salmon (1%). Three permits were issued to Pelican residents and three to Tenakee Springs residents; all were returned. Estimated harvests for these 2 communities were 63 salmon, a decrease of more than 100 salmon since 2011. Of that, 61 were sockeye salmon (97%), 1 was Chinook salmon (2%) and 1 was coho salmon (2%) (Table 13-4).

62. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. "Research and Analysis Homepage." Accessed July 2014. <http://labor.alaska.gov/research/pop/popest.htm>

Juneau Personal Use Area

Juneau fishers primarily harvest sockeye salmon from the Taku River and Sweetheart Creek in District 11, which is in the Juneau Nonsubsistence Area (Figure 13-1). These waters are under the management responsibility of the Division of Commercial Fisheries' Juneau Area office. Personal use regulations apply to salmon fishing for home uses in this area. Juneau area residents were the principal participants in the designated personal use fisheries in District 11. In 2012, the city and borough of Juneau had a population of 32,838, an increase of approximately 500 residents over 2011.⁶³

Regulations

The 2012 personal use permit conditions remained the same as 2011. Personal use regulations applied in the absence of positive customary and traditional use findings. The 2012 personal use permit for Juneau Management Area waters provided open seasons and limits for sockeye salmon at the following locations: in the Taku River from July 1–July 31, with a possession and annual limit of 5 sockeye salmon for a household of 1 person and 10 sockeye salmon for a household of 2 or more people; and in Sweetheart Creek from June 1–October 31, with a possession limit of 25 sockeye salmon and no annual limit. In all streams in the Juneau Management Area, except along the Juneau road system, the open season for pink salmon was June 1–September 30 with a 150 fish limit annually and in possession; for chum salmon, the open season was June 1–October 31 with an annual and possession limit of 50 fish.

Salmon could be taken under a personal use fishing permit by holders of a valid Alaska resident sport fishing license, by Alaska residents under the age of 16, or by Alaska residents with a permanent identification card (seniors and disabled veterans). Beach seines, cast nets, dip nets, gaffs, and spears were the gear allowed in the Juneau area. Set gillnets could not be used except in the Taku River, where set gillnets could not exceed 15 fathoms in length, had to be set from the Taku River Lodge upstream to the U.S.–Canada border, and could not be fished within 100 yd of the ADF&G fish wheels. In Sweetheart Creek, salmon could be taken for personal use only upstream from an ADF&G regulatory marker located near the stream mouth. Additionally, the permit holder had to be present at the net while it was in use. Snagging was prohibited in the personal use fisheries. Chinook and coho salmon, rainbow/steelhead trout, and Arctic char/Dolly Varden could be retained only if taken incidentally under a personal use permit. Salmon possession limits were 2 Chinook and 6 coho salmon.

Harvest Assessment Program

The total estimated salmon harvest for the Juneau Personal Use Area fisheries in 2012 was 6,875 salmon, consisting of 6,390 sockeye salmon (93%), 290 pink salmon (4%), 148 coho salmon (2%), 40 Chinook salmon (1%), and 6 chum salmon (<1%) (Table 13-3). This was a higher harvest than 2011. Sockeye salmon harvest increased by almost 4,000 fish, while harvests of all other species declined. An estimated 367 permits were fished in the Juneau Personal Use area, as compared to 2011 when there were an estimated 300 permits.

The estimated salmon harvest for the community of Juneau (including the communities of Douglas and Auke Bay), based on 659 permits issued and 610 returned (93%), totaled 9,220 salmon, including 8,453 sockeye salmon (92%), 371 coho salmon (4%), 293 pink salmon (3%), 52 chum salmon (1%), and 51 Chinook salmon (1%) and (Table 13-4). Compared to 2011, the 2012 harvest contained a much higher percentage of sockeye salmon (92% in 2012 vs 70% in 2011). While sockeye harvests increased by over 4,000 fish, harvests of all other salmon species either decreased or remained the same (Table 13-4). Total harvest in 2012 increased over 2011 by just over 3,000 fish.

63. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. "Research and Analysis Homepage." Accessed July 2014. <http://labor.alaska.gov/research/pop/popest.htm>

SITKA MANAGEMENT AREA

Sitka Subsistence and Personal Use Salmon Fisheries

Background and History

Subsistence and personal use salmon fisheries in the waters traditionally used by the residents of Sitka are under the management responsibility of the Division of Commercial Fisheries' Sitka Area office. In 1989, the BOF adopted a positive C&T finding for sockeye salmon in the waters of Section 13A south of the latitude of Cape Edward, in waters of Section 13B north of the latitude of Redfish Cape, and in waters of Section 13C (5 AAC 01.716 (a)(8)). At the March 1997 BOF meeting in Sitka, this finding was extended to include all other salmon species in all waters of District 13 (5 AAC 01.716 (a)(21)). Principal salmon waters and streams used by Sitka fishers include Klag Bay–Lake Anna, Lake Stream–Ford Arm, Necker Bay, Redoubt Bay, Salmon Lake, and Redfish Bay.

The residents of Sitka are the principal subsistence users of the salmon stocks in the area. In 2012, the city and borough of Sitka had a population of 9,058.⁶⁴ The Sitka Tlingit have traditionally used most of the Pacific coast of Baranof and Chichagof islands from Point Urey to Cape Ommaney, including the myriad islands lying off the coast, and up Peril Strait between Chichagof and Baranof islands into Hoonah Sound as far as Patterson Bay. Sitkans share the use of Yakobi Island and the sockeye salmon fisheries at Hoktaheen Cove and Surge Bay with the residents of Hoonah. Sitka residents' territory touches that of Angoon residents' in Peril Strait and Sitkoh Bay.

Regulations

Specific conditions on the 2012 subsistence–personal use salmon permit remained the same as 2011. The 2012 subsistence–personal use permit applied to the marine waters and anadromous lakes and streams within the Sitka Management Area. The season for sockeye salmon for all Sitka locations opened June 1 and closed between July 13 and August 31, depending on location. As stated on the permit, Falls Lake and Bay closed on July 13, but was open again from July 23 to August 15. On July 20, Hoktaheen Cove, Takanis Bay, and Gut Bay closed to sockeye salmon fishing. Leo's Anchorage closed on July 25. On July 31, Silver Bay, Politofski Lake, and other unlisted C&T areas closed. On August 15, Klag and Surge bays, Lake Anna, Ford Arm, and Hanus Bay (Lake Eva) closed. The last areas closed on August 31 and included Necker, Redfish, and Sitkoh bays.

Possession and annual limits for sockeye salmon varied from 10 fish in possession and annually at Leo's Anchorage and Silver Bay to 100 fish in possession and annually at Necker Bay. Sitkoh, Takanis, Surge, Klag, and Hanus bays, Hoktaheen Cove, and Politofski Lake had possession and annual limits of 50 sockeye salmon. Lake Anna, Ford Arm, Falls Lake, and Falls Bay had possession and annual limits of 25 fish. Redfish Bay had limits of 50 in possession and 100 fish annually. Gut Bay limits were 10 fish in possession and 20 fish annually. For subsistence–personal use locations not listed on the permit, the possession limit and annual limit was 10 sockeye salmon with a season of June 1–July 31. Redoubt Bay had a possession limit of 10 fish with an annual limit of 50, but this was altered inseason by an emergency order as discussed below.

In January 2003, the BOF adopted the Redoubt Bay and Lake Sockeye Salmon Management Plan (5 AAC 01.760). The plan provides a management approach for subsistence, sport, and commercial fisheries that target Redoubt Lake sockeye salmon based on an optimal escapement goal of 7,000–25,000 fish. As specified on the permit, by default the fishery is open from June 1–August 31 with a possession limit of 10 fish and an annual limit of 50 fish. If the projected run falls below 7,000 fish or above 10,000 fish, the season or limits will change inseason. In 2012, the projected escapement for the season exceeded

64. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. "Research and Analysis Homepage." Accessed July 2014. <http://labor.alaska.gov/research/pop/popest.htm>

30,000 sockeye salmon. In accordance with the plan, an emergency order was published on July 12, 2012 which increased the subsistence possession limit for sockeye to 25 fish with an annual limit of 100.

Salmon streams flowing across or adjacent to the Sitka road system were closed to subsistence–personal use fishing for coho and chum salmon. The season for chum salmon in other C&T waters within the Sitka Management Area, except for the listed sockeye salmon streams, was July 15–October 31, with a possession and annual limit of 50 fish. Pink salmon could be harvested from C&T areas within the Sitka Management Area, except those sockeye streams listed on the permit, from July 15–September 30, with a possession limit of 50 fish and annual limit of 150. Coho salmon could be taken in C&T areas within the Sitka Management Area from August 16–October 31 and in Redoubt, Necker, Redfish, and Sitkoh bays from September 1–October 31 with a possession limit of 20 fish and an annual limit of 40 fish.

The 2012 subsistence–personal use salmon permit for the Sitka Management Area stipulated that prior to August 16 Chinook salmon, rainbow/steelhead trout and Arctic char/Dolly Varden could only be taken incidentally by gear operated under the subsistence–personal use fishing guidelines of the permit. Allowable subsistence gear included hand purse seines, beach seines, drift gillnets, dip nets, gaffs, and spears. Drift gillnets could not exceed 50 fathoms. Cast nets were allowed in all areas except Redoubt Bay. In Redoubt Bay only, the use of rod and reel gear was allowed as subsistence gear and sport regulations applied to this gear. Portions of Falls Lake, Gut Bay, Silver Bay, and Indian River had closed areas and/or restricted gear types specified on the permit.

Harvest Assessment Program

As reported in Table 13-3, the estimated salmon harvest in the Sitka Management Area subsistence and personal use fisheries in 2012 was 15,650 salmon, consisting of 14,844 sockeye salmon (95%), 191 pink salmon (1%), 511 coho salmon (3%), 81 chum salmon (1%), and 24 Chinook salmon (<1%). This was an increase from the 2011 harvest estimate of 10,001 fish; contributions of each species to the overall harvest remained similar, but the harvest of most species increased, except for pink salmon. Sockeye salmon harvest increased from an estimated 9,464 salmon in 2011 and coho salmon harvests more than doubled the 2011 estimate. An estimated 449 permits were fished in the Sitka Management Area in 2012, compared to 308 permits in 2011.

As reported in Table 13-4, the estimated salmon harvest for the community of Sitka in 2012, based on 650 permits issued and 593 returned (91%), was 13,568 salmon, including 12,850 sockeye salmon (95%), 475 coho salmon (4%), 154 pink salmon (1%), 71 chum salmon (1%), and 17 Chinook salmon (<1%). The number of permits issued and returned increased, as did the estimated harvests. While the harvests of most species increased, the biggest increase, accounting for most of the overall harvest increases, was in sockeye salmon harvests, which increased from an estimated 8,890 salmon in 2011.

PETERSBURG MANAGEMENT AREA

The Petersburg Management Area includes the Kake Subsistence Area, the Petersburg–Wrangell Personal Use Area, the federal Stikine River subsistence fishery, and the Point Baker–Port Protection Subsistence Area. Overall, an estimated 185 state subsistence permits were fished in the Petersburg Management Area in 2012. The total estimated salmon harvest was 3,280 fish, with 76% of the harvest coming from sockeye salmon (Table 13-3). Fewer permits were fished in 2012 and fewer salmon were harvested. For any areas with a positive C&T determination not listed on the permit, the open season was June 1–July 31, with harvest limits for sockeye salmon of 10 in possession and 10 annually.

Kake Subsistence Area

Background and History

Subsistence salmon fisheries in the waters traditionally used by the residents of Kake are under the management responsibility of the Division of Commercial Fisheries' Petersburg Area office. In 1989, the BOF adopted a positive C&T finding for salmon in the waters of sections 9A and 9B north of the latitude

of Swain Point, in the waters of District 10 west of a line from Pinta Point to False Point Pybus, and in the waters of District 5 north of a line from Point Barrie to Boulder Point (5 AAC 01.716 (a)(10)). Principal salmon waters and streams used predominately by Kake fishers include Gut Bay and Falls Lake Creek on the southwest coast of Baranof Island, as well as Saginaw, Security (Salt Lake), Pillar (Kutlaku Creek), and Tebenkof (Alecks Creek) bays on Kuiu Island.

In 2012, Kake had a population of 598, an increase of about 20 residents over 2011.⁶⁵ Kake residents shared the use of the southern coastal waters of Admiralty Island with residents of Angoon and Petersburg. In recent years, principal subsistence salmon fishing by Kake residents has occurred in Gut Bay and Falls Creek on Baranof Island, and at Kutlaku Creek in Pillar Bay.

Regulations

Discretionary permit conditions did not change from 2011. The 2012 subsistence salmon permit for the Kake Area waters with a positive C&T finding provided for an open season for sockeye salmon in Alecks Creek and Shipley Bay of June 1–July 31. In the former stream, harvest limits were set at 50 salmon in possession and annually. In the latter area, the possession limit was 25 fish and the annual limit was 50 fish. In Bay of Pillars, the open season was from June 1–August 15 with annual and possession limits of 50 fish. For Falls Lake, the open seasons were June 1–July 13 and July 23–August 15 with possession and annual limits of 25 fish. The season for sockeye salmon in Gut Bay was June 1–July 20 with a limit of 10 fish in possession and 20 fish annually. Pink, chum, and coho salmon could be harvested in all streams in the Kake Subsistence Area, except for the sockeye salmon streams identified on the permits. The open season for pinks was between July 15–September 15 with a possession limit of 100 pink salmon and no annual limit. Chum salmon could be harvested from July 1–October 31 with a possession limit of 50 fish and no annual limit. The coho salmon season lasted from August 16–October 31 with a limit of 20 fish in possession and 40 fish annually.

Permitted subsistence gear included gaffs, spears, beach seines, dip nets, drift gillnets, and cast nets. Set gillnets could only be used in Shipley Bay within 100 yd of the terminus of Shipley Creek, and the permit holder was required to be physically present at the net while in operation. Drift and set gillnets could not exceed 50 fathoms.

Harvest Assessment Program

As reported in Table 13-3, the estimated salmon harvest in the Kake Subsistence Area in 2012 was 1,168 salmon, including 950 sockeye salmon (81%), 124 pink salmon (11%), 45 Chinook salmon (4%), 32 chum salmon (3%), and 16 coho salmon (1%). An estimated 53 permits were fished in the Kake Subsistence Area in 2011.

The estimated subsistence salmon harvest for the community of Kake in 2012, based on 142 permits issued and 115 returned (81%), was 1,235 salmon. The harvest consisted of 1,001 sockeye salmon (81%), 136 pink salmon (11%), 48 Chinook salmon (4%), 33 chum salmon (3%), and 16 coho salmon (1%), (Table 13-4). More permits were issued in 2012 than in 2011, but the total harvest declined from an estimated 2011 harvest of 2,146. The largest decrease was seen in sockeye salmon. .

Petersburg–Wrangell Subsistence–Personal Use Area

Background and History

Subsistence and personal use salmon fisheries in the waters traditionally used by the residents of Wrangell and Petersburg are under the management responsibility of the Division of Commercial Fisheries' Petersburg Area office. In 1989, the BOF adopted a positive C&T finding for salmon in many of the waters of Southeast Alaska. At that time, the BOF did not act on proposals requesting a positive

65. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. "Research and Analysis Homepage." Accessed July 2014. <http://labor.alaska.gov/research/pop/popest.htm>

C&T finding for salmon in the waters of districts 7 and 8, the principal waters used by Petersburg and Wrangell residents. In 2002, however, the BOF made a positive C&T finding for these districts (5 AAC 01.716 (a)(23)). These waters include Thoms Place, Harding River, Mill Creek, and the Stikine River.

Petersburg and Wrangell residents are the principal users of the salmon stocks of Salmon Bay on Prince of Wales Island, as well as Crystal Creek, Thoms Creek, Earl West Cove, Mill Creek, and the Stikine River. In 2012, the population of Petersburg was 2,968 and that of Wrangell was 2,448.⁶⁶ Both estimates are similar to the 2011 estimates.

Regulations

Few changes were made to the discretionary permit conditions from 2011. The 2012 subsistence–personal use salmon permit for the Petersburg Management Area was valid in the waters of districts 7 and 8. The permit provided an open season (June 1–July 31) for subsistence sockeye salmon in Shipley, Salmon, and Red bays, along with Thoms Place and Mill Creek. Limits for sockeye salmon were 25 in possession and 50 annually from Shipley Bay and 30 in possession and annually from Salmon Bay and Red Bay, combined. Thoms Place and Mill Creek had a combined possession limit of 20 fish and an annual limit of 40 fish.

For all streams in the Wrangell and Petersburg subsistence areas, except the sockeye salmon locations listed on the permit, fishing for pink and chum salmon was permitted. The open season for subsistence pink salmon fishing was July 15–September 15, with a daily possession limit of 100 pink salmon and no annual limit. The open season for subsistence chum salmon fishing was July 1–October 31, with a daily possession limit of 50 fish and no annual limit. Subsistence coho salmon was permitted in all the streams in the Wrangell and Petersburg subsistence areas from August 16–October 31, with a limit of 20 fish in possession and 40 annually.

Allowed subsistence gear included gaffs, spears, beach seines, dip nets, drift gillnets, and cast nets. Drift and set gillnets could not exceed 50 fathoms. Set gillnets could only be used in Shipley Bay within 100 yards of the terminus of Shipley Creek, and the permit holder was required to be physically present at the net while in operation. A federal subsistence permit was needed to fish the Stikine River.

Personal use permit conditions applied in the absence of positive C&T findings. Streams that crossed or were adjacent to the Petersburg or Wrangell road system were closed to personal use fishing. Both tips of the caudal fin (tail) had to be removed. Allowed personal use gear included beach seines, drift gillnets, cast nets, dip nets, spears, and gaffs. Gillnets could not exceed 50 fathoms in length.

Personal use regulations established a weekly sockeye salmon season at Hatchery Creek, which drains into Sweetwater Lake. The fishery was only open Thursdays through Sundays from June 1–June 30. In 2012, harvest limits were restricted to 3 fish daily and 9 annually. Personal use coho salmon fishing was open in Blind Slough and North Wrangell Narrows on Fridays from 6am to 8pm from August 10 to September 2 with possession and annual limits of 25 fish combined. The Anita Bay personal use permit allowed the harvest of Chinook, chum, and coho salmon May 1–October 31 with possession and annual limits of 25 fish in any combination. Outside of this area, the possession limit was 2 Chinook salmon and 6 coho salmon.

Harvest Assessment Program

The estimated salmon harvest in the Petersburg Subsistence–Personal Use Area in 2012 was 1,315 salmon, including 822 sockeye salmon (63%), 445 coho salmon (34%), 26 pink salmon (2%), and 22 chum salmon (2%) (Table 13-3). Compared to 2011, there was a slight increase in overall estimated harvest, coming from an increase in the estimated harvest of sockeye and coho salmon. The Petersburg

66. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. “Research and Analysis Homepage.” Accessed July 2014. <http://labor.alaska.gov/research/pop/popest.htm>

Subsistence-Personal Use Area shows a much lower reliance on sockeye salmon than any other management or use area. An estimated 73 permits were fished in 2012.

As reported in Table 13-4, the estimated subsistence salmon harvest for the community of Petersburg in 2012, based on 99 permits issued and 90 returned (91%), was 1,207 salmon, including 759 sockeye salmon (63%), 403 coho salmon (33%), 23 pink salmon (2%), and 22 chum salmon (2%). A similar number of permits was fished in 2012 as in 2011. Overall harvest increased slightly, as did harvests of sockeye and coho salmon. Harvests of chum, pink, and Chinook salmon all decreased slightly.

As shown in Table 13-3, the estimated salmon harvest in the Wrangell Subsistence-Personal Use Area in 2012 was 798 salmon, which included 709 sockeye salmon (89%), 12 pink salmon (2%), 63 chum salmon (8%), 8 Chinook salmon (1%), and 6 coho salmon (1%). Compared to 2011, the estimated overall harvest, as well as that of every species save chum salmon, decreased. An estimated 59 permits were fished in 2012.

The estimated subsistence salmon harvest for the community of Wrangell in 2012, based on 97 permits issued and 94 returned (97%), was 732 salmon, including 645 sockeye salmon (88%), 63 chum salmon (9%), 12 pink salmon (2%), 8 Chinook salmon (1%), and 3 coho salmon (<1%) (Table 13-4). Harvests were less than the estimated 2011 harvests, both in overall numbers and in numbers of each species harvested. Chum salmon was the only species that saw a slight increase in harvest level. Coho salmon experienced the most dramatic decline, from 196 salmon in 2011 to just 3 in 2012.

2012 Federal Stikine River Subsistence Salmon Fishery: Regulations

In January 2004, the U.S. and Canada negotiated a modified Pacific Salmon Treaty that allowed for a U.S. subsistence salmon fishery on the Stikine River. The Federal Subsistence Board implemented a Stikine River subsistence sockeye salmon fishery in 2004, followed by directed Chinook and coho salmon subsistence fisheries authorized in 2005. Regulatory changes implemented for the 2006 season included an increase in the mesh size of gillnets during the Chinook salmon fishery and an earlier starting date for the sockeye salmon fishery. In 2008, 2 additional regulatory changes were made: subsistence fishing permits became valid for the entire season (May 15–October 1); and the start date of the coho salmon fishery was moved up to August 1. The latter change allowed a continuous subsistence fishery throughout the season. There were no changes in subsistence fishing regulations or permit conditions for the 2012 fishing season. The inseason return estimate for Chinook salmon predicted the escapement goal would not be met. Therefore, the federal inseason manager issued a letter to permit holders requiring 48-hour reporting of any Chinook salmon harvested for the remainder of the Chinook salmon season. Post season total return estimates were above escapement goals for Chinook salmon.

Current Federal Regulations

The federal subsistence fisheries regulatory year begins April 1. Regulations are detailed in Subpart C of 36 CFR part 242. The sections relevant to the Stikine River are as follows:

§ __.24 Customary and traditional use determinations.

(2) Fish determinations. The following communities and areas have been found to have a positive customary and traditional use determination in the listed area for the indicated species:

Southeastern Alaska Area:

District 8 and waters draining into that District: Salmon, Dolly Varden, trout, smelt, and eulachon. Residents of drainages flowing into Districts 7 & 8, residents of drainages flowing into District 6 north of the latitude of Point Alexander (Mitkof Island), and residents of Meyers Chuck.

§ __.27 Subsistence taking of fish.

(i) Fishery management area restrictions.

(13) Southeastern Alaska Area.

(xv) You may take Chinook, sockeye, and coho salmon in the mainstem of the Stikine River only under the authority of a Federal subsistence fishing permit. Each Stikine River permit will be issued to a household. Only dip nets, spears, gaffs, rod and reel, beach seine, or gillnets not exceeding 15 fathoms in length may be used. The maximum gillnet mesh size is 5 1/2 inches, except during the Chinook season when the maximum gillnet mesh size is 8 inches.

(A) You may take Chinook salmon from May 15 through June 20. The annual limit is 5 Chinook salmon per household.

(B) You may take sockeye salmon from June 21 through July 31. The annual limit is 40 sockeye salmon per household.

(C) You may take coho salmon from August 1 through October 1. The annual limit is 20 coho salmon per household.

(D) You may retain other salmon taken incidentally by gear operated under terms of this permit. The incidentally taken salmon must be reported on your permit calendar.

(E) The total annual guideline harvest level for the Stikine River fishery is 125 Chinook, 600 sockeye, and 400 coho salmon. All salmon harvested, including incidentally taken salmon, will count against the guideline for that species.

The following conditions were included on the 2012 Stikine River subsistence fishing permit.

Seasons: Chinook - May 15 through June 20, sockeye - June 21 through July 31, coho - August 1 through October 1.

Harvest Limit: Chinook - annual limit is 5 per household, sockeye - annual limit is 40 per household, coho - annual limit is 20 per household.

C&T Determination: residents of drainages flowing into districts 7&8, residents of drainages flowing into district 6 north of the latitude of Point Alexander (Mitkof Island), and residents of Meyers Chuck.

Gear: Gillnets not exceeding 15 fathoms in length may be used. The maximum gillnet mesh size is 5 1/2 inches, except during the Chinook season when the maximum gillnet mesh size is 8 inches.

The total annual harvest level for the Stikine River is controlled by the in season manager and may be closed or expanded by special action.

Size: Chinook "jack" salmon is defined as less than 28 inches. Only Chinook equal to or greater than 28 inches are included in the annual harvest limit. Indicate both the number of Chinook taken that are greater than and less than 28 inches separately. See species legend for code difference.

Other: You may retain other salmon taken incidentally, however they must be recorded on your permit.

Harvest Assessment Program

For Chinook, coho, and sockeye salmon fisheries harvest assessment, a telephone-based monitoring program is used inseason, with permits and harvest reporting used for overall harvest assessment postseason. In 2012, 130 fishing permits were issued, with approximately 65% going to Wrangell households and the other 35% to Petersburg households. Year-end harvest reports were obtained from all but three permit holders. The Stikine River subsistence harvest totaled 1,546 salmon. The harvest

consisted of 1,302 sockeye salmon (84%), 112 coho salmon (7%), 47 chum salmon (3%), 53 Chinook salmon (3%), and 32 pink salmon (2%). There was also 1 Dolly Varden char harvested. Compared to 2011, a similar number of permit holders caught less salmon overall. Most of the decrease came from the harvest of sockeye salmon which dropped from a 2011 estimate of 1,741 sockeye, but each species excepting coho salmon experienced a decline in harvest in 2012. The proportion of the catch contributed by each species was similar, with sockeye salmon constituting 80% or more of the harvest in both years. Fishing patterns were similar to previous years in that most of the fishing effort and harvest occurred in the lower and middle portions of the river. Approximately 87% of the Chinook and sockeye salmon was harvested by July 28, a later date than in 2011; 90% of coho salmon was taken by late September.⁶⁷

Point Baker–Port Protection Subsistence Fisheries

Background and History

The Division of Commercial Fisheries' Petersburg Area office manages subsistence and personal use salmon fisheries in the waters used by fishers from the communities of Point Baker and Port Protection, which are also used by residents of Petersburg and Wrangell. These fisheries rely especially on the Salmon Bay and Red Bay sockeye salmon stocks at the northern end of Prince of Wales Island. In 1989, when the BOF adopted a positive C&T finding for salmon in some waters of Southeast Alaska, it did not act on proposals to make a similar finding for the principal waters used by Point Baker and Port Protection residents. In 1997, however, the BOF did adopt a positive C&T finding for salmon and other fishes in the waters of District 5 north of a line from Point St. Albans to Cape Pole, in the waters of Section 6A west of a line from Macnamara Point to Mitchell Point, and in the waters of Section 6B west of the longitude of Macnamara Point (5 AAC 01.716 (a)(20)). Harvests in these waters are included in the Petersburg Subsistence-Personal Use Area in Table 13-3.

In 2012, Point Baker had a population of 16 and Port Protection had a population of 42.⁶⁸

Regulations

The Point Baker drift gillnet subsistence sockeye salmon fishery occurs in the waters of Sumner Strait within 3 miles of the Prince of Wales Island shoreline north of Hole-in-the-Wall and west of the western side of Buster Bay. The fishery was open Wednesdays at 12:00 PM to Sundays at 12:00 PM, from June 13–July 31. Only drift gillnet gear, not to exceed 50 fathoms in length, was allowed. Harvest was limited to 25 sockeye salmon in possession and annually. Fishers could retain other species incidentally harvested during this fishery. Pink and chum salmon harvests were allowed in all streams within the Point Baker–Port Protection subsistence area, except for the sockeye salmon streams identified on the permit. There was a 100-fish possession limit for pink salmon, with no annual limit. For chum salmon, 50 fish were allowed in possession with no annual limit. Coho salmon could be harvested in all streams in the Point Baker–Port Protection subsistence area with a possession limit of 20 fish and annual limit of 40 fish.

Harvest Assessment Program

Port Protection households maintain either a Ketchikan or Point Baker post office address and receive mail via private carrier from Ketchikan. Port Protection harvests can be included in either the Point Baker or Ketchikan harvest estimates. In 2011, no salmon permits were issued to Port Protection residents. For Point Baker in 2012, 3 permits were issued and returned with no harvests recorded (Table 13-4).

KETCHIKAN MANAGEMENT AREA

The Ketchikan Management Area includes 3 subsistence areas where the BOF made positive C&T findings in 1989. Two of these areas are on the west coast of Prince of Wales Island: Hydaburg area

67. Robert Larson, USFS. Stikine River subsistence salmon fishery: 2012 season summary. United States Department of Agriculture Forest Service, unpublished report, 2012. Hereinafter referred to as (Larson 2012).

68. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. "Research and Analysis Homepage." Accessed July 2014. <http://labor.alaska.gov/research/pop/popest.htm>

waters and Craig–Klawock area waters. The third area is on the east coast of Prince of Wales Island, around the community of Kasaan. In addition, the Ketchikan Management Area includes the personal use fisheries in the Ketchikan area. All of these areas are under the management responsibilities of the Division of Commercial Fisheries’ Ketchikan Area office. There were an estimated 308 permits fished in the Ketchikan Management Area in 2012, slightly fewer than the 358 permits fished in 2011. The total estimated salmon harvest was 11,510 fish, slightly higher than the 2011 estimate of 11,296 salmon (Table 13-3). Sockeye salmon harvests contributed 85% of this harvest.

Craig, Klawock, and Hydaburg Subsistence Fisheries

Background and History

Hydaburg area waters with a positive C&T finding include Section 3A and the waters of District 2 in Nichols Bay north of lat. 54°42.12’ N (5 AAC 01.716 (a)(18)). Craig–Klawock area waters with a positive C&T finding include Section 3B east of a line from Point Ildefonso to Tranquil Point; Warm Chuck Inlet north of a line from a point on Heceta Island at lat. 55°44’ N, long. 133°25’ W to Bay Point; Section 3C in Karheen Passage north of lat. 55°48’ N and east of long. 133°20’ W; and Sarkar Cove and the Sarkar lakes (5 AAC 01.716 (a)(15)).

The communities of Hydaburg, Craig, and Klawock on the west coast of Prince of Wales Island primarily use the salmon stocks of sections 3A and 3B, with the main harvest locations at Hetta Inlet–Sukkwana Strait (Eek Creek), Big Salt–Trocadero Bay (Klawock River), and Sea Otter Sound (Sarkar River).

In 2012, Craig had a population of 1,242, Klawock had a population of 798, and Hydaburg had a population of 367.⁶⁹ All communities experienced a slight decrease in estimated populations since 2011.

Regulations

The 2012 subsistence sockeye salmon openings in Klawock were Mondays 8:00 AM to Fridays 5:00 PM from July 7–August 7, with a 20 sockeye salmon possession limit and no annual limit; in Hetta Inlet and Eek Creek, the season was June 1–August 31 with a possession limit of 20 sockeye salmon and no annual limit, and in Hugh Smith Lake fishing was open June 22–July 12 with a 12 sockeye salmon possession limit and no annual limit. Inseason, this system re-opened on July 20 through July 31 based on high sockeye salmon escapements. Other systems in the Ketchikan Management Area with C&T areas were open to sockeye salmon fishing June 1–July 31, with a 10 sockeye salmon possession limit and a 25 sockeye annual limit. Such streams had to be approved by ADF&G and listed on the permit. All streams in the Ketchikan Management Area with customary and traditional use areas were open for pink salmon July 1–September 30 with a 150 fish possession limit and no annual limit. Chum and coho salmon fishing was open in the same waters July 1–October 31 with a possession limit of 25 chum and 20 coho salmon. There was no annual limit for chum salmon, but there was a 40 coho salmon limit annually. Additional conditions on the 2012 subsistence–personal use salmon permit for the Ketchikan Management Area stipulated that hand purse seines, beach seines, spears, gaffs, cast nets, and dip nets were allowable subsistence–personal use gear allowed for general use. Chinook salmon, rainbow/steelhead trout, and Arctic char/Dolly Varden could only be taken incidentally by gear operated under the terms of the permit. Sockeye salmon could not be retained as incidental catch.

Harvest Assessment Program

The estimated salmon harvest for the Craig–Klawock–Hydaburg Subsistence Area in 2012 was 6,023 salmon, including 5,670 sockeye salmon (94%), 225 coho salmon (4%), 82 pink salmon (1%), and 46 chum salmon (1%)(Table 13-3). The 2012 harvest decreased from 6,023 fish in 2011. Sockeye and pink salmon decreased significantly, while coho salmon harvest increased from just 45 salmon in 2011.

69. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. “Research and Analysis Homepage.” Accessed July 2014. <http://labor.alaska.gov/research/pop/popest.htm>

As reported in Table 13-4, 161 permits were issued to residents of Craig and 117 (73%) were returned. The total estimated salmon harvest of Craig residents was 1,352, a decrease of 400 fish from 2011 estimates. By species, the harvest consisted of 1,153 sockeye salmon (85%), 105 pink salmon (8%), 67 coho salmon (5%), and 27 chum salmon (2%). The total estimated salmon harvest for Klawock, based on 121 permits issued and 89 returned (74%), was 2,900, a slight increase over the 2011 estimate of 2,567, consisting of 2,597 sockeye salmon (90%), 65 pink salmon (2%), 33 chum salmon (1%), and 205 coho salmon (7%). Sockeye and coho salmon harvests increased slightly, while chum and pink salmon harvests experienced a slight decline. The total estimated salmon harvest for Hydaburg, based on 55 permits issued and 43 returned (78%), was 1,797 salmon, the majority of which were sockeye salmon. An estimated 27 coho salmon and 9 pink salmon were also harvested. In 2011, the estimated harvest of salmon was 2,487, mainly sockeye salmon. Fewer permits were issued to each community in 2012, but response rates improved everywhere except for Craig.

Kasaan Subsistence Area

Background and History

The subsistence area on the east coast of Prince of Wales Island with a positive C&T finding for salmon includes the Kasaan area waters of District 2 north of the latitude of the northernmost tip of Chasina Point then west of a line from the northernmost tip of Chasina Point to the easternmost tip of Grindall Island to the easternmost tip of the Kasaan Peninsula (5 AAC 01.716 (a)(12)). Salmon fishing in all other marine waters along the east coast of Prince of Wales Island occurs under personal use or sport regulations. The principal waters used for personal use salmon fishing along the eastern coast of Prince of Wales Island are Kegan Lake, the Thorne River, and Hatchery Creek–Sweetheart Creek. The personal use fisheries are described in the Ketchikan Personal Use Area section.

In 2012, Coffman Cove had a population of 181, Edna Bay’s population was 39, Hollis had a population of 109, Kasaan’s population was 80, Thorne Bay’s population was 508, and the population of Whale Pass was 39.⁷⁰

Regulations

All streams in the Ketchikan Management Area with positive C&T findings not otherwise listed on the permit were open for subsistence sockeye salmon fishing June 1–July 31, with a 10 fish possession limit and a 25 fish annual limit. All streams with a positive C&T determination were open to pink salmon fishing July 1–September 30, with a limit of 150 fish in possession and no annual limit. Coho and chum salmon fishing was also open in these waters July 1–October 31, with a limit on coho salmon harvests of 20 fish in possession and 40 fish annually. The limit on chum salmon harvests was 25 fish in possession and no annual limit.

Allowable gear in the subsistence fishery included hand purse seines, beach seines, spears, gaffs, cast nets, and dip nets. Salmon could not be taken with a line attached to a rod or pole. Chinook salmon, rainbow/steelhead trout, and Arctic char/Dolly Varden could only be taken incidentally by gear operated under the terms of the permit. Sockeye salmon could not be retained as incidental catch.

Harvest Assessment Program

As reported in Table 13-3, in 2012 an estimated 76 permit holders fished in the Kasaan Subsistence Area with an estimated salmon harvest of 2,366 salmon. The harvest included 1,670 sockeye salmon (71%), 209 coho salmon (9%), 465 pink salmon (20%), and 23 chum salmon (1%). The total harvest increased from a 2011 estimate of 1,939. While sockeye salmon harvests decreased from an estimated 1,832 salmon in 2011, harvests of all other species increased. Fewer permits were fished in 2012 as compared to 2011.

70. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. “Research and Analysis Homepage.” Accessed July 2014. <http://labor.alaska.gov/research/pop/popest.htm>

Based on 15 permits issued to residents of Kasaan and 12 returned (80%) in 2012, an estimated 181 salmon were harvested, consisting almost entirely of sockeye salmon (153) as well as 28 coho and 1 pink salmon (Table 13-4). Thorne Bay residents were issued 26 permits, 22 of which were returned (85%), resulting in a harvest estimate of 125 salmon, including 122 sockeye salmon and 2 pink salmon (Table 13-4). Three permits were issued to Naukati Bay residents and all were returned. No salmon harvest was reported. In Hollis, 42 permits were issued and 36 were returned (86%). An estimated 1,054 salmon were harvested, including 610 sockeye salmon, 361 pink salmon, 81 coho salmon, and 2 chum salmon. In Coffman Cove 10 permits were issued and 7 were returned (70%). An estimated 35 sockeye were harvested. One permit was issued in Whale Pass and not returned, resulting in an estimate of 0 salmon harvested by Whale Pass residents.

Ketchikan Personal Use Area

Background and History

The Division of Commercial Fisheries' Ketchikan Area office is responsible for oversight of the subsistence and personal use salmon fisheries in districts 1, 2, 3, and 6. Some waters within sections 1A, 1C, 1D, 1E, 1F, and District 2 fall within the Ketchikan Nonsubsistence Use Area (Figure 13-2). The BOF made a positive C&T finding for salmon stocks in the waters traditionally used by the Tongass Tlingit of Saxman. These waters include the Naha River, Boca de Quadra in the waters of Sockeye Creek and Hugh Smith Lake, as well as the salt waters within 500 yards of the terminus of Sockeye Creek (5 AAC 01.716 (a)(19)).

The communities of Ketchikan and Saxman are the principal users of the fisheries in the Ketchikan area. In 2012, the population of the Ketchikan borough, excluding Saxman, was 13,472. Saxman, located within the Ketchikan Gateway Borough, had a population of 432.⁷¹

Regulations

The 2012 subsistence–personal use salmon permit for the Ketchikan Management Area provided for a July 1–August 30 open season for sockeye salmon at McDonald Lake (Yes Bay), with a possession and annual limit of 30 fish. Kegan Lake and Thorne River were open from June 1–July 31, with a possession limit of 12 sockeye salmon and an annual limit of 50 sockeye salmon. Other streams in the Ketchikan Management Area's personal use area, except the Ketchikan road system, were open June 1–July 31 with a limit of 10 sockeye salmon in possession and a 25 fish annual limit. Hatchery Creek was open June 1–June 30, Thursdays through Sundays, with a limit of 3 sockeye salmon in possession and 9 annually. Leask Creek and Mahoney Creek and Lake, and marine waters within 500 yards of the terminus of these streams, remained closed. For pink and chum salmon, all streams in the Ketchikan Management Area personal use area, except the Ketchikan road system, were open. The season for pink salmon ran from June 2–September 30 with a limit of 150 fish in possession and no annual limit. For chum salmon, the open season was from June 1–October 31 with a possession limit of 25 and no annual limit. The Ketchikan Creek personal use salmon fishery was opened on August 27 for 3 hours. A per person bag limit of 4 Chinook salmon and 4 coho salmon was in effect, with no size limit. Neets Bay, Nakat Inlet, Kendrick Bay, and Herring Bay personal use fisheries are open under regulations. Excluding the Ketchikan Creek personal use Chinook salmon fishery, coho and Chinook salmon, rainbow/steelhead trout, and Arctic char/Dolly Varden could only be taken incidentally by gear operated under personal use fishing guidelines of the permit. The possession limit for Chinook salmon was 2 fish and the possession limit for coho salmon was 6 fish. Sockeye salmon could not be retained as incidental catch. The legal gear types specified under the terms of this permit included hand purse seines, beach seines, spears, gaffs, cast nets, and dip nets. Salmon could not be taken with a line attached to a rod or pole. In Yes Bay, set or drift gillnets could only be used in specified locations from July 1 until August 30.

71. Alaska Department of Labor and Workforce Development (ADLWD), Juneau. n.d. "Research and Analysis Homepage." Accessed July 2014. <http://labor.alaska.gov/research/pop/popest.htm>

Harvest Assessment Program

The total estimated salmon harvest in the Ketchikan Personal Use Area in 2012 was 3,120 fish, including 2,437 sockeye salmon (78%), 340 chum salmon (11%), 319 pink salmon (10%), 19 coho salmon (1%), and 5 Chinook (<1%) (Table 13-3). An estimated 132 permits were fished in this area. The 2012 harvest is similar in composition to the 2011 harvest but is higher than the 2011 estimate (2,553 salmon). Harvests of all species except Chinook salmon remained static or increased.

As reported in Table 13-4, the total estimated salmon harvest for the community of Ketchikan (including Ward Cove), based on 265 permits issued and 221 returned (83%), was 3,526, including 2,863 sockeye salmon (81%), 309 chum salmon (9%), 297 pink salmon (8%), 53 coho salmon (2%), and 5 Chinook salmon (<1%). In Saxman, based on 21 permits issued and 18 returned (86%), a total of 371 salmon were harvested. Of the total, sockeye salmon constituted the largest proportion at 335 fish (90%) followed by pink salmon at 13 fish (4%) and then chum salmon at 23 fish (6%). Based on 3 permits issued and 2 returned (67%), in 2012 residents of Metlakatla harvested 110 salmon, including 92 sockeye salmon (84%), 14 chum salmon (13%), and 4 pink salmon (4%). Fewer permits were issued in Ketchikan/Ward Cove and Metlakatla in 2012, but harvests increased in all three communities.

Table 13-1.—Subsistence and personal use salmon harvests by district, Southeast region, 2012.

Fishing location	Name	Permits fished		Estimated salmon harvest					
		Reported	Estimated	Chinook	Sockeye	Coho	Chum	Pink	Total
District 1	Ketchikan-Behm Canal	160	192	5	2,437	19	340	319	3,120
District 2	Clarence Strait-East Prince of Wales Island	94	117	0	1,670	209	23	465	2,366
District 3	Inside Waters-West Prince of Wales Island	188	246	0	5,670	225	46	82	6,023
District 5	Sumner Strait	0	0	0	0	0	0	0	0
District 6	East Sumner Strait-North Frederick Sound	84	95	0	854	439	15	16	1,324
District 7	East Etolin Island-Wrangell Island-Ernest Sound	91	96	8	665	4	52	12	741
District 8	Stikine River	7	8	0	13	8	18	10	48
District 9	South Chatham Strait-West Frederick Sound	56	69	48	973	16	33	108	1,178
District 10	East Frederick Sound	2	2	0	29	0	0	26	55
District 11	Juneau-Taku Inlet-Stephens Passage	424	458	40	6,390	148	6	290	6,875
District 12	Angoon-North Chatham Strait-East Chichagof	50	63	0	1,161	59	0	165	1,385
District 13	Sitka-Outer Baranof and Chichagof-Peril Strait	640	706	21	14,793	511	80	181	15,585
District 14	Icy Strait-Glacier Bay	72	84	0	969	62	97	142	1,270
District 15	Lynn Canal-Chilkat Inlet	1,122	1,182	257	8,663	224	456	786	10,386
Yakutat Forelands	Yakutat Forelands	178	223	74	5,669	1,199	19	191	7,151
Yakutat Bay-Troll	Yakutat Bay-Troll	94	119	310	473	87	1	13	885
Subtotal, state permit fisheries		—	—	763	50,427	3,210	1,186	2,806	58,392
Stikine River	Stikine River Federal Fishery	53	53	53	1,302	112	47	32	1,546
Total		—	—	816	51,729	3,322	1,233	2,838	59,938

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

- Fishers with permits may fish at more than one location. As a result, the total number of permits cannot be derived simply by adding column values.

Table 13-2.—Historical subsistence and personal use salmon harvests, Southeast region, 1985–2012.

Year ^a	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
1985	ND	1,271	19	20,006	360	2,951	2,136	25,472
1986	ND	1,354	29	21,974	277	2,840	971	26,091
1987	ND	1,322	34	25,405	117	3,878	1,474	30,908
1988	ND	1,013	94	19,898	97	3,013	1,145	24,247
1989	ND	1,479	580	32,860	1,381	3,113	3,664	41,598
1990	ND	1,543	524	36,376	1,615	3,433	3,529	45,477
1991	ND	1,554	262	37,765	766	3,271	1,741	43,805
1992	ND	1,860	614	53,131	4,939	3,201	2,942	64,827
1993	ND	2,121	537	56,249	3,515	2,583	2,143	65,027
1994	ND	2,239	800	57,097	3,607	4,211	3,639	69,354
1995	ND	2,005	1,203	45,087	3,702	3,370	3,215	56,577
1996	4,172	3,341	1,170	69,216	3,090	5,553	3,204	82,233
1997	4,211	3,529	780	58,782	2,701	4,515	4,080	70,858
1998	4,273	3,629	1,082	62,551	3,264	6,442	3,910	77,250
1999	4,308	3,717	1,393	56,618	1,933	5,557	3,280	68,782
2000	3,771	3,170	1,359	52,867	2,151	3,414	2,619	62,411
2001	3,605	3,116	1,457	55,157	3,266	3,968	4,230	68,080
2002	3,326	2,732	1,857	56,379	3,176	2,183	3,210	66,804
2003	3,595	2,924	1,543	64,670	3,052	6,275	3,894	79,434
2004	3,703	3,235	1,583	61,419	2,446	3,151	3,164	71,763
2005	3,304	2,772	887	39,694	2,283	1,831	4,959	49,655
2006	3,405	2,809	1,356	54,862	1,873	1,731	3,603	63,425
2007	3,156	1,622	1,199	43,100	1,444	721	3,273	49,737
2008	3,153	2,820	1,052	41,548	3,555	1,421	1,897	49,472
2009	3,421	3,097	1,208	49,507	3,616	2,006	3,290	59,627
2010	2,217	1,829	1,828	52,258	3,885	878	3,721	62,571
2011	3,315	2,918	916	41,733	3,060	1,147	5,494	52,350
2012	3,267	2,853	763	50,427	3,210	1,186	2,806	58,392
5-year average (2007–2011)	3,052	2,457	1,241	45,629	3,112	1,235	3,535	54,751
10-year average (2002–2011)	3,260	2,676	1,343	50,517	2,839	2,134	3,650	60,484
Historical average (1985–2011)	3,558	2,408	939	46,897	2,414	3,210	3,127	56,586

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

a. For years prior to 1996, only permits returned with harvest data were included, and harvests reported in these years were not expanded into estimates. Caution should be used if comparing pre-1996 data with later data.

ND = no data.

Table 13-3.—Estimated subsistence and personal use salmon harvests by management and use areas, Southeast region, 2012.

Area	Permits fished		Estimated salmon harvest					
	Reported	Estimated	Chinook	Sockeye	Coho	Chum	Pink	Total
Yakutat Management Area	92	116	384	6,142	1,286	20	204	8,036
Haines Management Area	370	384	257	8,663	224	456	786	10,386
Juneau Management Area	455	474	40	8,520	269	104	597	9,530
Juneau Personal Use Area	357	367	40	6,390	148	6	290	6,875
Angoon Subsistence Area	48	54	0	1,161	59	0	165	1,385
Hoonah Subsistence Area	50	54	0	969	62	97	142	1,270
Sitka Management Area	409	449	24	14,844	511	81	191	15,650
Petersburg Management Area	169	185	53	2,482	466	117	162	3,280
Petersburg Subsistence- Personal Use Area	65	73	0	822	445	22	26	1,315
Wrangell Subsistence- Personal Use Area	56	59	8	709	6	63	12	798
Kake Subsistence Area	48	53	45	950	16	32	124	1,168
Stikine River Federal Subsistence Fishery	53	53	53	1,302	112	47	32	1,546
Ketchikan Management Area	292	308	5	9,777	453	409	866	11,510
Ketchikan Personal Use Area	115	132	5	2,437	19	340	319	3,120
Kasaan Subsistence Area	67	76	0	1,670	209	23	465	2,366
Craig-Klawock-Hydaburg Subsistence Area	110	100	0	5,670	225	46	82	6,023
Total	—	—	816	51,729	3,322	1,233	2,838	59,938

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

- Fishers with permits may fish at more than one location. As a result, the total number of permits cannot be derived simply by adding column values.

Table 13-4.—Subsistence and personal use salmon harvests by community, Southeast region, 2012.

Community	Permits		Estimated salmon harvest					
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	Total
Anchorage	17	12	13	137	8	0	6	164
Angoon	98	77	0	1,308	51	0	51	1,410
Auke Bay	8	8	0	94	2	24	0	120
Barrow	2	2	0	248	0	0	8	256
Coffman Cove	10	7	0	35	0	0	0	35
Craig	161	117	0	1,153	67	27	105	1,352
Denali National Park	1	1	0	6	0	0	4	10
Douglas	56	53	5	381	35	1	42	465
Dutch Harbor	1	1	0	0	0	0	0	0
Eagle River	4	3	0	0	0	0	0	0
Elfin Cove	1	0	0	0	0	0	0	0
Excursion Inlet	3	3	0	1	18	0	0	19
Fairbanks	8	7	0	141	0	0	11	152
Gustavus	22	21	0	84	0	1	2	88
Haines	426	406	235	7,245	177	349	679	8,686
Hollis	42	36	0	610	81	2	361	1,054
Hoonah	122	88	0	1,543	44	103	326	2,016
Hydaburg	55	43	0	1,761	27	0	9	1,797
Juneau	595	549	45	7,978	334	27	251	8,635
Kake	142	115	48	1,001	16	33	136	1,235
Kasaan	15	12	0	153	28	1	0	181
Ketchikan	261	217	5	2,857	53	303	297	3,514
Klawock	121	89	0	2,597	205	33	65	2,900
Klukwan	10	9	1	414	39	72	20	547
Metlakatla	3	2	0	92	0	14	4	110
Naukati Bay	3	3	0	0	0	0	0	0
Palmer	1	1	0	9	0	0	0	9
Pelican	3	3	0	14	0	0	0	14
Petersburg	146	137	21	1,335	408	44	38	1,846
Point Baker	3	3	0	0	0	0	0	0
Port Alexander	7	7	1	270	27	0	0	298
Saxman	21	18	0	335	0	23	13	371
Sitka	650	593	17	12,850	475	71	154	13,568
Skagway	16	16	5	60	0	0	23	88
Tenakee Springs	3	3	1	47	1	0	0	49
Thorne Bay	26	22	0	122	0	0	2	125
Tok	1	1	0	50	0	0	0	50
Ugashik	1	1	0	0	0	0	0	0
Valdez	1	0	0	0	0	0	0	0
Ward Cove	4	4	0	6	0	6	0	12
Wasilla	4	4	9	83	0	0	0	92
Whale Pass	1	0	0	0	0	0	0	0
Wrangell	97	94	8	645	3	63	12	732
Yakutat	142	112	369	5,336	1,116	11	202	7,034
Total	3,397	2,983	816	51,729	3,322	1,233	2,838	59,938

Source ADF&G Division of Subsistence, ASFDB 2013 (ADF&G 2014).

Table 13-5.— Subsistence salmon harvests by community for the Federal Stikine River subsistence salmon fishery, Southeast region, 2012.

Year	Permits		Estimated salmon harvest					Total
	Issued	Returned	Chinook	Sockeye	Coho	Chum	Pink	
Petersburg	47	47	21	576	5	22	15	639
Wrangell	83	83	32	726	107	25	17	907
Total	130	130	53	1,302	112	47	32	1,546

Source Larson (2012).

Table 13-6.— Historical subsistence salmon harvests by community for the Federal Stikine River subsistence salmon fishery, Southeast region, 2004–2012.

Year	Permits	Estimated salmon harvest					Total
	issued	Chinook	Sockeye	Coho	Chum	Pink	
2004	40	12	243	0	11	22	288
2005	35	15	252	53	22	69	411
2006	48	37	390	21	20	23	491
2007	44	36	244	23	11	59	373
2008	50	25	428	42	12	18	525
2009	80	31	723	21	46	66	887
2010	107	61	1,653	135	37	60	1,946
2011	129	66	1,741	40	74	189	2,110
2012	130	53	1,302	112	47	32	1,546
5-year average (2007–2011)	82	44	958	52	36	78	1,168
Historical average (2004– 2011)	67	35	709	42	29	63	879

Source Larson (2012).




Figure 13-1.—Juneau Nonsubsistence Area map, 2012.




Figure 13-2.—Ketchikan Nonsubsistence Area map, 2012.




Figure 13-3.—Southeast region subsistence and personal use harvests by species, 2012.




Figure 13-4.—Total salmon harvested by management area, Southeast region, 2012.

ACKNOWLEDGEMENTS

Each year, thousands of Alaska residents who participate in subsistence and personal use fisheries take the time to provide harvest information to ADF&G. We acknowledge their support with profound gratitude, for without it, a report like this would be impossible to produce.

We also offer thanks to the numerous ADF&G staff in the divisions of Commercial Fisheries, Sport Fish, and Subsistence who conduct the programs that collect, analyze, and report the subsistence and personal use fisheries harvest data every year. They, too, made this report possible.

Many ADF&G personnel generously made time to allow Division of Subsistence staff to interview them about subsistence databases and harvest assessment programs. We have relied upon their numerous insights about these programs to develop the Alaska Subsistence Fisheries Database, as well as to evaluate the data that appear in this report. We sincerely appreciate their help.

This annual report for 2012 is the result of the work of a number of Division of Subsistence staff. Former division employees Dave Caylor, Jeannie Heltzel, and Brian Davis helped design and update the Alaska Subsistence Fisheries Database. Data for 2011 were compiled by Terri Lemons, with assistance from Dave Koster. Division personnel who authored report chapters were James A. Fall, Nicole M. Braem, Caroline L. Brown, Sarah S. Evans, Lisa Hutchinson-Scarborough, Bronwyn Jones, Robbin La Vine, Meredith Ann Marchioni, Elizabeth Mikow, Joshua T. Ream, and Lauren A. Sill. We also acknowledge the contributions of Eunice Dyasuk, who administers the division's subsistence salmon permit program for Bristol Bay in Dillingham, as well as Lisa Olson, Garrett Zimpelman, and Adam Knight, who reviewed and edited the report.

As noted in the report itself, this is the 14th in a series of statewide summaries of subsistence and personal use fisheries harvest data. We encourage those who use this report to offer ideas and suggestions to improve future volumes in this series.

REFERENCES CITED

ADF&G

1987a. *The salmon fishing people of the Yukon River. Wildlife use notebook series No. 1.* Alaska Department of Fish and Game Division of Subsistence: Juneau.

1987b. *Family fish camps of the Yukon River. Wildlife use notebook series No. 3.* Alaska Department of Fish and Game Division of Subsistence: Juneau.

1988. *Salmon fishing methods of the Yukon River. Wildlife use notebook series No. 2.* Alaska Department of Fish and Game Division of Subsistence: Juneau.

2002. *Customary and traditional use worksheet: salmon, Chignik management area.* Alaska Department of Fish and Game Division of Subsistence: Anchorage.

2003. *Customary and traditional use worksheet: salmon: Chitina subdistrict, Prince William Sound management area.* Alaska Department of Fish and Game Division of Subsistence: Cordova.

Alaska Board of Fisheries

2011a. *The Alaska Board of Fisheries 2010/2011 proposed changes in the Cook Inlet, Kodiak and Chignik finfish; and king and Tanner crab (statewide, except Southeast/Yakutat) regulations.* Alaska Department of Fish and Game, Board Support Section: Juneau.

2011b. *Chignik finfish and Adak cod, actions taken.* Alaska Department of Fish and Game, Boards Support Section: Juneau.

Alaska Department of Fish and Game

1985. *Alaska habitat management guide reference maps, Southwest region, volume IV: human use of fish and wildlife.* Alaska Department of Fish and Game Division of Habitat: Juneau.

Alaska Department of Fish and Game and Alaska Inter-Tribal Council

2000. *Recommendations for a Unified Subsistence Fisheries Harvest Monitoring Program.* Prepared by the Subsistence Fisheries Harvest Assessment Working Group for the Office of Subsistence Management, U.S. Fish and Wildlife Service: Anchorage.

Andersen, D.B.

1992. *The use of dog teams and the use of subsistence-caught fish for feeding sled dogs in the Yukon River drainage, Alaska.* Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 210. <http://www.subsistence.adfg.state.ak.us/techpap/tp210.pdf>

Andersen, D.B., C.L. Brown, R.J. Walker, and K. Elkin

2004. *Traditional ecological knowledge and contemporary subsistence harvest of non-salmon fish in the Koyukuk River drainage, Alaska.* Alaska Department of Fish and Game, Division of Subsistence, Technical Paper No. 282. <http://www.adfg.alaska.gov/techpap/tp282.pdf>

Anderson, J.L.

2003. *Estimation of late run sockeye and coho salmon escapement in the Clark River, a tributary to Chignik Lake, Alaska Peninsula National Wildlife Refuge, 2002.* U. S. Fish and Wildlife Service, King Salmon Fish and Wildlife Field Office, Alaska Fisheries Technical Report Number 64: King Salmon.

Anderson, T.J., C.W. Russell, and M.B. Foster

2013. *Chignik Management Area salmon and herring annual management report, 2012.* Alaska Department of Fish and Game, Division of Sport Fish Research and Technical Services, Fishery Management Report No. 13-29: Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/FMR13-29.pdf>

- Andrews, E. and M. Coffing
1986. *Kuskokwim River subsistence Chinook fisheries: An overview*. Alaska Department of Fish and Game, Division of Subsistence. <http://www.adfg.alaska.gov/techpap/tp146.pdf> (Accessed December 11, 2012)
- Andrews, E.F.
1989. *The Akulmiut: territorial dimensions of a Yup'ik Eskimo society*. Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 177. <http://www.subsistence.adfg.state.ak.us/techpap/tp177.pdf>
- Bacon, J.J., T.R. Hepa, H.K. Brower Jr., M. Pederson, T.P. Olemaun, J.C. George, and B.G. Corrigan
2011rev.. [2009] *Estimates of subsistence harvest for villages on the North Slope of Alaska, 1994–2003*. North Slope Borough, Department of Wildlife Management: Barrow.
- Barker, J.H.
1993. *Always getting ready: Upterrlainarluta: Yup'ik Eskimo subsistence in southwest Alaska*. University of Washington Press: Seattle. ISBN 0295972343
- BBNA (Bristol Bay Native Association) and ADF&G
1996. *The harvest and use of freshwater fish in Togiak and Manokotak, 1994–95*. Natural Resource Department Bristol Bay Native Association, and Alaska Department of Fish and Game Division of Subsistence: Dillingham.
- Borba, B.M. and H.H. Hamner
2001. *Subsistence and personal use salmon harvest estimates, Yukon Area, 2000*. Alaska Department of Fish and Game Division of Commercial Fisheries Regional Information Report No. 3A01-27: Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/RIR.3A.2001.27.pdf>
- Bouwens, K.A.
2004. *An overview of the Chignik Management Area herring and salmon fisheries and stock status: report to the Alaska Board of Fisheries, November 2004*. Alaska Department of Fish and Game Division of Commercial Fisheries: Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fmr04-09.pdf>
2005. *Chignik management area salmon fisheries and stock status with particular reference to the cooperative fishery management plan: a report to the Alaska Board of Fisheries, November 2005*. Alaska Department of Fish and Game Division of Commercial Fisheries: Anchorage.
- Braem, N.M., J.S. Magdanz, D.S. Koster, and P. Fox
2013. *Subsistence harvests in Northwest Alaska: Selawik, 2010–2011*. Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 389: Fairbanks.
- Brannnian, L. and J. Fox
1996. *Upper Cook Inlet subsistence and personal use fisheries*. Alaska Department of Fish and Game Division of Commercial Fisheries Management and Development, Regional Information Report No. 2A96-03: Anchorage.
- Braund, S.R.
1982rev.. [1980] *Cook Inlet subsistence salmon fishery*. Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 54. <http://www.adfg.alaska.gov/techpap/tp054.pdf>
- Brown, C., J. Burr, K. Elkin, and R.J. Walker
2005. *Contemporary subsistence uses and population distribution of non-salmon fish in Grayling, Anvik, Shageluk, and Holy Cross*. Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 289: Fairbanks. <http://www.adfg.alaska.gov/techpap/tp289.pdf> (Accessed December 10, 2012)
- Brown, C., D. Koster, and P. Koontz
2010. *Traditional Ecological Knowledge and Harvest Survey of Nonsalmon fish in the Middle Yukon River Region, Alaska 2005–2008*. Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 358: Fairbanks. <http://www.adfg.alaska.gov/techpap/TP358.pdf>

- Brown, C.L., H. Ikuta, D.S. Koster, and J.S. Magdanz
2013. *Subsistence harvests in 6 communities in the Kuskokwim River drainage, 2010*. Alaska Department of Fish and Game Division of Subsistence Technical Paper No. 379: Fairbanks. <http://www.adfg.alaska.gov/techpap/TP%20379.pdf>
- Brown, C.L., J.S. Magdanz, D.S. Koster, and N.S. Braem
2012. *Subsistence harvests in 8 communities in the central Kuskokwim River drainage, 2009*. Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 365: Fairbanks. <http://www.adfg.alaska.gov/techpap/TP%20365.pdf>
- Brown, C.L., L.J. Slayton, A. Trainor, D.S. Koster, and M. Kostick
2014. *Wild resource harvests and uses, land use patterns, and subsistence economies in Manley Hot Springs and Minto, Alaska, 2012*. Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 400: Fairbanks.
- Bue, B.G., K.L. Schaberg, Z.W. Liller, and D.B. Molyneaux
2012. *Estimates of the historic run and escapement for the Chinook salmon stock returning to the Kuskokwim River, 1976–2011*. Alaska Department of Fish and Game, Fishery Data Series No. 12-49: Anchorage.
- Carroll, H.C. and T. Hamazaki
2012a. *Subsistence salmon harvests in the Kuskokwim area, 2008 and 2009*. Alaska Department of Fish and Game, Fishery Data Series No. 12-35 Anchorage: Anchorage. <http://www.adfg.alaska.gov/FedAidpdfs/FDS12-35>
- 2012b. *Subsistence salmon harvests in the Kuskokwim area, 2010*. Alaska Department of Fish and Game, Fishery Data Series No. 12-38 Anchorage: Anchorage. <http://www.adfg.alaska.gov/FedAidpdfs/FDS12-38>
- Coffing, M.W.
1991. *Kwethluk subsistence: Contemporary land use patterns, wild resource harvest and use and the subsistence economy of a lower Kuskokwim River area community*. Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 157: Juneau. <http://www.adfg.alaska.gov/techpap/tp157.pdf> (Accessed December 11, 2012)
- Coiley-Kenner, P., T.M. Krieg, M.B. Chythlook, and G. Jennings
2003. *Wild resource harvests and uses by residents of Manokotak, Togiak, and Twin Hills, 1999/2000*. Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 275. <http://www.adfg.alaska.gov/techpap/tp275.pdf>
- Conrad, S. and W. Davidson
2013. *Overview of the 2012 southeast Alaska and Yakutat commercial, personal use, and subsistence salmon fisheries*. Alaska Department of Fish and Game Division of Commercial Fisheries, Fishery Management Series Report No. 13-03: Anchorage.
- Davis, B.
2005. *Subsistence fisheries harvest assessment and traditional ecological knowledge, lower Alaska Peninsula and Aleutian islands*. Alaska Department of Fish and Game, Division of Subsistence.
- Dunker, K.J.
2010. *Upper Cook Inlet personal use salmon fisheries, 2007–2009*. Alaska Department of Fish and Game Divisions of Sport Fish and Commercial Fisheries, Fishery Data Series No. 10-89: Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/FDS10-89.pdf>
- Evans, S., M. Kukkonen, D. Holen, and D.S. Koster
2013. *Harvests and uses of wild resources in Dillingham, Alaska, 2010*. Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 375.
- Fall, J.A.
2006a. *Update of the status of subsistence uses in Exxon Valdez oil spill area communities, 2003*. Alaska Department of Fish and Game, Division of Subsistence. <http://www.adfg.alaska.gov/techpap/tp312.pdf>

- 2006b. *Update of the status of subsistence uses in Exxon Valdez oil spill area communities, 2003*. Alaska Department of Fish and Game, Division of Subsistence, Technical Paper No. 312. <http://www.adfg.alaska.gov/techpap/tp312.pdf>
2014. *Subsistence in Alaska: a year 2012 update*. Alaska Department of Fish and Game Division of Subsistence: Anchorage. http://www.adfg.alaska.gov/static/home/subsistence/pdfs/subsistence_update_2012.pdf
- Fall, J.A., D.B. Andersen, L. Brown, M. Coffing, G. Jennings, C. Mishler, A. Paige, C.J. Utermohle, and V. Vanek 1993. *Noncommercial harvests and uses of wild resources in Sand Point, Alaska, 1992*. Alaska Department of Fish and Game, Division of Subsistence, Technical Paper No. 226: Juneau. <http://www.subsistence.adfg.state.ak.us/techpap/tp226.pdf>
- Fall, J.A., N. Braem, S. Evans, D. Holen, T. Krieg, R. LaVine, T. Lemons, M.A. Marchioni, D.M. Runfola, L. Hutchinson-Scarborough, L. Sill, A. Trainor, and J.M. VanLanen 2012a. *Alaska subsistence salmon fisheries 2009 annual report*. Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 373: Anchorage. <http://www.adfg.alaska.gov/techpap/TP373.pdf>
- 2012b. *Alaska subsistence salmon fisheries 2009 annual report*. Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 373: Anchorage.
- Fall, J.A., A.R. Brenner, S.S. Evans, L. Hutchinson-Scarborough, B. Jones, R. La Vine, T. Lemons, M.A. Marchioni, E. Mikow, J.T. Ream, L.A. Sill, and A. Trainor 2013. *Alaska subsistence and personal use salmon fisheries 2011 annual report*. Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 387: Anchorage. <http://www.adfg.alaska.gov/techpap/TP387.pdf>
- Fall, J.A., D. Caylor, M. Turek, C. Brown, T. Krauthoefer, B. Davis, and D. Koster 2007. *Alaska subsistence fisheries 2004 annual report*. Alaska Department of Fish and Game Division of Subsistence. <http://www.adfg.alaska.gov/techpap/tp317.pdf> (Accessed December 7, 2012)
- Fall, J.A. and M.B. Chythlook 1997. *An overview of the subsistence fisheries of the Bristol Bay management area*. Alaska Department of Fish and Game Division of Subsistence: King Salmon.
- Fall, J.A., M.B. Chythlook, T.M. Krieg, and G. Jennings 2001. *Overview of the subsistence sockeye salmon fishery of the Kvichak River watershed, Bristol Bay, Southwest Alaska. Report to the Alaska Board of Fisheries, January 2001*. Alaska Department of Fish and Game, Division of Subsistence: Anchorage.
- Fall, J.A., M.B. Chythlook, J.C. Schichnes, and J.M. Morris 1996. *An overview of the harvest and use of freshwater fish by the communities of the Bristol Bay region, Southwest Alaska*. Alaska Department of Fish and Game, Division of Subsistence. <http://www.adfg.alaska.gov/techpap/tp166.pdf> (Accessed December 7, 2012)
- Fall, J.A., D.J. Foster, and R.T. Stanek 1984. *The use of fish and wildlife resources in Tyonek, Alaska. Tubughna Ch'adach' Elnen Ghuhdilt'a*. Alaska Department of Fish and Game, Division of Subsistence, Technical Paper No. 105: Anchorage.
- Fall, J.A., D.L. Holen, B. Davis, T. Krieg, and D. Koster 2006. *Subsistence harvests and uses of wild resources in Iliamna, Newhalen, Nondalton, Pedro Bay, and Port Alsworth, Alaska, 2004*. Alaska Department of Fish and Game, Division of Subsistence. <http://www.adfg.alaska.gov/techpap/tp302.pdf>
- Fall, J.A., L.B. Hutchinson-Scarborough, and P.A. Coiley 1995. *Fish and wildlife harvest and use in five Alaska Peninsula communities, 1989. Subsistence uses in Chignik Bay, Chignik Lagoon, Chignik Lake, Ivanof Bay and Perryville*. Alaska Department of Fish and Game, Division of Subsistence. <http://www.adfg.alaska.gov/techpap/tp202.pdf> (Accessed December 10, 2012)

- Fall, J.A. and D.S. Koster
2014. *Subsistence harvests of Pacific halibut in Alaska, 2012*. Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 388: Anchorage. <http://www.adfg.alaska.gov/techpap/TP388.pdf>
- Fall, J.A., T. Krieg, and M. Chythlook
2003. *An overview of the subsistence fisheries of the Bristol Bay management area*. Alaska Department of Fish and Game, Division of Subsistence. Report to the Alaska Board of Fisheries, December 2003: Anchorage.
- Fall, J.A., T.M. Krieg, and D.L. Holen
2009. *Overview of the subsistence fisheries of the Bristol Bay Management Area*. Alaska Department of Fish and Game Division of Subsistence Special Publication No. BOF 2009-07: Anchorage. http://www.subsistence.adfg.state.ak.us/specialpubs/SP2_SP2009-007.pdf
- Fall, J.A., R. Mason, T. Haynes, V. Vanek, L. Brown, G. Jennings, C. Mishler, and C. Utermohle
1993. *Noncommercial harvests and uses of wild resources in King Cove, Alaska, 1992*. Alaska Department of Fish and Game, Division of Subsistence. <http://www.adfg.alaska.gov/techpap/tp227.pdf> (Accessed December 11, 2012)
- Fall, J.A., J.C. Schichnes, M. Chythlook, and R.J. Walker
1986. *Patterns of wild resource use in Dillingham: Hunting and fishing in an Alaskan regional center*. Alaska Department of Fish and Game, Division of Subsistence. <http://www.adfg.alaska.gov/techpap/tp135.pdf> (Accessed December 11, 2012)
- Fall, J.A. and R. Shanks
2000. *Subsistence Fisheries Harvest Monitoring Strategy*. U.S. Fish and Wildlife Service, Office of Subsistence Management, Fisheries Resource Monitoring Program Final Report (Study No. 00-017). Alaska Department of Fish and Game, Division of Subsistence: Anchorage. <http://www.subsistence.adfg.state.ak.us/download/download/ssfhms.pdf>
- Fall, J.A. and R.T. Stanek
1990. *An overview of subsistence and personal use salmon fisheries in the Cook Inlet area: a report to the Alaska Board of Fisheries*. Alaska Department of Fish and Game Division of Subsistence: Anchorage.
- Fall, J.A., M.F. Turek, N.M. Braem, J.J. Simon, W.E. Simeone, D.L. Holen, L.C. Naves, L. Hutchinson-Scarborough, T. Lemons, V. Ciccione, T.M. Krieg, and D.S. Koster
2009. *Alaska subsistence salmon fisheries 2007 annual report*. Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 346: Anchorage. <http://www.subsistence.adfg.state.ak.us/techpap/TP346.pdf>
- Fienup-Riordan, A.
1983. *The Nelson Island Eskimo: social structure and ritual distribution*, no. 40 book in the The Alaskan book series. Alaska Pacific University Press: Anchorage, Alaska. ISBN 0935094091

1990. *Eskimo essays: Yup'ik lives and how we see them*. Rutgers University Press: New Brunswick. ISBN 0813515882

1995. *Boundaries and Passages: Rule and Ritual in Yup'ik Eskimo Oral Tradition*. University of Oklahoma Press: Norman, OK.
- Frothingham, A.
2011. *TCC Works with PSMFC on Net Exchange*. The Council: A Report to the Member Tribes of the Tanana Chiefs Conference 36(3), page 5.
- Hamazaki, T.
2011. *2011 reconstruction of subsistence harvests in the Kuskokwim areas, 1990–2009*. Alaska Department of Fish and Game, Fishery Manuscript No. 11-09: Juneau. www.adfg.alaska.gov/FedAidpdfs/FMS11-09.pdf
- Hammarstrom, L.F. and M.S. Dickson
2006. *2005 Lower Cook Inlet annual finfish management report*. Alaska Department of Fish and Game Divisions of Sport and Commercial Fisheries: Anchorage.

- Hammarstrom, L.F. and E.G. Ford
2011. *2010 Lower Cook Inlet annual finfish management report*. Alaska Department of Fish and Game, Fishery Management Report No. 11-26: Anchorage.
- Harritt, R.K.
2010. *Recent work at Difchahak, a center of Norton culture in Eastern Norton Sound, Alaska*. Arctic Anthropology 47(2), pages 80–89.
- Himmelheber, H.
1987. *Eskimo artists (fieldwork in Alaska, June 1936 until April 1937). [English translation of Eskimokünstler from the original German]*. Museum Rietberg: Zürich.
- Hochhalter, S.J. and P.A. Hansen
2011. *Summary of the noncommercial pot shrimp fishery by permit for Prince William Sound, Alaska, 2002–2005 and 2009–2010*. Alaska Department of Fish and Game, Fishery Data Series No. 11-67: Anchorage.
<http://www.adfg.alaska.gov/FedAidPDFs/FDS11-67.pdf>
- Holen, D.L., S.S. Evans, and B. Jones
2012. *The subsistence harvest of herring spawn on kelp in the Togiak District, Alaska, 2011 and 2012*. Alaska Department of Fish and Game, Division of Subsistence Special Publication No. BOF 2012-06: Anchorage.
- Holen, D.L. and J.A. Fall
2011. *Overview of subsistence salmon fisheries in the Tyonek Subdistrict and Yentna River, Cook Inlet, Alaska*. Alaska Department of Fish and Game, Division of Subsistence Special Publication No. BOF 2011-01: Anchorage.
- Holen, D.L., T.M. Krieg, and T. Lemons
2011. *Subsistence harvests and uses of wild resources in King Salmon, Naknek, and South Naknek, Alaska, 2007*. Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 360: Anchorage.
<http://www.adfg.alaska.gov/techpap/TP360.pdf>
- Holen, D.L. and T. Lemons
2012. *An overview of the subsistence fisheries of the Bristol Bay Management Area*. Alaska Department of Fish and Game Division of Subsistence Special Publication No. BOF 2012-05: Anchorage.
- Holen, D.L., J. Stariwat, T.M. Krieg, and T. Lemons
2012. *Subsistence harvests and uses of wild resources in Aleknagik, Clark's Point, and Manokotak, Alaska, 2008*. Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 368: Anchorage.
<http://www.adfg.alaska.gov/techpap/TP%20368.pdf>
- Hollowell, G., T. Otis, and E. Ford
2012. *2011 Lower Cook Inlet area finfish management report*. Alaska Department of Fish and Game Divisions of Sport Fish and Commercial Fisheries, Fishery Management Report No. 12-30: Anchorage.
- Hutchinson-Scarborough, L. and J.A. Fall
1996. *An overview of subsistence salmon and other subsistence fisheries of the Chignik management area, Alaska Peninsula, Southwest Alaska*. Alaska Department of Fish and Game, Division of Subsistence.
<http://www.adfg.alaska.gov/techpap/tp230.pdf> (Accessed December 7, 2012)
- Hutchinson-Scarborough, L., T. Lemons, J.A. Fall, D.L. Holen, and L. Olson
2010. *Chignik area subsistence salmon fisheries research report to the Alaska Board of Fisheries, January 2011*. Alaska Department of Fish and Game, Division of Subsistence Special Publication No. BOF 2010-06: Anchorage.
- Ikuta, H.
2012. *Options for amounts reasonably necessary for subsistence uses of salmon: Kuskokwim area; prepared for the January 2013 Anchorage Alaska Board of Fisheries meeting*. Alaska Department of Fish and Game Division of Subsistence, Special Publication No. 2012-07: Fairbanks.
http://www.adfg.alaska.gov/specialpubs/SP2_SP2012-007.pdf

- Ikuta, H., A.R. Brenner, and A. Goddhun
2013. *Socioeconomic patterns in subsistence salmon fisheries: historical and contemporary trends in five Kuskokwim River communities and overview of the 2012 season*. Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 382: Fairbanks. <http://www.adfg.alaska.gov/techpap/TP%20382.pdf>
- Ikuta, H., C.L. Brown, and D.S. Koster
2014. *Subsistence harvests in 8 communities in the Kuskokwim River drainage and lower Yukon River, 2011*. Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 396: Fairbanks. <http://www.adfg.alaska.gov/techpap/TP396.pdf>
- Jackson, J., J. Dinnocenzo, G. Spalinger, and M. Keyse
2012. *Kodiak Management Area commercial salmon fishery annual management report, 2011*. Alaska Department of Fish and Game, Fishery Management Report No. 12-48: Anchorage. www.adfg.alaska.gov/fedaidpdfs/fmr12-48.pdf
- Jackson, J.V.
2009. *Chignik Management Area Commercial Salmon Fishery Harvest Strategy, 2009*. Alaska Department of Fish and Game, Fishery Management Report No. 09-15: Anchorage. <http://www.cf.adfg.state.ak.us/FedAidPDFs/FMR09-15.pdf>
- Jallen, D.M., S.K.S. Decker, and T. Hamazaki
In prep. *Subsistence and personal use salmon harvests in the Alaska portion of the Yukon River drainage, 2012*. Alaska Department of Fish and Game, Fishery Data Series No. 14-XX: Anchorage.
2012. *Subsistence and personal use salmon harvests in the Alaska portion of the Yukon River drainage, 2011*. Alaska Department of Fish and Game, Fishery Data Series No. 12-72: Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/FDS12-72.pdf>
- Jones, M., T. Sands, C. Brazil, G. Buck, F. West, P. Salomone, S. Morstad, and T. Krieg
2014. *2013 Bristol Bay area annual management report*. Alaska Department of Fish and Game, Fishery Management Report No. 14-23: Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/FMR14-23.pdf>
- Jones, M., T. Sands, S. Morstad, P. Salamone, T. Baker, G. Buck, and F. West
2009. *2008 Bristol Bay area annual management report*. Alaska Department of Fish and Game, Fishery Management Report 09-30: Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/FMR09-30.pdf>
- Jones, M., T. Sands, S. Morstad, P. Salomone, G. Buck, F. West, T. Baker, and T. Krieg
2012. *2011 Bristol Bay area annual management report*. Alaska Department of Fish and Game, Fishery Management Report No. 12-21: Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/FMR12-21.pdf>
- Jones, M., T. Sands, S. Morstad, P. Salomone, G. Buck, F. West, C. Brazil, and T. Krieg
2013. *2012 Bristol Bay area annual management report*. Alaska Department of Fish and Game, Fishery Management Report No. 13-20: Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/FMR13-20.pdf>
- JTC (Joint Technical Committee of the Yukon River US/Canada Panel)
2012. *Yukon River salmon 2011 season summary and 2012 season outlook*. Alaska Department of Fish and Game, Division of Commerical Fisheries, Regional Information Report No. 3A12-01: Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/RIR.3A.2012.01.pdf>
2013. *Yukon River salmon 2012 season summary and 2013 season outlook*. Alaska Department of Fish and Game, Division of Commerical Fisheries, Regional Information Report No. 3A13-02: Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/RIR.3A.2013.02.pdf>
- Koskey, M. and K. Mull
2011. *Traditional Ecological Knowledge and Biological Sampling of Nonsalmon Fish Species in the Yukon Flats Region, Alaska. Final report for study 06-252*. Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 362: Fairbanks. <http://www.subsistence.adfg.state.ak.us/TechPap/TP362.pdf>

- Krauthoefer, T., J. Simon, M. Coffing, M. Kerlin, and W. Morgan
2007. *The harvest of non-salmon fish by residents of Aniak and Chuathbaluk, Alaska, 2001–2003*. Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 299: Juneau.
<http://www.adfg.alaska.gov/techpap/tp299.pdf>
- Krieg, T., M. Chythlook, P. Coiley-Kenner, D. Holen, K. Kamletz, and H. Nicholson
2005. *Freshwater fish harvest and use in communities of the Kvichak watershed, 2003*. Alaska Department of Fish and Game, Division of Subsistence. <http://www.adfg.alaska.gov/techpap/tp297.pdf> (Accessed December 10, 2012)
- Krieg, T.M., D.L. Holen, and D. Koster
2009. *Subsistence harvests and uses of wild resources in Igiugig, Kokhanok, Koliganek, Levelock, and New Stuyahok, Alaska, 2005*. Alaska Department of Fish and Game Division of Subsistence Technical Paper No. 322.
<http://www.adfg.alaska.gov/techpap/TP322.pdf>
- Lingnau, T. and P. Salomone
2003. *Informational Letter. Preliminary 2003 Yukon area Chinook and summer chum salmon fishery summary*.
- Magdanz, J.S.
1992. *Subsistence salmon fishing by permit in the Nome subdistrict and portions of the Port Clarence district, 1975–91*. Alaska Department of Fish and Game Division of Subsistence: Juneau.
- Magdanz, J.S., N.S. Braem, B.C. Robbins, and D.S. Koster
2010. *Subsistence harvests in Northwest Alaska, Kivalina and Noatak, 2007*. Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 354: Kotzebue.
www.subsistence.adfg.state.ak.us/techpap/tp354.pdf
- Magdanz, J.S., H. Smith, N. Braem, P. Fox, and D.S. Koster
2011. *Patterns and trends in subsistence fish harvests, Northwest Alaska, 1994–2004*. Alaska Department of Fish and Game Division of Subsistence, Technical Paper 366: Kotzebue.
<http://www.adfg.alaska.gov/techpap/TP%20366.pdf>
- Menard, J.
2010. *2010 Norton Sound salmon fisheries management plan*. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report No. 3A10-03: Anchorage.
<http://www.sf.adfg.state.ak.us/FedAidPDFs/RIR.3A.2010.03.pdf>
- Menard, J. and D.J. Bergstrom
2006. *Norton Sound Golovin and Moses Point subdistricts chum salmon stock status and action plan, 2007; a report to the Alaska Board of Fisheries*. Alaska Department of Fish and Game Division of Commercial Fisheries: Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/sp06-32.pdf>
- Menard, J. and S. Kent
2007. *2007 Norton Sound salmon season summary*. Alaska Department of Fish and Game Division of Commercial Fisheries [press release]: Nome.
- Menard, J., J. Soong, and S. Kent
2011. *2009 annual management report Norton Sound, Port Clarence, and Kotzebue*. Alaska Department of Fish and Game, Fishery Management Report No. 11-46: Anchorage.
<http://www.adfg.alaska.gov/FedAidPDFs/FMR11-46.pdf>
2012. *2010 annual management report Norton Sound, Port Clarence, and Kotzebue*. Alaska Department of Fish and Game, Fishery Management Report No. 12-31: Anchorage.
<http://www.adfg.alaska.gov/FedAidPDFs/FMR12-31.pdf>
- Menard, J., J. Soong, S. Kent, and A. Brown
2013. *2012 Annual management report Norton Sound-Port Clarence Area, and Arctic-Kotzebue*. Alaska Department of Fish and Game, Fishery Management Report No. 13-28: Anchorage.
<http://www.adfg.alaska.gov/FedAidPDFs/FMR13-28.pdf>

- Morris, J.M.
1987. *Fish and wildlife uses in six Alaska Peninsula communities: Egegik, Chignik, Chignik Lagoon, Chignik Lake, Perryville, and Ivanof Bay*. Alaska Department of Fish and Game, Division of Subsistence. <http://www.adfg.alaska.gov/techpap/tp151.pdf> (Accessed December 10, 2012)
- Nelson, D.
1994. *1993 Area Management Report for the Recreational Fisheries of the Kenai Peninsula*. Alaska Department of Fish and Game Division of Sport Fish, Fishery Management Report No. 94-7: Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fmr94-07.pdf>

1995. *Area Management Report for the Recreational Fisheries of the Kenai Peninsula, 1994*. Alaska Department of Fish and Game Division of Sport Fish, Fishery Management Report No. 94-7: Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fmr95-04.pdf>
- Nelson, D., D. Athons, P. Berkhahn, and S. Sonnichsen
1999. *Area management report for the recreational fisheries of the Kenai Peninsula, 1995–1997*. Alaska Department of Fish and Game, Fishery Management Report No. 99-3: Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fmr99-03.pdf>
- Oswalt, W.H.
1963a. *Napaskiak: an Alaskan Eskimo community*. University of Arizona Press: Tuscon.

1963b. *Mission of change in Alaska: Eskimos and Moravians on the Kuskokwim*. Huntington Library: San Marino.

1990. *Bashful No Longer: An Alaskan Eskimo Ethnohistory, 1778–1988*. University of Oklahoma Press: Norman, OK.
- Pete, M.C.
1993. “Foreword: Always getting ready: Upterrlainarluta: Yup’ik Eskimo subsistence in southwest Alaska” [in] J.H. Barker *Always getting ready: Upterrlainarluta: Yup’ik Eskimo subsistence in southwest Alaska*. University of Washington Press: Seattle. ISBN 0295972343
- Poetter, A.D., M.D. Keyse, and A.R. Tiernan
2012. *South Alaska Peninsula salmon annual management report*. Alaska Department of Fish and Game, Fishery Management Report No. 12-42: Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/FMR12-42.pdf>
- Ray, L., C. Brown, A. Russell, T. Krauthoefer, C. Wassillie, and J. Hooper
2010. *Local knowledge and harvest monitoring of nonsalmon fishes in the lower Kuskokwim River region, Alaska, 2005–2009. Final report to the U.S. Fish and Wildlife Service, Office of Subsistence Management, to fulfill obligations for Study No. FIS 06-351*. Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 356: Fairbanks. <http://www.adfg.alaska.gov/techpap/TP356.pdf>
- Scheaffer, R.L.
1990. *Elementary survey sampling*, 4th ed edition. PWS-Kent: Boston. ISBN 053492185X
- Schichnes, J. and M. Chythlook
1988. *Use of fish and wildlife in Manokotak, Alaska*. Alaska Department of Fish and Game, Division of Subsistence. <http://www.adfg.alaska.gov/techpap/tp152.pdf>

1991. *Contemporary use of fish and wildlife in Ekwok, Koliganek, and New Stuyahok, Alaska*. Alaska Department of Fish and Game, Division of Subsistence. <http://www.adfg.alaska.gov/techpap/tp185.pdf> (Accessed December 10, 2012)
- Scott, C.L., L.A. Brown, G.B. Jennings, and C.J. Utermohle
2001. *Community Profile Database*. Alaska Department of Fish and Game Division of Subsistence: Juneau.

- Senecal-Albrecht, D.
1998. *"Don't wait for Boldt": building co-management from the ground up: the success of salmon fishermen's groups in western Alaska*. "Crossing Boundaries," the seventh annual conference of the International Association for the Study of Common Property. Vancouver, British Columbia, Canada, June 10–14, 1998: Vancouver, B.C.
- Senecal-Albrecht, D.E.
1990. *Co-management as transaction: the Kuskokwim River Salmon Management Working Group*. McGill University: Montreal.
- Shelden, C.A., T. Hamazaki, M. Horne-Brine, G. Roczicka, M. Thalhauser, and H.C. Carroll
2014. *Subsistence salmon harvests in the Kuskokwim Area, 2011 and 2012*. Annual Report for Study 10-352 USFWS Office of Subsistence Management, Fisheries Resource Monitoring Program. ADF&G Division of Sport Fish and Commercial Fisheries: Anchorage. <http://www.adfg.alaska.gov/FedAidpdfs/FDS14-20>
- Shields, P. and A. Dupuis
2012. *Upper Cook Inlet Commercial Fisheries Annual Management Report, 2011*. Alaska Department of Fish and Game, Fishery Management Report No. 12-25: Anchorage. <http://www.adfg.alaska.gov/FedAidpdfs/FMR12-25>
- Simeone, W.E. and J.A. Fall
1996. *Patterns and trends in the subsistence salmon fishery of the Upper Copper River, Alaska*. Alaska Department of Fish and Game, Division of Subsistence: Cordova.
- Simeone, W.E. and J.M. Kari
2005. *The harvest and use of non-salmon fish species in the Copper River Basin, Alaska*. Alaska Department of Fish and Game, Division of Subsistence, Technical Paper No. 292. <http://www.adfg.alaska.gov/techpap/tp292.pdf>
- Simon, J., T. Krauthoefer, D. Koster, and D. Caylor
2007. *Subsistence salmon harvest monitoring report, Kuskokwim fisheries management area, Alaska, 2004*. Alaska Department of Fish and Game, Division of Subsistence, Technical Paper No. 313. <http://www.adfg.alaska.gov/techpap/tp313.pdf>
- Simon, J., T. Krauthoefer, D. Koster, M. Coffing, and D. Caylor
2007. *Bethel subsistence fishing harvest monitoring report, Kuskokwim fisheries management area, Alaska, 2001-2003*. Alaska Dept. of Fish and Game, Division of Subsistence. <http://www.adfg.alaska.gov/techpap/tp330.pdf> (Accessed December 10, 2012)
- Smith, E.A. and J.C. Linderman Jr.
2008. *Activities of the Kuskokwim River salmon management working group, 2007*. Alaska Department of Fish and Game, Fishery Management Report No. 08-74: Anchorage. <http://sf.adfg.state.ak.us/FedAidPDFs/FMR08-74.pdf>
- Smith, H.L. and J. Vreeman
1995. *Cultural resource inventory in the Bendeleben and Darby Mountains, Seward Peninsula, Alaska*. U.S. Department of the Interior, Bureau of Land Management, Alaska State Office: Anchorage.
- Soong, J., A.O. Banducci, S. Kent, and J. Menard
2008. *2007 annual management report Norton Sound, Port Clarence, and Kotzebue*. Alaska Department of Fish and Game Division of Commercial Fisheries: Anchorage. <http://www.adfg.alaska.gov/FedAidPDFs/FMR11-46.pdf>
- Stanek, R.T.
1985. *Patterns of wild resource use in English Bay and Port Graham, Alaska*. Alaska Department of Fish and Game, Division of Subsistence. <http://www.adfg.alaska.gov/techpap/tp104.pdf> (Accessed December 11, 2012)
- Stanek, R.T., D.L. Holen, and C. Wassillie
2007. *Harvest and uses of wild resources in Tyonek and Beluga, Alaska, 2005-2006*. Alaska Department of Fish and Game, Division of Subsistence. <http://www.adfg.alaska.gov/techpap/TP321.pdf> (Accessed December 10, 2012)

Stichert, M.A.

2007a. *2005 Chignik management area salmon and herring annual management report*. Alaska Department of Fish and Game, Fishery Management report No. 07-15: Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fmr07-15.pdf>

2007b. *Chignik management area salmon and herring annual management report, 2006*. Alaska Department of Fish and Game, Fishery Management report No. 07-56: Anchorage. <http://www.sf.adfg.state.ak.us/FedAidPDFs/fmr07-56.pdf>

Stickman, K., A. Balluta, M. McBurney, and D. Young

2003. *Nondalton Traditional Ecological Knowledge of Freshwater Fish*. U. S. Fish and Wildlife Service of Subsistence Management, Fisheries Resource Monitoring Program, Final Report (Study No. 01-075): Anchorage.

Stickney, A.

1984. *Coastal ecology and wild resource use in the central Bering Sea Area: Hooper Bay and Kwigillingok*. Alaska Department of Fish and Game, Division of Subsistence, Technical Paper No. 85. <http://www.adfg.alaska.gov/techpap/tp085.pdf> (Accessed December 10, 2012)

Tingley, A.C. and B. Davidson

2011. *Overview of the 2010 southeast Alaska and Yakutat commercial, personal use, and subsistence salmon fisheries*. Alaska Department of Fish and Game Division of Commercial Fisheries, Fishery Management Series Report No. 11-39: Anchorage.

Turek, M., N. Ratner, W.E. Simeone, and D.L. Holen

2009. *Subsistence harvests and local knowledge of rockfish *Sebastes* in four Alaska communities: final report to the North Pacific Research Board*. Alaska Department of Fish and Game Division of Subsistence Technical Paper No. 337: Juneau. <http://www.adfg.alaska.gov/techpap/TP337.pdf>

Walker, R.J. and M.W. Coffing

1993. *Subsistence salmon harvests in the Kuskokwim area during 1989*. Alaska Department of Fish and Game, Division of Subsistence Technical Paper No. 189: Juneau. <http://www.subsistence.adfg.state.ak.us/TechPap/tp189.pdf>

Wilburn, D.M. and N.W. Nichols

2013. *Annual summary of the commercial and subsistence salmon fisheries and salmon escapements in the Alaska Peninsula, Aleutian Islands, and Atka-Amlia Islands Management Areas, 2012*. Alaska Department of Fish and Game, Fishery Management Report No. 13-26: Anchorage.

Wolfe, R.J., J.J. Gross, S.J. Langdon, J.M. Wright, G.K. Sherrod, L.J. Ellanna, and V. Sumida

1984. *Subsistence-based economies in coastal communities of Southwest Alaska*. Alaska Department of Fish and Game Division of Subsistence, Technical Paper No. 89; Minerals Management Service, Alaska Region, U.S. Department of the Interior. <http://www.adfg.alaska.gov/techpap/tp089.pdf>

Wolfe, R.J., C. Stockdale, and C. Scott

2012. *Salmon harvests in coastal communities of the Kuskokwim Area, southwest Alaska*. Arctic-Yukon-Kuskokwim Sustainable Salmon Initiative: Anchorage.

Wright, J.M. and M.B. Chythlook

1985. *Subsistence harvests of herring spawn-on-kelp in the Togiak district of Bristol Bay*. Alaska Department of Fish and Game, Division of Subsistence. <http://www.adfg.alaska.gov/techpap/tp116.pdf>

Wright, J.M., J.M. Morris, and R. Schroeder

1985. *Bristol Bay regional subsistence profile*. Alaska Department of Fish and Game, Division of Subsistence. <http://www.adfg.alaska.gov/techpap/tp114.pdf> (Accessed December 10, 2012)