



U. S. Department of the Interior
Bureau of Land Management


BLM-Alaska Open File Report 52
BLM/AK/ST-94/021+6700+070
June 1994



Alaska State Office
222 West 7th, #13
Anchorage, Alaska 99513

Waterfowl Production on the Central Seward Peninsula, Alaska: Final Report

R. R. Jandt and A. E. Morkill



Authors

A. E. Morkill is a wildlife biologist with the Kobuk District, Fairbanks, Alaska.

R. R. Jandt is a wildlife biologist with the Kobuk District, Fairbanks, Alaska.

Open File Reports

Open File Reports identify the results of inventories or other investigations that are made available to the public outside the formal BLM-Alaska technical publication series. These reports can include preliminary or incomplete data and are not published and distributed in quantity. The reports are available at BLM offices in Alaska, the USDI Resources Library in Anchorage, various libraries of the University of Alaska, and other selected locations.

Copies are also available for inspection at the USDI Natural Resources Library in Washington, D.C., and at the BLM Service Center Library in Denver.

Waterfowl Production on the Central Seward Peninsula, Alaska: Final Report

INVESTIGATORS: R.R. Jandt and A.E. Morkill

Open File Report 52
June 1994

Table of Contents

Abstract	ii
Introduction	1
Study Area	1
Methods	2
Results	3
Kuzitrin River Basin: 1993	3
Discussion	4
Literature Cited	7

Figures

Figure 1. Study area locations on the Seward Peninsula, Alaska	1
Figure 2. Species composition of duck broods in the Kuzitrin River wetlands, Alaska, 1993	4
Figure 3. Number of broods in the three major age classes for dabblers and divers in the Kuzitrin River study area, Alaska, 1993	4
Figure 4. Estimated number of young ducks, geese and swans produced in Kuzitrin River valley, Alaska, from brood survey	4
Figure 5. Comparison of young ducks/km ² produced on three areas where BLM brood surveys were conducted	6

Tables

Table 1. Adult ducks observed in the Kuzitrin River study area, Alaska, 1993	2
Table 2. Species, type of broods, numbers of broods and young ducks observed in the Kuzitrin River study area, Alaska, 1993	3
Table 3. Comparison of density of young ducks obtained during July brood surveys on several federally-managed wetland areas in Alaska	3

Abstract

Abstract : Waterfowl brood surveys were conducted on public lands in the interior region of the Seward Peninsula, Alaska during 1989-1993. Production was monitored for four consecutive years on McCarthys Marsh and for three consecutive years in the upper Kuzitrin River basin. The 1993 estimated production of young ducks from the Kuzitrin flats was $4,888 \pm 1,857$. Northern pintails (*Anas acuta*), and greater scaup (*Aythya marila*) were the most common breeders. Multi-year production averages are given for the two production areas and management implications are discussed.

INTRODUCTION

Alaska's public wetlands are becoming an increasingly valuable resource as continental waterfowl breeding habitat diminishes because of development and drought conditions. Of the original 87 million acres of wetland habitat in the conterminous United States, less than half remains (Tiner 1984). During drought years in the prairie states and provinces, more than half of all northern pintails and one third of all wigeons (*Anas americana*) breed in Alaska (Lensink and Derksen 1990). Inventories in 1989-1993 have helped to identify important wetlands on Bureau of Land Management (BLM) lands in the Kobuk District and to gain an understanding of species composition, habitat selection, and habitat quality. Such baseline data is essential to determine the impacts of development and various land management policies on breeding waterfowl.

Ground brood counts have been used in several places in Alaska to estimate waterfowl production based upon the number and size of broods counted during the peak of breeding season in July (Brubaker and Witmer 1989, Doyle 1989, FWS 1991). The Kobuk District BLM conducted brood surveys on the Seward Peninsula beginning in 1989. From 1989 to 1992, BLM conducted waterfowl brood surveys in McCarthys Marsh (Anderson and Robinson 1991a, 1991b). From 1991 to 1993, BLM also surveyed production on the wetlands along the upper Kuzitrin River, from near Bunker Hill to Lava Lake, with the cooperation of the National Park Service (NPS)-Bering Land Bridge National Preserve (Brown and Jandt 1992, Brown *et al.* 1993). Results from the 1993 field surveys and an evaluation of the multi-year inventory are included in this final report.

STUDY AREAS

McCarthys Marsh is located in the Fish River valley and is 520 km² in size (Fig. 1). The Bendeleben

Mountains border the river flats to the north, and the Darby Mountains surround them on the east and southeast. The Fish River drains the basin, flowing south to Golovnin Bay off Norton Sound. McCarthys Marsh ranges in elevation from 15-90 m. Death Valley was also included in the McCarthys Marsh survey area, adding 182 km² of waterfowl habitat to the survey. Death Valley, east of the Darby Mountains, is drained by the Tubutulik River and its elevation averages 160 m.

The Kuzitrin River survey encompasses about 835 km² of the upper river basin between Bunker Hill and Lava Lake (Fig. 1). About one third of this area lies within the Bering Land Bridge National Preserve and most of the remainder is managed by the BLM-Kobuk District. The boundaries of this survey area were based on waterfowl production areas defined by the U.S. Fish and Wildlife Service (USF&WS), Office of Migratory Bird Management. The Kuzitrin, Noxapaga and Kougarok Rivers flow into the wetlands from the west and the north. The area is drained by the Kuzitrin River, which flows west to Imuruk Basin. In the northeastern part of the Kuzitrin study area there are extensive lava fields covering >200 km². The elevation of the Kuzitrin River survey area ranges from 15-90 m.

Plant communities in the two survey areas are similar. Forest cover is confined primarily to riparian zones, and is composed of white spruce (*Picea glauca*), black spruce (*Picea mariana*) and balsam poplar (*Populus balsamifera*), as well as willows (*Salix* spp.) and alders (*Alnus crispa*). Elsewhere, treeless tundra prevails with patches of various shrub communities, including dwarf birch (*Betula nana*), willows and alders. The predominant vegetation type in McCarthys Marsh is a complex of lichen tussock tundra and drainage ways/lake systems dominated by water sedge (*Carex aquatilis*) and white cottongrass (*Eriophorum scheuchzeri*) (Swanson *et al.* 1985). The upper Kuzitrin River flats are primarily a low shrub-tussock tundra with Bigelow sedge (*C. bigelowi*), tussock cottongrass (*E. vaginatum*), Labrador tea (*Ledum decumbens*), lowbush cranberry (*Vaccinium vitis-idaea*) and cloudberry (*Rubus chaemaemorus*) (Swanson *et al.* 1985). Some parts of the Kuzitrin valley also have lichen tussock tundra. Parts of both areas burned in the 1970s: 18 km² in McCarthys Marsh in 1977 (Wagon Wheel burn) and approximately 90 km² in the Kuzitrin study area (combination of five fires in 1971).

Ponds in these study areas are generally divided into oxbow or flood-formed lakes, or those of thermokarst origin. They vary widely in the amounts of associated aquatic and emergent vegetation. The most common types of emergent and





Figure 1. Study area locations on the Seward Peninsula, Alaska.



R. Brown prepares to conduct duck brood surveys in McCarthys Marsh.

submergent vegetation observed were buckbean (*Menyanthes trifoliata*), marsh fivefinger (*Potentilla palustris*), marsh marigold (*Caltha palustris*), yellow pond lily (*Nuphar polysephalum*), various sedges (*Carex*), water hemlock (*Cicuta mackenzieana*), bur reed (*Sparganium*), horsetail (*Equisetum*), bladderwort (*Utricularia vulgaris*), duckweed (*Lemna trisulca*) and pondweed (*Potamogeton*).

METHODS

Both McCarthys Marsh and the Kuzitrin River survey areas were divided into 2.6 km² plots defined by section lines on U.S. Geological Survey (USGS) 1:63,360 scale topographic maps. Each plot was stratified as "no habitat," "poor" or "other" (good) habitat based on water surface area, number of waterbodies and the presence of stream connections appearing on the maps. Lakes with water exchange from streams and that fluctuate with river systems are more productive than stag-

nant lakes and those with stable water levels (Lensink and Derksen 1990, Murphy *et al.* 1984). A random sampling of 13-15 plots in the "poor" and "other" strata were surveyed in each area, using the same plots in subsequent survey years. A detailed description of the stratification and sampling techniques can be found in the 1991 progress report (Brown and Jandt 1992).

A floatplane and a helicopter provided access to individual plots. Field methods followed those described in the FWS Standard Operating Procedures for ground brood surveys (1991). Survey teams walked or canoed around the ponds in sample plots using binoculars and spotting scopes to observe waterfowl. Duck broods were classified as age class I, II or III, according to Bellrose (1976). Priority was given to identifying, quantifying and aging broods, but all observed waterfowl were recorded. For more complete descriptions of the methods, see Anderson and Robinson (1991b). In some years, physical characteristics of ponds, such as pH, surface temperature, depth, and water clarity were recorded and plant specimens were collected.

We would like to express our gratitude to D. Chase and the NPS, Bering Land Bridge Park and Preserve staff for helping with the logistics of this survey. Biotechnician R. Brown played an integral role in helping collect and compile the data. Several others helped collect field data, including H. Brownell, R. Corbray, W. Gregg, H. Huntington, R. Jandt, and A. Morkill from BLM-Kobuk District, and N. Olson, B. Outwater, and T. Rodgers from the NPS. We thank pilot Buck Maxson of Arctic Air

Species	Broody Hens*	Other Adults	Total Adults	% of Total
Green-winged teal	13	41	54	5
American wigeon	14	62	76	7
Black scoter	7	70	77	7
Bufflehead	0	1	1	0
Common merganser	0	5	5	0
Greater scaup	25	501	526	46
Mallard	1	5	6	1
Northern pintail	32	263	295	26
Northern shoveler	8	2	10	1
Oldsquaw	7	34	41	4
Ruddy duck	0	1	1	0
Unidentified duck	0	16	16	1
White-winged scoter	2	22	24	2
Grand total	109	1023	1132	100

*With or without an observed brood

Table 1. Adult ducks observed in the Kuzitrin River study area, Alaska, 1993.

Species	NUMBER OF YOUNG				NUMBER OF BROODS		
	GANG BROOD	ORPHAN BROOD	BROOD W/HEN	TOTAL YOUNG	BROODY HENS	BROODS*	TOTAL BROODS
Green-winged teal	0	1	51	52	3	9	13
American wigeon	30	0	44	74	2	11	19.5
Black scoter	0	0	30	30	0	7	7
Greater scaup	0	0	98	98	4	19	23
Mallard	0	1	6	7	0	2	2
Northern pintail	10	9	60	79	16	17	35.7
Northern shoveler	0	1	33	34	1	8	9
Oldsquaw	0	5	33	38	1	7	8
Unidentified duck	0	1	0	1	0	1	1
White-winged scoter	0	0	12	12	0	2	2
Grand total	40	18	367	425	27	81	120.2

* Does not include gang broods. In calculating total broods, number of young in gang broods was divided by the average historical brood size for that species.

Table 2. Species, type of broods, numbers of broods and young ducks observed in the Kuzitrin River study area, Alaska, 1993.

Guides for providing safe and efficient transportation. Also, we are grateful to our supervisor, H. Brownell, and the staff at the Alaska Interagency Fire Coordination Center for bearing with us through numerous contingency plans.

RESULTS

Waterfowl Production and Species Composition

Kuzitrin River Basin: 1993

The Kuzitrin River production area contains roughly 475 km² of wetland habitat, with 130 km² in the "other" stratum and 345 km² in the "poor" stratum. Fifteen plots were surveyed in 1993, in-

cluding 10 plots in the "other" strata and 5 "poor" plots. Observers recorded 1,132 adult ducks with 425 young in 94 broods. In addition, 26 "broody" hens were observed exhibiting distraction displays although their broods were hidden or not observed. Observations other than ducks included 144 white-fronted geese (*Anser albifrons*), 191 Canada geese (*Branta canadensis*) with 8 young, 26 tundra swans (*Cygnus columbianus*) with 13 young, 45 red-necked grebes (*Podiceps grisegena*) with 10 young, 2 yellow-billed loons (*Gavia adamsii*), 18 Pacific loons (*G. pacifica*) with 2 young, 3 red-throated loons (*G. stellata*), and 14 sandhill cranes (*Grus canadensis*).

American wigeons, green-winged teals (*Anas crecca*), northern shovelers (*A. clypeata*) and northern pintails were the predominant dabbling ducks seen in the Kuzitrin River study area (Table 1). Wigeon broods composed 25% and pintails 45% of

Location	Agency	Young/km ²	Broods/km ²	Years	Reference
Kuzitrin River	BLM/NPS	10.9	2.5	91-93	This report
McCarthy's Marsh	BLM	9.7	2.2	89-92	This report
Pah River	BLM	2.1	0.5	1989	Anderson and Robinson 1991a
Pah River	BLM	4.5	0.8	1993	This report
Koyukuk	FWS	11.0	2.2	1990	Hodges and Conant 1990
Yukon Delta	FWS	9.8	2.4	1990	Hodges and Conant 1990
Selawik	FWS	4.5	—	1989	Brubaker and Witmer 1989
Selawik	FWS	7.5	1.6	1990	Hodges and Conant 1990

Table 3. Comparison of density of young ducks obtained during July brood surveys on several federally-managed wetland areas in Alaska.




Figure 2. Species composition of duck broods in the Kuzitrin River wetlands, Alaska, 1993.

the dabbler broods observed (Table 2, Fig. 2). Greater scaups, oldsquaws (*Clangula hyemalis*), and black scoters (*Melanitta nigra*) were the most common diving ducks, with scaups accounting for over half (58%) of the diver broods.

The average brood size across age classes (considering only observed broods of known size) for northern pintails was 3.1 (n=14), for American wigeons was 4.3 (n=16), and for greater scaups was 4.8 (n=17). These values are slightly lower than statewide multi-year averages of 4.34 for northern pintails, 4.87 for American wigeons and 6.22 for greater scaups (Hodges and Witmer 1990).

DISCUSSION

Timing of the 1993 brood survey may have been a little early for detecting the maximum number of diver broods because most were still in the downy stage (Class I) with none fully fledged (Class III) (Fig. 3). This indicates that some diver broods may not have hatched yet, and may account for the large number of scaup broody hens observed compared to other years. Timing should have been nearly optimal to see the largest number of dabbler broods, as 42% of them were in the partially-fledged stage (Class II).

The four-year average number of broods/km² produced in McCarthys Marsh was 9.7, while the three-year average for the Kuzitrin was 10.9. The number of young produced varies considerably from year to year, so it is useful to have more than one year's data to use in estimating productivity of wetlands for waterfowl. Figure 4 shows the variation in production among the three survey years in the Kuzitrin study area. In Table 3, the density of young waterfowl produced on BLM-




Figure 3. Number of broods in the three major age classes for dabblers and divers in the Kuzitrin River study area, Alaska, 1993.

managed wetlands are compared with some of the interior Alaska wildlife refuges. These averages are based on the entire basin, or management area, which of course contains some areas unsuitable for waterfowl. It is interesting to note that in spite of their small size, the Seward Peninsula wetlands are roughly comparable on a per unit basis to the much larger refuges. The Seward Peninsula has one of the highest densities of waterfowl breeding pairs (determined by aerial surveys in early spring) in the state, after the Copper River Delta and the Yukon Flats (Conant and Hodges 1985). Of course, the overall production of these two areas is dwarfed by larger wetland areas of the state; for example, the Yukon-Kuskokwim delta which encompasses 29,556 mi² (76,845 km²).

We had little success in attempting to relate observations to habitat characteristics such as pH, water hardness and temperature. Dabbler broods did not show a correlation with any of these variables, although diver broods seemed to correlate positively with pH (Brown and Jandt 1992). Other




Figure 4. Estimated number of young ducks, geese and swans produced in Kuzitrin River valley, Alaska, from brood survey data obtained in 1991-1993.



R Jandt examines a fledgling duck from the Kuzitrin survey.

investigators have attempted to correlate brood use and production with physical characteristics, and for the most part, it has been very difficult to obtain consistent results. Bertram (1992) looked at a number of chemical and physical variables in relation to brood density and found waterbody area to be the only variable with a reasonably good correlation to brood density. However, Murphy *et al.* (1984) found that while larger ponds had greater species richness, the smaller ponds supported the greatest density of ducks, and felt that hydrologic connection to a creek system, with resultant higher levels of nutrients such as phosphate and nitrite, was the key factor in determining habitat use by ducks. Seppi (1993) also looked at a number of different variables, and concluded that shoreline length of ponds had the greatest positive association with brood use. Heglund (1992) failed to produce a good predictive model of habitat use by broods on the Yukon Flats, after intensive study. She attributed part of the difficulty to the tendency of ducks to select ponds based on factors other than the habitat quality, such as the presence of other ducks, called the "decoy effect." Thus, although research into the factors that influence habitat preference for waterfowl at the individual pond level continues, it has been difficult to develop hard and fast criteria for these factors. Biologists attempting to evaluate habitat quality for waterfowl may have to con-

tinue to rely on their general knowledge of limiting factors and realize that a variety of things can influence habitat selection under differing conditions, both spatially and temporally.

Conclusions and Management Implications

Both the McCarthys Marsh and Kuzitrin waterfowl production areas are contained in the Bendeleben Mountains Planning Subunit of the Northwest Management Framework Plan (BLM 1982). Although the McCarthys Marsh area was identified as an important waterfowl nesting area in this plan and in the Seward 1008 Study Decision Record (BLM 1983), the Kuzitrin basin was not described as such. In future plans, the value of both areas for waterfowl production should be recognized and considered in Environmental Assessments, Environmental Impact Statements, and the permitting of authorized uses, such as reindeer herding and mining. The North American Waterfowl Management Plan (FWS 1986) recommends that "... public land management agencies should ... regulate land uses to prevent the destruction or degradation of waterfowl habitats." The baseline production data obtained during 1989-1993 provides the foundation for monitoring adverse impacts to waterfowl nesting habitat due to future developments and land uses on the Seward Peninsula.

In the larger perspective of ecosystem management, waterfowl serve as one of the measurable indicators of the health of the pond riparian ecosystems in this area. Riparian areas are lands adjacent to creeks, streams, rivers, and lakes where vegetation is strongly influenced by the presence of water. Riparian areas are considered some of the most biologically diverse habitats, providing food and cover for a variety of wildlife and fish. Wetland habitats support the highest population densities of both breeding and non-breeding birds on the Seward Peninsula (Kessel 1989), including songbirds, shorebirds, waterfowl and raptors. Incidental observations have been recorded of a number of mammalian species in the wetland areas during the waterfowl surveys, including moose, muskox, porcupine, muskrat, otter and beaver. BLM is mandated to conserve and maintain the long-term productivity of riparian areas, and to restore damaged areas to a level of production comparable to that prior to the disturbance. In order to do this, it is essential to know pre-disturbance production values, which was one of the goals of these studies.

The BLM's Fish and Wildlife 2000 plan (BLM 1987) directs each District to identify and rank by importance key waterfowl habitat areas on public lands. In addition, a strategy plan has been developed which specifically instructs the Bureau to conduct habitat inventories and evaluations to determine condition and improvement potential of 170 waterfowl habitat management areas covering nearly 12 million acres by the year 2000 (BLM 1989). The production data from these studies on the Seward Peninsula has provided a quantitative

and biologically-based means for accomplishing this ranking. Three waterfowl production areas have been surveyed using the ground brood count methodology in the Kobuk District. Based on the total production of waterfowl, it appears that the Kuzitrin River basin is the most productive area, followed by the McCarthys Marsh area, and the Pah River Flats (Anderson and Robinson 1991a, Jandt and Morkill 1994 is the least important as far as overall production (Fig. 5).




Figure 5. Comparison of young ducks/km² produced on three areas where BLM brood surveys were conducted. Kuzitrin River and McCarthys Marsh are three- and four-year averages, respectively, while the value given for Pah River is the average of 1989 and 1993 results.

Literature Cited

- Anderson, R. and S. Robinson. 1991a. Ground brood counts to estimate waterfowl populations on two habitats in western Alaska. The Wildlife Society: Proc. of the Western and Northwestern Sections Meeting in Reno, NV, Feb. 22-24, 1990.
- . 1991b. Ground brood counts to estimate waterfowl populations in BLM's Kobuk District, Alaska. 1990 progress report. United States Department of the Interior, Bureau of Land Management, Fairbanks, AK. Open File Report 32. 10pp.
- Bertram, M. 1992. Wetland ecology and sightability correction for waterfowl productivity surveys on the Koyukuk and northern unit of the Innoko National Wildlife Refuges (draft). Final Rep. 92-3, Koyukuk/Nowitna NWR, USDI, Fish and Wildl. Serv., Galena, AK. 29pp.
- Brown, R.J. and R.R. Jandt. 1992. Ground brood counts to estimate waterfowl production on BLM's Kobuk District, Alaska, 1991 progress report. USDI, Bur. of Land Manage., Fairbanks, AK. Open File Report 43. 18pp.
- Brown, R.J., A.E. Morkill, and R.R. Jandt. 1993. Ground brood counts to estimate waterfowl populations on BLM's Kobuk District, Alaska, 1992 progress report. USDI, Bur. of Land Manage., Fairbanks, AK. Open File Report 47. 11pp.
- Brubaker, R.A., and D.E. Witmer. 1989. Review of waterfowl inventory surveys at Selawik National Wildlife Refuge, 1985-1988, and proposed changes for future surveys (draft). Progress Rep. 89-1, Selawik NWR, USDI, Fish and Wildl. Serv., Kotzebue, AK. 15pp.
- Bureau of Land Management. 1982. Northwest management framework plan. Unpubl. Rep. USDI, Bur. Land Manage., Washington, D.C. 30 pp.
- . 1983. Seward 1008 study, decision record. USDI, Bur. Land Manage., AK830407-02, Fairbanks, Alaska. 66 pp.
- . 1987. Fish and Wildlife 2000, a plan for the future. USDI, Bur. Land Manage., Washington, D.C. 30 pp.
- . 1989. Waterfowl habitat management on public lands, a strategy for the future. USDI, Bur. Land Manage., Washington, D.C. 43 pp.
- Conant, B. and J.I. Hodges. 1985. Alaska-Yukon waterfowl breeding population survey, May 15 to June 21, 1985. Unpubl. rep., U.S. Fish and Wildlife Service, Juneau. 23 pp.
- Doyle, T.J. 1989. The use of one versus two brood surveys to estimate duck brood production in Alaska. (unpublished draft) U.S. Fish and Wildl. Serv., Tetlin National Wildl. Refuge, Tok, AK. 13pp.
- Fish and Wildlife Service. 1986. North American Waterfowl Management Plan. USDI, Fish and Wildl. Serv., Washington, D.C. 19 pp.
- . (1991). Draft duck production survey standard operating procedures manual (unpublished document), Anchorage, Alaska. 34pp.
- Heglund, P.J. 1992. Patterns of wetland use among aquatic birds in the interior boreal forest region of Alaska. PhD Thesis, University of Missouri, Columbia, MO. 396pp.
- Hodges J. and D. Witmer. 1990. Alaska duck production survey data analysis software. Produced in association with USDI, FWS, Migratory Bird Management, Juneau, AK.
- Jandt, R.R. and A.E. Morkill. 1994, in press. Waterfowl production and effects of fire on wetlands in the Pah River Flats. USDI, Bur. of Land Manage., Fairbanks, AK. Open File Report 50. 9pp.
- Kessel, B. 1989. Birds of the Seward Peninsula, Alaska. Univ. Alaska Press, Fairbanks, AK. 330pp.
- Lensink, C.J. and D.V. Derksen. 1990. Evaluation of Alaskan wetlands for waterfowl. Pp 45-84 in Alaska: regional wetland functions, Proceedings of a workshop, Anchorage, Alaska, May 28-29, 1986. The Environmental Institute, Univ. of Massachusetts at Amherst.
- Murphy, S.M., B. Kessel and L.J. Vining. 1984. Waterfowl populations and limnologic characteristics of taiga ponds. J. Wildl. Manage. 48(4):1158-1163.

Seppi, B. E. 1993. Use of wetlands by waterfowl broods in relation to habitat quality in the lower Innoko River area, Alaska. Master's Thesis, University of Alaska, Fairbanks, AK. 65pp.

Swanson, J.D., M. Schuman, and P.C. Scorup. 1985. Range Survey of the Seward Peninsula Reindeer Ranges, Alaska. USDA Soil Conservation Service, Anchorage, Alaska: 77 pp. and 63 maps.

Tiner, R.W. Jr. 1984. Wetlands of the United States: current status and recent trends. U.S. Fish and Wildl. Serv., National Wetlands Inventory, USGPO, Washington, DC 430pp.

Sources for Scientific Nomenclature

Bellrose, F. C. 1976. Ducks, geese and swans of North America, 3rd edition. Stackpole Books, Harrisburg, PA. 540pp.

Hultén, E. 1968. Flora of Alaska and neighboring territories. Stanford University Press, Stanford, CA. 1008pp.

BLM-Alaska Open File Report Series

1. Upper Colville River channel depth survey. Jack Mellor, June 1983. 16pp.
2. Archaeological investigations of the Delong mountains, northwest Alaska. Howard Smith, June 1983. 26pp.
3. Archaeological reconnaissance in the Central Arctic Management Area. Howard Smith, June 1983. 19pp.
4. Results of the 1981 bald eagle nest survey. Laurence Byrne, David Daum, Michael Small and Julie Henderson, June 1983. 20pp.
5. Results of the 1982 bald eagle nest survey. Laurence Byrne, Julie Henderson, Michael Small, June 1983. 12pp.
6. Results of the 1982 trumpeter swan survey. Laurence Byrne, Julie Henderson, Michael Small, June 1983. 7pp.
7. The 1982 peregrine falcon/raptor survey in the Central Arctic. Ted Swem, Robert Dittrock and James Silva, June 1983. 40pp.
8. Buckland reindeer/caribou conflict study final report. Layne Adams and Bruce Connery, June 1983. 60pp.
9. Denali Mine on Valdez Creek, southcentral Alaska. Beth Walton and Cheryl McCaffrey, November 1984. 9pp.
10. Effects of fire on a dwarf shrub-sedge tussock community. K. Van Waggoner and Marianne See, March 1985. 16pp.
11. The 1983 peregrine falcon/raptor survey. James Silva, April 1985. 23pp.
12. Status of the Ray Mountain caribou herd. Scott Robinson, June 1985. 11pp.
13. Landsat enhancement procedures and key, central Yukon area. Melanie Miller, June 1985. 10pp.
14. Fire occurrence in the Central Yukon Planning Area, 1956-1982. Melanie Miller, November 1985. 40pp.
15. Fire occurrence in the Northwest Planning Area, 1956-1982. Melanie Miller, November 1985. 37pp.
16. The 1984 fire season, Northwest Planning Area. Melanie Miller and Scott Robinson, November 1985. 21pp.
17. Wildlife of the Titna, Tozitna and Kateel water-sheds. Scott Robinson, December 1985. 28pp.
18. Wildlife of the Koyuk watershed. Scott Robinson, December 1985. 14pp.
19. Beaver Creek National Wild River cultural resources inventory. Susan Will, June 1986. 48pp.
20. Wildlife of the Squirrel River, Alaska. Scott Robinson, September 1987. 20pp.
21. Movements and distribution of the western arctic caribou herd across Buckland Valley and Nulato Hills, winter 1986-87. Scott Robinson and Larry Field, September 1987. 7pp.
22. Historical fire data: BLM-Alaska 1959-1985. Russell E. Hanson, September 1987. 21pp.
23. Movement and distribution of the western arctic caribou herd across Buckland Valley and Nulato Hills, winter of 1987-1988. Scott Robinson, December 1988. 11pp.
24. Fisheries investigations in the Beaver Creek drainage, 1988. Louis Carufel, June 1989. 16pp.
25. Bird communities of recently burned and unburned forest and scrub habitats in interior Alaska. Michael T. Hinkes and Kate Engles, June 1989. 33pp.
26. Movement and distribution of radio-collared caribou across Selawik Valley, Buckland Valley and Nulato Hills, winter 1988-89. Scott Robinson and Michael Spindler, November 1989. 25pp.
27. Ecology of moose in the White Mountains National Recreation Area, Alaska, 1985-1988. Winston Hobgood and Bruce M. Durtsche, August 1990. 16pp.
28. Three years of natural revegetation on the 1977 Bear Creek burn in interior Alaska. Russell Hanson, September 1990. 42pp.
29. Distribution, movements and seasonal use areas of caribou in the White Mountains Recreational Area, Alaska. Winston Hobgood and Bruce M. Durtsche, January 1991. 9pp.
30. Dall sheep in the White Mountains. Bruce M. Durtsche, Winston Hobgood and Jan Burris, October 1990. 10pp.
31. Fisheries investigations in the Beaver Creek drainage, White Mountains National Recreation Area, Alaska, Louis Carufel, January 1991. 19pp.
32. Ground brood counts to estimate waterfowl populations in BLM's Kobuk District, Alaska: 1990 progress report. Randi R. Anderson and Scott R. Robinson, March 1991. 10pp.
33. Movements and distribution of radio-collared caribou in the Buckland Valley and Nulato Hills, winter of 1989-1990. Scott R. Robinson and Mary Leykom, March 1991. 10pp.
34. Oil and gas development and Alaska's north slope: past results and future prospects. Arthur C. Banet, Jr., March 1991. 42pp.
35. Leasable mineral resource assessment of the southcentral planning area, Alaska. William R. Diel, December 1991. 68pp.
36. Petrographic survey and appraisal of reservoir quality and potential, National Petroleum Reserve-Alaska. Thomas C. Mowatt and Joseph A. Dygas, September 1991. 22pp.
37. Platinum and palladium in mafic-ultramafic igneous rocks, northwestern Alaska. Thomas C. Mowatt, November 1991. 21pp.
38. The Red Dog Deposit, northwestern Alaska: discovery, delineation, and development implications. Thomas C. Mowatt, Joseph A. Dygas and Christopher Gibson, November 1991. 15pp.
39. Mineral resources of western arctic Alaska. Thomas C. Mowatt, Joseph A. Dygas, Christopher Gibson and Aden L. Seidlitz, November 1991. 27pp.
40. Diagenetic relationships and reservoir quality implications in Brookian clastic sequences, National Petroleum Reserve, Alaska. Thomas C. Mowatt and June C. Mowatt, December 1991. 40 pp.
41. Oil and gas assessment of the Utukok Special Management Area, National Petroleum Reserve in Alaska. Robert J. Bascle and Richard L. Foland, March 1992. 28pp.
42. Petrographic analyses of selected horizons, Aurora 089 No. 1 OCS-Y-0943 well, offshore northeast Alaska (Part 1). Thomas C. Mowatt, Arthur C. Banet, Jr. and John W. Reeder, May 1992. 48pp.
43. Ground Brood counts to estimate waterfowl populations in BLM's Kobuk District, Alaska: 1991 progress report. R.J. Brown and R. R. Jandt, December 1992. 11pp.
44. Innoko National Wildlife Refuge oil and gas resource assessment. Robert Bascle, Aden Seidlitz and James Borkoski. July 1993.
45. Kodiak National Wildlife Refuge oil and gas resource assessment. Robert Bascle, Aden Seidlitz and James Borkoski. July 1993.
46. Teshekpuk Lake Special Management Area oil and gas resource Assessment, National Petroleum Reserve-Alaska. Robert Bascle. July 1993.
47. Ground Brood counts to estimate waterfowl populations in BLM's Kobuk District, Alaska: 1992 progress report. R.J. Brown, A.E. Morkill and R R Jandt July 1993. 8pp.
48. Moose Population Parameters and Distribution in the Squirrel River, Alaska, November 1992. A. E. Morkill and J. Dau.
49. Ruby-Poorman Mining District, Ruby Quadrangle, Alaska, November 1993, Rosalie E. L'Ecuyer. 41pp
50. Waterfowl production and effects of fire on wetlands in the Pah River Flats. April 1994, R.R. Jandt and A.E. Morkill. 9pp
51. Aquatic Resources of Salmon Fork Black River, Alaska, Jon Kostohrys, Brian Liubinski and Nathan Collin, April 1994. 18pp.

The Bureau of Land Management is responsible for the stewardship of our public lands. It is committed to manage, protect, and improve these lands in a manner to serve the needs of the American people for all times. Management is based on the principles of multiple use and sustained yield of our nation's resources within a framework of environmental responsibility and scientific technology. These resources include recreation, range, timber, minerals, watershed, fish and wildlife, wilderness, air, scenic, scientific and cultural values.
