

P

**REVIEW AND CRITICAL SYNTHESIS
OF THE LITERATURE ON RECOVERY OF ECOSYSTEMS FOLLOWING
MAN-INDUCED AND NATURAL-PHENOMENA-RELATED
DISTURBANCES: HARBOR SEALS AND KILLER WHALES**

by

Brent S. Stewart, Ph.D.
Staff Scientist

Pamela K. Yochem, D.V.M.
Staff Scientist

Joseph R. Jehl, Jr., Ph.D.
Staff Scientist, Research Director

Hubbs-Sea World Research Institute (HSWRI)
1700 South Shores Road
San Diego, CA 92109

Principal Investigator: Joseph R. Jehl, Jr., Ph.D.

FINAL REPORT

to

Restoration Planning Work Group
EXXON Valdez Oil Spill Office
645 "G" Street
Anchorage, Alaska 99501

30 June 1992

EXECUTIVE SUMMARY

Populations of marine mammals have suffered large reductions, sometimes to near extermination, by aboriginal and commercial harvests, incidental or indiscriminate killing, and epizootics during the past two centuries. After killing ended, many populations increased at annual rates varying from 7-21% in pinnipeds and 2-12% in cetaceans. The causes for recent steady declines, following population recoveries, of northern fur seals, northern sea lions, and harbor seals in the Bering Sea, Aleutian Islands, and western Gulf of Alaska, and of southern elephant seals in most of the southern ocean, remain unexplained.

Epizootics.--Recent epizootics killed over 18,000 seals (mostly harbor seals) in Europe, and an estimated several thousand at Lake Baikal; population responses following those reductions are undocumented. Historical occurrences of epizootics and the prevalence of antibodies to various viruses in current seal populations suggest that seals that survive these challenges provide nuclei for population recovery.

Climate.-- Seal and sea lion populations in the Pacific were reduced by the 1982-83 El Niño Southern Oscillation (ENSO) event. Recent studies have indicated only temporary demographic consequences. Historical, large-scale fluctuations in ocean conditions related to ENSOs may have influenced population changes in Antarctic pinnipeds, though not to the extent of affecting population persistence.

Overall long-term population data demonstrate the potential of pinnipeds and cetaceans to sustain high rates of growth following population reduction, even to very low abundance, so long as breeding and foraging habitats are not degraded.

Pollution.-- Fouling of pinnipeds and cetaceans by oil has evidently had insignificant effects on populations; substantial mortality has never been observed, even following catastrophic spills. The

effects of oiling depended on whether oil coated the body surface, was ingested, or aromatic hydrocarbons were inhaled. Most reports have been based on casual observations; results of systematic experiments have often been ambiguous.

Vulnerability of cetaceans is highest for species with small ranges, coastal/ice-dwelling/riverine habitats, limited diets, poor behavioral flexibility, and small populations. Species with large ranges, oceanic distribution, diverse prey, adaptable behavior, and large populations are least vulnerable. For pinnipeds, stressed or nursing animals, and recently-weaned pups are potentially vulnerable. But marine mammals are long-lived and even the loss of an entire cohort would have insignificant long-term demographic effects.

Prolonged inhalation of hydrocarbon vapors appears to pose the greatest risk to the viability of individuals. Animals with parasitic lung disease, which is relatively common in pinnipeds, would be especially vulnerable to respiratory challenges. Yet, for most pinnipeds, particularly in northern habitats, it is unlikely that petroleum vapors could become sufficiently concentrated to represent a threat.

Contaminants in food.--Pinnipeds are unlikely to directly ingest hydrocarbons, and their prey seem unlikely to accumulate residues. Thus, toxicity is not expected to be a significant health risk, except possibly in bearded seals, walruses, or harbor seals foraging in heavily contaminated benthic environments. Of greater significance is the potential direct effects of fouling on benthic communities, which may be transmitted to other parts of the food chain; for example, a reduction in octopus abundance might depress the recovery of harbor seals.

Killer whales consume a wide variety of prey, including fish, birds and mammals. They are unlikely to ingest toxic hydrocarbons, unless they prey on species that have accumulated residues.

Future research.--Because there are few data on pre-EVOS abundance of harbor seals and killer whales in the EVOS area, it is impossible to use simple counts of animals to decide whether a

population has recovered. For harbor seals, it may be possible to use early post-spill data on abundance, distribution, and pup production as a reference point for future assessments. However, other recovery criteria (e.g., habitat occupation; an arbitrarily-established, desired local population size; physical or physiological condition of individuals) need to be developed. Evaluation of the recovery process will require long-term monitoring of population abundance and seasonal distribution. Future research should document the movement patterns of harbor seals and killer whales in Prince William sound and their seasonal use of habitats in the EVOS area using satellite-linked or conventional VHF telemetry and intensive photo-identification studies (primarily killer whales). Surveys should cover a larger area and be conducted at all seasons of the year; this is especially needed for killer whales. Monitoring should be conducted at several year intervals--and at a level to provide statistically valid results-- to permit long-term, cost-effective evaluation.

Table of Contents

	Page
1.0. Introduction	6
1.1. Background	6
1.2. Objectives	6
2.0. Technical Approach	7
2.1. Information retrieved and sources of data	7
3.0. Review of available information of recovery of marine mammal populations from anthropogenic and natural disturbances	7
3.1. Rate, duration, and recovery following disturbance	7
3.1.1. Pinnipeds	7
A. Harbor seals	7
B. Other pinnipeds	9
3.1.2. Cetaceans	10
A. Killer whales	10
B. Other cetaceans	12
3.2. Dependency of recovery on habitat protection, changes in management practices and other restoration approaches	13
3.3. Indicators of recovery that are the most practical and cost-effective to measure	14
3.4. Approaches and strategies for determining how indicators of recovery are best monitored and tested to determine when recovery has occurred	15
4.0. Acknowledgements	16
5.0. References	16
6.0. Tables	28
7.0. Appendix 1. Annotated Bibliography.	35

1.0. Introduction

1.1. Background

On 24 March 1989, around 11 million gallons of North Slope crude oil spilled into Prince William Sound from the grounded oil tanker EXXON Valdez. About 60% of the oil was not recovered and drifted or was blown southwest along the Kenai Peninsula toward Shelikof Strait, resulting in the fouling of over 1200 miles of mainland and island coastline and an unknown area of ocean bottom. Resident populations of harbor seals and killer whales may have been affected during the spill by inhalation of volatile, short-chain hydrocarbons, ingestion of oil, immediate destruction of prey resources and long-term food chain contamination. Evidently, substantial numbers of harbor seals became oiled in the EXXON Valdez oil spill (EVOS) area. Some were likely exposed to toxic aromatic hydrocarbons in areas very near the spill source. Killer whale numbers have declined in the EVOS area since 1989; known (photo-identified) whales have been reported missing from well-studied killer whale pods in nearby areas of Prince William Sound. Additional studies have been conducted on the distribution and abundance of killer whales in Prince William sound to determine the relationship of the EVOS to changes in whale abundance but results of those studies have not yet been published. When abstracts or summaries were included in source documents we quoted them directly in our annotated bibliography. When no abstracts or summaries were present in the documents reviewed we constructed new abstracts.

1.2. Objectives

Here we summarize, in the form of an annotated bibliography, published information on the population effects of oil spills on harbor seals and other relevant pinnipeds and killer whales and other relevant cetaceans throughout their ranges. We also summarize demographic information on the responses of pinniped and cetacean populations to other anthropogenic and natural disturbances and on rates and patterns of population recovery. We use this data base as a guide to understanding

population growth rates of harbor seals and killer whales, particularly in the Gulf of Alaska. We include comparative data for cetaceans and pinnipeds and summarize their population responses to anthropogenic (especially oil spills) and natural disturbances.

2.0. Technical Approach

2.1. Information retrieval and sources of data

Computerized literature searches were made through DIALOG (accessing BIOSIS, AQUATIC SCIENCES AND FISHERIES ABSTRACTS and OCEANIC ABSTRACTS) and MELVYL (accessing all University of California book and periodical holdings). Direct searches were made of current scientific literature at libraries at Scripps Institute of Oceanography, San Diego State University, UCLA, and UC Davis. Finally, our personal and Institute libraries were the most productive sources of information on pinniped and cetacean biology. The literature recoveries from these initial searches were used in a hierarchical way to provide additional key words for additional searches and additional reference lists of previously published literature.

3.0. Review of available information of recovery of marine mammal populations from anthropogenic and natural disturbances

3.1. Rate, duration, and degree of recovery following disturbance.

3.1.1. Pinnipeds

A. Harbor seals

Harbor seals are relatively abundant residents of Prince William Sound and the Gulf of Alaska. Little is known of their daily and seasonal hauling patterns, absolute abundance, movements, life history parameters and diet within the EVOS area, but detailed information does exist for local populations elsewhere. Daily terrestrial abundance of harbor seals is greatest at mid-day or during

daytime low tides and seasonal terrestrial abundance is greatest during the molt in spring or summer and least in winter; breeding occurs from late winter through early spring or summer depending on latitude (e.g., Schneider and Payne 1983, Stewart 1984, Terhune and Almon 1983, Thompson et al. 1989, Yochem 1987). Terrestrial abundance at a large haulout area on Tugidak Island near the EVOS area declined substantially (about 85%) from 1976 through 1988 for unknown reasons, although large numbers of pups were harvested annually from 1964 through 1972 (Pitcher 1990). The trend in Prince William Sound was not documented. The decline in abundance at Tugidak Island is sharp contrast to the steady increases in harbor seal populations in most other parts of the species' range during the past several decades (e.g., Harvey et al. 1990, Heide-Jorgensen and Harkonen 1988, Olesiuk et al. 1990a, Stewart et al. 1988, Stewart et al. 1992).

Seasonal site-fidelity and short- and long-distance movements of harbor seals have been documented in some areas (e.g., Brown and Mate 1983, Pitcher and MacAllister 1981, Yochem et al. 1987) as have seasonal, sexual, and age-class segregation (e.g., Allen et al. 1988, Godsell 1988, Kovacs et al. 1990, Thompson et al. 1990). No comparable data are available for the EVOS-Prince William Sound area. The diet of harbor seals is relatively broad with benthic and epibenthic species of cephalopods and fish generally predominating (e.g., Brown and Mate 1983, Harkonen 1987, Olesiuk et al. 1990b, Pierce et al. 1991, Pitcher 1980a, 1980b, Thompson et al. 1991).

Harbor seal populations have been increasing in most areas where they have been studied in recent years where commercial or subsistence harvesting is low or absent (e.g., Harvey et al. 1990, Heide-Jorgensen and Harkonen 1988, Olesiuk et al. 1990a, Stewart et al. 1988, Stewart et al. 1992). Documented rates of population increase are relatively high, around 5-22% per year (Table 1). Most of the increases have occurred after bountied and indiscriminate killing and harvesting were outlawed. Degree of recovery is generally impossible to judge as pre-exploitation abundances are unknown. In a few other areas, however, populations have declined or fluctuated at low levels. In

some cases chronic pollution is believed to be responsible for reproductive failures and depressed populations of harbor and other seals (Helle et al. 1976, Reijnders 1978, Zakharov and Yablokov 1990). There has also been a persistent decline in the western Gulf of Alaska around Tugidak Island (Pitcher 1990), and perhaps in Prince William Sound. Causal factors may include 1) degradation of habitat (reduction of prey resources, natural environmental changes, virulent pathogens, etc.) or 2) substantial undocumented mortality associated with commercial fishing operations or native subsistence harvest.

In 1988 an epizootic killed over 18,000 seals, mostly harbor seals in European waters. In Swedish and Danish waters of the Kattegat and Skagerak more than 5300 harbor seals died; the population had previously numbered about 9100 and had increased from 1978-1988 at more than 12% per year (Dietz et al. 1989, Heide-Jorgensen and Harkonen 1988). An epizootic in the Soviet Union's Lake Baikal in 1987 killed several thousand Baikal seals (Grachev et al. 1989). Disease outbreaks in other species in the western Atlantic, Pacific, and Antarctic were less severe (Borst et al. 1986, Geraci et al. 1982, Hinshaw et al. 1984, Laws and Taylor 1957, Smith et al. 1974, Vedros et al. 1971), but there is no evidence of long-term demographic consequences in those areas. There are no published data on population responses following the 1987 and 1988 disease outbreaks. No long-term population effects of oil pollution on harbor seals or any other pinnipeds have been documented; documentation of chronic effects of oil pollution on individuals has been equivocal (Geraci and St. Aubin 1987, St. Aubin 1990).

B. Other pinnipeds

Throughout the world, populations of many pinniped species have been increasing at relatively high rates. Northern elephant seals (Mirounga angustirostris), for example, have been increasing at about 14% per year for nearly one hundred years (Stewart 1992). The duration of increases for other species varies according to the time at which commercial harvesting ended; pre-exploitation

abundance of any of those species is unknown. Following sustained population growth in the early 1900s, northern fur seals (Callorhinus ursinus) in the Bering Sea declined substantially, for unknown reasons, from the 1960s through the late 1980s. Northern sea lions have decreased steadily during the past two decades throughout the Aleutian Islands and western Gulf of Alaska, whereas their populations in the eastern Gulf of Alaska, Canada and Oregon and Washington have remained relatively stable or increased slightly. Southern elephant seals have also been declining in most areas of the Southern Ocean in recent years, following a period of recovery from commercial harvesting (Laws 1992).

Low reproductive success and high pup mortality among several species of pinnipeds in the Pacific in 1982 or 1983 coincided with the 1982/83 El Niño Southern Oscillation (ENSO; De Long and Antonelis, 1991; DeLong et al., 1991; Guerra C. and Portflitt K., 1991; Majluf, 1992; Stewart and Yochem, 1991; Trillmich and Dellinger, 1992). These results were evidently related to reduction, redistribution or disappearance of prey populations near rookeries. There is little evidence of substantial adult mortality nor in long-term demographic effects from that intense oceanographic disturbance, except perhaps at the Galapagos Islands.

3.1.2. Cetaceans

A. Killer Whales

Killer whales are widely distributed in the world's oceans (Dahlheim 1981). They occur in deep pelagic waters and in coastal areas, along ice edges, and in pack ice as well as in the tropics (Mitchell and Reeves 1988). Local movements and distribution appear to be largely dictated by distribution and availability of prey (Dahlheim 1981, Braham and Dahlheim 1982, Heimlich-Boran 1988). A partial list of prey items by geographic area was presented by Anon. (1982). Killer whales consume a variety of marine vertebrates and invertebrates, including fish, cephalopods and mammals. There are differences in food habits between sympatric populations in some areas: resident pods in

British Columbia and Washington consume mainly fish (especially salmon) whereas transients feed mostly on marine mammals, especially harbor seals (Heimlich-Boran 1988).

Using photo-identification, Olesiuk et al. (1990c) calculated a number of population parameters for killer whales off British Columbia and Washington. They reported an annual rate of increase of 2.92%; the percentage of mature females pregnant varied from 2.7-4.1%. Neonate mortality was 43%. The mean life expectancy was 50.2 years for females and 29.2 years for males, with predicted maximum life spans of 80-90 and 50-60 years, respectively. From computer simulations the authors predicted that the killer whales in this region could sustain a maximum non-selective harvest of 2.84%. They further predicted that a stationary population at carrying capacity would comprise 37% juveniles, 20% mature males, 14% reproductive females, and 29% post-reproductive females. Leatherwood et al. (1990) reported the following age structure among Prince William Sound killer whales: 22.41% adult males, 9.48% adult females (defined as females in close association with a calf), 3.9% calves, and 64.22% immatures and others (this group includes immature animals, adult females not associated with calves, and recently matured males that lack a prominent dorsal fin).

From 1962-1977, a total of 66 killer whales was removed from a few pods in British Columbia and Washington by a live-capture fishery to supply captive whales for oceanaria. Since then, the cropped pods have had higher birth rates (4.56%), lower mortality rates (bulls, 2.5%; cows 0.46%; juveniles, 1.99%) and have increased in number faster (pod growth rate = 3.01%) than uncropped pods (birth rate = 3.15%, pod growth rate = 1.67%) in the same areas (Bigg 1982, Balcomb et al. 1982).

Leatherwood et al. (1990) documented a minimum of 221 killer whales in Prince William sound in 1987 from photographs of their dorsal fins and color patterns. Those whales belonged to nine "resident" and eight "transient" pods, as defined by Bigg (1982). Recent DNA research has supported the hypothesis that these pods are genetically distinct (Hoelzel and Dover 1991). The

combined mortality rate for all ages and both sexes from 1984-86 was 1.9% in three pods, but 7.4% in another (AB pod). The latter pod has been interfering with the blackcod (Anoplopoma fimbria) longline fishery since 1985 and bullet wounds have been observed on some of its members.

Leatherwood et al. (1990) did not report an annual rate of population increase for killer whales but noted that 9 calves were born in 1986 and 1987. In British Columbia and Washington, where killer whales have been studied using the same techniques, annual rates of increase ranged from 1.67 to 3.01% (Balcomb et al. 1982, Bigg 1982, Olesiuk et al. 1990c) and annual mortality rates from 0.7% (adult females) to 2.81% (adult males).

Geraci and St. Aubin (1987) and Geraci (1990) reviewed the effects of oil on cetaceans and included a table of reports of cetaceans associated with oil. Only one incident involving killer whales was found, in which two whales (one sick, one dead) were observed in association with diesel fuel (quantity unknown) off the Alaskan peninsula.

Aside from occasional reports of mass die-offs or strandings (e.g., Oritsland and Christensen 1982, Christensen 1990), the most significant cause of killer whale mortality has been commercial whaling. For example, Christensen (1982) reported that 2399 killer whales were killed in Norwegian coastal waters between 1938 and 1980. This represented a mean annual catch of 57 whales. Christensen (1982) noted, however, that the length (and therefore presumably the age structure) of the catch did not change during that period. Although no population growth rates are available, the percentage of pregnant females ranged from 12-32.8%, as determined by catch data (Anon. 1982). Similar percentages of pregnant females have been calculated from Antarctic catch data (12.72-18.97%). Off Marion Island in the southern Indian Ocean, 36.3% of adult females observed had calves (Condy et al. 1978), although some may not have been young-of-the-year.

B. Other cetaceans

Population growth rates and related parameters have been measured in other species that have

experienced significant human disturbance, usually in the form of harvesting (either as target species, right whales for example; or incidental catch, dolphins in the Eastern Tropical Pacific (ETP) for example).

The relatively low birth and death rates of killer whales are mirrored by another large odontocete, the sperm whale. Females produce a calf only every 3-6 years, and the natural mortality rate is less than 1% per year (Gosho et al. 1984). A decrease in calving interval (from 6 to 5.2 years) has been documented in an exploited population off Durban, South Africa (Best et al. 1984).

Reilly and Barlow (1986) estimated that dolphins could approach a population growth rate of 9%, but they thought that rate was unlikely to be attained under most conditions. Barlow (1985) reported the following differences among a more intensively fished dolphin population in the ETP: smaller percent pregnant, larger percent lactating, and larger percent immature than less-exploited dolphin populations in the ETP. The highest rates of annual population increase in baleen whales are reported for southern right whales and range from 7.6% (Payne et al. 1990) to 11.7% (population as a whole) or 13% (cow-calf pairs) (Bannister 1990) (Table 2). Gray whales have increased at annual rates of about 4% or greater since the early 1900s, despite a harvest rate of about 1.2% per year (Reilly et al. 1983) and Bowhead whales, which also are harvested for subsistence purposes, increased at an annual rate of around 3% from 1978 through 1988 (Zeh et al. 1991). Moderate rates of increase for other whales were summarized by Best (1990). Reproductive rates have been reported for humpback whales; the mean calving rate (calves per mature female per year) is about 0.4 (Perry et al. 1990, Clapham and Mayo 1990). The mean calving interval for gray whales is 2.11 years and the birth rate (ratio of calves to adults) is about 0.14 (Reilly 1984).

3.2. Dependency of recovery on habitat protection, changes in management practices, and other restoration approaches.

In virtually all cases, recent population recoveries of pinnipeds and cetaceans has been

due to the termination of commercial harvesting or indiscriminate or incidental killing. Many species were reduced to very low levels during the harvesting periods and several were believed to have been exterminated. Presumably, foraging and breeding habitats were not degraded by the harvesting. The presence of abundant prey resources and good quality breeding habitat are probably the most important factors that allow sustained population growth, as soon as commercial exploitation ceases.

Quick resumption of population growth of eastern North Pacific pinnipeds (i.e., California sea lions, northern elephant seals, harbor seals) following the 1982/83 ENSO was evidently due to rapid recovery of prey resources; i.e., the degradation of habitat and reduction of carrying capacity was short-lived (Stewart, 1992; Stewart et al. 1992; Stewart and Yochem, unpubl; R. L. DeLong, pers. comm.). A consensus of recent literature on population modelling is the recognition that rapid and large population changes can occur with only moderate increases in adult mortality; population growth is less sensitive to changes in juvenile survival. Thus, if adult mortality is high during, after, or both, a population reduction (e.g., because of subsistence harvests or undocumented killing), the recovery may be delayed or a continued decline may also occur. Changes in harbor seal management practices (i.e., documenting all subsistence takes with respect to age and sex composition of harvest in and near the EVOS area, reducing and strictly regulating subsistence harvests) would probably be the most effective means of stimulating rapid population recovery.

3.3. Indicators of recovery that are the most practical and cost effective to measure

There are few data available on the pre-EVOS status of killer whales and harbor seals in the affected EVOS area. For harbor seals, relative abundance and distribution and relative annual production of young would be indicators that could be directly compared with early post-spill data and with similar data from comprehensive data bases from other regions. However, collection of data on haulout patterns, movements, and diet would be useful for determining whether changes in local abundance of seals might be due to lowered reproduction among resident seals or simply to

movements of surviving seals to more favorable breeding or foraging habitats or to changes in haulout patterns related to dietary shifts.

Photo-identification studies (perhaps in combination with VHF or satellite telemetry) of killer whales should be continued to document relative pod sizes and composition, home range (of residents) and large-scale movements (of residents and transients), and reproductive rates. Those studies should be made over a broader area in Prince William Sound and during more seasons than previous studies. Monitoring in alternate years or every three years would probably be most efficient as the studies should be continued for 15 years or more to provide any useful information on population trends.

Bigg (1982) and Balcomb et al. (1982) measured birth rates, mortality rates and net population change in cropped versus uncropped pods with relatively good success.

3.4. Approaches and strategies for determining how indicators of recovery are best monitored and tested to determine when recovery has occurred

First, "recovery" must be defined for killer whales and harbor seals because there are few or no pre-EVOS data to compare with post-EVOS data. One guideline for evaluating "recovery" might be whether or not animals have regained the ability to maintain self-replicating or growing populations. To determine whether or not and when these abilities have been regained would require long-term studies of abundance coupled with an assessment of seasonal movements of animals in and out of the area and of the magnitude of immigration and emigration. The case of harbor seals in Prince William Sound is further complicated by a probable declining trend prior to the EVOS (cf. Pitcher 1990). To evaluate the health or demographic trends of local Prince William Sound populations of these species, a combination of approaches would be most productive and should be conducted every two or three years. Combinations of satellite and VHF telemetry, aerial and boat surveys, ground observations, dietary studies (for harbor seals) and photo-identification studies (for killer whales) should be used but should be planned carefully to give statistically valid results and to

avoid the possibility of the studies themselves (i.e., disturbance) complicating interpretations of movements, reproduction and trends in abundance.

These studies need to be integrated with research by other groups on benthic, epibenthic, and mid-water column fish and invertebrate communities to determine the effects of their recoveries on local killer whale and harbor seal distribution.

4.0. Acknowledgements

We thank J. Francine, R. Pytesky, S. Bond, F. T. Awbrey, and K. Miller-McLune for assisting with literature collection, summary and preparation of the annotated bibliographic data base and S. Senner, J. Strand, D. Costa, and D. Sinniff for their reviews of the draft report.

5.0. References

- Allen, S. G., C. A. Ribic, and J. E. Kjelmyr. 1988. Herd segregation in harbor seals at Point Reyes, California. *Calif. Fish Game* 74:55-59.
- Allen, S. G., H. R. Huber, C. A. Ribic, and D. G. Ainley (1989). Population dynamics of harbor seals in the Gulf of the Farallones, California. *Calif. Fish Game* 75:224-232.
- Anon. 1982. Report of the workshop on identity, structure and vital rates of killer whale populations, Cambridge, England, June 23-25, 1981. *Rep. int. Whal. Commn* 32:617-631.
- Balcomb, K. C. III, J. R. Boran, S. L. Heimlich. 1982. Killer whales in Greater Puget Sound. *Rep. int. Whal. Commn* 32:681-685.
- Bannister, J. L. 1990. Southern right whales off Western Australia. *Rep. int Whal. Commn Spec. Iss.* 12:279-288.
- Barlow, J. 1985. Variability, trends, and biases in reproductive rates of spotted dolphins,

Stenella attenuata. U. S. Fish. Bull. 83:657-669.

Best, P. B. 1990. Recovery rates in whale stocks that have been protected from commercial whaling for at least 20 years. Appendix 6 of Annex F, Report of the International Whaling Commission 40:129-130.

Best, P. B., P. A. S. Canham, N. Macleod. 1984. Patterns of reproduction in sperm whales, Physeter macrocephalus. Rep. int. Whal. Commn Spec. Iss. 6:551-79.

Bester, M. N. 1980. Population increase in the Amsterdam Island fur seal, Arctocephalus tropicalis, at Gough Island. S. Afr. J. Zool. 15:229-234.

Bigg, M. 1982. An assessment of killer whale (Orcinus orca) stocks off Vancouver Island, British Columbia. Rep. int. Whal. Commn 323:655-666.

Borst, G. H. A., H. C. Walvoort, P. J. H. Reijnders, J. S. van der Kamp, and A. D. M. E. Osterhaus. 1986. An outbreak of a herpesvirus infection in harbor seals (Phoca vitulina). Journal of Wildlife Diseases 15:593-596.

Braham, H. W. 1984. Review of reproduction in the white whale Delphinapterus leucas, narwhal, Monodon monoceros, and Irrawaddy dolphin, Oracella brevirostris, with comments on stock assessment. Rep. int. Whal. Commn. Spec. Iss. 6:81-89.

Braham, H. W. and M. E. Dahlheim. 1982. Killer whales in Alaska documented in the Platforms of Opportunity Program. Rep. int. Whal. Commn. 32:643-646.

Brown, R. F. and B. R. Mate. 1983. Abundance, movements, and feeding habits of harbor seals, Phoca vitulina, at Netarts and Tillamook Bay, Oregon. U. S. Fish. Bull. 81:291-302.

Butterworth, D. S., J. H. M. David, L. H. McQuaid, and S. S. Xulu. 1987. Modeling the population dynamics of the South African fur seal Arctocephalus pusillus pusillus. NOAA Tech. Rep. NMFS 51:141-164.

- Calambokidis, J., B. L. Taylor, S. D. Carter, G. H. Steiger, P. K. Dawson and L. D. Antrim.
1987. Distribution and haul-out behavior of harbor seals in Glacier Bay, Alaska.
Can. J. Zool. 65:1391-1396.
- Chapman, D. G. 1981. Evaluation of marine mammal population models. In: C. W. Fowler
and T. D. Smith (eds.). Dynamics of large mammal populations. John Wiley and
Sons, New York.
- Christensen, I. 1982. Killer whales in Norwegian coastal waters. Rep. int. Whal. Commn
32:633-641.
- Christensen, I. 1990. A note on recent strandings of sperm whales (Physeter macrocephalus)
and other cetaceans in Norwegian waters. Rep. int. Whal. Commn 40:513-515.
- Clapham, P. J., and C. A. Mayo. 1990. Reproduction of humpback whales (Megaptera
novaeangliae) observed in the Gulf of Maine. Rep. int. Whal. Commn Spec. Iss.
12:171-175.
- Condy, P. R. 1978. Distribution, abundance, and annual cycle of fur seals (Arctocephalus
spp.) on the Prince Edward Islands. S. Afr. J. Wildl. Res. 8:159-168.
- Condy, P. R., R. J. van Aarde, and M. N. Bester. 1978. The seasonal occurrence and
behaviour of killer whales Orcinus orca, at Marion Island. J. Zool., Lond. 184:449-
464.
- Cooper, C. F. and B. S. Stewart. 1983. Demography of northern elephant seals, 1911-1982.
Science 219:969-971.
- Croxall, J. P. and R. L. Gentry (eds). 1987. Status, Biology, and Ecology of fur seals. NOAA
Tech. Rept NMFS 51:212 pp.
- Dahlheim, M. E. 1981. A review of the biology and exploitation of the killer whale, Orcinus
orca, with comments on recent sightings from Antarctica. Rep. int. Whal. Commn

31:541-546.

DeLong, R. L. and G. A. Antonelis. 1991. Impact of the 1982-1983 El Niño on the northern fur seal population at San Miguel island, California. pp. 75-83. In: F. Trillmich and K. Ono (eds.). Pinnipeds and El Niño: Responses to environmental stress. Springer-Verlag, Berlin.

DeLong, R. L., G. A. Antonelis, C. W. Oliver, B. S. Stewart, M. C. Lowry, and P. K. Yochem. 1991. Effects of the 1982-83 El Niño on several population parameters and diet of California sea lions on the California Channel Islands. pp. 166-184. In: F. Trillmich and K. Ono (eds.). Pinnipeds and El Niño: Responses to environmental stress. Springer-Verlag, Berlin.

DeMaster, D. P., D. J. Miller, D. Goodman, R. L. DeLong, and B. S. Stewart. 1982. Assessment of California sea lion fishery interactions. Trans. N. Amer. Wildl. Nat. Res. Conf. 47:

Dietz, R., M. P. Heide-Jorgensen, and T. Harkonen. 1989. Mass deaths of harbour seals (Phoca vitulina) in Europe. Ambio 18:258-264.

Felleman, F. L., J. R. Heimlich-Boran, and R. W. Osborne. 1991. The feeding ecology of killer whales (Orcinus orca) in the Pacific Northwest. pp. 113-141. In: Dolphin Societies (K. Pryor and K. S. Norris, eds.). Dolphin Societies. U.C. Press, Berkeley, CA. 397 pp.

Geraci, J. R. 1990. Physiologic and toxic effects on cetaceans. In: J. R. Geraci and D. J. St. Aubin (eds.). Sea Mammals and Oil: Confronting the Risks. Academic Press, New York.

Geraci, J. R. and D. J. St. Aubin. 1987. Effects of offshore oil and gas development on marine mammals and turtles. pp. 587-617. In: D. F. Boesch and N. N. Rabalais (eds).

Long-term environmental effects of offshore oil and gas development. Elsevier Applied Science, London and New York.

Geraci, J. R., D. J. St. Aubin, I. K. Barker, R. G. Webster, V. S. Hinshaw, W. J. Bean, H. L. Ruhnke, J. H. Prescott, G. Early, A. S. Baker, S. Madoff and R. T. Schooley. 1982. Mass mortality of harbor seals: pneumonia associated with influenza A virus. Science 215:1129-1131.

Godsell, J. 1988. Herd formation and haul-out behavior in harbour seals (Phoca vitulina). J. Zool., Lond. 215:83-98.

Gosho, M. E., D. W. Rice, and J. F. Breiwick. 1984. The sperm whale, Physeter macrocephalus. Mar. Fish. Rev. 46:54-64.

Grachev, M. A., V. P. Kumarev, L. V. Mamaev, V. L. Zorin, L. V. Baranova, N. N. Denikina, S. I. Belikov, E. A. Petrov, V. S. Kolesnik, R. S. Kolesnik, V. M. Dorofeev, A. M. Beim, V. N. Kudelin, F. G. Nagieva and V. N. Sidorov. 1989. Distemper virus in Baikal seals. Nature 338:209.

Guerra, C., C. G., and G. Portflitt K. 1991. El Niño effects on pinnipeds in northern Chile. pp 47-54. In: F. Trillmich and K. Ono (eds.). Pinnipeds and El Niño: Responses to environmental stress. Springer-Verlag, Berlin.

Harkonen, T. 1987. Seasonal and regional variations in the feeding habits of the harbour seal, Phoca vitulina, in the Skagerrak and the Kattegat. J. Zool., Lond. 213:535-543.

Harkonen, T. and M. P. Heide-Jorgensen. 1990. short-term effects of the mass dying of harbour seals in the Kattegat-Skagerrak area during 1988. Z. Saugetierk. 55:233-238.

Harkonen, T. and M. P. Heide-Jorgensen. 1990. Comparative life histories of east Atlantic and other harbour seal populations. Ophelia 32:211-235.

- Harvey, J. T., R. F. Brown, and B. R. Mate. 1990. Abundance and distribution of harbor seals (Phoca vitulina) in Oregon, 1975-1983. Northwest. Nat. 71:65-71.
- Harwood, J. 1990. The 1988 seal epizootic. J. Zool., Lond. 222:349-351.
- Heide-Jorgensen, M. P. and T. Harkonen. 1988. Rebuilding seal stocks in the Kattegat-Skagerrak. Mar. Mamm. Sci. 4:231-246.
- Heimlich-Boran, J. R. 1988. Behavioral ecology of killer whales (Orcinus orca) in the Pacific Northwest. Can. J. Zool. 66:565-578.
- Helle, E., M. Olsson, and S. Jensen. 1976. PCB levels correlated with pathological changes in seal uteri. Ambio 5:261-263.
- Hes, A. D., and G. P. Rouse. 1983. Population increase in the sub-Antarctic fur seal Arctocephalus tropicalis at Amsterdam Island. S. Afr. J. Antarct. Res. 13:29-34.
- Hinshaw, V. S., W. J. Bean, R. G. Webster, J. E. Rehg, P. Fiorelli, G. Early, J. R. Geraci, and D. J. St. Augin. 1984. Are seals frequently infected with avian influenza viruses? J. Virology 51:863-865.
- Hoelzel, A. R., J. K. Ford, and G. A. Dover. 1991. A paternity test case for the killer whale (Orcinus orca) by DNA fingerprinting. Mar. Mamm. Sci. 7:35-43.
- Hoover, A. A. 1988. Harbor seal. In: J. W. Lentfer (ed.). Selected marine mammals of Alaska. Marine Mammal Commission, Washington, D.C.
- Kerley, G. I. H. 1983. Relative population sizes and trends, and hybridization of fur seals Arctocephalus tropicalis and A. gazella at the Prince Edward Islands, Southern Ocean. S. Afr. J. Zool. 18:388-392.
- Kovacs, K. M., K. M. Jonas, and S. E. Welke. 1990. Sex and age segregation by Phoca vitulina concolor at haul-out sites during the breeding season in the Passamaquoddy

- Bay region, New Brunswick. *Mar. Mamm. Sci.* 6:204-214.
- Laws, R. M. 1992. History and present status of southern elephant seal populations. In: B. J. Le Boeuf and R. M. Laws (eds.). *Elephant seals*. University of California. In Press.
- Laws, R. and R. J. F. Taylor. 1957. A mass dying of crabeater seals, *Lobodon carcinophagus* (Gray). *Proceedings of the Zoological Society of London* 129:315-324.
- Leatherwood, S., C. O. Matkin, J. D. Hall, G. M. Ellis. 1990. Killer whales, *Orcinus orca*, photo-identified in Prince William Sound, Alaska, 1976 through 1987. *Can. Field Nat.* 104:362-371.
- Loughlin, T. R. and R. V. Miller. 1989. Growth of the northern fur seal colony on Bogoslof Island, Alaska. *Arctic* 42:368-372.
- Majluf, P. 1991. El Niño effects on pinnipeds in Peru. pp 55-74. In: F. Trillmich and K. Ono (eds.). *Pinnipeds and El Niño: Responses to environmental stress*. Springer-Verlag, Berlin.
- McLaren, I. A. 1990. Pinnipeds and Oil: Ecologic Perspectives. In: J. R. Geraci and D. J. St. Aubin (eds.). *Sea Mammals and Oil: Confronting the Risks*. Academic Press, New York.
- Merrick, R. L., T. R. Loughlin, and D. G. Calkins. 1987. Decline in abundance of the northern sea lion, *Eumetopias jubatus*, in Alaska, 1956-1986. *U. S. Fish. Bull.* 85:351-365.
- Mitchell, E. and R. R. Reeves. 1988. Records of killer whales in the western North Atlantic, with emphasis on eastern Canadian waters. *Rit Fiskideildir* 11:161-193.
- Olesiuk, P. F., M. A. Bigg, and G. M. Ellis. 1990a. Recent trends in the abundance of harbour seals, *Phoca vitulina*, in British Columbia. *Can. J. Fish. Aquat. Sci.* 47:992-1003.

- Olesiuk, P. F., M. A. Bigg, G. M. Ellis, S. J. Crockford, and R. J. Wigen. 1990b. An assessment of the feeding habits of harbour seals (Phoca vitulina) in the Strait of Georgia, British Columbia, based on scat analysis. Can. Tech. Rept. Fish. Aquat. Sci. 1730:1-135.
- Olesiuk, P. F., M. A. Bigg, and G. M. Ellis. 1990c. Life history and population dynamics of resident killer whales (Orcinus orca) in the coastal waters of British Columbia and Washington State. Rept. int. Whal. Commn., Special Issue 12:209-243.
- Oritsland, T. and I. Christensen. 1982. A mass stranding of killer whales at Lofoten, northern Norway, in June 1981. Rep. int. Whal. Commn. 323:642.
- Payne, M. R. 1977. Growth of a fur seal population. Philos. Trans. Royal Soc. Lond. 279(Ser B):67-79.
- Payne, P. M. and D. C. Schneider. 1984. Yearly changes in abundance of harbor seals, Phoca vitulina, at a winter haul-out site in Massachusetts. U. S. Fish. Bull. 82:440-442.
- Payne, R., V. Rowntree, J. S. Perkins, J. G. Cooke, and K. Lankester. 1990. Population size, trends and reproductive parameters of right whales (Eubalaena australis) off Peninsula Valdes, Argentina. Rep. int. Whal. Commn Spec. Iss. 12:271-278.
- Perry, A., C. S. Baker, L. M. Herman. 1990. Population characteristics of individually identified humpback whales in the central and eastern North Pacific: a summary and critique. Rep. int. Whal. Commn Spec. Iss. 12:307-317.
- Pierce, G. J., P. M. Thompson, A. Miller, J. S. Diack, D. Miller, and P. R. Boyle. 1991. Seasonal variation in the diet of common seals (Phoca vitulina) in the Moray Firth area of Scotland. J. Zool., Lond. 223:641-652.
- Pitcher, K. W. 1980a. Stomach contents and feces as indicators of harbor seal, Phoca vitulina, foods in the Gulf of Alaska. U. S. Fish. Bull. 78:797-798.

- Pitcher, K. W. 1980b. Food of the harbor seal, Phoca vitulina richardsi, in the Gulf of Alaska. U. S. Fish. Bull. 78:545-549.
- Pitcher, K. W. 1990. Major decline in number of harbor seals, Phoca vitulina richardsi, on Tugidak Island, Gulf of Alaska. Mar. Mamm. Sci. 6:121-134.
- Pitcher, K. W. and D. C. McAllister. 1981. Movements and haulout behaviour of radio-tagged harbour seals, Phoca vitulina. Can. Field-Nat. 95:292-297.
- Reijnders, P. J. H. 1978. Recruitment in the harbour seal (Phoca vitulina) population in Dutch Wadden Sea. Neth. J. Sea Res. 12:164-179.
- Reilly, S. B. 1984. Assessing gray whale abundance: a review. pp. 203. In: M. L. Jones, S. L. Swartz, and S. Leatherwood. The Gray Whale, Eschrichtius robustus. Academic Press, N. Y. 600 pp.
- Reilly, S. B. and Barlow, J. 1986. Rates of increase in dolphin population size. U. S. Fish. Bull. 84:527-533.
- Reilly, S. B., Rice, D. W. and A. A. Wolman. 1983. Population assessment of the gray whale, Eschrichtius robustus, from California shore censuses, 1967-1980. U. S. Fish. Bull. 81:267-281.
- Schneider, P. M. and P. M. Payne. 1983. Factors affecting haul-out of harbor seals at a site in southeastern massachusetts. J. Mammal. 64:518-520.
- Shaugnessy, P. D. and S. D. Goldsworthy. 1990. Population size and breeding season of the Antarctic fur seal Arctocephalus gazella at Heard Island--1987-1988. Mar. Mamm. Sci. 6:292-304.
- Smith, A. W., R. J. Brown, D. E. Skilling, and R. L. DeLong. 1974. Leptospira pomona and reproductive failure in California sea lions (Zalophus californianus californianus). J. Amer. Vet. Med. Assoc. 165:996-998.

- St. Aubin, D. J. 1990. Physiologic and toxic effects on pinnipeds. In: J. R. Geraci and D. J. St. Aubin (eds.). *Sea Mammals and Oil: Confronting the Risks*. Academic Press, New York.
- Stewart, B. S. 1984. Diurnal hauling patterns of harbor seals at San Miguel Island, California. *J. Wildl. Manage.* 48:1459-1461.
- Stewart, B. S. 1992. Population recovery of northern elephant seals on the southern California Channel Islands. In: D. R. McCullough and R. H. Barrett (eds.). *Wildlife 2001*. Elsevier Press. In Press.
- Stewart, B. S., G. A. Antonelis, R. L. DeLong, and P. K. Yochem. 1988. Abundance of harbor seals on San Miguel Island, California, 1927 through 1986. *Bull. So. Calif. Acad. Sci.* 87:39-43.
- Stewart, B. S. and P. K. Yochem. 1991. Northern elephant seals on the Southern California Channel Islands and El Niño. pp. 234-243. In: F. Trillmich and K. Ono (eds.). *Pinnipeds and El Niño: Responses to environmental stress*. Springer-Verlag, Berlin.
- Stewart, B. S., P. K. Yochem, R. L. DeLong, and G. A. Antonelis. 1992. Status and trends in abundance of pinnipeds on the Southern California Channel Islands. In: F. G. Hochberg (ed). *Proceedings of the Third California Islands Symposium*, Santa Barbara Museum of Natural History, Santa Barbara, CA. In Press.
- Terhune, J. M., and M. Almon. 1983. Variability of harbor seals numbers on haul-out sites. *Aquat. Mamm.* 10:71-78.
- Testa, J. W., G. Oehlert, D. G. Ainley, J. L. Bengtson, D. B. Siniff, R. M. Laws, and D. Rounsevell. 1991. Temporal variability in Antarctic marine ecosystems: Periodic fluctuations in the phocid seals. *Can. J. Fish. Aquat. Sci.* 48:631-639.
- Thompson, P. M., M. A. Fedak, B. J. McConnell and K. Nicholas. 1989. Seasonal and sex-

- related variation in the activity patterns of common seals (Phoca vitulina). J. Appl. Ecol 26:521-536.
- Thompson, P. M., G. J. Pierce, J. R. Hislop, D. Miller, and J. S. W. Diack. 1981. Winter foraging by common seals (Phoca vitulina) in relation to food availability in the inner Moray Firth, N. E. Scotland. J. Anim. Ecol. 60:283-294.
- Trillmich, F. and D. Limberger. 1985. Drastic effects of El Niño on Galapagos pinnipeds. Oecol. 67:19-22.
- Vedros, N. A., A. W. Smith, J. Schonewald, G. Migaki, and R. C. Hubbard. 1971. Leptospirosis epizootic among California sea lions. Science 172:12450-1251.
- Wilkinson, I. S. and M. N. Bester. 1988. Is onshore human activity a factor in the decline of the southern elephant seal? S. Afr. J. Antarct. Res. 18:14-17.
- Wursig, B. 1990. Cetaceans and oil: Ecologic Perspectives. In: J. R. Geraci and D. J. St. Aubin (eds.). Sea Mammals and Oil: Confronting the Risks. Academic Press, New York.
- Yochem, P. K. 1987. Haul-out patterns and site fidelity of harbor seals at San Nicolas and San Miguel Islands, California. M.S. thesis, San Diego State University, San Diego, CA.
- Yochem, P. K., B. S. Stewart, R. L. DeLong, and D. P. DeMaster. 1987. Diel hauling patterns and site fidelity of harbor seals (Phoca vitulina richardsi) on San Miguel Island, California, in autumn. Mar. Mamm. Sci. 3:323-332.
- York, A. E. 1987. On comparing the population dynamics of fur seals. NOAA Tech. Rep. NMFS 51:133-140.
- Zakharov, V. M. and A. V. Yablokov. 1990. Skull asymmetry in the Baltic grey seal: effects of environmental pollution. Ambio 19:266-269.

Zeh, J. E., J. C. George, A. E. Raftery, and G. M. Carroll. 1991. Rate of increase, 1978-1988, of bowhead whales, Balaena mysticetus, estimated from ice-based census data. Mar. Mamm. Sci. 7:105-122.

6.0. Tables

Table 1. A summary of growth rates expressed as percent increase per annum in various pinniped populations.

SPECIES	AREA	RATE	PERIOD	NOTES	SOURCE
<u>Phoca vitulina richardsi</u>	Alaskan peninsula	-3.5	1976-85	E	1
"	Tugidak I.	-19.0	1976-79	E	2
"	"	-7.0	1982-88	E	2
"	British Columbia	12.5	1973-88	E	3
"	Oregon	8.1	1975-83	E, I, D	4
"	Gulf of Farallones, Double Pt.	7.6	1976-87	E, I	5
"	Gulf of Farallones, S. Farallon I.	17.0	1974-86	E, I	5
"	San Miguel I.	22.0	1958-76	E, I	6
"	"	5.0	1976-86	E, I	6
<u>Phoca vitulina concolor</u>	Massachusetts	11.9	1972-83	E, D	7
"	Kattegat-Skagerrak	12.0	1979-86	E	8
<u>Callorhinus ursinus</u>	Pribilof Is.	8.0	1911-24	E	9
"	"	0.0	1950-55	E	10

"	"	-6.0	1955-65	E	10
"	"	0.0	1965-75	E	10
"	"	-7.8	1975-81	E	10
"	"	-1.8	1981-86	E	10
"	Commander I.	0.0	1974-82	E	10
"	Robben I.	-5.8	1974-82	E	10
"	Bogoslof I.	57.0	1980-88	E	11
<u>Eumetopias jubatus</u>	Alaska	-2.7	1956-86	E	12
<u>Arctocephalus</u> <u>tropicalis</u>	Gough I.	15.9	1955-77	E	13
"	Marion I.	10.5	1951-74	E, I	14
"	"	12.9	1974-89	E, I	15
"	"	15.0	1974-81	E, I	16
"	Amsterdam I.	11.0	1956-81	E	14
"	"	7.8	1955-69	E	17
"	"	16.5	1969-81	E	14
"	Prince Edward I.	9.7	1982-87	E, I	15

<u>Arctocephalus gazella</u>	Heard I.	20.7	1962-88	I	18
"	Bird I.	13.1	1958-75	E	19
"	Marion Is.	15.1	1974-81	E	16
"	Prince Edward Is.	11.3	1981-89	E, I	15
<u>Arctocephalus pusillus</u> <u>pusillus</u>	Southern Africa, mainland colonies	7.5	1971-83	E	20
"	Southern Africa, island colonies	-3.5	1971-83	E	20
"	Southern Africa	5.8	1971-80	E	21
<u>Arctocephalus australis</u>	All stocks	11.0	1953-72	E	22
<u>Arctocephalus townsendi</u>	Isla de Guadalupe	7.5	1954-77	E	23
<u>Mirounga angustirostris</u>	San Miguel I.	13.6	1964-81	E, I	24
"	San Nicolas I.	16.5	1959-81	E, I	24
"	Año Nuevo	15.8	1968-80	E, I	24
"	Farallon I.	53.3	1974-80	E, I	24
"	Isla de Guadalupe	5.4	1965-77	E	24
"	Islas San Benito	5.9	1965-77	E	24

SPECIES	AREA	RATE	PERIOD	NOTES	SOURCE
<u>Mirounga leonina</u>	South Georgia	0.0	1951-85	E	25
"	Patagonia	5.1	1975-82	E	25
"	"	3.2	1982-90	E	25
"	Iles Kerguelen	-4.6	1970-77	E	25
"	Heard I.	-2.4	1949-85	E	25
"	Marion I.	-4.8	1974-83	E	25
"	"	-1.9	1983-89	E	25
"	Macquarie I.	-2.1	1949-85	E	25
<u>Zalophus californianus</u>	California	8.7	1927-46	E	23
"	"	6.7	1947-70	E	23
"	San Miguel I.	5.0	1971-81	E	26
<u>Halichoerus grypus</u>	United Kingdom	7.0	Early 1960s-late 1970s	E	27

NOTES: D = Relaxation from disturbance; E = Exploited population; I = Immigration

SOURCE: 1 = Pitcher 1986, cited in Hoover 1988; 2 = Pitcher 1990; 3 = Olesiuk et al. 1990; 4 = Harvey et al. 1990; 5 = Allen et al. 1989; 6 = Stewart et al. 1988; 7 = Payne and Schneider 1984; 8 = Heide-Jorgensen and Harkonen 1988; 9 = Lander 1981; 10 = York 1987; 11 = Loughlin and Miller 1989;

Table 1, continued

12 = Merrick et al. 1987; 13 = Bester 1980; 14 = Condry 1978; 15 = Wilkinson and Bester 1990; 16 = Kerley 1983; 17 = Hes and Rouse 1983; 18 = Shaughnessy and Goldsworthy 1990; 19 = York 1987, after Payne 1977; 20 = Butterworth et al. 1987; 21 = Shaughnessy and Butterworth 1981, cited in York 1987; 22 = Vaz-Ferreira 1982, cited in York 1987; 23 = Chapman 1981; 24 = Cooper and Stewart 1983; 25 = Laws In Press; 26 = DeMaster et al. 1982; 27 = Harwood 1981

Table 2. A summary of growth rates expressed as percent increase per annum in various cetacean populations.

SPECIES	AREA	RATE	PERIOD	NOTES	SOURCE
<u>Balaenoptera musculus</u>	Iceland	4.8	1969-88	E	1
"	"	5.2	1979-90	E	2
<u>Megaptera noveangliae</u>	Iceland	11.5	1970-88	E	1
"	"	13.8	1979-88	E	1
"	"	14.8	1979-90	E	2
"	Western Australia	4.8	1963-88	E	1
"	Eastern Australia	10.0	1983-87	E	1
"	NW Atlantic	9.4	1979-86	E	1
<u>Eubalaena glacialis</u>	Argentina	7.6	1974-86	E	1
"	Western Australia	11.7	1977-87	E	3
"	South Africa	6.8	1971-87	E	1
<u>Balaena mysticetus</u>	Bering/Beaufort/ Chuckchi Seas	3.1	1978-88	E	4
"	Bering/Chukchi Seas	3.0- 4.5	1978-89	E	1

SPECIES	AREA	RATE	PERIOD	NOTES	SOURCE
<u>Balaenoptera musculus</u>	Iceland	4.8	1969-88	E	1
"	"	5.2	1979-90	E	2
<u>Megaptera noveangliae</u>	Iceland	11.5	1970-88	E	1
"	"	13.8	1979-88	E	1
"	"	14.8	1979-90	E	2
<u>Eschrichtius robustus</u>	California stock	2.5	1967-80	E	5
<u>Orcinus orca</u>	British Columbia	3.01	1973-81	E	6
"	"	1.67	1973-81	U	6
"	Puget Sound	2.3	1973-81	E	7

NOTES: E = Exploited population; U = Unexploited population

SOURCE: 1 = Best 1990; 2 = Sigurjonsson and Gunnlaugsson 1990; 3 = Bannister 1990; 4 = Zeh et al. 1991; 5 = Reilly 1984; 6 = Bigg 1982; 7 = Balcomb et al. 1982.

7.0. Appendix 1.

Annotated bibliography of the literature on recovery of ecosystems following man-induced and natural-phenomena-related disturbances: Harbor seals and killer whales.

Allen, S. G.; Ainley, D. G.; Page, G. W.; Ribic, C. A. (1984)

The effect of disturbance on harbor seal haul out patterns at Bolinas Lagoon, California

U.S. Fish. Bull. 82 (3). 493-500.

ABSTRACT

We studied harbor seals at Bolinas Lagoon, California, from May 1978 to June 1979. Field observation and two time lapse motion picture cameras were used to monitor the numbers of seals and of disturbances, and to provide information on tidal height. Peak numbers occurred during the summer. During nonbreeding seasons, high numbers occurred at low tides, and during the breeding season they occurred in early afternoon except when haul out areas were flooded. Seals were disturbed by humans on 71% of days monitored; people in canoes were the primary source of disturbance. Human activities closer than 100 m caused seals to leave haul out sites more than activities at greater distances.

Allen, S. G.; Ribic, C. A.; Kjelson, J. E. (1988)

Herd segregation in harbor seals at Point Reyes, California

Calif. Fish Game 74 (1). 55-59.

ABSTRACT

A mixed class aggregation of harbor seals resides at Double Point, California. Seals are present year round but are more abundant during the March through June breeding season. Seals were segregated by sex and age class within the herd, and patterns changed during the season. Early in the 1984 and 1985 breeding seasons the herd was equally composed of males, females, and immatures, but mid-way was composed of mostly females and pups. Late in the season more males and immatures were present. We hypothesize that changes in segregation patterns are related to intolerance of males and immature by pregnant females and females with pups.

Allen, S. G.; Huber, H. R.; Ribic, C. A.; Ainley, D. G. (1989)

Population dynamics of harbor seals in the Gulf of the Farallones, California

Calif. Fish Game 75 (4). 224-232.

ABSTRACT

We surveyed harbor seals, *Phoca vitulina*, in the Gulf of the Farallones, California, at all known haul-out sites from March 1982 through February 1984, and studied them intensively at two haul-out sites, Double Point and the South Farallon Islands, from 1976 to 1986. Though present year round, seals were most abundant onshore during the breeding/molt season (March-July). The relative abundance of seals onshore at Double Point during the 1987 breeding season was double the number in 1976, and at the South Farallon Islands, numbers in 1986 were four times higher than in 1974. Individual females observed at Double Point during two 3-year sets had a 0.89 and 0.92 probability of parturition in successive years.

Anonymous (1982)

Report of the Workshop on Identity, Structure and Vital Rates of Killer Whale Populations, Cambridge, England, June 23-25, 1981

Rep. int. Whal. Commn 32. 617-631.

ABSTRACT

Results of the workshop are summarized in this report. Killer whale pregnancy rates (% of mature females pregnant) from catch data were corrected for a 15-month gestation period as follows: Norway 12%-32.8%, Antarctic 13.74%-18.97%. A partial list of food items by geographic area is included in the appendices.

Baker, J. R.; McCann, T. S. (1989)

Pathology and bacteriology of adult male Antarctic fur seals, Arctocephalus gazella, dying at Bird Island, South Georgia

Br. vet. J. 145. 263-275.

ABSTRACT

A high mortality rate occurs in Antarctic fur seal males on the breeding beaches of Bird Island, South Georgia. The main causes of death were infections of fighting wounds and pneumonias. The bacteria involved appear to be opportunistic pathogens, predominantly various strains of streptococci.

Baker, J. R. (1989)

Natural causes of death in non-sucking grey seals (Halichoerus grypus)

Vet. Rec. 125. 500-503.

ABSTRACT

Thirty-four grey seals which died of natural causes were examined. They ranged in age from aborted fetuses to adults, but suckling pups were excluded from the study. The commonest primary cause of death was pneumonia and a variety of parasitoses occurred as secondary lesions.

Balcomb, K. C., III; Boran, J. R.; Heimlich, S. L. (1982)

Killer whales in Greater Puget Sound

Rep. int. Whal. Commn 32. 681-685.

ABSTRACT

From 1974 to 1980 killer whales (Orcinus orca) in a previously exploited population in Greater

Puget Sound, North America increased in number at a rate of 2.3% per year. The average adult female reproductive rate was observed to be 0.089 calves per year, and the average mortality rate for all whales was calculated to be 0.010 per year.

Bannister, J. L. (1990)

Southern right whales off Western Australia

In: Individual recognition of cetaceans: Use of photo-identification and other techniques to estimate population parameters. Report of the International Whaling Commission (Special Issue 12)

Hammond, P. S.; Mizroch, S. A.; Donovan, G. P., eds. Pages 279-288

International Whaling Commission, Cambridge

ABSTRACT

Southern right whale aerial counts and head callosity photographs from southern Western Australia, over some eleven years to 1987, have provided information on: a significant population increase, at least in cow-calf pairs; differences in some body characters (lip callosities, some body markings) compared with animals off South Africa and Argentinian; patterns of distribution and dispersal along the coast; and reproduction (most mating activity unlikely to be taking place on the coast, one animal first seen as a probable yearling giving birth nine years later, calving interval averaging three years). Up to 81 individuals (including 21 calves) have been identified in one year along some 1,100 km of coastline. Point estimates of rates of population increase ranged from 11.7% (all animals to 13.0% (cow-calf pairs).

Barham, E. G. (1982)

Marine mammals in Monterey Bay, California, during the years 1950-1955

Calif. Fish Game 68 (4). 213-223.

ABSTRACT

Over about a 5-year period, 180 sightings of 12 marine mammal species and three unidentified mammal categories were made during 239 weekly sea trips. Most frequently sighted were the Pacific white-sided dolphins and Dall's porpoise. Mean herd sizes were 13.90 and 5.98 animals, respectively. The white-sided dolphin was absent from early May to early September; the majority of sighting occurred from December to March. Dall's porpoise was seen around the calendar, but more frequently during the last half of the year. Killer, short-finned pilot, and sperm whales were noted only in that same period. The majority of gray whale sightings were correlated with their winter migrations, but a few stragglers were seen as late as July. No seasonality is evident from the humpback whale sightings. Northern fur seals were observed only from January to June. Two northern elephant seals and one sea otter were noted.

Barlow, J. (1985)

Variability, trends, and biases in reproductive rates of spotted dolphins, Stenella attenuata

U.S. Fish. Bull. 83 (4). 657-669.

ABSTRACT

Temporal changes were examined in three parameters that affect reproduction of spotted dolphin populations in the eastern Pacific. Of mature females, percent pregnant decreased markedly from the period 1971-1973 to the period 1974-83. Within the period 1974-83, percent pregnant remained relatively constant. Of pregnant females, percent lactating increased during the period 1971-83. The percentage of sexually mature females did not change. Potential biases in the measurement of the three parameters were identified by examining the effects of sampling conditions. The percentage of mature females that are pregnant and the percentage of pregnant females that are lactating were found to be robust to sampling conditions. The percentage of mature females in a sample was found to depend significantly on the number of dolphins killed per set, and annual variability was too large to be explained by random sampling error. Comparisons between two populations show that the more exploited population has a lower percent pregnant, although the opposite might be expected from density compensatory effects. Percent lactating and percent immature were higher in the more exploited population.

Bayer, R. D. (1985)

Six years of harbor seal censusing at Yaquina Estuary, Oregon

Murrelet 66. 44-49.

ABSTRACT

Harbor seals regularly hauled out at Yaquina Estuary only during tides lower than about +0.3m. Few seals (and no pups) were present during the pupping season in May and June. The greatest numbers hauled out in August or September; a smaller secondary peak appeared in February into two of six years. Seal abundance was about the same from 1977-1983, with the summer maximum ranging from 40 to 72 seals.

Berkson, J. M.; DeMaster, D. P. (1985)

Use of pup counts in indexing population changes in pinnipeds

Can. J. Fish. Aquat. Sci 42. 873-879.

ABSTRACT

A series of population simulations were used to test the accuracy of estimating the discrete rates of population change (RPC) from annual pup counts. The simulations indicate that pup counts can give a biased estimate of RPC, and the magnitude and direction of bias depends on which life history parameters are density dependent and on the maximum rate of population change. In general, if pre-census pup survival is density dependent the estimated RPC using pup counts is too

low. If post-census pup survival is density dependent, the estimated RPC is too high. If adult survival is density dependent, there is very little bias in the estimate. The results indicate that pup counts can be reliable indicators of population growth, but caution should be used in interpreting the results unless density feedback mechanisms have been identified.

Best, P. B.; Canham, P. A. S.; Macleod, N. (1984)

Patterns of reproduction in sperm whales, Physeter macrocephalus

In: Reproduction in whales, dolphins and porpoises. Reports of the International Whaling Commission (Special Issue 6)

Perrin, W. F.; Brownell, R. L., Jr.; DeMaster, D. P., eds. Pages 51-79

International Whaling Commission, Cambridge

ABSTRACT

From ovarian activity the peak period of breeding in the sperm whale seems to occur between October and December in the Southern Hemisphere and between April and June in the Northern Hemisphere. While large males (over 13.7m in length) may be the prime breeding animals, the density of spermatozoa in seminal fluid suggests that males may be physiologically fertile at an average length of 12.5m. Gestation (estimated from the difference between peaks of mating and calving) may last 15 to 16 months, though a comparison of neonatal and adult brain weights indicates a longer period (18.9 months). Equal numbers of male and female calves are born at an average length (from an examination of 15 neonates) of 4.00 ± 0.13 m; sexual dimorphism in size at birth cannot be demonstrated. Sperm whale milk is composed on average of $35.5 \pm 1.3\%$ total solids, $24.4 \pm 1.2\%$ fat, $9.1 \pm 0.3\%$ protein and $0.7 \pm 0.04\%$ ash. Diving ability of newborn animals seems to be relatively poor. Neonates demonstrate a strong 'following response' that wanes rapidly with age. An interdependence of school members is suggested from behavioural observations. Calves grow to about 6.1 m in length at one year of age and weigh 2,698 kg, or an increase of 1,673 kg since birth. Tests for lactose in stomach contents were positive in males up to 13 years of age and in females up to 7 1/2 years. Solid food is taken for the first time before the age of one year. Juvenile sperm whales tend to eat smaller and younger squid than adults. Heart weight relative to body weight may be about twice that of adults, suggesting a higher metabolic and thus feeding rate. Criteria for accurate measurement of reproductive (=pregnancy) rates are discussed. Mean calving interval for Donkergat (west coast of South Africa) is estimated as 5.2 years and 6.0-6.5 years for Durban (east coast of South Africa). A decrease in the calving interval from 6 to 5.2 years at Durban was observed between 1962-65 + 1967 and 1973-75. The mean duration of lactation may increase with the age of the female. Some of the older juveniles found with lactose in their stomach may represent offspring of older females in an extended period of lactation. The benefits of possible communal suckling by sperm whale calves are discussed.

Best, P. B. (1990)

Trends in the inshore right whale population off South Africa, 1969-1987

Mar. Mamm. Sci. 6 (2). 93-108.

ABSTRACT

Results of annual aerial surveys of the right whale population along the southern coast of South Africa from 1971 to 1987 are analysed. About 91.5% of cows with calves and 82% of unaccompanied adults on the South African coast in spring are found within the standard survey area, with some indications that the range is expanding up the west coast. In the nearshore region, most right whales (90%) are found within 1.85 km, and all cows with calves within 0.93 km of the coast. Within the standard survey area, specific areas of concentration can be predictably identified, both for cows with calves and unaccompanied adults. Although total counts (and counts expressed per hour flown) have increased overall, by a best estimate of 6.8% (95% CL 4.6, 9.0) per year from 1971 to 1987, some concentration areas for both classes have failed to show an increase over the same time period. Photographic flights since 1979 have permitted the movements of individually identified adult females to be monitored between successive calves. In at least one concentration area from which no overall increase between 1971 and 1987 was apparent (Mossel Bay), a large net emigration rate was found, nearly all of which involved a shift to the main nursery area to the west, off De Hoop. Reasons for the dissimilar dynamics of different concentration areas are not yet known.

Best, P. B. (1990)

Recovery rates in whale stocks that have been protected from commercial whaling for at least 20 years

Rep. int. Whal. Commn 40. 129-130.

ABSTRACT

Population growth rates are presented for five whale species (gray, right, bowhead, humpback, blue) that have been protected from commercial harvest for at least 20 years.

Bester, M. N. (1990)

Population trends of Subantarctic fur seals and southern elephant seals at Gough Island

S. Afr. J. Antarct. Res. 20 (1). 9-12.

ABSTRACT

The Subantarctic fur seal (*Arctocephalus tropicalis*) population at Gough Island in the South Atlantic Ocean is continuing to increase rapidly since its recovery from exploitation. The intrinsic rate of increase is however slowing down on established breeding colony beaches in the western sector as congested conditions develop. The rate of increase on the more recently colonised breeding colony sites on the east coast is high but some beaches here remain unexploited by

breeders despite the increased density on the west coast. The small breeding population of southern elephant seals (Mirounga leonina) either remained stable, or declined very slowly, over the past 17 years.

Bigg, M. (1982)

An assessment of killer whale (Orcinus orca) stocks off Vancouver Island, British Columbia

Rep. int. Whal. Commn 32. 655-666.

ABSTRACT

A study of photographically identifiable individual killer whales was undertaken during 1973-81. In all, 30 pods were found, containing about 260 whales. A pod is a long term family or kinship group which periodically joins with others to form communities. Around Vancouver Island there are two resident communities and one transient community. The three communities do not associate with one another. Resident and transient killer whales differ in pod size and behaviour. The coastal range of movements for most resident pods is probably about 300 nm. Transient pods appear to range further. Births occur mainly during fall and winter. At birth, lengths average about 8 ft. First pregnancies generally occur at 16 ft, or at an age of at least 6.7 years. Sexual maturity occurs in males at 19 ft, or at an age of at least 12 yrs. A long term stability in pod composition permits direct measurement of vital statistics. The rate of calf production to an average age of six months is 10.30% per cow. The minimum interval between calving is three years. Many cows apparently rarely give birth. Annual natural mortality rates average 2.80% for bulls, 0.70% for cows and 2.30% for juveniles. Pods increase at an average net rate of 2.52% per year. Exploited pods have a slightly higher productivity than unexploited pods.

Bjorge, A. (1991)

Status of the harbour seal Phoca vitulina L. in Norway

Biol. Conserv. 58. 229-238.

ABSTRACT

The current estimate of the number of harbour seal Phoca vitulina in Norwegian waters, including Svalbard, is 4129, based on actual counts of seals at the haul-out sites during the period 1977-88. Comparison with an estimate obtained during the early 1960s indicates that there has been little change in the overall number of harbour seals in Norway during the last 25 years. Three distinct types of habitats are utilized by harbour seals in Norway--open rocky coasts, deep fjords and estuarine sandbanks. The habitat types and the status of legislative protection of harbour seals and their habitats are described.

Blix, A. S.; Grav, H. J.; Ronald, K. (1979)

Some aspects of temperature regulation in newborn harp seal pups

Am. J. Physiol. 236 (3). R188-R197.

ABSTRACT

Harp seals are born on the drifting ice of the North Atlantic Ocean during arctic winter when temperatures of -20°C , occasionally in combination with wind of 10 m/sec, might prevail for days. At birth the pups lack subcutaneous blubber and the wet infantile fur has a conductance value of $30.0 \text{ W}\cdot\text{m}^{-2}\cdot^{\circ}\text{C}^{-1}$ as compared with only $2.0 \text{ W}\cdot\text{m}^{-2}\cdot^{\circ}\text{C}^{-1}$ when dry. While still wet immediately after birth the pups are nevertheless able to retain body core temperature by shivering. This activity leads to reduction of muscle fat and glycogen stores. Nonshivering thermogenesis commences in thermogenic adipose tissue by virtue of loosely coupled mitochondria. Thermogenic adipose tissue is found at birth both as a subcutaneous layer along the back and as internal deposits around venous plexuses in the neck, on the pericardium, on the kidneys, and the abdominal walls. After about 3 days of sucking the subcutaneous adipose tissue loses its thermogenic function being gradually transformed into blubber, whereas the internal deposits persist at least until the pups venture into water at the age of 3-4 wk.

Boal, J. (1980)

Pacific harbor seal (Phoca vitulina richardii) haul out impact on the rocky midtidal zone

Marine Ecol. Prog. Ser. 2 (4). 265-269.

ABSTRACT

A study of haul out and adjacent non-haul out rocks of the Pacific harbor seal Phoca vitulina richardii (Gray), on the Monterey Peninsula of California, revealed significant differences in algal morphology and per cent composition, and in numbers of animals present. It is suggested that the mechanical and chemical impact imposed by the seals on the haul out sites are responsible for these differences.

Boulva, J.; McLaren, I. A. (1979)

Biology of the harbor seal, Phoca vitulina, in eastern Canada

Bull. Fish. Res. Board Can. Need Pages.

ABSTRACT

The biology of harbour seals in eastern Canada is reviewed. Population surveys and estimates, rates of decrease of bounty kill and age structure of heavily vs. lightly exploited populations all suggest an overall decline of about 4% per year between 1950 and 1973 in the Maritimes. Best estimates of pre-weaning and post-weaning mortality rates, produce near equilibrium in population simulation. If females were fertile 1 year younger this would only allow a very small sustainable yield. The very

high natural mortality rates, possibly a result of shark predation and overexposed breeding sites in the reduced range, may make eastern Canadian harbor seals particularly vulnerable to hunting.

Braham, H. W.; Dahlheim, M. E. (1982)

Killer whales in Alaska documented in the Platforms of Opportunity Program

Rep. int. Whal. Commn 32. 643-646.

ABSTRACT

Sighting records of over 1,100 killer whale (Orcinus orca) groups from Alaskan waters obtained from 1958 to 1980 are now in the National Marine Mammal Laboratory's Platforms of Opportunity Program data base. From a preliminary analysis, and with as yet no correction for sighting effort, killer whales are distributed primarily over the continental shelf in water less than 200 m deep in the eastern North Pacific (southeast Alaska to the eastern Aleutian Islands) but seem to concentrate near the 200 m contour over the shelf slope in the southeastern Bering Sea. In the Gulf of Alaska over 60% of our sightings were within 20 km of shore, while in the North Pacific from Kodiak Island to Unimak Pass they were uniformly distributed from shore to beyond 56 km. Pod size ranged from 1-100 animals, with only 1% of the groups containing 20 or more individuals. From a review of the literature, and in light of our sightings of concentrations near areas of known high productivity of fishes and other marine mammals, killer whales appear to feed upon fish when locally abundant and to switch to marine mammals when fish are less available.

Braham, H. W. (1984)

Review of reproduction in the white whale, Delphinapterus leucas, narwhal, Monodon monoceros, and Irrawaddy dolphin, Orcaella brevirostris, with comments on stock assessment

In: Reproduction in whales, dolphins and porpoises. Reports of the International Whaling Commission (Special Issue 6)

Perrin, W. F.; Brownell, R. L. Jr.; DeMaster, D. P., eds. Pages 81-89

International Whaling Commission, Cambridge

ABSTRACT

Estimates of vital reproductive parameters for white whales, Delphinapterus leucas, have been made over the past 20 years as a result of data collected from a long, but intermittent history of exploitation. Based on two dentinal growth layers per year, reported in the literature, they may live to 25-30 years. Females become sexually mature at 5 years of age and males at 8 years; they begin active breeding 1-3 years later. Evidence is presented which suggests that estimating age using current methods may result in error in ages at which certain life history events occur. From an observed pregnancy rate of 0.41, and assuming an average gestation period of 14.5 months and a crude birth rate of 0.33-0.38, a gross annual reproductive rate of 0.09-0.12 is estimated. Observed rates are 0.09-0.14. Most vital reproductive rates are unknown for narwhals, Monodon monoceros. Almost no life history information is available on the Irrawaddy dolphin, Orcaella brevirostris.

Reevaluation and new estimates of stock abundance and vital rate parameters are needed for white whales because under present harvest levels 44% of the defined harvested stocks are being exploited at or above replacement yield.

Brown, R. F.; Mate, B. R. (1983)

Abundance, movements, and feeding habits of harbor seals, Phoca vitulina, at Netarts and Tillamook Bays, Oregon

U.S. Fish. Bull. 81. 291-301.

ABSTRACT

Patterns of seasonal abundance of harbor seals at Netarts and Tillamook Bays, Oregon, were documented by recording numbers of seals hauling out on tidally exposed sand flats in both bays. Harbor seal abundance at Tillamook Bay coincided with the annual return (October-November) of chum salmon, Oncorhynchus keta, to a hatchery on Whiskey Creek. Observations of seals preying on adult salmon resulted in estimated losses of 6.1, 7.2, and 1.5% of the total chum returns for 1978, 1979, and 1980, respectively, due to seal predation in the Whiskey Creek area. Other prey species of harbor seals at Netarts Bay were identified by the recovery of prey hard parts from seal feces collected on haul-out areas. The Pacific sand lance, Ammodytes hexapterus, was the most frequently identified prey item. Ten species of flatfish (Order Pleuronectiformes) were identified as harbor seal prey with five species (Parophrys vetulus, Glyptocephalus zachirus, Citharichthys sordidus, Microstomus pacificus, and Lyopsetta exilis) ranking among the seven most frequently occurring food items. In general, benthic and epibenthic fish appeared to be important in the harbor seal diet. Distributions, abundances, and estimated sizes of identified prey species indicated that harbor seals had fed both in Netarts Bay and in the nearshore ocean. Movements of radio-tagged harbor seals between Netarts Bay and Tillamook Bay were common (45.4% of tagged seals made at least one move between bays). Tagged harbor seals frequented at least four different estuaries and one coastal haul-out area, ranging from 25 to 550 km from the tagging area.

Budylenko, G. A. (1981)

Distribution and some aspects of the biology of killer whales in the South Atlantic

Rep. int. Whal. Commn 31. 523-525.

ABSTRACT

Killer whales (Orcinus orca) are widely distributed in the South Atlantic. Data are presented which were obtained from the whaling vessel Yuri Dolgorukiy for the 15 years 1960-75 in the months November-May. During this period killer whales were observed in warm, temperate and cold waters.

In temperate waters, the bulk of the diet consists of dolphins and fish, while in the colder water, minke whales are the dominant food species. Killer whales are most commonly seen in groups of about 10 animals. Groups of 10-20 animals are less common and groups of over 100 are occasionally seen. The size of the killer whale groups is determined by the hunting strategy.

Butterworth, D. S.; David, J. H. M.; McQuaid, L. H.; Xulu, S. S. (1987)

Modeling the population dynamics of the South African fur seal Arctocephalus pusillus pusillus

NOAA Tech. Rep. NMFS 51. 141-164.

ABSTRACT

Aerial survey and tag-recapture assessments of fur seal pup population numbers are considered at the 23 breeding colonies around the southeastern and western coasts of southern Africa during the period 1971-83. Exponential growth curves are fitted for each colony assuming a constant relative bias between the various assessment methods used. The pup population for all colonies combined is estimated to have grown at an average annual rate of 3.9% (SE 1.1%). The population is now dominated by four major mainland colonies which contribute 78% to the 1983 total pup population estimate of 310,000. Mainland colonies have increased over the period considered at an average annual rate of 7.5% (SE 1.5%), while island colonies have declined at 3.5% (SE 0.9%) per annum. Estimation of change in growth rate suggests that this rate has increased over the period though not significantly ($P=0.07$).

No direct assessments are available for adult and juvenile survival rates for the South African fur seal Arctocephalus pusillus pusillus, but limitations can be placed on possible ranges of values. An approach is suggested which imposes the constraint of a population dynamics model for adult females upon these ranges, estimates of total pup population size and growth rate, and knowledge of the average annual harvest of pups from 1971 to 1983. This provides refined probability distributions for various demographic parameters; the annual average pup harvesting rate is estimated to have been 38% (SE 5%), and the annual adult female survival rate 0.92 (SE 0.02%). An example is given of how the approach can be extended to provide estimates for the total population and to detect possible density-dependent effects, priority should be given to further assessments of the major mainland colonies, particularly those at Wold and Atlas Bays.

Chapman, D. G. (1981)

Evaluation of marine mammal population models

In: Dynamics of large mammal populations

Fowler, C. W.; Smith, T. D., eds. Pages 277-296

John Wiley & Sons, New York

ABSTRACT

Population models used for a variety of marine mammals are classified (growth models, yield models, multispecies and ecosystem models). Population size estimates and exponential growth rates are presented for several species.

Christensen, I. (1982)

Killer whales in Norwegian coastal waters

Rep. int. Whal. Commn 32. 633-641.

ABSTRACT

Observations reported by whalers and fishermen indicate that killer whales are present in all areas of the Norwegian coast throughout the year. No migration pattern has been discovered, but killer whales occur in the greatest numbers in the Lofoten and More areas and off the southern west coast. A mean catch of 57 killer whales per year over the period 1938-1980 has not changed the length distribution in the catch. Decreasing fatness from 1951 to 1968 may have been caused by a decreasing availability of food. Reported weights indicate that meat constitutes 40% of total body weight, blubber 29%, bones 22%, and viscera 9%. Feeding behaviour and pod organization is described from recorded observations. Female killer whales seem to reach sexual maturity at a length of 15 feet, and an age of about 6 years. Matings occur throughout the year, with a maximum in October-December. A gestation period of 12 months and one birth every second year is indicated. Preliminary age determinations indicate continued growth in groups of body length up to 20-25 years and a life span of at least 35 years.

Christensen, I. (1984)

Growth and reproduction of killer whales, Orcinus orca, in Norwegian coastal waters

In: Reproduction in whales, dolphins and porpoises. Reports of the International Whaling Commission (Special Issue 6)

Perrin, W. F.; Brownell, R. L., Jr.; DeMaster, D. P., eds. Pages 253-258

International Whaling Commission, Cambridge

ABSTRACT

Length data and reproductive material collected by coastal whalers during the period 1938-67 and 1978-81 are analysed. Female killer whales attain sexual maturity at a length of 15-16 ft (4.6-4.9 m) and an age of about 8 years, and males at about 19 ft (5.8 m) and 15 years. Mating occurs throughout the year, with a maximum in October-December. The birth rate seems to be one calf every three years. Preliminary age determinations indicate that physical maturity is attained at 20-25 years, with a life span of at least 35 years.

Christensen, I. (1990)

A note on recent strandings of sperm whales (Physeter macrocephalus) and other cetaceans in Norwegian waters

Rep. int. Whal. Commn 40. 513-515.

ABSTRACT

Strandings of 27 sperm whales on the coast of Norway and sightings of an additional 9 dead and drifting whales in Norwegian coastal waters in 1988 and 1989 are recorded. Also reported are the strandings of other cetaceans, including 6 killer and 7 pilot whales, which also occurred in these years. Factors accounting for the death of the stranded whales are discussed.

Clapham, P. J.; Mayo, C. A. (1990)

Reproduction of humpback whales (Megaptera novaeangliae) observed in the Gulf of Maine

In: Individual recognition of cetaceans: Use of photo-identification and other techniques to estimate population parameters. Report of the International Whaling Commission (Special Issue 12)

Hammond, P. S.; Mizroch, S. A.; Donovan, G. P., eds. Pages 171-175

International Whaling Commission, Cambridge

ABSTRACT

A high level of effort in the Gulf of Maine between the years 1979 and 1987 has produced detailed resighting histories of individual humpback whales, Megaptera novaeangliae, allowing us to conduct a study of reproduction in this population. The crude birth rate during this period ranged from 0.045 to 0.103 (mean=0.079), with no significant year-to-year variation. An alternative measure gave a mean reproductive rate of 0.41 calves per mature female per year. Sixty-five females were observed with a total of 120 calves. Thirty-one females were observed with more than one calf during the study. Observed calving intervals were: 1 year (n=2), 2 years (n=36), 3 years (n=14), 4 years (n=2) and 5 years (n=1); 52 of the 55 intervals involved complete resighting histories between years of known calving. Five calves born during the study period were observed with calves of their own in later years. Data from these whales, and from other mothers whose age can be estimated, suggest that the majority of females attain sexual maturity by the age of six. Of 94 calves born prior to 1987, 72 (76.6%) were resighted in at least one year after separation from their mothers, providing further support for the belief that the composition of a humpback whale feeding stock is determined matrilineally. Eleven of a possible 13 calves were observed to the age of 6 years and may therefore have survived to breeding age. Two others were probably lost to entanglements in commercial fishing gear, which may represent a significant source of mortality in this population. The value of using variation in the shape, size and scarring of the dorsal fin to identify individuals is discussed.

Clark, D. W. (1986)

Archaeological and historical evidence for an 18th-Century "Blip" in the distribution of the northern fur seal at Kodiak Island, Alaska

Arctic 39 (1). 39-42.

ABSTRACT

Recovery of fur seal Callorhinus ursinus remains from archaeological sites on Kodiak Island, Alaska, shows a low harvest prior to late prehistoric and early historic time. Then there is a pronounced increase in the frequency of fur seal bones in refuse layers. Russian records do not show any significant take of fur seals from Kodiak, but by the end of the 18th century and the beginning of the next century there are reports that this animal, formerly abundant in the area, had become rare. This may indicate that conditions had reverted to their earlier prehistoric state.

Colbourne, P. L.; Terhune, J. M. (1991)

Harbour seals (Phoca vitulina) do not follow herring movements in the Bay of Fundy, Canada

Ophelia 33 (2). 105-112.

ABSTRACT

Aerial surveys of hauled-out harbour seals (Phoca vitulina) were conducted along the mainland New Brunswick coast of the Bay of Fundy, 1-2 times monthly from January to mid-December 1987. A gradual summer increase and autumnal decrease in seal numbers were observed. Atlantic herring (Clupea harengus) in the Bay of Fundy move onshore and are caught in weirs between May and November. Comparison of the seals' distribution on haul-out sites with that of the Atlantic herring catch indicates that both species moved independently of each other. The seals' bi-weekly movements to different regions within the Bay appear to be random.

Condry, P. R.; van Aarde, R. J.; Bester, M. N. (1978)

The seasonal occurrence and behaviour of killer whales Orcinus orca, at Marion Island

J. Zool., Lond. 184. 449-464.

ABSTRACT

The paper describes the occurrence of Killer whales at Marion Island (Prince Edward group) in the south Indian Ocean from August 1973 to November 1976. They occur seasonally, being most numerous from October to December. Their occurrence is synchronized with the seasonal haul out of Southern elephant seals, but the seasonality of King, Rockhopper and Macaroni penguins is also likely to influence their occurrence. The largest herds occur in October, the month during which the mean group size is also largest. Sex and age composition are given, adult males being significantly more numerous than adult females, while 36.3% of the latter had calves. Hunting activity appears to be greatest between 15.00 and 17.00 hrs, and most Killer whales were seen within

100 m of the shore. Aspects of hunting, attacking, feeding and resting behaviour are discussed. The body measurements of a young male found on a beach are given.

Condy, P. R. (1978)

Distribution, abundance, and annual cycle of fur seals (Arctocephalus spp.) on the Prince Edward Islands

S. Afr. J. Wildl. Res. 8. 159-168.

ABSTRACT

The following annual growth rates are presented for A. tropicalis: 10.5% (1951-1974), Marion Island; 11.0% (1956-81), 16.5% (1969-81), Amsterdam Island.

Cooper, C. F.; Stewart, B. S. (1983)

Demography of northern elephant seals, 1911-1982

Science 219. 969-971.

ABSTRACT

Northern elephant seals (Mirounga angustirostris) were hunted to near extinction in the 19th century. Protection has allowed them to recolonize former habitat on islands off California, where the population is increasing more than 14 percent per year. Immigration of young pregnant females from Baja California initiated the California rookeries but is responsible for only a small part of recent population growth. Almost 25,000 northern elephant seal pups were born in the species' range in Mexico and the United States in 1982 in comparison with only six known births in 1911.

Costa, D. P.; Croxall, J. P.; Duck, C. D. (1989)

Foraging energetics of Antarctic fur seals in relation to changes in prey availability.

Ecol. 70 (3). 596-606.

ABSTRACT

This research examines the energy budget of breeding female Antarctic fur seals, both when food was plentiful and when it was scarce. The energy expenditure and change in body mass of lactating female Antarctic fur seals, Arctocephalus gazella, foraging at sea was measured in two years using doubly labeled water at South Georgia Island. There was no difference between years in mass gain, water influx, mass-specific field metabolic rate (FMR), or absolute FMR. Mean at-sea FMR over both years was 9.52 ± 0.55 W/kg ($n=22$), a value that is 6.7 times the predicted basal rate but only 1.9 times the FMR measured onshore. Comparable results have been reported for similar-sized northern fur seals.

Krill, the nearly exclusive prey of breeding females, were very scarce in 1984 at South Georgia. Fur

seal foraging trips were twice as long in 1984 as in 1985 and total mass-specific energy expended by females during these trips was significantly greater. In addition, females were significantly lighter at parturition in 1984, and both pup mortality and the proportion of pups that died from starvation were double the 1985 values. Female condition at parturition and average foraging-trip duration (i.e., offspring-provisioning rate) appear to reflect prey availability. The similarity between years in mass increase suggests that females do not return to feed their pups until they replenish their own reserves. Antarctic fur seal females may have a limited ability to increase the relative time spent foraging because even in normal years only 5% of their time at sea is spent resting. This contrasts with northern fur seals, Callorhinus ursinus, which typically spend 17% of their time at sea inactive. Apparently these northern seals can increase their foraging effort by increasing the proportion of time spent foraging. This would account for the observed between-year difference in at-sea FMR of C. ursinus while foraging-trip duration remained fairly constant.

Croxall, J. P.; Prince, P. A.; Ricketts, C. (1985)

Relationships between prey life-cycle and extent, nature and timing of seal and seabird predation in the Scotia Sea

In: Antarctic nutrient cycles and food webs

Siegfried, W. R.; Condy, P. R.; Laws, R. M., eds. Pages 516-533

Springer-Verlag, Berlin

ABSTRACT

The impacts of seals and breeding seabirds on prey resources in the Scotia Sea are assessed, and for seabirds seasonal changes are modelled. Although the data available are the best for any part of the Southern Ocean important deficiencies exist, which are summarized. Total consumption is estimated as 23×10^6 t, fairly evenly divided between seabirds and seals and between South Georgia and the rest of the area. Krill, Euphausia superba, (70%) is much more important than squid (16%) or fish (8%) but there are few data on winter diets. The main consumers of Krill are Crabeater Seals, Logodon carcinophagus, Macaroni, Eudyptes chrysolophus, and Chinstrap, Pygoscelis antarctica, penguins, with penguins accounting for 76% of the intake by birds. Peak demand is in February; consumption estimates are high compared with stock assessments and reasons for this are discussed. Despite variations between seasons, species, and sites, Krill (mostly sexually mature individuals, 35-45 mm long) taken by predators and by net-hauls in the southern Scotia Sea are generally similar, although the largest Krill are absent from predator samples. At South Georgia, however, most predators take 45-55 mm long Krill which are rare in net-hauls; possible causes and implications of this important discrepancy are discussed. The significance of differential predation on male and female Krill (bias favoring males in the south and females at South Georgia) is evaluated in terms of seasonal changes in energy content (due to lipogenesis), and it is suggested that the latter might be a factor influencing timing of breeding in certain predators. Other aspects of krill-predator interactions, including diving patterns and catching rates, are reviewed; more data on Krill demography and on seasonal variations in the age, sex and status of Krill taken by predators are required to assess critically predator-prey interactions. Elephant Seals, Mirounga leonina, take perhaps 75% of squid eaten by predators. A comparison of

quantitative data on squid diets of Sperm Whales, Physeter catodon, albatrosses and certain seals show the importance of Mesonychoteuthis, Moroteuthis and Kondakovia, although other taxa are as important for certain species (e.g., octopods for seals). All these squid are rarely caught by net-hauls and data from predators presently provide the best conspectus of the abundance of the Southern Ocean epi- and meso-pelagic squid of potential commercial importance. Fish are less important to predators than Krill and squid, although the total consumption (77% by seals, especially Elephant, Fur, Arctocephalus gazella, and Weddell, Lepotonychotes weddelli, seals) is a substantial proportion (e.g., at least 38% for Notothenia rossii at South Georgia) of estimated standing stocks, which may therefore be appreciably under-estimated. Seabirds (cormorants, penguins) mainly take immature nototheniids; Fur Seals mostly eat adult Champscephalus and are thus now in direct potential competition with commercial operations.

Croxall, J. P.; McCann, T. S.; Prince, P. A.; Rothery, P. (1988)

Reproductive performance of seabirds and seals at South Georgia and Signy Island, South Orkney Islands, 1976-1987: Implications for Southern Ocean monitoring studies

In: Antarctic Ocean and Resources Variability

Sahrhage, D., ed. Pages 261-285

Springer-Verlag, Berlin

ABSTRACT

Aspects of the reproductive performance over the last decade of Black-Browed, Grey-Headed and Wandering Albatrosses, Gentoo and Macaroni Penguins and Antarctic fur seals, at Bird Island, South Georgia and for Adelie and Chinstrap Penguins at Signy Island, south Orkney Islands, are summarized and reviewed. Breeding success of the Wandering Albatross, which breeds in winter and eats fish and squid, has remained constant, while population size has declined gradually but significantly. The other species at South Georgia, which breed in summer and feed extensively on krill, have shown major fluctuations in some or all of: breeding population size, breeding success, foraging trip duration and offspring growth rate. 1977-78 and 1983-84 were summers of particularly poor reproductive performance by almost all species; circumstantial evidence realint this to reduced availability of krill is discussed. The fluctuations in reproductive performance of the krill-eating, summer-breeding penguins at Signy Island are not synchronized with those at South Georgia; they correlate best (especially for Chinstraps, which suffered badly in 1980-81 and 1982-83) with the date of ice break-out in late spring. Numerous parameters of albatross, penguin and fur seal biology are reviewed in terms of their sensitivity and suitability for detecting changes in the marine environment.

Dahlheim, M. E. (1981)

A review of the biology and exploitation of the killer whale, Orcinus orca, with comments on recent sightings from Antarctica

Rep. int. Whal. Commn 31. 541-546.

ABSTRACT

Killer whales, Orcinus orca, are cosmopolitan in distribution. At present, a single species is recognized; however, various geographical races may exist. Population estimates are not available on a worldwide basis. Killer whales are usually found in pods and most activities appear to be group orientated. A population birth rate of 4-5% is suggested. Various other life history parameters of O. orca are discussed. Killer whales appear to be opportunistic feeders. Distributions of this species seems to be dependent upon the distribution and migration of prey items. A worldwide summary of the exploitation and utilization of this species is given for the years 1948-80.

Davis, M. B.; Renouf, D. (1987)

Social behaviour of harbour seals, Phoca vitulina, on haulout grounds at Miquelon

Can. Field Nat. 101 (1). 1-5.

ABSTRACT

A breeding colony of more than 700 harbour seals (Phoca vitulina) on the French island of Miquelon was observed during 1982. Behavioural interactions among different age and sex classes were documented, and the spatial organization of seals on the haulout grounds was assessed photographically. The study revealed a consistent hauling out pattern, and a predictable arrangement of animals on the beach, with a high degree of site tenacity. Our observations are discussed in light of existing theories of the social organization of harbour seal.

DeLong, R. L.; Antonelis, G. A. (1991)

Impact of the 1982-1983 El Nino on the northern fur seal population at San Miguel Island, California

In: Pinnipeds and El Nino: Responses to environmental stress

Trillmich, F.; Ono, K., eds. Pages 75-83

Springer-Verlag, Berlin

ABSTRACT

After 14 years of increasing production (following colonization), northern fur seal pup births declined at San Miguel Island in 1983 by 60% at Adams Cove and 64% at Castle Rock. Births increased at annual rates of 15 and 21% following the decline in 1983 but had not recovered to the 1982 level by 1987.

DeLong, R. L.; Antonelis, G. A.; Oliver, C. W.; Stewart, B. S.; Lowry, M. C.; Yochem, P. K. (1991)

Effects of the 1982-83 El Nino on several population parameters and diet of California sea lions on the California Channel Islands

In: Pinnipeds and El Nino: Responses to environmental stress

Trillmich, F.; Ono, K., eds. Pages 166-184

Springer-Verlag, Berlin

ABSTRACT

Births declined at all rookeries in the Southern California Bight in 1983. The magnitude of the decline in births varied among rookeries with decreases of 30% at San Miguel Island, 43% at San Nicolas Island, 62% at San Clemente Island, and 71% at Santa Barbara Island. In 1984 births increased on San Miguel Island, decreased further on San Nicolas Island, and remained low on Santa Barbara and San Clemente islands. In 1986 numbers of pups born on all islands were still 10% below the numbers born in 1982. The weights of male and female pups showed similar annual trends declining about 25% to 35% at San Miguel and San Clemente Islands from 1982 to 1983. Pups weights remained low in 1984 and did not return to 1982 values until 1985. It was not clear whether the declines in births resulted from reduced pregnancy rates among females, increased female mortality or both.

DeMaster, D. G.; Miller, D. J.; Goodman, D.; DeLong, R. L.; Stewart, B. S. (1982)

Assessment of California sea lion fishery interactions

Trans. North Am. Wild. Nat. Res. Conf. 47. 253-264.

ABSTRACT

California sea lions on the Southern California Channel Islands increased at rates of 4 to 6% from 1971 through 1981. A growth rate of about 5% per year is thought to be a minimum estimate of the maximum rate of population change for this species.

Engelhardt, F. R. (1982)

Hydrocarbon metabolism and cortisol balance in oil-exposed ringed seals, Phoca hispida

Comp. Biochem. Physiol. 72C (1). 133-136.

ABSTRACT

1. Ringed seals were exposed experimentally to oil contamination by feeding of a [¹⁴C] naphthalene marked crude oil in fish for up to 4 days at a rate of 5 ml/day. 2. Mixed function of oxygenase (MFO) activity, measured as aryl hydrocarbon hydroxylase in liver and kidney, was found to be induced, in particular in kidney tissue where the activity increased 3-fold. 3. MFO induction correlated with a high degree of conversion of crude oil hydrocarbons to water-soluble metabolites.

Most of the radioactivity was found in the polar fraction of plasma and urine. 4. Plasma cortisol levels were somewhat elevated by captive holding, and increased markedly after oil-exposure. Cortisol half-life decreased after oil exposure from 1 3/4 to 1 hr.

Everitt, R. D.; Braham, H. W. (1980)

Aerial Survey of Pacific harbor seals in the southeastern Bering Sea

Northwest Sci. 54 (4). 281-288.

ABSTRACT

Between June 1975 and June 1977, five aerial surveys were conducted along the eastern Aleutian Islands, and throughout Bristol Bay to study the distribution and abundance of the harbor seal (*Phoca vitulina richardsi*) during the breeding season. The number of group sightings and the total number of seals observed varied significantly with the tide height ($P < 0.01$). Fifty-seven percent more seals were observed on a low tide than in the same area surveyed near high tide. Three locations - Port Moller, Port Heiden, and Ginder River along the north side of the Alaska Peninsula - accounted for 78 percent of the study area population count, and for approximately 8.5 percent of the entire Alaska population estimate. A minimum abundance for the study area is estimated at 29,000 animals.

Fowler, C. W. (1988)

Population dynamics as related to rate of increase per generation

Evol. Ecol. 2. 197-204.

ABSTRACT

Inflection points in animal population growth curves (expressed as a fraction of equilibrium population levels) are correlated with rates of increase per generation. The inflection point declines with increasing rates of increase per generation. This apparent unifying principle of population dynamics is independent of phenomena related to body size since the rate of increase per generation is not correlated with body size. Species as diverse as whales and bacteria appear to conform to the pattern. More studies of the observed relationships are to be encouraged.

Gales, N. J.; Burton, H. R. Burton (1989)

*The past and present status of the southern elephant seal (*Mirounga leonina* Linn.) in Greater Antarctica*

Mammalia 53 (1).

ABSTRACT

Counts of southern elephant seals at the Vestfold Hills from 1958 to the present reveal that the population has declined by half to two thirds. The decline of this predominantly male, moulting

population is similar to reported declines of breeding populations of elephant seals in the Kerguelen province. The population of moulting seals at the Windmill Islands is larger than previously thought and its status is unknown. The Vestfold Hills' and Windmill Islands' populations are the only known aggregations of southern elephant seals in Greater Antarctica. The accessibility of suitable areas on the Antarctic coast for moulting during summer is shown to be the reason for selection of these sites.

Tagging studies show that elephant seals from the Vestfold Hills and the Windmill Islands migrate to and from the Kerguelen province, and that up to 40% of the seals moulting in the Vestfold Hills return to moult the following year. The Kerguelen Plateau and the Antarctic coast represent important foraging grounds for sub-adult, male elephant seals.

Geraci, J. R.; Smith, T. G. (1976)

Direct and indirect effects of oil on ringed seals (Phoca hispida) of the Beaufort Sea

J. Fish. Res. Board Can. 33. 1976-1984.

ABSTRACT

Ninety-six ringed seals (Phoca hispida) were taken from nets at Brown's Harbour, Northwest Territories in the fall of 1974. Comparison with two other new samples from 1971 and 1972 revealed a lower proportion of young-of-the-year and a lower mean weight of seals in all age-classes. Six seals immersed in Norman Wells crude oil for 24 h at the field netting site suffered only transient eye problems and minor kidney and possibly liver lesions; no permanent damage was observed. Three seals transported to the University of Guelph all died within 71 min after oil was introduced into their pool. Hematologic and blood chemical studies indicate that death was caused by oil superimposed on the stress of captivity. Six, 3-4 wk-old wild whitecoat harp seal (P. groenlandica) pups at the Magdalen Islands, Quebec, were coated with crude oil. No significant differences in core body temperatures were noted and no deleterious effects were observed. Five captive ringed seals at Guelph were subjected to a cumulative dosage of Norman Wells crude oil fed with their fish food. High dosage (75 ml) and low dosage (25 ml) of crude oil were also fed to two groups of six harp seal pups. No significant lesions or behavioral changes were noted. These experiments were of an acute nature and reflect the effects of a brief contact with oil only. Effects of longer contact as would probably be the case in an offshore oil well blowout situation are discussed. Possible effects of large-scale offshore oil fields are also considered.

Geraci, J. R.; St. Aubin, D. J.; Reisman, R. J. (1983)

Bottlenose dolphins, Tursiops truncatus, can detect oil

Can. J. Fish. Aquat. Sci 40. 1516-1522.

ABSTRACT

Two trained bottlenose dolphins, Tursiops truncatus, were tested for their ability to detect visually 12 different oils and 22 oil mixtures confined at the surface in small cylinders. Detection thresholds were established in tests using progressively lighter substances. The animals detected 6-mm slicks

of crude, residual, and refined motor oils, and diesel slicks thicker than 17 mm. They could not detect 6-mm thicknesses of leaded gasoline or transparent mineral oil. One dolphin's ability to detect oil improved with experience. While blindfolded, one dolphin could detect 12-mm-thick samples of two crude oils, Bunker C and mineral oil, but only when the latter two were churned and contained air bubbles. We conclude that dolphins detect the thicker concentrations of oil that occur near the source of an oceanic spill, but not lightly colored or refined products that tend to disperse into thin films.

Geraci, J. R.; St. Aubin, D. J. (1987)

Effects of offshore oil and gas development on marine mammals and turtles

In: Long-term environmental effects of offshore oil and gas development

Boesch, D. F.; Rabalais, N. N., eds. Pages 587-617

Elsevier Applied Science, London

ABSTRACT

During the past five years, studies on marine mammals have brought us closer to an understanding of basic behavioral and physiological responses to oil. For example, experiments have shown that dolphins can detect oil and, under certain circumstances, will avoid it. Oil can cause subtle damage to their skin, the full impact of which is still being assessed. The threat to otters and polar bears is unequivocal. Oiled fur is ineffective as an insulator, and attempts to groom can lead to oil ingestion. Fouling of baleen has short-term effects on water flow and feeding efficiency, although the consequences may not be as great as was predicted. Noise and disturbance associated with offshore production may be within the limits of tolerance for some species.

The full range of effects on turtles is poorly understood. Young turtles can eat tarballs which seal their mouths and interfere with normal feeding. Oil fouling of nests can lead to embryonic abnormalities and hatchling mortality. Turtles are particularly vulnerable to disturbances during the nesting season. The greatest impact of offshore oil and gas activities may result not from direct mortality, but rather through subtle alterations of habitat, in association with intrinsic stressors within the environment. We provide recommendations which reflect our interpretation of the most significant data gaps and emphasize the need for selective long-term monitoring.

Geraci, J. R.; Anderson, D. M.; Timperi, R. J.; St. Aubin, D. J.; Early, G. A.; Prescott, J. H.; Mayo, C. A. (1989)

Humpback whales (Megaptera novaeangliae) fatally poisoned by dinoflagellate toxin

Can. J. Fish. Aquat. Sci 46. 1895-1898.

ABSTRACT

During a 5-wk period beginning in late November, 1987, 14 humpback whales, Megaptera novaeangliae, died in Cape Cod Bay after eating Atlantic mackerel, Scomber scombrus, containing

saxitoxin (STX), a dinoflagellate neurotoxin responsible for paralytic shellfish poisoning in humans. We propose a line of evidence to explain how whales, by virtue of their diving adaptations, may be particularly vulnerable to this systemic neurotoxin. Absence of STX in New England waters and shellfish during the episode suggests that the mackerel, representing the northern stock which spawns in the Gulf of St. Lawrence, accumulated the toxin there and delivered it to the Gulf of Maine and Cape Cod Bay in the fall of 1987. These findings challenge common perceptions of the manner in which planktonic toxins move through the food chain, and offer new insights into natural mortality and strandings of marine mammals. It seems appropriate to search for STX and other phytotoxins when investigating marine mammal mortalities.

Geraci, J. R. (1990)

Physiologic and toxic effects on cetaceans

In: *Sea Mammals and Oil: Confronting the Risks*

Geraci, J. R.; St. Aubin, D. J., eds. Pages 167-198

Academic Press, New York

ABSTRACT

An oil spill at sea adds an element of risk to the environment of a whale or dolphin. Fresh crude oil or volatile distillates release toxic vapors that can damage sensitive tissues; harmful fractions may be swallowed or consumed through contaminated prey; and thicker tarry substances with entrapped debris may linger at the surface, plugging the vital baleen and digestive apparatus of whales that engulf them.

In spite of numerous observations of cetaceans in spills, none of these effects has been detected, or at least recorded with any certainty. Experimental evidence shows that dolphins can see oil at the surface and that they prefer to avoid it. Other cetaceans seem to be comparably equipped to detect oil. Yet in the wild, whales and dolphins have been observed swimming and feeding in its presence without apparent ill effect. Perhaps, in these instances, the stimulus was not noxious enough, or perhaps cetaceans disregard oil when they are engaged in more engrossing or important activities. Unlike furbearers, cetaceans do not lose heat through fouling of the skin. Furthermore, cetacean epidermis is nearly impenetrable, even to the highly volatile compounds in oil, and when skin is breached, exposure to these fractions does not impede the progress of healing. There is no evidence that oil or tar balls significantly foul the feeding apparatus of baleen whales; laboratory studies suggest that such fouling has only transient effects. Current technology provides the means to probe deeper- to the molecular level, if necessary- for damage by oil to cetaceans. Probing may satisfy our scientific curiosity, but would not bring us closer to an understanding of the central question. On the whole, it is quite improbable that a species or population of cetaceans will be disabled by a spill at sea, whatever the likelihood that one or a few animals might be affected or even killed. Yet some habitats, and therefore their residents, are more vulnerable than others. The ice-edge, refuge for bowheads, narwhals, and beluga whales, is a riskier trap for them than pelagic waters. And in coastal areas with bustling oil production activity, dolphins might be the unwitting sentinels of a deteriorating environment. The stage is now set for decisions to identify, wisely utilize, and monitor such habitats.

Gerrodette, T.; Gilmartin, W. G. (1990)

Demographic consequences of changed pupping and hauling sites of the Hawaiian Monk Seal

Conserv. Biol. 4 (4). 423-430.

ABSTRACT

During the last 30 years, changes in the size of Hawaiian monk seal populations at several locations have been associated with the amount and type of human disturbance. Recreational beach activities caused monk seals to alter their pupping and hauling patterns. Survival of pups in suboptimal habitats was low, leading to gradual population declines. During the last decade at Kure Atoll, the process has been reversed: human disturbance on beaches has decreased, and traditional pupping and hauling sites have been reestablished. Subsequently, high survival rates of young seals, coupled with two successful enhancement programs for female pups, have led to dramatic changes in the age and sex composition of the population. Based on these changes, the monk seal population at Kure Atoll soon should begin to increase. Apparently small behavioral changes in such vital activities as feeding and reproduction can have large demographic consequences. Therefore, monitoring of endangered species should include data on habitat use and age and sex composition, as well as estimates of abundance.

Glockner-Ferrari, D. A.; Ferrari, M. J. (1990)

Reproduction in the humpback whale (Megaptera novaeangliae) in Hawaiian water, 1975-1988: the life history, reproductive rates and behavior of known individuals identified through surface and underwater photography

In: Individual recognition of cetaceans: Use of photo-identification and other techniques to estimate population parameters. Report of the International Whaling Commission (Special Issue 12)

Hammond, P. S.; Mizroch, S. A.; Donovan, G. P., eds. Pages 161-169

International Whaling Commission, Cambridge

ABSTRACT

Humpback whales were studied in the waters off the west coast of Maui, Hawaii during the period 1975-88 using photographic techniques. Using surface and underwater photographs of color pattern characteristics, 584 adults and 268 calves were identified. Using photographs of the undersurface of the flukes, 210 adults and 2 calves were identified. Resighting histories were compiled for 69 individuals. Intervals between first and last sighting ranged from 1-13 years. One known male, observed in ten different years over a 13 year interval, was estimated to be a minimum of either 18 or 23 years of age (depending on certain assumptions) when last sighted. Of 34 resighted mothers, 31 produced more than one calf: 1 had seven calves, 1 had six calves, 2 had four calves, 6 had three calves and 21 had two calves. Maximum calving intervals were 1 year (n=7), 2 years (n=17), 3 years (n=8), 4 years (n=9), 5 years (n=5), 6 years (n=1), 7 years (n=1) and 9 years (n=2). Of the calves, 52.9% were males and 47.1% females. One male calf was later resighted at 6, 7 and 10 years of age. Evidence suggests that this individual reached sexual maturity by age

seven. A decrease in the occurrence of mothers and calves in nearshore waters off the west coast of Maui, Hawaii during the 1977-88 study period was demonstrated.

Godsell, J. (1988)

Herd formation and haul-out behavior in harbour seals (Phoca vitulina)

J. Zool., Lond. 215. 83-98.

ABSTRACT

Haul-out patterns of individually marked harbour seals Phoca vitulina during the breeding season indicated that herds which formed along the extensive beaches of Sable Island, Canada did not represent stable social units. There was large variation in the number, size and composition of herds and no segregation of seals by age, sex or breeding condition. Marked seals hauled out with many different individuals but not frequently nor consistently with the same ones. Pairs of seals which hauled out in the same herds for up to 30% of their times ashore generally occurred together in only one area, which suggests that associations between individuals were the consequence of similar site use rather than because of social cohesion. However, nursing females showed high levels of site fidelity; this behaviour probably serves as a safeguard against possible pup loss. Low levels of interactions between seals when ashore indicated that social relationships between adults were neither established nor maintained on land. Fights and displays may establish dominance relationships among breeding males while in the water but these did not appear to influence male haul-out patterns. Other than mother-pup relationships, social factors played little, if any, role in the formation or composition of harbour seal herds in a habitat where haul-out space was not limited.

Gosho, M. E.; Rice, D. W.; Breiwick, J. F. (1984)

The sperm whale, Physeter macrocephalus

Mar. Fish. Rev. 46 (4). 54-64.

ABSTRACT

This article reviews the status of the sperm whale, Physeter macrocephalus. The following topics are addressed: distribution and migration, life history and ecology, exploitation and population size, and management. Females attain sexual maturity at 9 yrs and produce a calf every 3-6 yrs. Males have a prolonged puberty, beginning at 9 yrs and ending at 20 yrs. Natural mortality rates range from 0.005-0.009/year.

Guerra C., C. G.; Portflitt K., G. (1991)

El Nino effects on pinnipeds in northern Chile

In: Pinnipeds and El Nino: Responses to environmental stress

Trillmich, F.; Ono, K., eds. Pages 47-54

Springer-Verlag, Berlin

ABSTRACT

The changes in abundance and distribution of the southern sea lion and the south American fur seal are documented along the northern Chilean coast during the 1982-83 El Nino.

Hamilton, P. K.; Mayo, C. A. (1990)

Population characteristics of right whales (Eubalaena glacialis) observed in Cape Cod and Massachusetts Bays, 1978-1986

In: Individual recognition of cetaceans: Use of photo-identification and other techniques to estimate population parameters. Report of the International Whaling Commission (Special Issue 12)

Hammond, P. S.; Mizroch, S. A.; Donovan, G. P., eds. Pages 203-208

International Whaling Commission, Cambridge

ABSTRACT

Right whale, Eubalaena glacialis, occurrence in Cape Cod and Massachusetts Bays was investigated from 1978 through 1986. Using photo-identification techniques, a total of 113 individual whales were identified with a maximum of 47 whales sighted during a calendar year. Although whales were sighted in all months of the year except December, peak abundance occurred from February through April. Mothers with calves consistently appeared in April or later. The mean calving interval of nine mature females was three years. Residency in the study area varied from one to 165 days. An unusual summer residency group, including nine mother/calf pairs, present in the study area from July through October 1986, is discussed.

Harkonen, T. (1987)

Seasonal and regional variations in the feeding habits of the harbour seal, Phoca vitulina, in the Skagerrak and the Kattegat

J. Zool., Lond. 213. 535-543.

ABSTRACT

Studies of the feeding of harbour seals have been carried out at the Tjarno Marine Biological Laboratory since 1977. The studies are based on fish otoliths found in faeces at seal haulouts. The

present paper compares feeding habitats at two different localities. Three families of fish, gadoids, pleuronectids and clupeoids were predominant in the seals' diet at a rocky shore habitat. Pleuronectids made up 75% of the diet at a sandy shore habitat. Temporal variations in feeding habits are also examined. The results indicate that harbour seals are opportunists in their choice of prey species, but some locally abundant species do not appear in the diet.

Härkönen, T.; Heide-Jorgensen, M. P. (1990)

Density and distribution of the ringed seal in the Bothnian Bay

Holarct. Ecol. 13. 122-129.

ABSTRACT

A modified strip census of basking ringed seals in the Bothnian Bay was carried out during the last week of April and the first week of May 1988. Of the total ice covered area, 23174 km², 3236 km² (14%) was covered by the transects. The mean density in the area was 0.101 ringed seals km², with substantial regional and local variation. Highest densities were found in the compact drift ice area. The estimated total population of ringed seals on the ice was 2093 in the Bothnian Bay proper and 248 in the North Quark. The 95% confidence limits of the estimates are $\pm 24\%$.

Härkönen, T. J.; Heide-Jorgensen, M. P. (1990)

Comparative life histories of east Atlantic and other harbour seal populations

Ophelia 32 (3). 211-235.

ABSTRACT

During the seal epizootic in 1988 tissue samples were collected from harbour seals, Phoca vitulina vitulina, that died in the Kattegat-Skagerrak area. The material is used for describing life history and estimating vital parameters of East Atlantic harbour seals. From a similar size at birth of 81 cm and 8.7kg, females and males become sexually mature at 127 and 130 cm, which is close to 85% of the asymptotic lengths of 146 and 156 cm. Asymptotic weight is 67 and 75 kg for females and males respectively, but shows considerable fluctuation during seasons of parturition, mating and moulting. During lactation females lose 26 kg (35%), and males decrease from 72 to 62 kg during the July mating period. Females have a mean age at sexual maturity of 3.72 years and a mean age at first parturition of 4.64. Males mature one year later. The overall pregnancy rate of females was 92% from age 3 to age 36, with lowered reproductive success after the age of 25 years. Ovulation occurs mainly during July, with a distinct peak during the last two weeks of the month. Maximum age for females was 36 and for males 34 years, and the sex ratio is skewed in favour of females in older segments. Based on an exponential rate of population increase ($r=0.11$) survival rates were calculated at 0.91 for adult males, whereas age-specific pregnancy rates were used to demonstrate possible combinations of juvenile and adult survival in females. The East Atlantic harbour seal is identical in the schedule of its reproductive cycle to the subspecies Phoca vitulina concolor and P. vitulina richardsi, but differs in seasonal timing of reproduction. Harbour seals in Alaska exhibit greater asymptotic length and weight, but no statistically significant differences in growth or reproduction could be detected between subspecies. The harbour seals in

the Kattegat-Skagerrak are slightly smaller than the other two subspecies and have a relatively high rate of reproduction.

Harvey, J. T.; Brown, R. F.; Mate, B. R. (1990)

Abundance and distribution of harbor seals (Phoca vitulina) in Oregon, 1975-1983

Northwest. Nat. 71. 65-71.

ABSTRACT

Harbor seals were observed on 32 haulout sites in Oregon during aerial counts conducted from 1975 to 1983; 90% were seen on 14 sites. The greatest number of seals seen on a haulout was 985 recorded at Cape Arago in July 1982. Counts of harbor seals in 1982 and 1983 were 38.6% greater than counts from 1975 to 1980. These data indicate an increase in numbers of harbor seals in Oregon, an increase corroborated by other information, namely, lower counts made before 1975 and increased use of new haulout sites since 1975. Between 1975 and 1983, numbers within bays increased, whereas numbers on most offshore rocks remained somewhat constant. Decreased harassment and mortality since implementation of the Marine Mammal Protection Act of 1972 doubtless allowed harbor seals to increase in numbers and to reoccupy protected haulout sites in bays.

Harwood, J. (1981)

Managing gray seal populations for optimum stability

In: Dynamics of large mammal populations

Fowler, C. W.; Smith, T. D., eds. Pages 159-172

John Wiley and Sons, New York

ABSTRACT

The gray seal population (as measured by pup production) in the United Kingdom increased 6-7% annually from the mid-1960s through the late 1970s. Data collected over the past several decades are used to derive life table parameters and population projection matrices for British grey seals.

Heide-Jorgensen, M. P.; Härkönen, T. J. (1988)

Rebuilding seal stocks in the Kattegat-Skagerrak

Mar. Mamm. Sci. 4 (3). 231-246.

ABSTRACT

The harbor seal (Phoca vitulina) population in the Kattegat-Skagerrak area has been dwindling for several centuries due to excessive hunting pressure. Corrected hunting statistics during 1890-1976

are used to estimate changes in population size over the past century. After protection was introduced in the 1960s and 1970s the harbor seal population in the area increased at an exponential rate of 0.12 and exceeded 5,000 animals in 1986. The present rate of population growth is used for modelling the influence of fertility and age-specific mortality. It is found that the observed high rate of increase is only realistic if female fertility rate is very high, the range of juvenile mortality rate is 0.33-0.52 and adult mortality is less than 0.15. Commonly cited higher mortality rates are not realistic in the Kattegat-Skagerrak area.

Heimlich-Boran, J. R. (1988)

Behavioral ecology of killer whales (Orcinus orca) in the Pacific Northwest

Can. J. Zool. 66. 565-578.

ABSTRACT

Killer whales (Orcinus orca) were found to use different physio-graphic regions of their habitat in unique ways. Resident whales fed more in areas of high relief subsurface topography along salmon migratory routes, and may use these geographic features to increase feeding efficiency. Transient whales fed in shallow protected areas around concentrations of their prey, harbor seals (Phoca vitulina). Whales traveled across deep, featureless areas in moving from one feeding area to another. Whales rested depending on the previous sequence of behaviors and played in open water areas or adjacent to feeding areas. The location of food resources and habitats suitable for prey capture appears to be the prime determining factor in the behavioral ecology of these whales. These patterns of behavior most likely represent cultural mechanisms that have been learned through trial and error experiences leading to successful foraging strategies.

Helle, E. (1980)

Lowered reproductive capacity in female ringed seals (Pusa hispida) in the Bothnian Bay, northern Baltic Sea, with special reference to uterine occlusions

Ann. Zool. Fennici 17. 147-158.

ABSTRACT

The reproduction of female ringed seals, Pusa hispida (Schreber), was studied in 225 specimens trapped with seal nets in October-November at Simo, on the northern Bothnian Bay (65°35'N, 25°00'E), and 28 specimens shot on the ice of the Bothnian Bay in April-May, 1973-79.

The ovulation frequency in mature specimens was 0.984. The sizes of the corpora lutea and c. albicantia were not dependent on maternal age. The c. luteum of the pregnant females was larger on average than the regressing c. luteum or c. albicans of the non-pregnant females in October-November, some 4 months after implantation. The proportion of normal pregnant females averaged 28%, decreasing from a maximum of 52% in the group aged 7-8 years to 8% in the females over 25 years of age. Females with normal ovulation but no macroscopic uterine signs of pregnancy, i.e. a missed pregnancy, were almost as common, reaching a maximum of 68% among the newly matured females aged 5-6 years, and averaging 23% thereafter. An average of 42% had

an occlusion in one or both of the uterine horns, the proportion of both single (26%) and bilateral occlusions (16%) increasing with age. This membranous occlusion sealed up the uterine tract, forming a closed chamber with varying amounts and types of fluid. The average position of the occlusion was just over 60 mm from the tip of the horn, ranging from near the tip to the bifurcation. Other reproductive failures averaged 5%. Major changes took place in the reproductive status even during the period 1974-79, with the proportion of pregnant females dropping from over 30% to under 20% and cases of occlusion increasing from 35% to 59%. The numerous reproductive failures, possibly linked with hormonal imbalance and/or severe infections, at present form the most serious threat to the future of the Baltic seal population.

Hellou, J.; Huang, Y. S.; Upshall, C.; Ni, I. H.; Payne, J. F. (1991)

Polycyclic aromatic hydrocarbons in harp seals (Phoca groenlandica) from the Northwest Atlantic

Arch. Environ. Contam. Toxicol. 21. 135-140.

ABSTRACT

There is virtually no information available on concentrations of polycyclic aromatic hydrocarbons (PAH) in seals from any of the world's oceans. The largest harp seal population in the world is found in Canadian waters of the Northwest Atlantic. Samples of muscle tissue obtained from twenty eight harp seals ranging in age from foetuses to animals 22 years old were analyzed for total PAH and lipid content. Concentrations were determined in terms of crude oil and chrysene equivalents in line with recommendations of the International Oceanographic Commission. Overall,, relatively low concentrations were found, the highest values being less than 1 ppm in terms of chrysene equivalents and 4 ppm in terms of petroleum hydrocarbon equivalents. There was no evidence of bioaccumulation with age, the concentrations in juvenile seals (1-5 years) being higher than concentrations in older animals (6-20 years). There was also no correlation between PAH concentration and fat content. There is a possibility that the animals having elevated levels of PAH are from the Gulf of St. Lawrence herd, indicating the importance of obtaining more information on PAH levels on marine mammals and other organisms from this and similar regionally contaminated seas.

Hes, A. D.; Rouse, G. P. (1983)

Population increase in the sub-Antarctic fur seal Arctocephalus tropicalis at Amsterdam Island

S. Afr. J. Antarct. Res. 13. 29-34.

ABSTRACT

The population of fur seals at Amsterdam Island increased 7.8% per year between 1955 and 1969.

Hindell, M. A. (1991)

Some life-history parameters of a declining population of southern elephant seals, Mirounga leonina

J. Anim. Ecol. 60. 119-134.

ABSTRACT

Mark-resight data were analysed for thirteen cohorts from a declining population of southern elephant seals branded at Macquarie Island between 1951 and 1965. First year survival was essentially stable during the 1950s at about 46% for females and 42% for males. There was a dramatic fall in first year survival during the 1960s, declining to less than 2% for both sexes in 1965. Post-year-1 survival did not change between the 1950s and the 1960s. Comparisons with a stable population of southern elephant seals at South Georgia indicated that both first year and adult survival were lower in the Macquarie Island population. There were no changes in the age at first breeding of the Macquarie Island seals during the study, but this was on average 1 year later than at South Georgia. It is hypothesized that the current decline in elephant seal numbers at several of their major breeding islands is due to the populations returning to pre-sealing levels after they had risen to abnormally high levels with the end of commercial exploitation early this century. Possible tests of the hypothesis include studying the diet and foraging behaviour of southern elephant seals to gain an understanding of the predator-prey relationships, continuing to census the Macquarie Island population to determine if the population levels out at around the estimated pre-sealing levels, and monitoring northern elephant seal populations which were also severely exploited but are currently increasing rapidly.

Hoelzel, A. R.; Ford, J. K. B.; Dover, G. A. (1991)

A paternity test case for the killer whale (Orcinus orca) by DNA fingerprinting

Mar. Mamm. Sci. 7. 35-43.

ABSTRACT

The minisatellite DNA profiles from four captive killer whales (two adult males, a female and her calf) were compared to determine paternity between two potential fathers. One of the males was clearly excluded, while the other shared all paternal bands with the calf. The background of this technique, and its potential applications in captive breeding programs and field studies are discussed.

Hoelzel, A. R.; Dover, G. A. (1991)

Genetic differentiation between sympatric killer whale populations

Hered. 66. 191-195.

ABSTRACT

The genetic variation within and between putative Killer whale (Orcinus orca) populations was examined by DNA fingerprinting nuclear genomes and sequencing the D-loop region of the

mitochondrial genome. Mitochondrial DNA variation indicated that two sympatric populations in the northeastern Pacific were as genetically distinct as North Pacific populations from a South Atlantic population. The two sympatric populations are known to pursue different foraging strategies. DNA fingerprinting showed very low levels of variation within populations relative to comparisons between allopatric populations, suggesting inbreeding. These results are consistent with predictions about the genetic structure of Killer whale populations based on behavioural observations and variation in colour morphology.

Hoover, A. A. (1988)

Harbor seal

In: Selected marine mammals of Alaska

Lentfer, J. W., ed. Pages 125-157

Marine Mammal Commission, Washington, D.C.

ABSTRACT

The life history and population biology of harbor seals in Alaska are reviewed and conservation, management, and research issues are discussed.

Jefferson, T. A. (1990)

Status of Dall's porpoise, Phocoenoides dalli, in Canada

Can. Field Nat. 104 (1). 112-116.

ABSTRACT

Dall's porpoise, Phocoenoides dalli, is one of the most commonly sighted cetaceans throughout its range in temperate waters of the North Pacific Ocean and surrounding seas. Dall's porpoises are common both offshore and in deep inshore waters of British Columbia. There appear to be few serious conservation problems in the eastern Pacific at present, although little is known of the behavior and ecology of this species. Probably the major threat facing this species in Canada is environmental contamination by such substances as organochlorines and heavy metals. Much more research is needed before these threats can be properly assessed but, in the meantime, a conservative approach to porpoise management is suggested.

Jones, M. L. (1990)

The reproductive cycle in gray whales based on photographic resighting of females on the breeding grounds from 1977-82

In: Individual recognition of cetaceans: Use of photo-identification and other techniques to estimate population parameters. Report of the International Whaling Commission (Special Issue 12)

Hammond, P. S.; Mizroch, S. A.; Donovan, G. P., eds. Pages 177-182

International Whaling Commission, Cambridge

ABSTRACT

Gray whales (Eschrichtius robustus) with distinctive natural markings were systematically photographed in San Ignacio Lagoon, Mexico from 1977 to 1982. In this paper, information is presented on breeding cycles for individually known females, including the range of values observed for length of calving interval and the relative frequencies of different length calving intervals (expressed in years). About 6,000 photographs were taken and 562 different gray whales were identified. Among these were 55 sexually mature females that were followed through 2 to 6 seasons on their winter breeding grounds; they produced a total of 115 calves over the 6-year period.

The length of time between the birth of consecutive calves was documented for 42 cows. Calving intervals ranged from 1-4 years, but were predominantly 2 years (1 calf every other year). The observed intervals were: 1 year (n=1), 2 years (n=48), 3 years (n=6) and 4 years (n=5). The mean length of the calving interval, or breeding cycle, for the population from 1977-82, was estimated as 2.11 (SD=0.403) years.

Jonsgard, A.; Lyshoel, P. B. (1970)

A contribution to the knowledge of the biology of the killer whale Orcinus orca (L.)

Norw. J. Zool. 18 (1). 41-48.

ABSTRACT

Biological data from 1413 killer whales caught by Norwegian whalers in northeastern North Atlantic waters in the period 1938-1967 are examined. Four more killer whales were examined by two Norwegian biologists in 1967. The distribution and migration of killer whales in these waters seem to be dependent upon the distribution and migration of the herring. Mammals are eaten by larger killer whales. Females and males seem to attain sexual maturity when about 16 and 19 feet long respectively. The young are very close to 7 feet in length at birth. The breeding season may stretch over several months, although there is some evidence that relatively more calves are born in late autumn and winter. In length, adult females and males seldom exceed 26 and 30 feet respectively.

Kannan, N.; Tanabe, S.; Ono, M.; Tatsukawa, R. (1989)

Critical evaluation of polychlorinated biphenyl toxicity in terrestrial and marine mammals: increasing impact of Non-ortho and Mono-ortho coplanar polychlorinated biphenyls from land to ocean

Arch. Environ. Contam. Toxicol. 18. 850-857.

ABSTRACT

Residues of potentially toxic non-ortho chlorine substituted coplanar 3,3',4,4'-tetra-3,3',4,4',5-penta-, 3,3',4,4',5,5'-hexachlorobiphenyl and their mono- and di-ortho analogs (2,3',4,4',5-penta-, 2,3,3',4,4'-penta-, 2,3,3',4,4',5-hexa and 2,2',3,3',4,4'-hexa-, 2,2',3,4,4',5-hexachlorobiphenyl) were determined in humans, dogs, cats (terrestrial), a finless porpoise (*Neophocoena phocaenoides-coastal*), Dall's porpoises (*Phocoenoides dalli, dalli*), Baird's beaked whales (*Berardius bairdii*) and killer whales (*Orcinus orca*-open ocean). Among the coplanar polychlorinated biphenyl (PCB) congeners, the concentration of the di-ortho congeners was the highest and the non-ortho congeners was the lowest. However, all three coplanar PCBs occurred at significantly higher levels than toxic polychlorinated dibenzo-p-dioxins (PCDDs) and polychlorinated dibenzofurans (PCDFs). The relative bioconcentration and metabolic capacity of terrestrial and marine mammals to these chemicals, suggest that the toxic threat of coplanar PCBs increases from land to ocean, but the reverse is true for PCDDs and PCDFs. The toxic threat of coplanar PCBs to higher aquatic predators such as cetaceans was principally assessed by 2,3,7,8-T4CDD Toxic Equivalent Analysis which is based on the induction of arylhydrocarbon hydroxylase (AHH) and ethoxyresorufin (E)-deethylase (EROD). Analysis indicates, in particular, that the bioaccumulation of toxic 3,3',4,4',5-penta- and 2,3,3',4,4'-pentachlorobiphenyls in carnivorous marine mammals is a cause for considerable concern.

Kasuya, T.; Marsh, H. (1984)

Life history and reproductive biology of the short-finned pilot whale, Globicephala macrorhynchus, off the Pacific coast of Japan

In: Reproduction in whales, dolphins and porpoises. Reports of the International Whaling Commission (Special Issue 6)

Perrin, W. F.; Brownell, R. L. Jr.; DeMaster, D. P., eds. Pages 259-310

International Whaling Commission, Cambridge

ABSTRACT

After a period of heavier exploitation in the 1940s, the short-finned pilot whale has been hunted at a lower rate of several hundreds per year by a small-type-whaling and drive fishery off the Pacific coast of Japan. Age data from 373 females and 170 males obtained from 27 schools which were stranded or driven during the 17-years period from 1965 cover all months except March, April, September and November. Breeding is diffusely seasonal, with a single parturition peak in July-August. A single calf is born at a mean body length of 140 cm after 14.9 months gestation and nursed for a minimum of about two years. Calves of older cows may be nursed for considerably longer than this. Females mature at 7-12 ($X=9.0$) years, produce an average of four to five calves,

and have their last calf before age 40 years, even though they may live up to 63 years. In contrast males have a maximum longevity of only 46 years and probably continue to be capable of reproduction until death. In males, puberty begins at 7 to 17 ($X=14.6$) years and social maturity at an average of 17 years.

The age composition suggests that the total mortality rate is lowest in the post-pubertal stage and that it increases after age 28 (male) or age 46 (female). Males have a higher total mortality rate than females at any given age. The juvenile total mortality rate is probably higher than that of post-pubertal animals. These differences in total mortality rates may reflect differences in natural mortality rates. Using an hypothetical stationary population model, we estimate that the total annual mortality rate over all age classes is 8.3% (male) and 4.5% (female). Thus there are more reproductive females than adult males. The mating system is polygynous. Males may migrate between schools after weaning. However, females probably stay in their mother's school for life, so that the breeding schools are essentially matrilineal kinship groups.

Kerley, G. I. H. (1983)

Relative populations sizes and trends, and hybridization of fur seals Arctocephalus tropicalis and A. gazella at the Prince Edward Islands, Southern Ocean

S. Afr. J. Zool. 18. 388-392.

ABSTRACT

Fur seals were counted at the Prince Edward Islands during the 1981/1982 austral summer. Classified counts, adjusted for pup undercounting and mortality, pregnancy rate and seasonal haulout patterns, of Arctocephalus tropicalis and A. gazella are presented. These populations have entered a phase of rapid growth, as indicated by higher rates of population growth than previously found. The possible role a A. gazella immigration is unknown. There has been an increase in the number of breeding localities used by both species. At present the extent of hybridization between these two species appears limited, possibly by behavioural, ecological and genetic processes.

Kovacs, K. M.; Jonas, K. M.; Welke, S. E. (1990)

Sex and age segregation by Phoca vitulina concolor at haul-out sites during the breeding season in the Passamaquoddy Bay region, New Brunswick

Mar. Mamm. Sci. 6 (3). 204-214.

ABSTRACT

The size and composition of groups of harbor seals at two haul-out sites were studied during the breeding season of 1989, in the Passamaquoddy Bay region of Atlantic Canada. Evidence of segregation both by age and sex was found in the distinct composition of the two groups. One group contained mainly males and no pups, and the other had a sex ratio not significantly different from one and contained pups. The proportion of females increased at the nursery site with the onset of

birthing in the region while the proportion of males increased through the breeding season at the other site. No increase in the number of adults, in total, was detected over the study period, suggesting that sexual segregation and not a change in haul-out frequency was responsible for the disparity in the sex structure of the two groups. The proportion of juveniles was significantly greater at the male dominated site than at the nursery site.

Krieber, M.; Barrette, C. (1984)

Aggregation behaviour of harbour seals at Forillon National Park, Canada

J. Anim. Ecol. 53. 913-928.

ABSTRACT

(1) In order to explain the aggregation behaviour of harbour seals (*Phoca vitulina*) at their landing sites, two possibilities were investigated: grouping in response to resting site distribution, or in response to changes in individual time-energy budget. (2) To verify if seals aggregated according to the distribution of their landing sites, we studied their site selection. Although seals select sites with distinctive features, they frequented only those whose area was sufficient or that were close enough to other suitable sites to hold a tight resting group. Hence, the distribution of landing sites was ruled out as the sole factor explaining aggregation. (3) By a closer look into time budgeting, we found that membership in a group generated an individual benefit. This takes the form of an increased time allocation to sleep in larger and tighter groups. While individual scanning rate decreased with group size and proximity, overall vigilance increased with group size and decreased with proximity. Although evidently desiring to join larger and tighter groups, seals saw their density regulated on each site by aggressive interactions. Rate of aggressive interaction did not vary with density, but the proportion leading to departures increase with density. No fighting advantage was found in larger size, but the first animal to land on the site was generally the winner. (4) Changes in the landing behaviour in relation to weather were consistent with other studies. However, seasonal weather trends could not fully explain the increase in group size from spring to autumn, during which time the proportion of used space remained roughly constant.

Lander, R. H. (1981)

A life table and biomass estimate for Alaskan fur seals

Fish. Res. 1. 55-70.

ABSTRACT

Data on growth rates (8% per year, 1911-1924), adult female survival, immature female survival, and fecundity are presented for *Callorhinus* at the Pribilof Islands.

Laws, R. M. (1992)

History and present status of southern elephant seal populations.

In: Elephant seals

Le Boeuf, B. J.; Laws, R. M., eds. Need Page Numbers

University of California Press, Berkeley

ABSTRACT

The population history (sealing) and recent trends are reviewed for southern elephant seal (*Mirounga leonina*) colonies. The colony at South Georgia has remained stable over the past several decades, one new colony on the South American Atlantic coast has increased, whereas colonies at many other locations have, for unknown reasons, declined.

Leatherwood, S.; Bowles, A. E.; Krygier, E.; Hall, J. D.; Ignell, S. (1984)

Killer whales (Orcinus orca) in southeast Alaska, Prince William Sound, and Shelikof Strait; a review of available information

Rep. int. Whal. Commn 34. 521-530.

ABSTRACT

Information on killer whales is summarized from a formal sightings network, 1976-1981 (Southeast Alaska), aerial surveys, 1976-1978 and small boat surveys, 1976-1982 (Prince William Sound), aerial surveys, 1982-1983 (Shelikof Strait), and interviews with marine scientists, fishermen and other knowledgeable mariners (all areas). Although present year-round, killer whales in each area increase in numbers and concentrate in specific areas in late summer through early fall, apparently in response to concentrations of salmon. In Prince William Sound, one group apparently remains within a limited home-range (less than 25 km radius) in Knight Island Passage each summer. Minimum counts from portions of the three areas are 93, 80 and 66 whales, respectively. Data indicate concurrent presence of killer whales in other unstudied portions of all three areas; so, 'populations' in each area certainly number in excess of 100 animals. Herds included an average of 4-6 animals (n=890) in Southeast Alaska. Of the three areas surveyed, Prince William Sound and Southeast Alaska appear most suitable as areas for long-term observational studies of killer whales of the sort pioneered in inland waters of Washington and British Columbia.

Leatherwood, S.; Matkin, C. O.; Hall, J. D.; Ellis, G. M. (1990)

Killer whales, Orcinus orca, photo-identified in Prince William Sound, Alaska, 1976 through 1987

Can. Field Nat. 104 (3). 362-371.

ABSTRACT

Individual Killer Whales, Orcinus orca, were identified from photographs taken incidental to other research (1976-1983) or during intensive studies of the species (1984-1987) in Prince William Sound (PWS), Alaska. The 232 identified animals were grouped into pods, based on observed associations, and assigned to the following conservatively defined age/sex categories: adult males (52, 22.41%), adult females (22, 9.48%), calves born in 1986 or 1987 (9, 3.9%) or 'immatures/others' of both sexes (149, 64.22%). Because of imprecision, this last category may include some adult females, as an animal was only classified as adult female if it was closely associated with a calf, and some recently matured adult males. Age/sex composition in PWS is similar to that noted for other areas of the eastern North Pacific and for Iceland, but caution is indicated in using such data. There were at least 13 births in PWS from 1985 through 1987. In 1984, calves were grouped with 'juveniles'. The 4 newborn calves first detected in 1985 are now classified as 'immature/others'. The combined mortality rate of 1.9% in three pods (AB, AE, AI) is similar to that in British Columbia and Washington (1.7 percent), while that in AB pod (7.4 percent) is significantly higher. Members of AB pod have been interfering with a longline fishery for Sablefish (Blackcod), Anoplopoma fimbria, since 1985. At least 10 (possibly 14) members of AB pod were reported wounded by gunshot between late 1984 and fall 1986, some apparently fatally. There were no new bullet wounds detected in 1987. Although no carcasses were found, from 1986 to 1987, there were apparently 7 deaths in AB pod, 2 in AE pod, and one each in AK and AN pods, leaving a documented minimum population of 221 whales in PWS by the end of the 1987 season.

Loughlin, T. R.; Rugh, D. J.; Fiscus, C. H. (1984)

Northern sea lion distribution and abundance: 1956-80

J. Wildl. Manage. 48 (3). 729-740.

ABSTRACT

The present distribution and abundance of the northern sea lion (Eumetopias jubatus) was determined from surveys made between 1975 and 1980 and compared to estimates made approximately 20 years earlier. The previous population estimate of 240,000-300,000 is similar to our estimate of 245,000-290,000. Declines appear to have occurred in the eastern Aleutian Islands, Pribilof Islands, and near Kodiak Island in the Gulf of Alaska. Increases may have occurred principally in the western and central Aleutians and west of Kodiak Island. Seasonal movements and indirect and direct effects by humans are considered the likely causes for regional changes in distribution and abundance.

Loughlin, T. R.; Nelson, R., Jr. (1986)

Incidental mortality of northern sea lions in Shelikof Strait, Alaska

Mar. Mamm. Sci. 2 (1). 14-33.

ABSTRACT

The incidental catch of northern sea lions (Eumetopias jubatus) in the walleye pollock (Theragra chalcogramma) joint-venture fishery in Shelikof Strait, Alaska, was studied during 1982-1984 to assess the nature and magnitude of the catch. Data were obtained by placing U.S. observers on foreign processing vessels. Dead sea lions recovered from trawl nets were counted, sexed and measured; teeth were removed for age determination by dental laminae; and stomach contents were analyzed. Although the fishery has continued to expand both in number of boats and estimated total catch (74,136 metric tons [t] in 1982 to 171,539 t in 1984), the estimated incidental catch of northern sea lions has declined (ranging from 958 to 1,436 in 1982, 216 to 324 in 1983 and 237 to 355 in 1984). Of the sea lions processed, 73 percent were caught between 2000 and 0500 h, probably during net retrieval. Most caught sea lions were females ranging in age from 1-25 yr with a mean age of 6.43 yr; 79 percent of the females were sexually mature and probably part of the reproducing population. Males had a mean age of 4.8 yr and only 12 percent were old enough to obtain and defend territories. Analysis of stomach contents showed that the sea lions consumed pollock the same size as that taken by the commercial fishery. The impact of the incidental catch on the Gulf of Alaska sea lion population is unknown.

Loughlin, T. R.; Merrick, R. L. (1988)

Comparison of commercial harvest of walleye pollock and northern sea lion abundance in the Bering Sea and Gulf of Alaska

In: Proceedings of the International Symposium on Biological Management (89-1)

Melteff, B., ed. Pages 679-700

Alaska Sea Grant, Anchorage

ABSTRACT

Northern sea lions are declining in most of Alaska and the synergistic effects of commercial fisheries have been mentioned as one plausible reason for the decline. We examined walleye pollock commercial fishing catch data for 1971-1986 to assess the possible role of the indirect effect of fish removal on northern sea lion abundance trends in the Bering Sea and Gulf of Alaska. The commercial catch in areas near major sea lion rookeries was compared with trends in sea lion abundance and correlation coefficients calculated. Results from the analysis were inconclusive although in some areas, such as the eastern Aleutian Islands and central Gulf of Alaska, high correlation between the amount of fish caught and sea lion abundance trends were apparent, while in other areas the correlation was equivocal.

Loughlin, T. R.; Miller, R. V. (1989)

Growth of the northern fur seal colony on Bogoslof Island, Alaska

Arctic 42 (4). 368-372.

ABSTRACT

Northern fur seal, Callorhinus ursinus, pups were first observed on Bogoslof Island, southeast Bering Sea, in 1980. By 1988 the population had grown at a rate of 57%/year to over 400 individuals, including 80+ pups, 159 adult females, 22 territorial males, and 188 subadult males. Some animals originated from rookeries of the Commander Islands, whereas others are probably from the Pribilof Islands. In 1983 and 1985 over 50% of the females were estimated to be >6 years of age, based on vibrissae color. The rookery is in the same location where solitary male fur seals were seen in 1976 and 1979 and is adjacent to a large northern sea lion rookery.

Lowry, L. F.; Frost, K. J. (1981)

Feeding and trophic relationships of phocid seals and walruses in the Eastern Bering Sea

In: The eastern Bering Sea shelf: oceanography and resources, Vol. 2

Hood, D. W.; Calder, J. A., eds. Pages 813-824

Office of Marine Pollution Assessment, NOAA, BLM, Seattle

ABSTRACT

Recent data on food habits of five species of phocid seals and walruses (Odobenus rosmarus) in the eastern Bering Sea are reviewed. Harbor seals (Phoca vitulina richardsi), spotted seals (Phoca largha), and ribbon seals (Phoca fasciata) all feed to a large extent on pelagic and semidemersal fishes. Demersal fishes are eaten by all three of these species but appear to be of greatest importance in the diet of ribbon seals. Crustaceans and octopus (Octopus spp.) are also eaten. Ringed seals (Phoca hispida) also feed on pelagic and semidemersal fishes. Crustaceans make up a considerable portion of the diet of ringed seals, especially young animals. Bearded seals (Erignathus barbatus) and walruses feed primarily on benthic organisms. Walruses feed almost exclusively on clams. Clams, crabs and shrimp make up the bulk of the diet of bearded seals. Geographical, seasonal, year-to-year, and age-related variations in feeding are evident in all species for which sufficiently large samples have been examined. Harbor, spotted, ribbon, and ringed seals all depend primarily on a pelagic food web and compete for food with one another and with fur seals (Callorhinus ursinus), sea lions (Eumetopias jubatus), cetaceans, and seabirds. Walruses and bearded seals compete for clams in some areas. Gray whales (Eschrichtius robustus) feed on benthic crustaceans, which are also eaten by bearded seals and walruses. Commercial fisheries harvest a portion of the food resource of Bering Sea pinnipeds which may influence populations of some species. Available data on foods of phocid seals and walruses are inadequate in all seasons and in all regions except the northern Bering Sea. Data are lacking for all species in southern and central regions during winter months, for walruses in all areas and seasons, and for harbor seals in the southeastern Bering Sea.

Lowry, L. F.; Frost, K. J.; Loughlin, T. R. (1988)

Importance of walleye pollock in the diets of marine mammals in the Gulf of Alaska and Bering Sea, and implications for fishery management

Proc. Int. Symp. Biol. Mgmt. Walleye Pollock Nov. 1988, Anchorage, Alaska.

ABSTRACT

Approximately 31 species of marine mammals occur at least seasonally in portions of the Gulf of Alaska and Bering Sea. Walleye pollock are eaten to some degree by euryphagous baleen whales, including fin, minke, humpback, and sei whales, and have been found in the stomachs of sperm whales and Dall's porpoise. Pollock may be a seasonally important food for belukha whales, harbor porpoises, and killer whales. Pollock are the most important food (35% of total energy intake) of northern fur seals in the eastern Bering Sea in summer and are also eaten in the Gulf of Alaska. Pollock composed 58% of the stomach contents of Steller sea lions and 21% of the stomach contents of harbor seals collected in the Gulf of Alaska, and are also important prey in the Bering Sea. Pollock are important foods for spotted and ribbon seals when they are associated with the Bering Sea ice front during March-June, and are sometimes eaten by bearded seals.

Foraging activities of marine mammals may affect walleye pollock populations by: (1) influencing abundance of certain size/age classes directly through predation; (2) influencing the productivity of pollock populations by feeding on the same prey base (e.g., copepods, euphausiids, and forage fishes); and (3) preying on species that are competitors or predators of pollock. Conversely, pollock fisheries may affect marine mammals by altering the abundance and age-class structures of pollock stocks and incidentally killing marine mammals during fishing activities. Available data are not adequate to accurately model or monitor interactions between marine mammals, pollock populations, and pollock fisheries. Clearly both fisheries and marine mammals remove large amounts of pollock from a complex and dynamic ecosystem. Changes in population status of marine mammals, whether or not they are directly caused by fishing activities, may have major consequences for fishery management.

Majluf, P. (1991)

El Nino effects on pinnipeds in Peru

In: Pinnipeds and El Nino: Responses to environmental stress

Trillmich, F.; Ono, K., eds. Pages 55-74

Springer-Verlag, Berlin

ABSTRACT

The foraging behaviour, diet, pup growth and mortality of the fur seals and sea lions in Peru under EN and non-EN conditions are compared. During the EN, female fur seals had to dive more often to greater depths and they spent longer times at sea foraging. They were still not able to obtain

enough food since they and their young were in poor condition and suffered higher mortality. Sea lions were similarly affected.

McLaren, I. A.; Smith, T. G. (1985)

Population ecology of seals: retrospective and prospective views

Mar. Mamm. Sci. 1 (1). 54-83.

ABSTRACT

This review focuses on population ecology, with critical accounts of past work and future possibilities in age determination, body growth and condition, estimating abundances, mortality rates and lifespans, reproduction, comparative life histories, population dynamics, population modelling and seals in ecosystems. We suggest ways to reduce errors in age determination and to improve methods of obtaining and presenting growth data. Generalized van Bertalanffy growth equations are promoted as a basis for analyzing species differences and intra-population variation in body lengths. Indices other than blubber thickness may be better for following body condition. Catch-effort and survival-index methods of estimating abundances have limited applicability, total counts are only locally useful, and sample counts may only be accurate for scattered, ice-breeding species. Some new techniques for population indices are promising. Pre-adult mortality remains difficult to assess. Although not always recognized, adult mortality rates do increase with age, as well described by Gompertz functions. Existing estimates of lifespans are unreliable, and a new approach is outlined. There are methodological problems in estimating ages of maturity. Corpora albicantia should not be used for back-extrapolation, and more study is needed of use of teeth annuli as indicators of maturity. Age-specific proportions of females parous based on reproductive tracts may disagree with proportions recruited in breeding groups, suggesting that the former may often be in error. Allometric relationships among body sizes and life-history variables need more reliable data, especially since the residuals of such relationships are of greater interest. Brain size may be a better scalar. Direct evidence of density dependence in population growth of seals is sparse. Early survival has been more widely shown to be density-dependent, but only among polygynous species where crowding on land may be a byproduct of sexual selection; there is as yet no good evidence of trophic restraints. Evidence of density dependence of ages of maturity is generally unconvincing. Predation, especially by sharks, may be critical in some species. Characteristics of equilibrium populations might profitably be sought in mass remains in middens and historic kill sites. More attention should be paid to the search for density-dependent influences. Supposed impacts of fisheries and pollutions are not wholly convincing. Natural epidemics may keep some populations below resource or space saturation, and some high-latitude species may show large year-to-year variations in recruitment and abundances. Evidence for such density-independent effects should be sought in residuals of growth curves and in teeth layers. Although surplus yield and production/biomass models have been tried, realistic pinniped models must be completely age-structured and time-dependent. Simple models have questionably assumed stationarity to derive life-history parameters. The best available estimates of density dependence of such parameters give no resolution when extrapolated toward equilibrium, and only limited efforts have been made to introduce stochasticity. Better data, not improved model structures, are needed for better understanding. Recent work has contradicted the assumed voraciousness of seals, but their system impacts and dependencies are not well understood. Extended Lotka-Volterra equations used to model Antarctic food webs, including seals, are merely heuristic. Fixed seal biomasses enter as

top-down, driving functions in a Bering Sea model, which accordingly cannot be used to analyze or manage their populations. Some Soviet models are tantalizing but ill-specified. The introduction of harbor seals in well-chosen lakes might give more insights into system roles than would more elaborate modelling. We wonder if pinniped ecology is well served by too many enthusiasts operating under too many restraints.

McLaren, I. A. (1990)

Pinnipeds and Oil: Ecologic Perspectives

In: Sea Mammals and Oil: Confronting the Risks

Geraci, J. R.; St. Aubin, D. J., eds. Pages 55-101

Academic Press, New York

ABSTRACT

Pinnipeds share many characteristics with other marine mammals, and indeed with large mammals in general, especially in demographic features, energetics, and social behavior. This allows us to draw on a wide range of empirical and theoretical literature to assess possible responses of pinnipeds to an environmental impact. Their amphibious nature poses special circumstances under which they face such threats. If we take as a premise that an oil pollution catastrophe has produced a large kill of pinnipeds, it is important to consider the rate at which the population is restored to its original level. Indeed, rate of return to "normality" is often an explicit component of environmental impact statements. The rate of recovery depends on the species and circumstances; the estimates available are for populations of pinnipeds under conditions that probably produce close to the maximum values. Those closer to equilibrium should show a less enhanced rate of increase following substantial reduction of numbers. A population already limited by the carrying capacity of the environment may not be capable of producing an immediate population response. Some species, such as the northern fur seal and Steller's sea lion, are indeed declining, presumably because the carrying capacity of their environments is in some way deteriorating. For these, a catastrophic kill might not be followed by population recovery. The rate of recovery of a seal population will also depend on the segments of the population that are killed. Loss of a year's offspring is more quickly compensated than is an equivalent loss of all age groups from the population, particularly of adults. Yet even the long recovery time for populations at equilibrium before a catastrophe might be optimistic if individuals in crowded environments fail to respond to population reduction.

Merrick, R. L.; Loughlin, T. R.; Calkins, D. G. (1987)

Decline in abundance of the northern sea lion, Eumetopias jubatus, in Alaska, 1956-1986

U.S. Fish. Bull. 85. 351-365.

ABSTRACT

Aerial, ship, and onshore surveys were conducted to assess the abundance of northern sea lions, Eumetopias jubatus, in southwestern Alaska, from the central Gulf of Alaska through the central

Aleutian Islands, during June-July of 1984-86. Counts of northern sea lions from these surveys were compared with counts made in 1956-62 and 1975-79. These data indicated that the number of adults and juveniles onshore declined 52% from 140,000 animals in 1956-60 to 68,000 in 1985 - an annual rate of decline of at least 2.7%. Numbers have declined throughout the region, with the greatest declines in the eastern Aleutian Islands (79%) and the least in the central Aleutian Islands (8%). This was not due to emigration because significant increases have not been noted elsewhere. Between the 1960s and mid-1970s, there were large decreases in the eastern Aleutian Islands and western Gulf of Alaska, and a major increase in the central Aleutian Islands. Beginning in the late 1970s declines occurred in all areas. The cause of the declines are unknown, but they may be associated with disease, prey availability or quality, or a combined effect of these and other factors. Factors which may contribute to the declines include the pre-1973 commercial harvests, entanglement of juveniles in marine debris, incidental takes in fisheries, and killing by fisherman.

Mikhalev, Y. A.; Ivashin, M. V.; Savusin, V. P.; Zelenaya, F. E. (1981)

The distribution and biology of killer whales in the Southern Hemisphere

Rep. int. Whal. Commn 31. 551-566.

ABSTRACT

Biological and distributional data for Southern Hemisphere killer whales, collected between 1961/62 and 1978/79 are analysed. It appears that killer whales are found in warm waters in winter and migrate into high latitudes in the summer. From the available data six populations of killer whales are proposed (classified by their winter distribution): Western American; Eastern American; Western African; Eastern African; Western Australian; Eastern Australian; although it is believed that further populations will be determined in the future for the open waters of the Atlantic, Indian and Pacific Oceans. The migration appears to be linked with that of its prey species, in particular the minke whale. Morphological and reproductive data are presented and discussed. A new species of killer whale, the dwarf killer whale Orcinus nanus is proposed.

Mitchell, E.; Reeves, R. R. (1988)

Records of killer whales in the western North Atlantic, with emphasis on eastern Canadian waters

Rit Fiskideildar 11. 161-193.

ABSTRACT

Published records of killer whales, Orcinus orca, in the western North Atlantic, from central Labrador (about 55°N) south to the West Indies, were compiled. We also compiled unpublished records, with emphasis on eastern Canadian waters, from statistics of Canada's east-coast shore whaling industry, data from Canadian whale sighting and tagging cruises in 1966-1973, the files of the Arctic Biological Station and other sources.

Although killer whales were taken only rarely by whalers in southern Labrador, Newfoundland and Nova Scotia, these whales were known to frequent the whaling grounds. In addition to the written and verbal reports of observations by the whalers, the wounds and scars on mysticetes landed at

the whaling stations provide evidence of encounters with killer whales (though there is no way of telling where or when the attempts at predation occurred). Jukker whales enter embayments along the coast of Newfoundland and are known from as far up the St. Lawrence as the Saguenay River confluence. In the past at least, they apparently preyed on the formerly large population of white whales, Delphinapterus leucas, in the St. Lawrence Estuary. There are few records from Nova Scotia and the Bay of Fundy. In New England, killer whales are sighted annually off Cape Cod, particularly in autumn, presumably as they follow migrating bluefin tuna, Thunnus thynnus, inshore. Winter records from the southeast U.S., the Bahamas and the West Indies support the hypothesis that at least some killer whales in the western North Atlantic undertake a north-south seasonal migration. However, there are also summer records in the Caribbean region. Therefore, two or more stocks may occur in the Northwest Atlantic or the migration scheme may be complicated.

Neff, J. M. (1990)

Effects of oil on Marine Mammal Populations: Model Simulations

In: Sea Mammals and Oil: Confronting the Risks

St. Aubin, J. R.; St. Aubin, D. J., eds. Pages 35-54

Academic Press, New York

ABSTRACT

The effect of an oil spill on a population of marine mammals can be assessed in a number of ways. One is to use a mathematical model that incorporates information on the behavior of spilled oil with data on the life history of the species in question. Also needed is some understanding of direct and indirect effects of oil on individuals--the kind of information usually generated from field observations and experimental studies. The advantage of a computer model is that it always produces a result: a prediction of impact. Modeling may be applied as part of risk assessment when developing offshore oil reserves. Models were created for the impacts of oil on marine mammals in the Southern California Bight, northern fur seals in the Bering Sea, bowhead and gray whales in Alaskan waters and sea otters in California. These modeling efforts are examples of the state of development of the approach. They can be quite sophisticated, incorporating a full range of environmental influences on oil and movements of animals that might contact it. When assumptions are made regarding mortality, the models can project the rate and degree to which a population recovers. A notable weakness in this scheme is that for most species, effects of contact are poorly understood, and the premise that contact is fatal is both extreme and results in gross overestimates of the impact of a spill. It is critical then to define more precisely how oil affects an animal. This can be done by combining direct observations and experimental studies with an understanding of life history, habitat use, social organization, demography, and other elements that can be used to measure the vulnerability of an animal to oil.

Newby, T. C. (1973)

Changes in the Washington State Harbor seal population, 1942-1972

Murrelet 54. 4-6.

ABSTRACT

A major decline of harbor seals was noted in the Nisqually Delta in southern Puget Sound from the 1940s through the 1960s due to extensive bounty hunting. The total Washington population declined from about 5,000 to 6,000 seals in the 1930s to fewer than 2,000 in 1971.

Nishiwaki, M.; Sasao, A. (1977)

Human activities disturbing natural migration routes of whales

Sci. Rep. Whales Res. Inst. 29. 113-120.

ABSTRACT

Human activities which disturbed natural migration routes of 2 whale populations were investigated. In Tokyo Bay (Japan) the annual catch of the Baird's beaked whale (Berardius bairdii) in the Boso whaling grounds is between 50-300. Although the whales showed no signs of average body length decrease, an apparent decrease of whales despite increasing catch per unit effort indicates that the maritime transportation has disturbed the migration routes of these whales. This also occurs on the northern coast of Kyushu. The reason for the catch decline cannot be damage by whaling as in the minke whale (Balaenoptera acutorostrata). Catch per unit effort is, as a whole, increasing, but the number of whales caught is decreasing without average body length decrease. Great numbers of vessels are coming into and going out of both areas, and the number is still increasing. Not only these whales, but also various species of marine mammals may be troubled by the bay traffic. Warning about the activities by man which disturb or expel wild animals from their natural habitats and migration routes are suggested.

Olesiuk, P. F.; Bigg, M. A.; Ellis, G. M. (1990)

Recent trends in the abundance of harbour seals, Phoca vitulina, in British Columbia

Can. J. Fish. Aquat. Sci 47. 992-1003.

ABSTRACT

Aerial censuses of harbour seals (Phoca vitulina) were conducted in the Strait of Georgia (1966-88), the lower Skeena River (1977-87), off the southwest coast of Vancouver Island (1976-87), off the northeast coast of the Queen Charlotte Islands (1986), in Jervis Inlet (1987), and at the entrance to Queen Charlotte Strait (1988). The estimated number of seals in the Strait of Georgia, the primary study area, increased from 2170 in 1973 to 15180 in 1988; the number in the lower Skeena River from 520 in 1977 to 1590 in 1987; and the number off southwestern Vancouver Island from 210 in 1976 to 1130 in 1987. The trends indicated that populations throughout British Columbia had been increasing at a rate of about 12.5%/year since 1973. Based on the density of seals in the areas surveyed, and the relative distribution of bounty and commercial kills, the total post-pupping

population in British Columbia was estimated to have numbered 75000-88000 in 1988, compared with 9000-10500 when the species was protected in 1970. Despite the recent increases, which probably reflect the recovery from historic kills, there was no evidence of density-dependent changes in the population growth rate.

Olesiuk, P. F.; Bigg, M. A.; Ellis, G. M.; Crockford, S. J.; Wigen, R. J. (1990)

An assessment of the feeding habits of harbour seals (Phoca vitulina) in the Strait of Georgia, British Columbia, Based on Scat Analysis

Can. Tech. Rep. Fish. Aquatic Sci 1730. 1-135.

ABSTRACT

Seasonal and regional variations in the diet of harbour seals in the Strait of Georgia were described based on 2841 scat samples collected from 58 sites (11 estuaries and 47 non-estuary haulouts) in all months, and the diet compared with the diet in other regions of the province based on 159 samples. Prey remnants were separated from other faecal matter using an elutriator and prey identified using a wide variety of structures including otoliths, teeth, vertebrae, cranial, appendicular and caudal elements, scutes and spines. Otoliths alone provided an incomplete and biased representation of the diet. We derived and employed a new index, referred to as a split-sample frequency of occurrence, to assess the relative importance of prey. The validity of the assumptions underlying the index, and scat analyses in general, were addressed. Annual prey consumption was estimated by combining dietary information with data on the abundance and distribution of seals and estimates of their daily food requirements. The diet in the Strait of Georgia was dominated by Pacific hake and herring, which comprised 42.6 and 32.4% of the overall diet respectively. The former was consumed primarily during April-November, which coincided with the post-spawning dispersal and movement of hake to shallower water, and the latter mainly during December-March, which coincided with the pre-spawning emigration of herring into the Strait of Georgia. Annual hake consumption in 1988 was estimated at 4214 tonnes, which represented 3.5% of the total stock biomass or 71% of mean recent commercial harvests. Annual herring consumption was estimated at 322.6 tonnes, which represented 3.2% of the total stock biomass or 27% of mean recent commercial harvests. Salmonids comprised 4% of the overall diet and consisted mainly of adult salmon that were taken as they returned to rivers to spawn, especially in estuaries. However, predation on trout also appeared to have been significant in localized areas. Annual salmonid consumption was estimated at 394 tonnes, which represented approximately 2.8% of mean recent annual escapement. Lingcod comprised 3.0% of the overall diet and were preyed upon mainly during November-April when males were defending nets. Annual lingcod consumption was estimated at 294 tonnes, which was roughly equivalent to the recent commercial and sport harvests combined. Other important prey in the Strait of Georgia were plainfin midshipman (3.4% of overall diet), surfperches (2.3%), cephalopods (2.1%), flatfishes (1.2%), sculpins (1.2%) and rockfishes (1.1%). The Strait of Georgia data, and the limited data collected from other regions, indicate the harbour seal is an opportunistic predator in that diets varied regionally and seasonally depending on the local availability of prey.

Olesiuk, P. F.; Biggs, M. A.; Ellis, G. M. (1990)

Life history and population dynamics of resident killer whales (Orcinus orca) in the coastal waters of British Columbia and Washington State

In: Individual recognition of cetaceans: Use of photo-identification and other techniques to estimate population parameters. Report of the International Whaling Commission (Special Issue 12)

Hammond, P. S.; Mizroch, S. A.; Donovan, G. P., eds. Pages 209-243

International Whaling Commission, Cambridge

ABSTRACT

Life history parameters are derived for the resident form of killer whale in the coastal waters of British Columbia and Washington State based on the demographic changes observed in two communities (closed to immigration and emigration) that were monitored between 1973-4 and 1987. Females have a mean life expectancy of 50.2 year, typically give birth to their first viable calf at 14.9 years of age, produce an average of 5.35 viable calves over a 25.2 year reproductive lifespan and have a maximum longevity of about 80-90 years. Calving is diffusely seasonal with most births occurring in October-March. Neonate mortality is approximately 43%. The estimated proportion of mature females pregnant varies from 0.274 in April to 0.411 in September. Males have a mean life expectancy of 29.2 years, typically attain sexual maturity at 15.0 years and physical maturity at 21.0 years of age, and have a maximum longevity of about 50-60 years. Mortality curves are U-shaped for both females and males, but the curve is narrower for males. There is no evidence of density dependence in the life history parameters as a result of cropping prior to the start of the study or as the populations increased during the study.

The derived life history parameters are used to develop a sex- and age-specific matrix population model and to calculate life tables. The model accurately emulates the demographic changes observed during the study. Population projections indicate that both communities represent stable populations below their carrying capacity. These populations had a finite annual rate of increase of 2.92% and were composed of 50% juveniles, 19% mature males, 21% reproductive females and 10% post-reproductive females. Discrepancies between the sex- and age-structure of the study populations and those of a stable population can be largely attributed to the selective cropping of pods prior to the start of the study. Simulations indicate that the population could sustain a maximum non-selective harvest of 2.84%; or maximum selective harvests of 4.70% of juveniles or 8.34% of adults, which represented total population harvest levels of 1.89% and 3.17% respectively. Sensitivity analyses reveal that populations are robust to changes in mortality rates, particularly adult mortality rates, which implies that density dependence is expressed primarily through changes in reproductive parameters. It is predicted that (1) a stationary population at carrying capacity will comprise 37% juveniles, 20% mature males, 14% reproductive females and 29% post-reproductive females; and (2) in a stationary population, females surviving to the end of their 14.0 year reproductive lifespan will produce an average of 2.0 calves.

Ono, K. A.; Boness, D. J.; Oftedal, O. T. (1987)

The effect of a natural environmental disturbance on maternal investment and pup behavior in the California sea lion

Behav. Ecol. Sociobiol. 21. 109-118.

ABSTRACT

Observed changes in maternal investment due to an environmentally induced decrease in food supply (the 1983 El Niño-Southern Oscillation) are compared with a priori predictions for the California sea lion (*Zalophus californianus*). Changes in behavior, growth and mortality of offspring were also examined. Data collected in the first two months postpartum for the years before (PRE), during (EN), and the two years after (POST1 and POST2) the 1983 El Niño indicate that females initiated postpartum feeding trips earlier during the food shortage, and spent more time away on individual feeding trips in both the El Niño year and the year after. Perinatal sex ratios (female:male) in the years PRE, EN, POST1 and POST2 were 1:1, 1.4:1, 1.1:1 and 1:1.4, respectively. Fewer copulations were observed during the El Niño year, but this difference was not statistically significant. Pups spent less time suckling in the food shortage year and the year following, but attempted to sneak suckle more. Pups were less active and played on land less in the El Niño and following year. Finally, maternal investment as measured by milk intake of offspring was decreased, pups grew more slowly, and suffered increased mortality during the food shortage year. Despite expected sex differences in maternal investment and pup behavior in response to food shortage, there were no sex-biased differences in response in either females or pups. As expected, the food shortage did not affect adult males since they migrate north during the non-breeding season where the environmental perturbation was less severe.

Oritsland, T.; Christensen, I. (1982)

A mass stranding of killer whales at Lofoten, northern Norway, in June 1981

Rep. int. Whal. Commn 32. 642.

ABSTRACT

A report of a mass stranding in which 14 killer whales out of a pod of about 30 ran ashore.

Payne, P. M.; Schneider, D. C. (1984)

Yearly changes in abundance of harbor seals, Phoca vitulina, at a winter haulout site in Massachusetts

U.S. Fish. Bull. 82. 440-442.

ABSTRACT

Harbor seals increased at a Massachusetts coastal haul-out site at around 11.9% per year from 1972 through 1980 mostly from dispersal of seals from rookeries in Maine. The growth was thought to have been facilitated by protection of seals from bountied hunting after the passage of the Marine Mammal Protection Act in 1972.

Payne, R.; Rowntree, V.; Perkins, J. S.; Cooke, J. G.; Lankester, K. (1990)

Populations size, trends and reproductive parameters of right whales (Eubalaena australis) off Peninsula Valdes, Argentina

In: Individual recognition of cetaceans: Use of photo-identification and other techniques to estimate population parameters. Report of the International Whaling Commission (Special Issue 12)

Hammond, P. S.; Mizroch, S. A.; Donovan, G. P., eds. Pages 271-278

International Whaling Commission, Cambridge

ABSTRACT

Photo-identification of individual whales, based on natural marking, has been used to study a population of southern right whales, Eubalaena australis, on their winter assembly grounds around Peninsula Valdes, Argentina. Right whales have raised patches of roughened skin (callosities) on their heads. The pattern of callosities differs between individuals. This enables known individuals to be monitored over time. Photographs of individual whales have been obtained from aerial surveys conducted each year between June and December from 1971 through 1986. A total of 909 distinct individuals were identified over the period, of which 481 were identified in more than one season. These data have been used to estimate various population parameters. The mean calving interval is estimated to be 3.6 years (95% confidence interval 3.3 to 4.1 years). It is estimated that there were 99 (SE 18) calvings in the population in 1986, which implies a total population of about 1,200 in that year. The population is estimated to be increasing at a rate of 7.6% p.a. (SE 1.7%). These estimates should be treated with caution until the validity of the underlying assumptions has been verified.

Pechmann, J. H. K.; Scott, D. E.; Semlitsch, R. D.; Caldwell, J. P.; Vitt, L. J.; Gibbons, J. W. (1991)

Declining amphibian populations: the problem of separating human impacts from natural fluctuations

Science 253. 892-895.

ABSTRACT

Reports of declining amphibian populations in many parts of the world are numerous, but supporting long-term census data are generally unavailable. Census data from 1979 to 1990 for three salamander species and one frog species at a breeding pond in South Carolina showed fluctuations of substantial magnitude in both the size of breeding populations and in recruitment of juveniles. Breeding population sizes exhibited no overall trend in three species and increased in the fourth. Recent droughts account satisfactorily for an increase in recruitment failures. These data illustrate that to distinguish between natural population fluctuations and declines with anthropogenic causes may require long-term studies.

Perrin, W. F.; Oliver, C. W. (1982)

Time/area distribution and composition of the incidental kill of dolphins and small whales in the U.S. purse-seine fishery for tuna in the eastern Tropical Pacific, 1979-80.

Rep. int. Whal. Commn 32. 429-444.

ABSTRACT

Data are presented for 4,642 spotted dolphins, Stenella attenuata; 1,745 spinner dolphins, S. longirostris; 99 striped dolphins, S. coeruleoalba; 1,535 common dolphins, Delphinus delphis; two bottlenose dolphins, Tursiops truncatus; one Pacific white-sided dolphin, Lagenorhynchus obliquidens; and one false killer whale, Pseudorca crassidens. Possible source of biases in the samples are discussed.

Perrin, W. F.; Henderson, J. R. (1984)

Growth and reproductive rates in two populations of spinner dolphins, Stenella longirostris, with different histories of exploitations

In: Reproduction in whales, dolphins and porpoises. Reports of the International Whaling Commission (Special Issue 6)

Perrin, W. F.; Brownell, R. L. Jr.; DeMaster, D. P., eds. Pages 417-430

International Whaling Commission, Cambridge

ABSTRACT

A model of density-dependent change in net reproductive rate (births minus deaths) has been used in assessing status of dolphin stocks in the eastern tropical Pacific. The eastern spinner population has been estimated to be at a lower fraction of original size (17-25%) than is the population of whitebelly spinners (58-72%). Higher reproductive rates would be expected in the former than in the latter on the basis of the density-dependent model, provided the latter is above its level of maximum net production. Based on analyses of over 4,000 specimens collected through 1978: (1) there is a relative paucity of fully adult males in the eastern spinner population (possibly resulting in lower average fertility), (2) the eastern spinner female attains sexual maturity about one tooth-layer unit (probably one year) earlier than does the whitebelly spinner, (3) ovulation rates in young females is lower in the eastern spinner population, and (4) the proportion of all females which are sexually mature is lower in the eastern spinner population. Gross annual reproductive rates (proportion female X proportion of females mature X pregnancy rate) are not different in the two populations (about 8-10% in both cases). This comparison does not confirm the hypothesis that a density-dependent increase in gross reproduction occurs in the spinner dolphin.

Perrin, W. F.; Reilly, S. B. (1984)

Reproductive parameters of dolphins and small whales of the family Delphinidae

In: Reproduction in whales, dolphins and porpoises. Reports of the International Whaling Commission (Special Issue 6)

Perrin, W. F.; Brownell, R. L. Jr.; DeMaster, D. P., eds. Pages 97-133

International Whaling Commission, Cambridge

ABSTRACT

The purposes of this review are to describe and critique methods used to estimate reproductive parameters, to summarize estimates in the literature and to examine patterns in the estimates and their implications. Reviewed are gestation period, fetal growth rate, size at birth, size and age at attainment of sexual maturity, average size and age of adults, maximum size, asymptotic length, ovulation rate, pregnancy rate, calving interval, length of lactation, weaning age, length of "resting" period, age and sex structure, and birth rates. Also discussed are the effects on the estimates of seasonality, schooling segregation, geographical variation and exploitations and the relationships between parameters.

Perry, A.; Baker, C. S.; Herman, L. M. (1990)

Population characteristics of individually identified humpback whales in the central and eastern North Pacific: a summary and critique

In: Individual recognition of cetaceans: Use of photo-identification and other techniques to estimate population parameters. Report of the International Whaling Commission (Special Issue 12)

Hammond, P. S.; Mizroch, S. A.; Donovan, G. P., eds. Pages 307-317

International Whaling Commission, Cambridge

ABSTRACT

Methods developed to obtain and record photographs of tail flukes and accompanying sighting data for humpback whales are described. Published descriptions of the migratory movement, abundance, reproductive histories and social organization of individually identified humpback whales based on our long-term studies in Hawaii and southeastern Alaska and other studies in the eastern and central North Pacific are reviewed. Biases and limitations associated with the collection and analysis of photographic data are discussed and additional methods that can be useful in describing population parameters for humpback whales and other cetacean species are suggested.

Humpback whales in the North Pacific, like those in the North Atlantic, appear to form geographically isolated feeding herds which intermingle on one or more wintering grounds. Mark-recapture analyses of resighting data suggest a seasonal population of 327 to 421 in the southeastern Alaska feeding region, and 1,113 to 1,701 on the Hawaiian wintering grounds. In Hawaii, multiple sightings of 18 sexually mature females provided an estimated calving rate

(calves/female/year) of 0.58, but this value may be inflated by sighting biases. In southeastern Alaska, multiple sightings of 41 mature females provided an estimated calving rate of 0.37, which we believe is a polygynous mating system involving male-male competition for mature females. In southeastern Alaska, the foraging strategies of humpback whales appear to be noncompetitive and, on occasion, cooperative.

Pierce, G. J.; Thompson, P. M.; Miller, A.; Diack, J. S. W.; Miller, D.; Boyle, P. R. (1991)

Seasonal variation in the diet of common seals (Phoca vitulina) in the Moray Firth area of Scotland

J. Zool., Lond. 223. 641-652.

ABSTRACT

Seasonal variation in the diet of common seals (Phoca vitulina) in the Moray Firth, north-east Scotland, was determined from analysis of faecal samples collected at haul-out sites during each month of 1988. Data on diet of common seals in 1987 are also presented. Limitations of the methods available for quantification of diet are discussed. Although some of the observed variation in diet from month to month may reflect changes in the sampling regime, a clear seasonal pattern was apparent, with clupeids predominating in the winter and sandeels in the summer. The trends observed are consistent with opportunistic feeding on the most abundant prey.

Pitcher, K. W. (1980)

Food of the harbor seal, Phoca vitulina richards, in the Gulf of Alaska

U.S. Fish. Bull. 78 (2). 544-549.

ABSTRACT

A total of 548 harbor seals were collected by rifle throughout the Gulf of Alaska from 1973 through 1978. Stomach contents were sorted and prey remains identified as an index of diet. Fishes composed 74.5%, cephalopods 21.5% and decapod crustaceans 4.0% of the prey identified.

A minimum of 27 species of fish were identified belonging to 13 families. Cephalopods included both octopus and squids of the family Gonatidae. The five top-ranked prey of harbor seals in the Gulf of Alaska were walleye pollock, octopus, capelin, eulachon, and Pacific herring. Either walleye pollock or octopus was the top-ranked food in all marine areas and eulachon was dominant in the estuarine and freshwater habitats of the Copper River Delta. Walleye pollock was the top-ranked item in the eastern areas: northeastern Gulf of Alaska, Prince William Sound, and the Kenai coast. In the western areas: Lower Cook Inlet, Kodiak, and the Alaska Peninsula, octopus had the highest ranking. In Lower Cook Inlet, octopus and shrimps made up over 60% of both total occurrences and volumes which was nearly twice the percentages in other areas. Salmon were found in the diet of harbor seals from both Prince William Sound and the Kodiak Island area only during the summer. In the Kodiak area, feeding on Pacific sand lance appeared to be greatest in the fall while use of capelin seemed to peak in the spring. Use of Pacific herring by harbor seals appeared greatest in the spring in Prince William Sound.

Pitcher, K. W.; McAllister, D. C. (1981)

Movements and haulout behaviour of radio-tagged harbour seals, Phoca vitulina

Can. Field Nat. 95. 292-297.

ABSTRACT

Movements, haulout area fidelity, and haulout frequency of harbor seals (Phoca vitulina) were studied in the Kodiak Island area, Alaska, by relocating radio-tagged animals captured on a large hauling area. Eight of 35 radio-tagged seals were found on hauling areas other than the capture site. The longest movement was 194 km and one seal crossed 74 km of open ocean. Movement rates up to 27 cm/d were recorded. There appeared to be considerable fidelity to one or two specific haulout location by individual radio-tagged animals. Resident, radio-tagged seals of a large hauling area were hauled out during 50% of the daily radio checks in June and 41% from 1 August to 5 September. On an individual basis, frequency of haulout ranged from 16 to 80% of the days.

Pitcher, K. W. (1986)

Variation in blubber thickness of harbor seals in southern Alaska

J. Wildl. Manage. 50 (3). 463-466.

ABSTRACT

Measurements of blubber thickness, sculp weights, and other morphometric data were obtained from 559 harbor seal (Phoca vitulina) collected along the southern Alaskan coast in order to evaluate seasonal fatness in relation to sex and age and to compare indices of fatness. In the older age classes females were fatter than males. Older seals of both sexes were fatter than younger animals. Adolescent and mature animals of both sexes were fatter during winter than during the reproductive and molt periods. Sculp weight divided by total body weight, sculp weight divided by standard length and condition index were all correlated positively with blubber thickness. These findings facilitate comparisons of fatness in populations regionally or over time.

Pitcher, K. W. (1990)

Major decline in number of harbor seals, Phoca vitulina richardsi, on Tugidak island, Gulf of Alaska

Mar. Mamm. Sci. 6 (2). 121-134.

ABSTRACT

Tugidak Island, located in the Gulf of Alaska, was once the site of one of largest local concentrations of harbor seals (Phoca vitulina richardsi) in the world. This population, which probably consisted of about 20,500 animals in the mid-1960s, declined by about 85% between 1976 and 1988. The population appeared to decline more rapidly during the late 1970s than during the 1980s. Causes for the decline are not apparent. There appear to be both similarities and dissimilarities between this decline and recent declines in abundance of northern fur seals

(Callorhinus ursinus) and Steller sea lions (Eumetopias jubatus) in the Bering Sea and Gulf of Alaska.

Ray, G. C.; Dobbin, J. A.; Salm, R. V. (1978)

Strategies for protecting marine mammal habitats

Oceanus 21 (2). 55-67.

ABSTRACT

The protection of marine mammal species requires identification of areas of special biological significance, or 'critical habitats'. Recent legislation and research work have emphasised the value of a habitat approach to management, which the authors have attempted to illustrate by means of a management model, based on walrus (Odobenus rosmarus divergens) populations in the Bering Sea. The model integrates data on : 1) the hydrography of the study area, 2) the distribution, life history, food habits and community structure of the walrus, 3) existing and proposed socioeconomic activities, 4) legal and jurisdictional matters. Areas of high vulnerability are identified by mapping the mating grounds, pupping grounds, migration routes and food supply, and then overlaying additional data in a colour-coded system. This gives a visual presentation of potential conflicts and responsibility for action, which may be used as a strategic planning tool for protection measures such as the establishment for sanctuaries. The article concludes with a discussion of the usefulness of this and other approaches to management problems.

Reijnders, P. J. H. (1983)

The effect of seal hunting in Germany on the further existence of a harbour seal population in the Dutch Wadden Sea

Zeit. Saugetierkunde 48. 50-54.

ABSTRACT

Calculated influence of dispersal and hunting pressure on seals in the Wadden Sea. Due to pollution effects pup production in the Dutch seal population is too low compared to the stable population in Schleswig-Holstein. Nevertheless, aerial surveys showed that the total number of seals remained fairly stable at about 500 specimens since 1974. It is demonstrated that since hunting was stopped in Niedersachsen and Schleswig-Holstein unrestricted dispersal in the Wadden Sea area could take place. It is calculated that the Dutch seal population in 1980 contained 41% animals originating from outside the area. Besides another 15% of the animals were repatriated by seal nursery stations. Abstinence of hunting in the whole area is of vital importance for the further existence of a harbour seal population in the Dutch Wadden Sea.

Reijnders, P. J. H. (1985)

On the extinction of the southern Dutch harbour seal population

Biol. Conserv. 31. 75-84.

ABSTRACT

The harbour seal population in the southern part of the Netherlands, the Delta area, has decreased sharply since about 1953. Causes for the observed decrease, such as hunting pressure, habitat limitation through construction works, disturbance and water pollution, are discussed. As in the Wadden Sea, where the seal population has been studied more intensively, the initial sharpest decrease was caused by overhunting, probably followed by the impact of pollution. Construction works have had a secondary effect, as by the time these began, numbers were already very much reduced.

Reilly, J. J.; Fedak, M. A.; tedak, M. A. (1991)

Rates of water turnover and energy expenditure of free-living male common seals (Phoca vitulina)

J. Zool., Lond. 223. 461-468.

ABSTRACT

The water and energy metabolism of free-living male common seals (Phoca vitulina) during the mating season was investigated using labelled water methods. All three seals, which were captured on two occasions, were in negative energy balance during the study. The daily energy expenditure of one animal, estimated using doubly-labelled water was 52.5 MJ. This is equivalent to six times the basal metabolic rate predicted from Kleiber's (1975) allometric equation. Rates of water turnover were slightly lower than predicted from the allometric equation of Richmond, Langham and Trujillo (1962). The observed rates of water turnover and energy expenditure are considerably higher than those of seals which fast during the mating season, and are consistent with the observed differences in behaviour between males of the common seal and other pinniped males during mating.

Reilly, S. B. (1984)

Observed and maximum rates of increase in gray whales, Eschrichtius robustus

In: Reproduction in whales, dolphins and porpoises. Reports of the International Whaling Commission (Special Issue 6)

Perrin, W. F.; Brownell, R. L. Jr.; DeMaster, D. P., eds. Pages 389-399

International Whaling Commission, Cambridge

ABSTRACT

Shore station censuses indicate that the California stock of gray whales increased at an annual rate

of 2.5% during 1967-1980, concurrent with an annual exploitation of approximately 1.2%, i.e. net reproduction was near 3.7% per year. In light of this net reproduction, vital-rate estimates for the circa 1967 population were re-evaluated. The most likely values during that period were: pregnancy rate of 0.467, mean age at sexual maturity of 8 years, adult survival of 0.945 and juvenile survival of 0.899. To estimate maximum net rate of increase, biologically defined limits of pregnancy rate, age at sexual maturation and juvenile survival were used to generate a Leslie matrix, with adult survival held constant at the 1967 level. The dominant eigenvalue of the matrix was calculated as 1.069. If a stable age structure can be assumed in populations at a very low level, this indicates a possible maximum net rate of increase approaching 6.7% per year.

Reilly, S. B.; Barlow, J. (1986)

Rates of increase in dolphin population size

U.S. Fish. Bull. 84 (3). 527-533.

ABSTRACT

Annual finite rates of increase in dolphin population size were estimated to vary up to a maximum of 1.09, using simulation, based on ranges in vital rates. Vital rate ranges were defined from values reported in the literature where possible, otherwise by making assumptions about biological or logical limits. Given information on current values, or limits, of one or more vital rate, one can use the figures presented to determine ranges of possible rates of increase in population size. The highest rates estimated here (up to 1.09) are probably unrealistic, because of the unlikely combinations of high fecundity and low mortality needed to achieve them.

Reilly, S. B. (1987)

Reanalysis of rate of change in the California-Chukotka gray whale stock, 1967/68 - 1979/80

Rep. int. Whal. Commn 37. 347-349.

ABSTRACT

A statistical analysis of the relationships among annual population estimates, mean annual visibility and mean distance offshore of passing whales from the 1967-1980 Monterey gray whale censuses indicates that there was in fact a significant positive rate of change in gray whale abundance during those years.

Reilly, S. B. (1990)

Seasonal changes in distribution and habitat differences among dolphins in the eastern tropical Pacific

Marine Ecol. Prog. Ser. 66. 1-11.

ABSTRACT

Large-scale patterns of dolphin distribution and oceanography were studied from research-vessel

surveys conducted in the pelagic eastern tropical Pacific during June to November 1982, 1986 and 1987. Substantial changes were observed in relation to previously reported winter distributions for spotted and/or spinner dolphin schools (*Stenella attenuata* and/or *S. longirostris*) and for striped dolphin schools (*S. coeruleoalba*). These dolphin species were sighted in abundance west of 120°W along 10°N coincident with seasonal shoaling of a thermocline ridge. No seasonal distribution changes were observed for common dolphin schools (*Delphinus delphis*); as in the winter, they occupied upwelling-modified waters of the region. Highest-density areas for the 3 school types were statistically different between spotted/spinner dolphin school and common dolphin schools. Striped dolphin schools could not be discriminated from the other 2 types based on these habitat variables, indicating other factors or processes contribute to the observed spatial separation of the 3 distributions.

Renouf, D.; Gaborko, L.; Galway, G.; Finlayson, R. (1981)

The effect of disturbance on the daily movements of harbour seals and grey seals between the sea and their hauling grounds at Miquelon

Appl. Anim. Ethol. 7. 373-379.

ABSTRACT

The movements of harbour seals and grey seals through a narrow channel connecting their hauling grounds with the sea were recorded during the daylight hours from 14 to 27 June 1980. There was only a slight increase in seabound travel after the seals were disturbed by humans, and the animals did not necessarily go to sea when the sand flats they hauled out on were flooded by the high tide. There was no recognizable diurnal pattern to their movements, and no relationship between the direction and intensity of the traffic and various meteorological factors.

Schneider, D. C.; Payne, P. M. (1983)

Factors affecting haul-out of harbor seals at a site in southeastern Massachusetts

J. Mamm. 64. 518-520.

ABSTRACT

Several factors affected the number of seals appearing near shore at Manomet, but only tide and disturbance had any significant effect on the percentage hauling out. Occasional disturbance did not affect the total number of seals in the study area, but may nonetheless be important if the number of hours spent out of water is critical.

Shaughnessy, P. D.; Goldsworthy, S. D. (1990)

Population size and breeding season of the Antarctic fur seal Arctocephalus gazella at Heard Island-1987/88

Mar. Mamm. Sci. 6 (4). 292-304.

ABSTRACT

Breeding colonies of the antarctic fur seal Arctocephalus gazella on Heard Island (53°10'S, 73°30'E) are situated on the sheltered northern and eastern coasts on flat vegetated terrain near streams and pools. Pupping in the 1987/88 summer began on 21 November, with 90% of births in 26 d. The median birth date was 11 December. Pup counts at Heard Island made in seven breeding seasons from 1962/63 to 1987/88 show an exponential rate of increase of 21%, which may be inflated due to undercounting in early years. The total of 248 births in 1987/88 represents an exponential increase of 37% since the previous year, but pups may have been undercounted then. Based on the number of pups born, the breeding population is estimated at 870-1,120. During the breeding season, the largest number of animals ashore was 835. Many non-breeding fur seals began hauling out from early January and 15,000 animals were estimated to be ashore by late February, a far larger number than expected from the size of the breeding population. Both the breeding and non-breeding components of the population may be augmented by immigration. The source of immigrants may be undiscovered breeding colonies of this species in the northwestern sector of the Kerguelen Archipelago or the concentration at South Georgia. Further censuses are required at Heard Island to monitor the population growth.

Slater, L. M.; Markowitz, H. (1983)

Spring population trends in Phoca vitulina richardi in two central California coastal areas

Calif. Fish Game 69. 217-226.

ABSTRACT

Two areas separated by about 17km with comparable numbers of harbor seals were surveyed in Spring 1980. Population trends in the two areas were significantly different with a decline at the San Mateo County sites. In contrast to this area which had a low pupping rate, nursery herds were observed at the two Santa Cruz County sites. A group composed almost exclusively of males was observed in San Mateo County. Since there is no indication of movement between the areas, these contrasting distributions are interpreted as resulting from differences in habitat characteristics and frequency of human encroachment.

Slouten, E.; Lad, F. (1991)

Population biology and conservation of Hector's dolphin

Can. J. Zool. 69. 1701-1707.

ABSTRACT

During the past decade, Hector's dolphins, Cephalorhynchus hectori, have suffered an alarming level of mortality due to entanglement in commercial and amateur gill nets. In this paper we study two Leslie matrix population models that incorporate known features of dolphin fertility and mortality, focussing on the information they provide regarding age distributions and maximum population growth rates. The simplest model specifies constant survival rates over many age-classes. The second model uses more realistic curves of age-specific survival rates. The results indicate that Hector's dolphin, like most other small cetaceans, has a low potential for population growth. Growth rates of 1.8-4.9% per year are likely to be the maximum possible for Hector's dolphin population, and C. hectori (and C. commersonii) populations are likely to be declining under recent levels of net entanglement. Survival rate estimates from free-living populations, subject to natural and net-entanglement mortality, showed decreasing populations. Even with the most optimistic reproductive parameters, survival rates would need to be some 5-10% higher than those observed in populations subject to gill-net entanglement before population growth could occur. The likely consequences of a reduction in entanglement mortality through conservation management are explored using the survivorship curve model. These simulations show that the age structure of the population can have an important effect on changes in the size and growth rate of the population during the recovery phase following a reduction in entanglement mortality.

Smith, R. I. L. (1988)

Destruction of Antarctic terrestrial ecosystems by a rapidly increasing fur seal population

Biol. Conserv. 45. 55-72.

ABSTRACT

The terrestrial environment of Signy Island, South Orkney Islands, maritime Antarctic, is undergoing rapid and possibly irreversible change caused by a natural biological agent. During the past decade there has been a dramatic increase in the number of Antarctic fur seals Arctocephalus gazella coming ashore on the island during the short summers. It is not known whether significant numbers of seals were present on the island prior to the initiation of commercial hunting in the early 1820s. The impact that the continuing increase of these seals had made on the island's terrestrial and freshwater environments has been sudden and locally devastating. The fragile cryptogram-dominated vegetation has suffered physical damage from which it may be impossible to recover. These seals are also frequenting several of the island's freshwater lakes which are becoming increasingly eutrophic. The long-term implications of this impact are causing serious concern for the future of the lowland terrestrial and freshwater ecosystems on Signy Island if the fur seal population continues to increase.

Smith, T. D. (1983)

Changes in size of three dolphin (Stenella spp.) populations in the eastern tropical Pacific

U.S. Fish. Bull. 81. 1-13.

ABSTRACT

Dolphins from three populations, one of Stenella attenuata and two of S. longirostris, have been killed incidentally in the yellowfin tuna purse seine fishery in the eastern tropical Pacific, two populations since about 1959 and the other since about 1969. Size changes in these populations are estimated from numbers killed each year, population size estimates in 1979, and net recruitment rates. Ranges of values for some parameters are considered, accounting for some uncertainties. Assuming central values of the ranges of maximum net recruitment rate (3%) and the population level giving maximum net productivity (65%), one S. longirostris population, the eastern spinner dolphin, is near 20% of pre-exploitation levels: the S. attenuata population, the northern offshore spotted dolphin, is between 35 and 50%; and the second S. longirostris population, the whitebelly spinner dolphin, is between 58 and 72% of pre-exploitation levels.

Smith, T. G.; Geraci, J. R.; St. Aubin, D. J. (1983)

Reaction of bottlenose dolphins, Tursiops truncatus, to a controlled oil spill

Can. J. Fish. Aquat. Sci 40. 1522-1525.

ABSTRACT

Three captive bottlenose dolphins, Tursiops truncatus, were observed in an ocean pen measuring 14 x 11 m divided into three equal areas by oil-containment booms. Each dolphin was placed in the pen alone for 4 d with no oil present, and 2 d with tar-colored mineral oil in one of the areas. We noted their area of surfacing, underwater movements, dive times, and reactions. After a few brief contacts with the oil, dolphins completely avoided surfacing in the slick. The oil also acted as a temporary barrier restricting their underwater movement.

St. Aubin, D. J. (1990)

Physiologic and toxic effects on pinnipeds

In: Sea Mammals and Oil: Confronting the Risks

Geraci, J. R.; St. Aubin, D. J., eds. Pages 103-107

Academic Press, New York

ABSTRACT

Oil fouling has been implicated in the deaths of pinnipeds, though much of the evidence has been circumstantial. Large-scale mortality has never been observed, even after some of the more catastrophic spills. In general, the prediction that spilled oil would have its greatest impact on

young pinnipeds in cold, ice-bound waters has been borne out following the discharge of residual oil in the Gulf of St. Lawrence in 1969.

Pinnipeds show little behavioral or physiologic reactions to the noxious characteristics of oil. Incidental ingestion during feeding, exposure to vapor concentrations that might be expected under natural conditions at sea, and limited surface fouling with relatively fresh oil do not appear to cause significant distress. Pinnipeds trapped near the source of a spill, or forced to emerge in heavy accumulations of oil in leads and around rookeries, exhibit the most severe effects. For fur seals, experimental studies indicate that surface fouling will decrease the insulative value of the pelt, possibly leading to thermal and energetic stress. Individuals of all species and groups that are compromised by pre-existing disease, or stressed by an unfavorable habitat, intra-specific competition, or unusual environmental conditions may be the most vulnerable to the effects of oil exposure.

Stewart, B. S. (1984)

Diurnal hauling patterns of harbor seals at San Miguel Island, California

J. Wildl. Manage. 48. 1459-1461.

ABSTRACT

Harbor seals haul-out on a variety of substrates (tidal sand and mud bars, intertidal rocks and reefs, ice floes and glacial drift, sand and cobble beaches, and man-made floats and rafts) in various parts of their range. In some areas, hauling areas are accessible only at or near low tide and consequently surveys of seal populations are timed with respect to tide only. Using time-lapse photography the author documented year-round patterns of terrestrial abundance of harbor seals at haul-outs on San Miguel Island in southern California where beaches are accessible at all tides. Seals exhibited strong diurnal patterns; few were ashore in early morning or late afternoon and peak abundance was between 1200 and 1400 hrs throughout the year. Tide did not influence the timing of haulout but seals were more abundant ashore at afternoon low tides compared to afternoon high tides when corrected for hour of the day. Human intrusion (persons on site, aircraft and helicopters, etc.) disrupted hauling patterns on 21% of the days (n=272). Maximum yearly terrestrial abundance was in late May and early June when seals were molting. The author concluded that estimates of population size on the southern California Channel Islands can be made more reliable by standardizing survey times to early afternoon. In other parts of the species' range where supralittoral hauling space is available and used by seals, studies of diurnal influences on hauling patterns may be desirable before estimates of population size and growth are made based on low tide censuses.

Stewart, B. S.; Karl, S. A.; Yochem, P. K.; Leatherwood, S.; Laake, J. L. (1987)

Aerial surveys for cetaceans in the former Akutan, Alaska, whaling grounds

Arctic 40 (1). 33-42.

ABSTRACT

Randomized aerial surveys were flown between 26 July and 26 August 1984 to search for cetaceans in two areas of southwestern Alaska: one on both Bering Sea and Pacific Ocean sides of the Aleutian islands near the defunct Akutan shore-whaling station, which operated from 1912 through 1939, the other overlapping continental slope and shallow continental shelf waters between the Aleutians and the Pribilof Islands. Surveys were made at altitudes between about 150m and 245m from a Partenavia P68 Observer with a plexiglass nose bubble, which permitted center-line viewing. Searches covered 3940 nautical miles (nm), including some 2403 nm of random transects. Sightings were made of gray whales (10 sightings, 14 individuals), fin whales (3,11), minke whales (1,1), unidentified beaked whales (1,6), Dall's porpoises (47,131), killer whales (8,26) and harbor porpoises (4,7). A Fourier series model was used to estimate density of Dall's porpoises as 115 individuals (CV = 0.263) per 1000 nm² on the whaling grounds and 16.6 individuals (CV = 0.0) per 1000nm² in the Bering Sea north of the whaling grounds. These estimates are comparable to those previously reported for the same general areas (97.2 animals per 1000nm², SD = 49.5). There were too few sightings of other cetaceans to permit calculation of meaningful density estimates. At least four species of great whales (blue, fin, humpback and sperm) were sufficiently abundant during the first four decades of this century to support significant whaling activities within about 100nm of Akutan (more than 5300 whales were caught during 23 years of whaling, 1912-39). Although previous studies of the fisheries showed a downward trend in catch per unit of effort and an increase in distance traveled to take whales, whales were still being taken at relatively high rates (0.28-0.51 whales per gross catcher day) at the end of the fishery in 1939. Populations of fin, humpback, blue and sperm whales were probably significantly reduced by shore and pelagic whaling conducted widely in the North Pacific since 1939. The low number of sightings on the present surveys probably means that populations on and near the whaling grounds remain depressed from such activities.

Stewart, B. S.; Antonelis, G. A., Jr.; DeLong, R. L.; Yochem, P. K. (1988)

Abundance of harbor seals on San Miguel Island, California, 1927 through 1986

Bull. So. Cal. Acad. Sci. 87 (1). 39-43.

ABSTRACT

Counts of harbor seals ashore were made each year at San Miguel Island in southern California from 1973 through 1986. Historical counts were available for 1927, 1958, 1964, 1965, and 1969. The data indicated that the colony increased about 22% per year from 1958 through 1986 overall, but that the rate of increase may have declined somewhat after 1976. Counts of terrestrial abundance of seals were corrected using radiotelemetry data to yield estimates of absolute abundance. Hauling patterns may have changed during the 1982/83 El Niño Southern Oscillation which makes interpretations of trends in abundance after 1983 problematic. The authors concluded that accurate determinations of absolute abundance of harbor seals on the Channel Islands and elsewhere will

require further studies of temporal, geographic, and habitat influences on the proportions of seals ashore.

Stewart, B. S. (1989)

The ecology and population biology of the northern elephant seal, Mirounga angustirostris Gill 1866, on the Southern California Channel Islands

Ph.D. dissertation, University of California, Los Angeles, CA

ABSTRACT

The patterns of seasonal abundance, behavior, and population biology of northern elephant seals (Mirounga angustirostris) on the Southern California Channel Islands were studied from 1979 through 1987 to investigate the influences of crowding on colony growth and neonatal survival. Four primary peaks in terrestrial abundance of elephant seals are related to reproduction in winter and age- and sex-related differences in timing of the molt in spring and summer. The magnitudes of these peaks on San Nicolas Island varied among years. The magnitudes of these peaks on San Miguel varied among years. Non-breeding season peaks in abundance are unreliable for assessing absolute abundance or trends in colony size, but annual fluctuations at one rookery may be useful indicators of temporary or long-term oceanographic or environmental changes or suggestive of demographic changes on other rookeries. An understanding of the demographics of all colonies is, however, essential for interpreting seasonal changes in abundance on each. Births continued to increase exponentially at average annual rates of 11.5% and 7.8% at San Nicolas and San Miguel islands, respectively. Crowding during the breeding season but insignificant effects on increases in births in most habitats and the greatest rates of increase in births occurred at densities very near the carrying capacities (K) of the breeding habitats (i.e., at about 0.75 K). Pre-weaning pup mortality was independent of densities on rookeries and remained low at about 4% on San Nicolas Island and generally less than 6% on San Miguel Island. Crowding on some beaches where female distribution did, however, influence the tenure of lactating females and growth of their pups. Females were evidently in poorer physical condition in 1983 and 1984 than in other years presumably because of decreased food abundance or altered prey distribution which may have required them to range further when at sea from late 1982 through early 1984 during an intense ocean warming event. Consequently, parous females arrived on rookeries later in 1983, nursed their pups fewer days in 1983 and 1984 which resulted in pups being weaned in poorer condition and were at sea feeding longer in spring 1983 than in other years. They also conceived a significantly greater proportion of male progeny in 1984 than in other years.

Stewart, B. S.; Leatherwood, S.; Yochem, P. K.; Heide-Jorgensen, M. P. (1989)

Harbor seal tracking and telemetry by satellite

Mar. Mamm. Sci. 5. 361-375.

ABSTRACT

We tested a satellite Platform Transmitter Terminal (PTT) in the laboratory (on a float and on captive seals) and on a free-ranging harbor seal in the Southern California Bight to investigate the

utility of satellite telemetry in documenting seals' at-sea behavior and movements. We used records from a microprocessor-based time-depth recorder (TDR) to interpret location and diving records from the PTT. For the free-ranging harbor seal, we obtained at least one uplink during 70% (while the seal was at sea) to 82% (while she was ashore) of satellite passes and at least one location each day. Of 62 locations determined by Service Argos for the free-ranging seal, 20 were verified from TDR records to have been at sea; these indicated that the seal may have ranged up to 48 km from the haul-out site, although most locations were within 5 km. The accuracies of locations calculated when the seal was at sea were substantially less than when it was ashore (1.5km), thus limiting at-sea tracking of seals by satellite to rather gross movements. Fewer transmissions were detected and locations calculated when the seal was actively diving than when it was swimming near the surface as it departed from or returned to the haul-out site. Consequently, average dive durations indicated by the PTT were substantially shorter than those calculated from TDR records. Documentation of foraging areas and detailed at-sea movements using satellite technology may not be possible for pinnipeds unless PTT-transmission rates are increased substantially from the 1 per 45 sec maximum rate now permitted by Service Argos.

Stewart, B. S.; Yochem, P. K. (1991)

Northern elephant seals on the Southern California Channel Islands and El Nino

In: Pinnipeds and El Nino: Responses to environmental stress

Trillmich, F.; Ono, K., eds. Pages 234-243

Springer-Verlag, Berlin

ABSTRACT

Pup mortality of northern elephant seals increased at the Channel Islands in winter 1983 because of intense storms, astronomically high tides, and elevated sea level coincident with El Nino. Pregnant females arrived later in 1983, nursed their pups fewer days in 1983 and 1984 and were at sea feeding longer in spring 1983 than in other years.

Stewart, B. S.; Yochem, P. K.; DeLong, R. L.; Antonelis, G. A. (1992)

Status and trends in abundance of pinnipeds on the Southern California Channel Islands

In: Recent advances in California islands research

Hochberg, F. G., ed. Pages In Press

Santa Barbara Museum of Natural History, Santa Barbara, California, Santa Barbara

ABSTRACT

Pinnipeds were relatively uncommon on the Southern California Channel Islands in the early 1900s because of overexploitation by commercial sealers and indiscriminate killing. Populations of four of the species recorded have increased greatly during the past several decades. Northern sea lions

once bred on the Channel Islands in small numbers but they have not been seen since 1984. Guadalupe fur seals wander to the Channel Islands from their sole rookery in Mexico and may soon colonize them. Populations of northern elephant seals, California sea lions, harbor seals, and northern fur seals will probably continue to increase, although at temporarily reduced rates because of greater mortality and slower growth of pups and juveniles during the 1982-83 El Nino Southern Oscillation event. In addition, competition for food resources or breeding space, or both, will restrict further growth and some populations may stabilize abruptly by the end of the century.

Summers, C. F. (1978)

Trends in the size of British gray seal populations

J. Appl. Ecol. 15. 395-400.

ABSTRACT

Trends in size of the more important stocks of British Grey seals are presented. Together with their West Atlantic counterparts they are increasing exponentially. Attempts to control stock size have indicated that they are best reduced by adult culls and not by pup hunting.

Terhune, J. M.; Almon, M. (1983)

Variability of harbour seal numbers on haul-out sites

Aquat. Mamm. 10. 71-78.

ABSTRACT

The high variability in numbers of seals hauled out were interpreted as evidence for high population mobility. Most seals hauled out during low tide. Human disturbance affected the numbers of seals hauled out which limited the researchers abilities to estimate population size.

Testa, J. W. (1987)

Long-term reproductive patterns and sighting bias in Weddell seals (Leptonychotes weddelli)

Can. J. Zool. 65. 1091-1099.

ABSTRACT

The reproductive performance of tagged Weddell seals (Leptonychotes weddelli) was monitored at McMurdo Sound, Antarctica, from 1970 to 1984. An age-specific reproductive schedule revealed the major onset of pupping at age 6 years, and a mean age of first birth of 7.1 years. The average asymptotic pupping rate of 0.61 is reached by age 10. The cost of pupping in a given year is reflected in a 0.05 drop in the probability of pupping the following year. This cost is not evident in females over 7 years old, suggesting that postweaning condition affects newly mature females more than those that are fully mature. Annual adult reproductive rates ranged from 0.46 to 0.79, with a possible periodicity of 5 to 6 years. Simulations were conducted to determine the impact on

reproductive estimates of sighting biases associated with seals having had at least one pup (Parous) or having pupped that season (With-Pup). Age at first reproduction as deduced from an age-specific pupping schedule is strongly affected by both forms of sighting bias, but bias in sighting Parous females was the more important. Estimates of adult reproductions were affected minimally. Comparisons of reproductive estimates with those of Weddell seals at Signy Island are discussed with regard to the effects of sighting biases.

Testa, J. W.; Siniff, D. B. (1987)

Population dynamics of Weddell seals (Leptonychotes weddelli) in McMurdo Sound, Antarctica

Ecòl. Monog. 57 (2). 149-165.

ABSTRACT

Populations dynamics of Weddell seals in McMurdo Sound, Antarctica, were studied from 1970 to 1984 using mark-recapture estimation, aerial surveys, age structure and magnitude of harvests, and direct counts of pup production. Similar data from earlier studies were used to reconstruct the history of the population during the period of human presence after 1956. Jolly-Seber estimates of population size indicated a general decline from 1970 to 1976, low numbers in 1976 and 1978, and relative stability from 1979 to 1984 at about 1500 adult seals. From 1970 to 1983, mean annual survival of adult females and male Weddell seals was estimated at 0.85 and 0.76. Counts made in aerial surveys in the summers of 1982/1983 and 1983/1984 were substantially below similar survey counts in the 1960s. Pup production in the breeding colonies also declined from 1967 to 1976, but has recovered and been stable since 1979. Collections spanning 28 yr indicated significant shifts in age structure characteristic of populations undergoing a rapid increase and then decline. The age structure of both sexes indicated a constant age of recruitment to the adult population at 5 yr, which corresponds closely with age of maturity in females. Generalized least squares and standard regression analyses failed to detect any correlation of adult survival or female reproductive estimates with population size or ice conditions, but there were significant trends in survival estimates from 1970 to 1983 that may be due to shifting age structure.

The most plausible interpretation of these data is that heavy harvests of seals to feed dog teams in the mid-1905s severely depleted the resident population of adult Weddell seals. When population studies began in the early 1960s the population was expanding rapidly, probably as a result of immigration by juveniles. The population declined to low levels in 1976-1978 and has been fairly stable since 1979, probably at a level lower than before harvesting began. Survival and reproductive parameters of these Weddell seals are low relative to other pinnipeds, suggesting that the adult population is at an equilibrium with its environment.

Testa, J. W.; Siniff, D. B.; Croxall, J. P.; Burton, H. R. (1990)

A comparison of reproductive parameters among three populations of Weddell seals (Leptonychotes weddellii)

J. Anim. Ecol. 59. 1165-1175.

ABSTRACT

(1) Tagged populations of Weddell seals at three sites in Antarctica were compared to resolve earlier differences in estimated reproductive parameters and to examine interannual patterns in adult reproduction. (2) The estimated reproductive rate from McMurdo Sound (0.68) was lower than that reported from the Vestfold Hills (0.80). The estimate from Signy Island was found to be biased upward and not comparable to the other two sites. (3) Average age at first sighting with pup was 6 years at McMurdo Sound, 7 years at Signy Island and 8 years at the Vestfold Hills. (4) Sampling at Signy Island and the Vestfold Hills was earlier in the breeding season and less intensive than at McMurdo Sound. Subsets of data from McMurdo were used to simulate the sampling regime at the other two sites. The results were higher estimates of both reproductive parameters (0.80 pups female⁻¹ and 7 years) such that there were no significant differences between sites. A similar sampling bias probably occurs at Signy and the Vestfold Hills and would account for the differences between sites. (5) Probability of sighting an adult female with a pup was found to correlate well with adult reproductive rate at McMurdo Sound and was used as an index to adult reproduction at all three sites. This reproductive index appears to have fluctuated in approximate synchrony at Signy Island and McMurdo Sound, sites separated by over 145° of longitude, but not at the Vestfold Hills.

Testa, J. W. (1991)

Temporal variability in Antarctic Marine Ecosystems: Periodic fluctuations in the phocid seals

Can. J. Fish. Aquat. Sci 48. 631-639.

ABSTRACT

Three species of seals around Antarctica have shown quasi-cyclic patterns in some aspect of their biology: the age structure of crabeater seals (Lobodon carcinophagus) around the Antarctic Peninsula has shown strong cohorts separated by 4- to 5-yr intervals; juvenile leopard seals (Hydrurga leptonyx) have appeared in unusually large numbers at Macquarie Island, also at 4- to 5-yr intervals; and Weddell seals (Leptonychotes weddellii) in McMurdo Sound have undergone fluctuations in reproductive rate every 4-6 yr. Complex demodulation was used to compare patterns among these three data sets and with the Southern Oscillation Index (SOI). All of the the seals data sets showed evidence of cyclical behavior when demodulated at a period of 5 yr. The Weddell seals were generally in phase with the SOI since the Weddell series began in 1970. The leopard seals and SOI were in phase in the 1960s, but thereafter the SOI series led the leopard seal series by about one quarter of a cycle. The crabeater series was more complicated, but similarities with the other data sets also were seen. If these tentative observations are confirmed, they point to large-scale oceanographic variation, possibly related to the El Niño-Southern Oscillation (ENSO), as an important mechanism in Antarctic marine ecosystems.

Thompson, P. M.; Rothery, P. (1987)

Age and sex differences in the timing of moult in the common seal, Phoca vitulina

J. Zool., Lond. 212. 597-603.

ABSTRACT

This study followed the progress of the annual moult within a population of common seals in Orkney, Scotland. Moulting seals were seen over a three-month period, from 7 June until 16 September. Yearlings were first to start moulting. Amongst older seals, females completed their moult an average of seven days earlier than immature males and 19 days earlier than mature males. Differences in the timing of moult appeared to be related to the age or reproductive status of the animals, and may be the result of differential changes in levels of the sex hormones.

Thompson, P. M. (1989)

Seasonal changes in the distribution and composition of common seal (Phoca vitulina) haul-out groups

J. Zool., Lond. 217. 281-294.

ABSTRACT

Seasonal changes in the distribution and composition of common seal haul-out groups were followed in a study area in Orkney, Scotland. A marking programme was also undertaken, using both conventional and radio-tags, to study individual movements between sites and seasonal changes in site-use. Certain haul-out sites were used only in the breeding season, while others were used during the winter. Seals were seen at one site all year round and at another during only the pre-pupping and moult period. On one island where two sites were used during the summer, there were significant differences in the sex ratio of groups at the two sites: at one site males predominated and few pups were seen; on another, nearby, mothers and pups were regularly seen, although the site was also used by males. There was also evidence for segregation of the sexes outside the breeding season. Repeated observations of marked seals showed that seals used several different haul-out sites throughout the year, and that the seasonal changes in abundance at different sites resulted from individual changes in site-use. These changes in site-use are discussed in relation to feeding movements, breeding requirements and the physical characteristics of different sites.

Thompson, P. M.; Fedak, M. A.; McConnell, B. J.; Nicholas, K. (1989)

Seasonal and sex-related variation in the activity patterns of common seals (Phoca vitulina)

J. Appl. Ecol. 26. 521-536.

ABSTRACT

Radio-telemetry was used to study common seal activity patterns around Orkney, and to assess the degree of individual, seasonal and sex-related variation in haul-out behaviour in 1984-86. During the summer, both males and females hauled-out regularly, in a marked diurnal pattern, but there were differences in other aspects of their haul-out behaviour. Males showed a marked change in

their haul-out behaviour at the beginning of moult. At this time, they hauled-out every day and, consequently, male behaviour became more synchronous. The change in behaviour of the male followed in 1985 was closely related to changes in abundance at the haul-out site that he used over this period. In contrast, females showed no sign of a similar change in behaviour at the beginning of moult, but appeared to spend more of their time at sea after lactation. During the winter, seals appeared to spend more time in offshore waters, although they regularly returned to the inshore study area to haul-out. At this time of year there was no marked diurnal haul-out pattern. There was a high degree of individual variation in the effect of the tidal cycle on activity patterns, which appeared to be related to tide-related changes in food and habitat availability. Overall, however, the tidal cycle had less effect on haul-out behaviour in summer, when strong circadian patterns were more important. The variations in activity patterns found in this study are discussed in relation to seasonal changes in feeding activity and the demands of breeding and moult.

Thompson, P. M.; Harwood, J. (1990)

Methods for estimating the population size of common seals, Phoca vitulina

J. Appl. Ecol. 27. 924-938.

ABSTRACT

This study evaluates the use of two different survey techniques for providing an estimate of the size of the common seal population in Orkney, U.K. In August 1985, an aerial survey was made over the coast of Orkney during the seals' moult. These results were compared with those from a series of boat surveys made over a sample of this area during the pupping season, in June and July. Over twice as many seals were found hauled-out on the survey made during the moult, and it is recommended that future estimates of population size should be based on surveys made at this time of year. Time-lapse photography was used to look at changes in the number of common seals hauled-out in relation to the tidal cycle and the time of day. These data, together with data collected on the activity patterns of radio-tagged individuals, were used to provide correction factors to compensate for seals which were in the water at the time of the survey. The application of these correction factors to the survey total of 6616 produced a provisional estimate 9331 (95% C.L.s 8147-10515) for the size of the Orkney common seal population. This estimate is discussed in relation to previous estimates of the size of both the Orkney and the total British common seal populations.

Thompson, P. M.; Pierce, G. J.; Hislop, J. R. G.; Miller, D.; Diack, J. S. W. (1991)

Winter foraging by common seals (Phoca vitulina) in relation to food availability in the inner Moray Firth, N.E. Scotland

J. Anim. Ecol. 60. 283-294.

ABSTRACT

Parallel studies of fish distribution and the diet and activity of common seals were made to assess the relationship between the seals' winter feeding activity and the distribution and abundance of their prey. Echosounder and trawling surveys revealed that a large part of the fish biomass was

sprat and small herring, while faecal analyses showed that >90% of common seal prey (by weight) were clupeoid fish. During the day, clupeoids concentrated in trenches and holes more than 12 m deep. Radio-tagged seals were located regularly over these areas. At night, clupeoid shoals rose in the water column and became more dispersed. Diel changes in seal activity patterns suggest that seals fed more often during the day. Prey sizes were estimated from the size of otoliths retrieved from seal faeces. Estimated sizes of clupeoids taken by seals were similar to the sizes of fish caught in trawls, even though estimates were not corrected to allow for partial digestion of otoliths. This suggests that the rapid otolith digestion rates previously reported from captive seals may have been artificially high, or that the Moray Firth seals selected fish larger than those caught in trawls.

Trillmich, F.; Limberger, D. (1985)

Drastic effects of El Niño on Galapagos pinnipeds

Oecol. 67. 19-22.

ABSTRACT

Population dynamics of pinnipeds living in the tropical upwelling ecosystem of the Galapagos were strongly influenced by the 1982-83 Southern Oscillation-El Niño (EN) event which was the strongest recorded in this century. The Galapagos fur seal (Arctocephalus galapagoensis) population lost the four youngest year classes (1980-1983) almost entirely and approximately 30% of the adult females and non-territorial males. Mortality of large territorial males was almost 100%. Most of the 1982 year class of Galapagos sea lions (Zalophus californianus wollebaeki) died and there was a much lower pup production in the breeding season following EN. Recurrent EN events must strongly influence age structure and average population size of these and other otariid species depending on tropical upwelling ecosystems.

Tsukagoshi, T. (1983)

Some peculiar phenomenons on killer whale, Orcinus orca.

Bull. Jap. Soc. Fish. Oceanogr. 44. 127-131.

ABSTRACT

LANGUAGE: Japanese. The author describes his experience of sighting killer whales (Orcinus orca), with [sic] happened only 3 times during his 30 years seafaring life. First he saw a single one attacking a sperm whale 200 miles south off the Shionomisaki Cape in April, 1953. The second time was in July, 1959, at 157-30E, 32-30W. The last time was in April, 1974, 20 miles south off Bali Island in Indonesia. On the second and third occasions a good amount of tuna and bonito were caught. He concludes that killer whales feed on tuna and bonito in warm areas.

Waters, S.; Whitehead, H. (1990)

Population and growth parameters of Galápagos sperm whales estimated from length distributions

Rep. int. Whal. Commn 40. 225-235.

ABSTRACT

The length distributions of female and immature sperm whales, *Physeter macrocephalus*, off the Galápagos Islands were studied using data collected between February-April 1985 and January-June 1987. Sperm whales were measured using a photographic technique. There were few small whales and most animals were between 8.6-10.2m. A simple population model estimated population and growth parameters for female and immature sperm whales off the Galápagos by comparing the Galápagos length distribution with that expected given certain population and growth parameters. The derived parameters were not always consistent with those currently accepted. The Galápagos data were most consistent with a significantly reduced pregnancy rate (2-4% of mature females per year) and smaller lengths of females at physical maturity (10.2m), and suggested a low mean mortality of mature females if the population is near equilibrium.

Watkins, W. A. (1986)

Whale reactions to human activities in Cape Cod waters

Mar. Mamm. Sci. 2 (4). 251-262.

ABSTRACT

A review of whale observations of more than 25 years indicated that each of the species commonly observed within 35 km of Cape Cod reacted differently to stimuli from human activities, and that these responses have gradually changed with time. These reactions appeared to result mostly from three types of stimuli: primarily underwater sound, then light reflectivity, and tactile sensation. The whale reactions were related to their assessment of the stimuli as attractive, uninteresting or disturbing, their assessment of the movements of the sources of the stimuli as expected or unexpected. Whale reactions were modified by their previous experience and current activity: habituation often occurred rapidly, attention to other stimuli or preoccupation with other activities sometimes overcame their interest or wariness of stimuli, and inactivity seemed to allow whales to notice and react to stimuli that otherwise might have been ignored. The changes over time in the reactions of whales to stimuli from human activities were gradual and constantly varying with increased exposure to these activities.

Weigle, B. (1990)

Abundance, distribution and movements of bottlenose dolphins (Tursiops truncatus) in lower Tampa Bay, Florida

In: Individual recognition of cetaceans: Use of photo-identification and other techniques to estimate population parameters. Report of the International Whaling Commission (Special Issue 12)

Hammond, P. S.; Mizroch, S. A.; Donovan, G. P., eds. Pages 195-201

International Whaling Commission, Cambridge

ABSTRACT

Boat surveys of bottlenose dolphins (Tursiops truncatus) inhabiting lower Tampa Bay, Florida, were conducted between April and October in 1983 and 1984. Objectives included: (1) examining the abundance and distribution of dolphins over 230km² in southern Tampa Bay and the adjacent Gulf of Mexico; (2) identifying individual animals using photographs of scars and other natural markings on the dorsal fin; and (3) characterizing the range of movement and interactions among recognizable dolphins.

A total of 70 surveys were carried out using a 5m or 12m vessel. School size and number of calves was evaluated, behavior recorded and individuals photographed. Photographs were classified based on location of fin notches and cataloged. Mean monthly dolphin school size was lowest in April (2.8) and highest in September (6.1). The largest schools were observed around the mouth of Tampa Bay where it joins the Gulf of Mexico. Zones with highest density were also around the Bay mouth. Mean density (dolphins/km²) was highest from July (0.38) through September (0.36). Calves constituted 9.7% of all dolphins observed. Dolphins with distinct, naturally marked fins were recognizable in 142 of 319 schools; 246 animals were cataloged with 75 being sighted two to seven times. Thirty seven dolphins photographed three or more times were classified into three herds based on location of the sightings; a Tampa Bay herd, a Pinellas herd, and a Sarasota Bay herd. Members of the Tampa Bay herd were observed on both sides of the Bay and interacted with members of the Sarasota Bay and Pinellas herds. The large number of recognizable animals sighted only once (171) suggests that transient dolphins, perhaps nearshore or offshore animals, use lower Tampa Bay for foraging in the summer months, probably following schools of mullet inshore. An apparently open population of dolphins used the study area, creating a high potential for genetic mixing. Dolphin ranges within the study area were calculated to be up to 166km² and may represent only a portion of the total home range of the animals studied.

Wells, R. S.; Scott, M. D. (1990)

Estimating bottlenose dolphin population parameters from individual identification and capture-release techniques

In: Individual recognition of cetaceans: Use of photo-identification and other techniques to estimate population parameters. Report of the International Whaling Commission (Special Issue 12)

Hammond, P. S.; Mizroch, S. A.; Donovan, G. P., eds. Pages 407-415

International Whaling Commission, Cambridge

ABSTRACT

Field studies begun in 1970 and continuing to date have identified at least three adjacent resident populations (or communities) of bottlenose dolphins along the central west coast of Florida. We have used photo-identification, mark-recapture techniques, behavioral observations, radio-tracking and brief captures for biological sampling to examine the structure and dynamics of these populations. Population designations are based on consideration of individual home ranges, social association patterns and genetics. Although the populations are relatively discrete in terms of ranges and associations, electrophoretic analyses of blood samples indicate that genetic exchange occurs between populations. Males travelling between populations appear to be the probable vectors for genetic exchange.

Most field effort has concentrated on the Sarasota dolphin population. Most of its members are identifiable from natural marks or tagging efforts over the last 20 years. This population consists of about 100 individuals. For the analyses presented here, we considered 116 dolphins identified during 1980-87. Of these, the sex was known for 90 dolphins and the age has been estimated for 79 dolphins. The long time span of the study and the high proportion of identifiable individuals has allowed us to estimate vital rates for this population. An annual recruitment rate to age 1 of 0.048 was countered by a minimum mortality rate of 0.910 and a mean annual loss rate from other causes of 0.029 (e.g., emigrations, mortalities for which carcasses were not recovered or undocumented changes in identifying characteristics). Immigration was infrequent, with a mean annual rate about 0.02. The mean fecundity rate was 0.144. Knowledge of maternal relationships allowed comparisons of the percentage of calves observed in the field vs the percentage of young of the year. Because of the prolonged period of association between mothers and calves, there were nearly six times as many mother-calf pairs as mothers with young of the year. To test the effectiveness of photo-identification techniques, we compared the number of correct identifications made in the field 'by eye' against the number identified from photographs. We identified 89% of these well-marked dolphins correctly by eye. Because virtually all the dolphins were marked in the most-heavily surveyed portion of the study area, it was also possible to test the accuracy and precision of mark-recapture methods. Both the Petersen and Schnabel methods underestimated the known population size, although the Schnabel estimate was less biased. This bias was likely due to heterogeneity of sighting probabilities; different age-sex classes were shown to have different sighting probabilities.

Wilkinson, I. S.; Bester, M. N. (1988)

Is onshore human activity a factor in the decline of the southern elephant seal?

S. Afr. J. Antarct. Res. 18 (1). 14-17.

ABSTRACT

Comparison of areas of high and low human activity on Marion Island shows no difference in rates of decline of elephant seal numbers. Spatial distribution of births also shows no change in the period 1976 - 1986, suggesting that no shift in breeding population distribution has occurred in the period as a result of the level of human activity on Marion Island. Furthermore, comparisons of Marion Island with other breeding sites of elephant seals, where human activity is lower, show no significant differences in the rates of decline of the species. Direct onshore human disturbance is therefore rejected as a significant factor in the decline of the species.

Wilkinson, I. S.; Bester, M. N. (1990)

Continued population increase in fur seal, Arctocephalus tropicalis, and A. gazella, at the Prince Edward Islands

S. Afr. J. Antarct. Res. 20 (2). 58-83.

ABSTRACT

Population estimates of fur seals were obtained at the Prince Edward Islands during the 1987/88 and 1988/89 austral summers. The populations of Arctocephalus tropicalis at Marion Island and Prince Edward Island are increasing at a rate of 12.9% and 9.7% respectively. The Marion Island population is still in the recolonisation phase of rapid growth while the Prince Edward Island population is either still in the slow establishment phase or approaching maturity. The lack of previous data for Prince Edward Island prevents a comparison of growth rates. An apparent slowing of the rate of population growth in A. gazella at Marion Island is conceivably the result of undercounts in 1974, giving an overestimate of growth between 1974 and 1981. As a result of this undercount the importance of immigration to A. gazella numbers at Marion Island may have been overemphasized previously. The apparent failure of the A. gazella population to move past the slow establishment phase may be a result of the absence of krill (Euphausia superba) in the vicinity of the Prince Edward Islands. Hybridization between the species is continuing.

Wittelman, G. J.; Redfearn, A.; Pimm, S. L. (1990)

The extent of complex population changes in nature

Evol. Ecol. 4. 173-183.

ABSTRACT

Many models of animal populations show complete yet predictable patterns of density changes under simple and plausible assumptions. Yet one previous attempt to determine the extent and importance of complete dynamics concluded that they were likely only in some laboratory

populations, but not in field populations. Ecologists have treated changes more complex than a return to a simple equilibrium, such as the cyclical changes in populations of lynx and voles in the arctic, as special cases. Highly variable populations, such as insects, are usually thought to be driven by unpredictable changes in the weather. Here, we assemble 71 populations counted for over 50 years, and suggest that complex yet predictable population changes are more common than previously thought.

Woodley, T. H.; Read, A. J. (1991)

Potential rates of increase of a harbour porpoise (Phocoena phocoena) population subjected to incidental mortality in commercial fisheries

Can. J. Fish. Aquat. Sci 48. 2429-2435.

ABSTRACT

We estimated the potential intrinsic rate of increase (r) of the harbour porpoise (Phocoena phocoena) population in the Bay of Fundy and Gulf of Maine using empirical data on reproductive rates and several hypothetical survival schedules. Schedules of survival to maximum ages of 12 and 15 yr, were calculated from two potential natural mortality schedules combined with several schedules of incidental mortality estimates. The most realistic results were obtained when natural mortality of non-calves were calculated from Caughley's (1966. Ecology 47:906-918) smoothed age-frequency equation for Himalayan thar (Hemitragus jemlahicus) and applied in conjunction with a range of calf natural mortality estimates. This model indicates that harbour porpoises have a limited capacity for population increase, and populations are unlikely to sustain even moderate levels of incidental mortality (4% of the population per year). Extending the maximum age used in the models from 12 to 15 yr does little to increase estimates of r for the harbour porpoise population, and hence their susceptibility to incidental mortality.

Wursig, B. (1990)

Cetaceans and Oil: Ecologic Perspectives

In: Sea Mammals and Oil: Confronting the Risks

Geraci, J. R.; St. Aubin, D. J., eds. Pages 129-165

Academic Press, New York

ABSTRACT

Many aspects of behavior, diet, and habitat use may lead cetaceans into contact with spilled oil. Given the host of interacting variables, it is difficult to state precisely which species or individuals might be most vulnerable. However, several assumptions can be made.

Encounters with oil are likely to be prolonged in species that frequent restricted areas such as bays and estuaries. Examples are breeding and feeding humpback, gray, right, bowhead, and beluga whales, narwhals, bottlenose dolphins, harbor porpoises, and river dolphins. Cetaceans that range

widely may contact some oil as they move quickly through a fouled area, but are less likely to suffer long-term exposure. After a spill, oil is distributed primarily at the surface and on the seabed. Consequently, cetaceans that feed in these areas are more likely to contact oil than those that feed in the water column. These include skim-feeding right and bowhead whales, surface-lunging porpoises, and bottom-feeding gray whales- in other words, all mysticetes, except possibly the minke whale. Harbor porpoises and some dolphins may contact oil when they feed on flatfish and other bottom-dwelling prey. Dolphins that habitually force schools of prey to the surface may also be at risk. As a group, baleen whales appear to be the most vulnerable in view of their generally low numbers, their peculiar feeding strategies, and their dependence on selected, localized habitats for feeding and reproduction. Among the odontocetes in North American waters, restrictive habitat increases the risk of exposure for belugas, narwhals, harbor porpoises, and bottlenose dolphins. Most other odontocetes are too mobile and wide-ranging for oil to present much of a threat.

Yablokov, A. V.; Olson (Editors), M. (1989)

Influence of human activities on the Baltic ecosystem

Gidrometeoizdat, Leningrad

ABSTRACT

Present volume is a result of studies of a group of Soviet and Swedish researchers united under the project "Research of toxicant influence on the dynamics of seal populations" of Soviet-Swedish cooperation in the field of environment protection sponsored by the USSR State Committee of Hydrometeorology and Swedish National Environment Protection Board. The volume contains studies presented at the Soviet-Swedish Symposium held in Moscow in 1986.

Cooperation between Soviet and Swedish scientists on the problems relating to seals started in 1982 on the initiative of the Swedish side who had obtained alarming data concerning an abrupt decrease of the number of grey and ringed seals as a result of the accumulation of polychlorobiphenyls in the seal organism. The Swedish side was represented by researchers from the Department of Vertebrate Zoology, Swedish National Museum of Natural History (Stockholm); Department of Pathology, Faculty of Veterinary Medicine, Swedish University of Agricultural Sciences (Uppsala) and Special Analytical Laboratory, Swedish National Environment Protection Board (Solna). The Soviet side involved researches from N.K. Koltzov Institute of Developmental Biology, USSR Ac. Sci. (Moscow); Institute of Experimental Meteorology (Obninsk); Institute of Applied Geophysics, USSR State Committee of Hydrometeorology (Moscow); Atlant NIRO (Kaliningrad); Institute of Biology, Ac. Sci. Latv. SSR (Salaspils) and Lymnological Institute, USSR, Ac. Sci. (Irkutsk). Joint work was related to three major aspects of the problem: 1) study of distribution and numbers of the Baltic seal; their nutrition and ecology; 2) investigation of pathological changes in the seal organisms; 3) chemical analysis of seal tissue samples for the content of pollutants. These three aspects are elucidated in the present volume. Summing up the studies carried out it should be said that important comparative evidence was obtained as to the content of pollutants in the Baltic, Ladoga and Baykal seals. It turned possible to discover earlier unknown and extremely alarming evidence related to the damage of skin and some skeleton elements (probably, as a result of disturbances of the organism hormonal balance). It was shown for the first time for Pinnipedia that the analysis of asymmetry of morphological structures could be used for estimating alterations in

the state of populations. Known forms of histopathology were added up by discovered changed patterns of seal spermatogenesis. During the last years as a result of efforts undertaken by different countries to limit pollutants waste the situation with pollution of the Baltic Seas was somewhat improved. However, the state of seal populations, as the last link of food chains, still causes grave worries. The following fact alone shows the gravity of the situation: at present the number of seals in all the Baltic Sea is similar to that in the Ladoga Lake alone. Investigations carried out under this project are only a part of national and international efforts of scientists aimed at investigations and improvement of the situations in the Baltic Sea, restorations of its seal population. In this respect experiments of the Swedish scientists on seal reproduction in captivity are of great interest (results of these studies are also here presented). The present book sums up only the first stage of cooperation between Soviet and Swedish scientists in the studies of Baltic seals. This cooperation must get more fundamental and extensive.

Yochem, P. K.; Stewart, B. S.; DeLong, R. L.; DeMaster, D. P. (1987)

Diel haul-out patterns and site fidelity of harbor seals (Phoca vitulina richardsi) on San Miguel Island, California in autumn

Mar. Mamm. Sci. 3. 323-332.

ABSTRACT

We studied the haul-out patterns and movements of harbor seals (Phoca vitulina richardsi) on San Miguel Island, California, from 23 October through 6 December 1982 by attaching a radio transmitter to each of 18 seals and monitoring their presence ashore with continuously scanning receivers. Seals hauled out at all hours although, on average, the largest proportion of tagged seals was ashore between 1300 and 1500 h. Median durations of haul-out bouts of individual seals ranged from 4.7 to 21.8 h; 81% of all haul-out bouts were less than 12 h and 3% were longer than 24 h. Eighty-one percent of the seals that were resighted at least twice used only the sites where they were tagged; two seals used two sites and one seal used three. Most seals were hauled out on fewer than 51% of the days sampled. On average, about 41% of tagged seals hauled out each day whereas an average of about 19% was hauled out during peak afternoon hours. Using telemetry data to correct a count of 412 seals made during an aerial survey, we estimated absolute abundance at about 2,168 seals; a modified Peterson mark-recapture model produced an estimate of about 1445 seals.

Yochem, P. K. (1987)

Haul-out patterns and site fidelity of harbor seals at San Nicolas and San Miguel Islands, California

Master of Science Thesis, San Diego State University, San Diego, CA

ABSTRACT

I used radiotelemetric and observational data to study haul-out behavior and site fidelity of harbor seals at San Nicolas and San Miguel islands, California, from 1982 through 1984. Thirty-seven harbor seals were radio-tagged and flippers-tagged and an additional 36 were marked with

flipper-tags only. The proportion of days seals hauled out varied by time of day, period, and year. The largest proportion was ashore in early afternoon; some seals tended to haul out during daylight hours (0600-1800) and some at night (1800-0600). In 1982, individuals hauled out most often (i.e., the largest proportion of days) during the early molt period; the proportion of radio-tagged seals hauled out per day was also greatest during this period. The proportion of seals hauled out declined during late molt and post-molt periods. Thus, changing haul-out patterns are at least partially responsible for the observed seasonal changes in the number of seals hauled out (number ashore is highest in late spring, lowest in winter). Haul-out patterns in 1983 differed from those in 1982, which may have been due to the different age- and sex- class composition of the two samples. Alternatively, haul-out patterns in 1983 may have reflected a change in harbor seal feeding behavior in response to the effects of the 1982-1983 El Niño-Southern Oscillation event on prey abundance and availability. Factors such as parturition and molt also affected haul-out behavior. Adult females ashore outnumbered adult males during the reproductive period, perhaps because females needed to spend more time on land suckling their pups and resting. Seals also hauled out more often during the molt. The annual peak in the number of seals ashore which occurs during the early molt period is apparently caused by a temporary overlap of molting males and females; the sex ratio ashore is 1:1 during this period. Females molt earlier than males, however, and by the late molt period adult males ashore outnumbered adult females. Seals exhibited a high degree of site fidelity. No individual used more than four of the harbor seal haul-out sites at either island; seals that used more than one site had primary or "preferred" sites and were usually seen there. Primary sites were consistent between years during the molt period, when mating occurs, suggesting that mating is not random.

Yochem, P. K.; Stewart, B. S.; Mina, M.; Zorin, A.; Sadovov, V.; Yablokov, A. (1990)

Non-metrical analyses of pelage patterns in demographic studies of harbor seals

In: Individual recognition of cetaceans: Use of photo-identification and other techniques to estimate population parameters. Report of the International Whaling Commission (Special Issue 12)

Hammond, P. S.; Mizroch, S. A.; Donovan, G. P., eds. Pages 87-90

International Whaling Commission, Cambridge

ABSTRACT

Phenotypic variation in pelage patterns of North Pacific harbor seals has been observed and described by several researchers. We developed a classification scheme of non-metrical pelage characters (e.g., presence or absence of spots in various body areas) and tested intra- and inter-observer variability in scoring black-and-white photographs of harbor and largha seals. Observer agreement was good overall, but some observers disagreed when scoring the more subjective characters (e.g., spot density). We also compared pelage patterns of harbor seals from San Miguel, Santa Rosa and San Nicolas islands and found inter-island differences in two characters.

York, A. E.; Hartley, J. R. (1981)

Pup production following harvest of female northern fur seals

Can. J. Fish. Aquat. Sci 38. 84-90.

ABSTRACT

Female northern fur seals (Callorhinus ursinus) were harvested commercially from 1956 to 1968 and pelagic collections were taken for research purposes from 1958 to 1974. Early survival rates (birth to age 2) for males increased from an average of 0.32 before the harvest to 0.38 afterwards. Numbers of female pups whose births were precluded by the harvest are estimated for the years 1956-79; these account for about 70% of the difference between the numbers of pups actually born and the level of pup births before 1956. Estimates of the increased numbers of pups due to the increase in the early survival rates are presented for the years 1958-74.

York, A. E. (1987)

On comparing the population dynamics of fur seals

NOAA Tech. Rep. NMFS 51. 133-140.

ABSTRACT

A relatively simple age-structured model applicable to most species of fur seals was constructed. Using the model and available data on vital parameters and observed rates of increase or decrease of the various populations of fur seals, I investigated the interrelationships among the vital parameters and their effect on the rate of increase of the population. There are some similarities among the populations: (1) all must have high adult survivorship, i.e., greater than 85% per year; (2) changes in age at first reproduction alone do not greatly affect the rate of increase of the population; and (3) small changes (not statistically detectable without very large sample sizes) in any combination of vital parameters can significantly change the rate of increase of the population. There are also two important dissimilarities: (1) the observed rates of population increase for the southern species (as high as 15-16% per year) are much higher than the maximum rate of increase observed for the Pribilof Island population of northern fur seals (8% per year); and (2) many of the southern populations of fur seals are increasing, whereas most of the populations of northern fur seals have decreased recently or have remained stable. The first suggests that scientists must be circumspect in applying vital parameters estimated for Callorhinus to Arctocephalus; the second implies that comparisons of population dynamics must take into account the environmental differences which affect the vital parameters. The model also allows one to estimate adult survival if the growth rate and the average age of the breeding females are known.

Zakharov, V. M.; Yablokov, A. V. (1990)

Skull asymmetry in the Baltic Grey Seal: effects of environmental pollution

Ambio. 19. 266-269

ABSTRACT

The most dramatic increase in DDT and PCB levels in the Baltic occurred after 1955. The present study investigates whether morphological changes resulting from the disturbance in developmental stability can be found in the grey-seal populations born during the most significant period of pollution that occurred after 1960. Skull characteristics were examined in animals born before 1940 (pre-pollution group) and in animals born after 1960 (pollution group). It was shown that the pollution group had sharply increased levels of asymmetry in almost every character analyzed. The findings suggest a dramatic change in the developmental stability of the Baltic grey seal during the period of heavy pollution after 1960.

Zeh, J. E.; George, J. C.; Raftery, A. E.; Carroll, G. M. (1991)

Rate of increase, 1978-1988, of bowhead whales, Balaena mysticetus, estimated from ice-based census data

Mar. Mamm. Sci. 7 (2). 105-122.

ABSTRACT

The number of bowhead whales, Balaena mysticetus, passing within viewing range of the ice-based census at Point Barrow, Alaska, during spring migrations from 1978 to 1988 is estimated from the visual census data. The trend in the annual numbers yields an estimated rate of increase of 3.1% per year with a 95% confidence interval ranging from 0.1% to 6.2% for the Bering-Chukchi-Beaufort Seas bowhead stock during this period. Alternative treatments of the data suggest less precise or somewhat lower estimates, but all results indicate that the stock was increasing.