

HARZA-EBASCO

Susitna Joint Venture
Document Number

2131

Please Return To
DOCUMENT CONTROL

Temperature Tolerance in Young Pacific Salmon, Genus *Oncorhynchus*¹

BY J. R. BRETT

*Pacific Biological Station, and
Department of Zoology, University of Toronto
(Received for publication September 19, 1951)*

CONTENTS

Introduction	266
Acknowledgements	267
Materials	267
Methods	272
Results	277
Upper limits of temperature tolerance	277
Lower limits of temperature tolerance	283
Preferred temperatures	291
Comparison of temperature resistance	293
Zones of thermal tolerance	296
Discussion and Conclusions	298
Time and temperature	298
Comparison with some other salmonoids	302
Some ecological relations	302
Summary	305
References	307
Appendix I. Statistical procedure	311
Appendix II. Tables IV-XIV	315

ABSTRACT

Lethal limits of high and low temperatures were determined for the young of five species of Pacific salmon, the spring (*Oncorhynchus tshawytscha*), the pink (*O. gorbuscha*), the sockeye (*O. nerka*), the chum (*O. keta*) and the coho (*O. kisutch*).

For acclimation temperatures ranging from 5° to 24°C. significant differences between species in their resistance to high temperatures was obtained. The spring and coho were most resistant. The pink and chum salmon were least resistant, and the sockeye was distinguishable from the latter two by greater resistance for prolonged exposure to high temperatures. No species could tolerate temperatures exceeding 25.1°C. when exposed for one week.

A fanning-out of the opercula was shown to be directly correlated with the onset of death from a low temperature. By use of this criterion mixed lethal effects at low temperatures were demonstrated and found to be influenced by the size of the fish and by the salinity of the water. None of the species could withstand temperatures lower than 4°C. when acclimated to 20°C. and above. When taken from holding troughs as low as 5°C., coho and sockeye could not tolerate long exposure (four days) to 0°C.

¹Based upon portions of a thesis accepted by the Faculty of Graduate Studies, University of Toronto, in partial fulfillment of the requirements for the degree of Doctor of Philosophy.

In a vertical gradient little difference in preferred temperature was observed, either between species or for different acclimation temperatures. The 12° to 14°C. stratum was the region of greatest concentration.

Specific differences in temperature responses are in keeping with taxonomic and ecological distinctions.

INTRODUCTION

NO ACTIVITY of an animal escapes the effect of temperature. Distribution, development, propagation and mere existence, each is influenced strongly in some manner by temperature. This influence must be met and surmounted either through resistance or adaptation, external avoidance or internal control. It is known that the upper and lower limits of temperature-tolerance in fish are extended through both adaptation and resistance (Fry, 1947a), and the varying degrees of these two attributes separately and collectively distinguish the species in this respect. By conducting experiments on tolerance to high and to low temperatures among the young of the five North American species of Pacific salmon, the relative abilities of these species to cope with extremes of temperature have been described in the following analysis.

Among earlier experiments, interest in the ability of fish to survive temperatures in the region of the freezing point of water and slightly below was expressed by Regnard (1895). In 1899, Maurel and Lagriffe while investigating both upper and lower levels of temperature-tolerance, chiefly in fresh-water fish, concluded that these species were better adapted for resisting low than high temperatures. Later investigators, dealing mainly with mortality from high temperatures, expressed the resistance in terms of the temperature reached before death when heated at a constant rate (Huntsman and Sparks, 1924), or averaged either the times to death (Loeb and Wasteneys, 1912) or the number of fish dead following a given exposure (Hathaway, 1927) at various constant temperatures. This quantitative expression of temperature-tolerance has been developed to provide a more inclusive treatment, borrowing from the methods of pharmacological procedure concerning dosage-mortality (Fry *et al.*, 1946).

The phenomenon of thermal adaptation in relation to previous temperature history provides the organism with greater scope for environmental experience. The term "acclimation" has been used to describe this effect, although "acclimatization" is apparently synonymous (Doudoroff, 1942; Heilbrunn, 1943; Brett, 1944). The importance of temperature-acclimation in nature and in experimental work has been stressed with significant emphasis (Doudoroff, 1945; Fry, 1947a). By working systematically with different temperature-acclimations the various levels of both upper and lower thermal tolerance can be determined within sufficient statistical limits to permit accurate prediction. The development of precise methods of physiological measurement has set the stage for physiological description.

The close taxonomic relation of the Pacific salmons, genus *Oncorhynchus*, (Milne, 1948) coupled with fairly distinct ecological habits, provides interest in the affinities which might be revealed by a rigorous analysis of their temperature-tolerances. Similar work on other salmonoids is mounting (Fry *et al.*, 1946;

Fry, 1947b: C attributes on the role of temperature animals will be

I am greatly both directed wide variety of has been the s

The math of Dr. D. B. E

The experimology which is of Lands and ment of Land laboratory and fish by Messrs measure to the

The salmon Pacific Biological of Fisheries, V tional shipmen

The Fish extended leave tance. It is a

SOURCE

The five America were

Each lot insulated box, Successful ship

TABLE I. The h

Species
1. Spring
2. Pink
3. Sockeye
4. Chum
5. Coho

Fry, 1947b; Graham, 1949). With the gradual recording of such physiological attributes on a comparable basis within and between taxonomic groups of fishes, the role of temperature in the ecology, and possibly in the evolution, of these animals will become increasingly apparent.

ACKNOWLEDGMENTS

I am greatly indebted to Dr. F. E. J. Fry who has for a number of years both directed and stimulated research on temperature relations in fish. His wide variety of interests and knowledge in the problems of experimental biology has been the source of a wealth of suggestions and a constant inspiration.

The mathematical analysis of the data has been under the helpful guidance of Dr. D. B. DeLury of the Ontario Research Foundation.

The experiments were conducted in the Laboratory for Experimental Limnology which is operated jointly by the University of Toronto and the Department of Lands and Forests, located at the Southern Research Station of the Department of Lands and Forests, Maple, Ontario. The excellent facilities of the laboratory and the attention devoted to feeding and maintaining the stocks of fish by Messrs. D. Cucin, G. Stofa and W. Sanderson contributed in a large measure to the successful execution of the research.

The salmon eggs were obtained through the courtesy of Mr. F. Neave, Pacific Biological Station, British Columbia, and of Mr. C. H. Ellis, Department of Fisheries, Washington. They have each responded to urgent pleas for additional shipments when transportation problems resulted in minor catastrophies.

The Fisheries Research Board of Canada has not only seen fit to grant extended leave of absence to the author, but has also given him financial assistance. It is a pleasure to acknowledge this support.

MATERIALS

SOURCE

The five species of Pacific salmon common to the west coast of North America were obtained as eyed eggs from three hatcheries (Table I).

Each lot of approximately two thousand eggs was shipped in a fiberglass-insulated box, packed with ice and perforated to permit air exchange for the eggs. Successful shipment by air was possible if not more than thirty-six to forty-eight

TABLE I. The hatchery locations and dates of fertilizing, shipping and 50 per cent hatch for the five species of Pacific salmon.

Species	Hatchery	Fertilized	Shipped	50% Hatch
1. Spring	Dungeness, Wash.	30/8/49	20/10/49	9/11/49
2. Pink	" "	28/9/49	7/12/49	12/12/49
3. Sockeye	Issaquah, "	10/10/49	25/11/49	6/1/50
4. Chum	Nile Creek, B.C.	28/10/49	7/1/50	4/2/50
5. Coho	" " "	14/11/49	7/1/50	27/1/50

Fry, 1947b; Graham, 1949). With the gradual recording of such physiological attributes on a comparable basis within and between taxonomic groups of fishes, the role of temperature in the ecology, and possibly in the evolution, of these animals will become increasingly apparent.

ACKNOWLEDGMENTS

I am greatly indebted to Dr. F. E. J. Fry who has for a number of years both directed and stimulated research on temperature relations in fish. His wide variety of interests and knowledge in the problems of experimental biology has been the source of a wealth of suggestions and a constant inspiration.

The mathematical analysis of the data has been under the helpful guidance of Dr. D. B. DeLury of the Ontario Research Foundation.

The experiments were conducted in the Laboratory for Experimental Limnology which is operated jointly by the University of Toronto and the Department of Lands and Forests, located at the Southern Research Station of the Department of Lands and Forests, Maple, Ontario. The excellent facilities of the laboratory and the attention devoted to feeding and maintaining the stocks of fish by Messrs. D. Cucin, G. Stofa and W. Sanderson contributed in a large measure to the successful execution of the research.

The salmon eggs were obtained through the courtesy of Mr. F. Neave, Pacific Biological Station, British Columbia, and of Mr. C. H. Ellis, Department of Fisheries, Washington. They have each responded to urgent pleas for additional shipments when transportation problems resulted in minor catastrophies.

The Fisheries Research Board of Canada has not only seen fit to grant extended leave of absence to the author, but has also given him financial assistance. It is a pleasure to acknowledge this support.

MATERIALS

SOURCE

The five species of Pacific salmon common to the west coast of North America were obtained as eyed eggs from three hatcheries (Table I).

Each lot of approximately two thousand eggs was shipped in a fiberglass-insulated box, packed with ice and perforated to permit air exchange for the eggs. Successful shipment by air was possible if not more than thirty-six to forty-eight

TABLE I. The hatchery locations and dates of fertilizing, shipping and 50 per cent hatch for the five species of Pacific salmon.

Species	Hatchery	Fertilized	Shipped	50% Hatch
1. Spring	Dungeness, Wash.	30/8/49	20/10/49	9/11/49
2. Pink	" "	28/9/49	7/12/49	12/12/49
3. Sockeye	Issaquah, "	10/10/49	25/11/49	6/1/50
4. Chum	Nile Creek, B.C.	28/10/49	7/1/50	4/2/50
5. Coho	" " "	14/11/49	7/1/50	27/1/50

hours elapsed between packing and receiving. An initial mortality not exceeding 2 per cent inevitably followed the handling necessitated by shipment. With this exception, egg losses from other causes were exceptionally low in all cases where subsequent experiments were performed.

FEEDING AND CARE OF YOUNG

The transition stage from alevin to free-swimming, feeding fry is a precarious one for young salmonids. The habit of feeding must be developed and encouraged, usually by frequent presentation of small particles of food. By directing a jet of water into a small aluminum screened basket containing finely ground beef or hog liver, adequate dispersal of the food over periods of 15 to 20 minutes, four times daily, was achieved. This routine was maintained for the first month of feeding, followed by reduction in feeding frequency and a change of diet, mainly in accordance with fish-cultural procedures for salmon currently practised in Washington State (Burrows, no date). The diet selected was a slight modification of one reported by Ellis (1948) which had been found to give best growth and least mortality for young spring salmon when tested on a variety of diets. A mixture of 50 per cent beef or hog liver together with 48 per cent ground "fish-pack" (haddock and cod fillet waste) and 2 per cent yeast was provided, up to the second month, followed by a reduction in liver to 30 per cent for the balance of the experimental period.

Mortality in the stock tanks with a constant temperature of $8.8^{\circ} \pm 0.2^{\circ}$ was virtually negligible. No prophylactic treatments were introduced. At higher temperatures, 20°C. and above, up to 5 per cent mortality was observed in all species, and infrequent treatments (two to three times per month) with a 1:4,000 solution of Roccal were applied (Burrows). Two cases where disease became significant were encountered; one with five-month-old chum salmon, from a single tank, which necessitated discarding the remaining fish as well as one series of obviously discordant data; the other, with three-month-old sockeye, raised by stages to 24°C. and apparently incapable of deriving adequate nutrition from their diet at such an elevated temperature.

The pH of the well water supplying the laboratory was 7.3 with total solids amounting to 254.3 parts per million (Table II).

RETAINING TROUGHS

The retaining troughs were each supplied with running water tapped from hot- and cold-water sources of relatively constant temperature and pressure. Adjustment of these with regular inspection permitted setting the temperature of a trough (above 9°C.) to within $\pm 0.1^{\circ}\text{C.}$ of any desired temperature. Usually two, sometimes three, species were cultured in a divided trough.

Below 9°C. a refrigeration unit reduced the temperature in a single holding tank in which the five species were retained separately in cages of fine aluminum wire screening. Thus, different levels of temperature-acclimation were readily obtained with a high degree of accuracy and constancy.

SIZE AND AGE

Uniformity was maintained at the same time to ensure abundance of subsequent hatchlings to specific differences.

The time (about two months) from hatching to the time when the knowledge of their first stages.

TABLE II. Mineral constituents of the water supply used in these experiments, as reported by the Chief Provincial Analyst, Sept. 24, 1947.

	<i>parts per million</i>
Total solids	254.3
Alkalinity	202.0*
Total hardness	181.0*
Iron and Aluminum (oxides)	4.7
Iron	0.2
Calcium	89.5
Magnesium	14.1
Potassium and Sodium	36.8
Sulphates	45.1
Chlorides	13.0

*As carbonate of lime.

SIZE AND AGE OF EXPERIMENTAL FISH

Uniformity of conditions in every feature of the history of the young fish was maintained as far as possible. Keeping them in the same or similar troughs, at the same temperature, and presenting the same diet in sufficient quantity to ensure abundance, were the first precautions. The variations in response to subsequent high or low temperatures might then be considered as attributable to specific differences only.

The time for commencing experiments was set at three months after hatching (about two months after feeding commenced), and then continued for an additional two to three months. The fish from higher temperatures were used first. The knowledge that the young chum and pink salmon move to salt water early in their first year motivated making comparison of the species in the very young stages.

TABLE III. Mean fork-lengths, weights and ages of the salmon fry used in temperature-tolerance tests.

Species	Length (cm.)	Weight (g.)	Age (months)
Upper temperature tolerance			
Spring	4.44 ± 0.40	1.03 ± 0.27	3.6
Pink	3.81 ± 0.29	0.30 ± 0.15	3.8
Sockeye	4.49 ± 0.84	0.87 ± 0.45	4.7
Chum	5.44 ± 0.89	1.62 ± 1.03	4.9
Coho	4.78 ± 0.60	1.37 ± 0.62	5.2
Lower temperature tolerance			
Spring	4.72 ± 0.41		7.7
Sockeye	4.50 ± 0.53		5.8
Chum	5.00 ± 0.51		5.2
Coho	4.83 ± 0.45		5.4

In Table III mean fork-lengths and weights (with standard deviations) and the average age from hatching of all samples used in the temperature-tolerance tests are compiled. No weights are included for the fish used in low-temperature tests. These samples were not removed from the lethal tanks until some time after death, and water absorption had affected their weight.

LETHAL BATHS

The lethal baths, six in number and measuring 22 inches square by 11 inches deep (Figure 1) were each constructed of galvanized iron coated inside with aluminum paint and adapted for use in either upper or lower thermal-tolerance tests. The addition of complete insulation with fiberglass of one-inch thickness was of value in reducing temperature variation to a minimum. Thermostatically controlled, 120-watt heater-coils in pyrex tubing counterbalanced a steady loss of heat, mainly from aeration and from a very slow exchange of water, equal to the volume of the tank every twenty-four hours. In the low-temperature lethals the heat loss was augmented by the addition of a layer of crushed ice, partitioned off on three sides of the tank by a removable galvanized iron sheet. A standardized

thermometer
ments. Often
in general, v

Under
never reduced
high lethals
experiments

PREFERRED

Two pro
concerning
variously ad
36 inches hi
tively. Plate
complete th
the bottom
holes, the




FIGURE 1. Apparatus for determining upper and lower lethal temperatures being assembled. Two units, one with four, the other with two lethal baths are shown with connections for thermal control, aeration and water exchange. (Photograph by Mr. W. P. Rice.)

FIGURE 2. A
under low illu
of

thermometer calibrated in intervals of 0.1°C . was used throughout all experiments. Often no detectable change in the tank temperatures was apparent and, in general, variation did not exceed a range of more than 0.1°C . from any set level.

Under maximum loads of fish per tank (40) oxygen concentrations were never reduced below 93 per cent saturation (5.24 cc. O_2/l ., 26.5°C .) in tests on high lethals, and not below 81 per cent (6.91 cc. O_2/l ., 6.0°C .) in low-lethal experiments.

PREFERRED-TEMPERATURE TANKS

Two preferred-temperature tanks were used in a limited series of experiments concerning the region most frequented in a vertical temperature gradient by variously acclimated salmon fry. These tanks, illustrated in Figure 2, stand 36 inches high, with length and width measurements of 36 and 20 inches respectively. Plate glass facings held in angle iron edging and bolted to the main frame complete the outer structure. Water, usually of low temperature, is introduced at the bottom through a metal tube perforated uniformly over its length with small holes, the displaced water being drained off at the top. A coiled copper tube




FIGURE 2. A preferendum tank divided into ten cells by white cord for recording of fish positions under low illumination. Thermometers are situated in the front right of each cell. A faint outline of the coiled copper tube for carrying hot water is visible within the tank.

forming a closed system conducts hot water in a downward spiral around the inside periphery of the tank in contradirection to the rising "cold front". By adjusting the temperature and rate of flow of the introduced cold water as well as that of the hot water within the copper coil, any desired temperature gradient can be obtained.

METHODS

The approach to the problem of describing the temperature-tolerance of young Pacific salmon has been to hold samples of each species at different non-lethal temperatures but otherwise similar conditions. These variously acclimated samples were later tested for their tolerance to high and to low temperatures, ranging from rapidly lethal to sublethal levels. The data were treated graphically or mathematically to distinguish such differences as might occur.

The application of these methods are considered below.

ACCLIMATION

As early as 1895 Davenport and Castle reported on the "acclimatization of organisms to high temperature", and threads of this principle have been variously woven around the theme of temperature relations in fish by Loeb and Wasteneys (1912), Hathaway (1927), Binet and Morin (1934), Sumner and Doudoroff (1938), and more generally by Weigmann (1929, 1930, 1936), Ogle and Mills (1933), Heilbrunn (1943), and others. Yet a great deal of experimental work has been done without adequate regard for the conditioning effects of temperature in the past-history of animals. A study of changes in heat-tolerance for the goldfish, *Carassius auratus*, from both low (4°C.) and moderately high (20°C.) temperature-acclimations (Brett, 1946) led to the general conclusion that rate of acclimation was related to metabolic rate. Thus, at low temperatures, acclimation proceeded at a slow rate but increased to a very rapid rate at high temperatures, probably in geometric progression. Conversely the loss of acclimation to any level of temperature was a comparatively slow process under most conditions.

The thermal history for the egg, alevin and early fry stage for each of the salmon species was 8.8°C. with very little variation. When the fry were about two months old the process of moving them through a series of temperature-acclimations was initiated. The *minimum* standards set for acclimation from the holding-trough temperature of 8.8°C. to any one of the following temperatures were:

To 5°C.	4 weeks at 5°C.
To 10°C.	3 weeks at 10°C.
To 15°C.	3 weeks at 15°C.
To 20°C.	1 week at 15°C., 2 weeks at 20°C.
To 23°C.(or 24°C.)	1 week at 15°C., 1 week at 20°C., 1 week at 23°C.

Repetition of some of the upper lethal-temperature experiments after one and even five months' lapse of time without further change in acclimation gave no significant change in heat-tolerance.

The fish indicated by the holding temperature fully cultured they prove to be more resistant when acclimated from 15°C. to five species next in line lethal-temperature 24°C. show after prolonged exposure to divergent temperatures more resistance about 40% obtained in though at almost completely, the chum and provide com

UPPER LETHAL

The method associates with the method which produces of the data 1914; Doudoroff tracing the temperature is necessary

Each 10 0.5°C. difference chosen by previous non-lethal tests by close inspection plotted graph of an asymmetrical various temperatures fry from 10°C. (1937) that

The first species to be investigated was the spring salmon. Preliminary tests indicated that a maximum acclimation temperature without significant loss in the holding troughs was close to 24°C. This was later substantiated by successfully culturing over 200 young spring salmon at 24°C., at which temperature they proved to be very active and to be good feeders, but had reduced growth rate when compared with groups from lower temperatures, particularly those from 15°C. A level of 24°C. was introduced, therefore, as the highest standard acclimation temperature for all species. By chance one of the most hardy of the five species had been used to set the standard for the others. The pink salmon, next in line by age for progressive acclimation to 24°C., fed poorly. Immediate lethal-temperature experiments following minimum acclimation standards for 24°C. showed a breaking away from the usual temperature-time mortality curve after prolonged exposure to a temperature of 24.5°C. Extrapolation of this divergent trend indicated that 24°C. bordered on 50 per cent lethality for prolonged exposure. Evidence of the unsatisfactory nature of such a high acclimation temperature was convincing in the sockeye fry. The latter species, while apparently more resistant than the pink salmon, showed a complete aversion to feeding in about 40 per cent of cases which later appeared as typical "pin heads", often obtained in hatcheries when young fish do not develop the feeding habit (although at an equable temperature). The growth of the remainder was curtailed almost completely and their activity was quite apparently reduced. Consequently, the acclimation temperature was lowered to 23°C. for the sockeye, the chum and the coho. Insufficient numbers of spring and pink salmon remained to provide comparison at the new acclimation level.

UPPER LETHAL TEMPERATURE

The method of lethal-temperature determination as conducted by Fry and associates has remained basically consistent from its inception (Brett, 1941), with the marked exception of the duration of exposure to given temperatures which produce some but not complete mortality. The analysis and interpretation of the data have changed and expanded considerably (see Fry *et al*, 1942; Brett, 1944; Doudoroff, 1945; Fry *et al*, 1946; Fry, 1947a; Hart, 1947, 1949). Without tracing the history of these changes, a discussion of the present treatment of temperature data and the current terminology used to describe the observations is necessary.

Each lethal bath is regulated to a constant temperature almost exactly 0.5°C. different from the temperature of an adjacent bath and appropriately chosen by preliminary tests to span the conditions from rapid to slow, partial, or non-lethal temperature effects. Records of the times to death for all fish are kept by close inspection. These latter have been called the *resistance times* which, if plotted graphically on normal axes in order of occurrence, take on the appearance of an asymmetrical S-curve. A series of such curves can be plotted, including the various temperatures investigated, as has been done for a sample of spring salmon fry from 10°C. acclimation in Figure 3a. It has been demonstrated (Bliss, 1935a, 1937) that many dosage-mortality curves can be resolved into straight-line

relations if the proper derivative of time is applied and the variation of response is normally distributed within the sample. By converting the axes into probability units as one variable (order of death) and logarithm of time as the other variable (time to death) a linear progression of points is frequently obtained (Figure 3b). The application of this principle to lethal-temperature experiments with fish was shown to be quite appropriate by Fry *et al.* (1946). The hundreds of observations which have since been made on many species of fish (Hart, 1947, 1949; present paper) and found to adhere closely to the above interpretation have added convincingly to the validity of the relation. The normality of the distribution permits application of standard statistical treatments. In addition the mean, median and mode, all coincide in a normal distribution, so the descriptive value of the single figure (50 per cent point) is evident.

From each lethal-temperature experiment a series of median resistance times may be plotted for the corresponding levels of temperature, in the manner of Figure 4a. It is apparent that for every state of acclimation the possibility of a series of such points exists. Thus, an overall picture of the effect of temperature can be constructed. Conveniently enough these curves, in the case of high-temperature tolerance, can be resolved into straight lines by using the logarithm of time against temperature (Figure 4b). A distinct break in the semilogarithmic plot, not otherwise evident, occurs at a progressively earlier point of time in the lines for lower levels of acclimation (usually below 20° to 15°C. for Pacific salmon). The discovery of this break (Fry *et al.*, 1946) and its variable occurrence with acclimation was most significant and has constituted the main difference in experimental procedure from that of earlier investigations. The definition of *lethal temperature* has been that temperature at which 50 per cent of the population is dead after indefinite exposure. The stumbling block in the past has been the duration of the experimental test. Doudoroff (1945) questioned the 14-hour period used by Fry *et al.* (1942) and shortened to 12 hours by Brett (1944). The answer, as indicated above, was provided when breaks in the resistance time-temperature relations showed that mortality from temperature as a primary cause had ceased. The duration, even as long as the seven-day period used for Pacific salmon should be governed by this factor since it varies for different species and different acclimations. As long as the resistance times continue to be finite the fish are considered to be in a *zone of resistance*. Beyond this lies the *zone of tolerance* (Fry, 1947a).

At one acclimation there are any number of resistance times but only one lethal temperature. To distinguish indices derived from high- and low-temperature experiments the terms *upper* and *lower* lethal temperatures are applied respectively.

LOWER LETHAL TEMPERATURE

Temperatures distributed from 0°C. ($0.1^\circ \pm 0.1^\circ\text{C.}$) to 7°C. at one degree intervals were used in lower lethal-temperature determinations. One or two instances of experiments at fractional degrees are reported, but the use of 20 fish per tank from a limited total sample precluded carrying the investigation to a finer point.

There seem to be no reports of mortality from high temperature in the literature. Cessation of resistance and complete loss of equilibrium (Often no reference to lethal-temperature) has been recorded for many species, by reference to the literature. In the case of low temperature, the source of constant mortality from this cause. Usually from a large sample of fish, the equable temperature has been performed. A sample of fish, 1942; Brett, 1944; operations in

Close inspection of the death the chilled fish could be a symptom before covering, a correlation $r = 0.9$ highly significant. The correlation $r = 0.9$ almost entirely from direct observations, a temperature, a

In practice, long, 3 inches, samples for each test temperature regardless of the container was recorded. The container varied, but the mortalities, by percent. The number of fish showing

The median (similar to upper) pattern which any further experiments of line of reason curves to be

There seems to be relatively little trouble in deciding when a fish has died from high temperature, except in a few instances (Brett, 1944; Hart, 1949). A cessation of respiratory movements and muscular contraction accompanied by complete loss of response to stimuli have been regarded as quite decisive criteria. Often no reference concerning such end points is included in reports on upper lethal-temperature experiments. A check on 180 Pacific salmon, including all species, by removing the fish to a lower temperature immediately after "death" was recorded, resulted in no recoveries. On the other hand the depressing effects of low temperatures, producing a type of "suspended animation", have been the source of considerable trouble in establishing satisfactory criteria of death from this cause. Usually groups of fish at a single low temperature have been removed from a large sample and tested for mortality by immersion in water of a more equable temperature, the recoveries being noted over the first 24 hours. This has been performed at intervals throughout the experiment, or, when only a small sample of fish was available, at the end of a given exposure time (Fry *et al.*, 1942; Brett, 1944). The lack of a more direct criterion of death has restricted operations in this field.

Close inspection of Pacific salmon revealed that with the approach of cold-death the characteristically immobile and closely compressed opercula of the chilled fish commence to fan out perceptibly. By systematically recording this symptom before removal of each sample to a testing tank (at 12°C.) for recoveries, a comparison of the "predicted" and "actual" mortalities was made. A highly significant correlation between the two was obtained (coefficient of correlation $r = +.90$, $P_{.01} = .37$), the cases of non-agreement being scattered almost entirely on the side of greater "actual" mortality. Unpredicted recovery was virtually non-existent. Consequently, the resistance times could be tabulated from direct observation of the fish in the lethal baths as in the upper lethal experiments, and the median resistance times plotted for different degrees of low temperature, a system hitherto not employed in low temperature work.

In practice, lots of ten fish were placed in small plastic cylinders (6 inches long, 3 inches in diameter) capped at either end with plastic screening. Two such samples for each species from a given acclimation were inspected at each of the test temperatures. When the number of predicted dead in the first cylinder—regardless of the number dead in the second, had reached 50 per cent, the container was removed to running water at 12°C. The treatment of the second container varied in order to test the prediction value over a greater range of mortalities, but usually contained estimated deaths of between 50 and 100 per cent. The number of *actual* dead was recorded 24 hours later and excluded all fish showing perceptible activity—rarely a questionable category.

The median resistance times, determined by plotting the order of death (similar to upper temperature resistance), when transposed to a graph follow a pattern which has not been possible to convert into a more convenient form by any further resolution of the data. A solution for the problem of duration in the experiments on effects of high temperature was presented. Following the same line of reasoning, though not apparent in this case, a flattening-out of the curves to become asymptotic with the time axis, is indicative of continued, pro-

longed survival of the sample under the corresponding temperature conditions. This latter inflection occurs by at least 5,000 minutes (three and a half days) in practically all cases, and considerably earlier for the *higher* acclimations. Consequently, a limit of 5,500 minutes was set for the duration of all low-temperature work.

A further use of the lethal temperature to delineate the biokinetic range of fishes has been illustrated by the construction of a trapezium relating upper and lower lethal temperatures to acclimation temperature (Fry *et al.*, 1942, and later references). For every stage of acclimation there is a corresponding lethal temperature. In the upper temperature region the lethal and acclimation temperatures approach each other, finally providing an *ultimate upper lethal temperature* (Fry *et al.*, 1946) beyond which no extension of temperature-tolerance is possible for the species as we know it. Such relations for the Pacific salmon have been illustrated in Figures 20 to 24.

PREFERRED TEMPERATURE

The specific aim of the investigation was to work out in some detail the limits of tolerance; it was also possible to carry out research on temperature selection but of a preliminary nature and consequently presented as such.

Measurements of the aggregation of fish in horizontal temperature gradients have been conducted effectively by Doudoroff (1938) and Sullivan (1949). Emphasis must be placed on the horizontal nature of these gradients since *vertical* gradients were employed for the Pacific salmon. One highly significant difference is apparent, namely, that a gravity gradient is inextricably involved in a vertical tank. It has been customary to reduce as far as possible all interfering factors when recording the responses of an organism to a gradient of a given identity. The methods employed in the present instance were to habituate ten previously acclimated fry to feeding freely in the preferendum tank for one week at the same acclimation temperature. No other control was instituted. Live *Daphnia* regularly introduced with finely ground food, and slowly swept around the tank by currents produced through aeration, served to scatter the fish in an irregular manner throughout the tank. Lighting during the habituation period was from overhead 150-watt bulbs and from sunlight through side windows (Figure 2).

On the day of an experiment, feeding was reduced (excessively fed fish tend to sink to the bottom when inactive) and only between 10 p.m. and midnight were observations on distribution in a temperature gradient recorded. At this time lighting, sufficient only to record positions accurately, was produced by two 3-candle-power sources, placed 35 to 40 inches on either side of a middle point of the tank. Any defence of territory which had been exhibited under full illumination was never displayed in the greatly reduced lighting of an experiment. The distribution in various thermal gradients was then noted by counting the number in each cell of known temperature.

STATISTICS

The
and the st
the calcul

The a
in response
These are
the five s

UPPER L

SPRIN
times and
times at
axes, and
acclimate
during th
The letha
(Table X)
has been
mortality

The
the level
parallel w
series of p
that for t
the points
results (u
acclimate
times for
resulted
equalled
served re
by Fry *et*
or lesser

PINK
species to
length sh
acclimati
perature

Comp
ment of Zo

STATISTICAL TREATMENT

The various sources of variability in response to extremes of temperature and the statistical treatment of the data are presented in Appendix I. Most of the calculations were made by the method of analysis of variance.¹

RESULTS

The actual levels of tolerance to extremes of temperature and the differences in response among the five species have constituted the main theme of study. These are presented by dealing with one aspect of temperature and considering the five species collectively under each heading.

UPPER LIMITS OF TEMPERATURE TOLERANCE

SPRING SALMON. Upper limits of temperature-tolerance include the resistance times and lethal temperatures for each acclimation. A typical series of mortality times at different test temperatures has been depicted in Figure 3a on normal axes, and on probit and logarithmic axes in Figure 3b for young spring salmon acclimated to 10°C. At 24.0°C. and below no deaths were recorded for that sample during the 10,000-minute (one week) duration of the experiment (Table IV). The lethal temperature therefore lies somewhere between 24.0° and 24.5°C. (Table XI). In the latter graph the mean of the logarithms of the times to death has been calculated. The very close approximation of the median and mean mortality times is apparent and in agreement with the findings of Fry *et al.*, 1946.

The median times to death have been plotted further in Figure 4a, illustrating the levelling-off of the median resistance times at lower temperatures to become parallel with the time axis. In Figure 4b the resolution of these data into a linear series of points has been achieved. The line A—B drawn almost at right angles to that for the median resistance times for 5°, 10° and 15°C. acclimations denotes the points of time at which continuation of the experiment provides no change in results (up to 10,000 minutes). Although the resistance times of the higher acclimated fish (20° and 24°C.) showed an increased tolerance at comparable times for periods of 1,000 minutes (about 17 hours) and less, continued exposure resulted in continued mortality to a level of death (line B—C) which finally equalled that of the lower acclimation of 15°C. This phenomenon has been observed repeatedly over the higher levels of acclimation for every species studied by Fry *et al.*, (1942, 1946) and Hart (1947, 1949). It is a characteristic, to a greater or lesser extent, of all the species considered in the following presentation.

PINK SALMON. The pink salmon were decidedly the most difficult of the five species to handle in fresh water after the first month of feeding, and for their length showed the least weight (Table III). The difficulties experienced in acclimating this species to 24°C. were noted earlier. Their intolerance to a temperature as high as 24.0°C. is also apparent in the distribution of their resistance

¹Complete tables of data are included in a Ph.D. thesis, 1951, in the libraries of the Department of Zoology, University of Toronto, and the Pacific Biological Station, Nanaimo, B.C.




FIGURE 3a. Times to death at different lethaltest temperatures among young spring salmon acclimated to 10°C. Plotted on arithmetic axes.




FIGURE 3b. Times to death at different lethaltest temperatures among young spring salmon acclimated to 10°C. Plotted on probit X logarithmic axes. Calculated geometric mean resistance times coincide with the median resistance times (at probit 5.0).

TEMPERATURE - °C.

FIGURE

TEMPERATURE - °C.

FIGURE 4b
acclimated




FIGURE 3a. Times to death at different lethaltest temperatures among young spring salmon acclimated to 10°C. Plotted on arithmetic axes.




FIGURE 3b. Times to death at different lethaltest temperatures among young spring salmon acclimated to 10°C. Plotted on probit X logarithmic axes. Calculated geometric mean resistance times coincide with the mean resistance times (at probit 5.0).

TEMPERATURE - °C.
28
26
24
22

FIGURE

TEMPERATURE - °C.
28
26
24
22
10

FIGURE 4b
acclimated




FIGURE 4a. Median resistance times to high temperatures among young spring salmon acclimated to temperatures indicated. Plotted on arithmetic axes.




FIGURE 4b. Median resistance times to high temperatures among young spring salmon acclimated to temperatures indicated. Plotted on arithmetic X logarithmic axes. (See text for further explanation.)

times following a minimum period of acclimation to this level of temperature (Table IV). After 2,000 minutes' exposure to 25.5°, 25.0° and 24.5°C., no increase in their tolerance over that for 20°C. acclimated samples was obtained.

The results for lower acclimations (Figure 5) were in accordance with general expectations except for the 5°C. group. Mortality was so rapid in lethal tanks with temperatures of 22.5° and 23.0°C. that it was decided to test the majority of the remainder of a limited sample at 21.0° and 19.0°C. Only four fish finally




FIGURE 5. Median resistance times to high temperatures among young pink salmon acclimated to temperatures indicated. The encircled point for 5°C. acclimated pinks denotes the use of four fish instead of the normal ten in other points.

remained for an intermediate test at 22.0°C. The reliability of these data is therefore not as great as that for the more orderly points of higher acclimations.

SOCKEYE SALMON. Sockeye salmon, although held at a maximum acclimation of 23°C. (thus 1°C. lower than the highest for pink salmon), also showed an inherent intolerance for such levels of temperature as 24.0°C. and 24.5°C. after prolonged exposure (Table IV, Figure 6). However, of the five species, the sockeye are probably the best adapted to fresh water, often spending two years in lakes before migrating seaward. A completely fresh-water subspecies is also commonly reported (Dymond, 1936). The responses of the young sockeye to high temperatures were quite similar in pattern to those of a fresh-water species of Salmonidae, *Salvelinus fontinalis* (Fry *et al.*, 1946). The lines connecting median resistance times for acclimations of 20°C. and below are fairly regularly spaced and are not significantly different in slope to be distinguished as not parallel (Table X).

The app
what mi
of the to
As a res
drawn b
most lik
mately p

FIGURE
acclim
and 20

CHU
in Table
for expe
1,000 mi
in life al
Chin and
after 9,00
half the s
to 24°C.

COH
the stock
ted some

The apparent variation in slope for the 15°C. acclimated group is not beyond what might be expected from chance. This has been derived from a consideration of the total data mustered in the tables of analysis of variance (Tables VI to X). As a result, in this instance only, the line for the 15°C. acclimation has not been drawn by inspection as the best straight line for the plotted points, but as the most likely relation to be expected on the basis of the total data, that is, approximately parallel.




FIGURE 6. Median resistance times to high temperatures among young sockeye salmon acclimated to temperatures indicated. A somewhat lowered resistance among the 23° and 20°C. acclimated groups for prolonged exposure to 24.5°C., below the expected level, is indicated by the dotted line on the extreme right-hand side.

CHUM SALMON. The results of experiments with chum salmon are recorded in Table VII and Figure 9. This species showed the greatest amount of variability for experiments in which the mean resistance time approached or exceeded 1,000 minutes. Like the pink salmon, the chum normally move to sea quite early in life although some have been maintained in fresh water up to two years by Chin and Kuroda (1935). Acclimation to 23°C. was quite successful. However, after 9,000 minutes' (6 days') exposure to 24°C. in a temperature-tolerance test, half the sample had died, confirming the impossibility of acclimating this species to 24°C.

COHO SALMON. An accident in the early history of the coho salmon eliminated the stock of eggs held in one of two troughs. This loss, while unselective, necessitated some curtailment in the programme of study. At the time of experimentation




FIGURE 7. Median resistance times to high temperatures among young chum salmon acclimated to temperatures indicated.




FIGURE 8. Median resistance times to high temperatures among young coho salmon acclimated to temperatures indicated.

with upper
coho were
the spring
permit mor
In retrospe
between co
temperature

The re
PROLO
high temper
5°C. These
exposure we
In this resp
sockeye ren
salmon, it
conditions
tures have
one week).
The results
Figures 9 an
3,000 minut
that for the
first day (1
period a ra
chum salmon
direction of
perimentati

LOWER LIM

SPRING
lethal levels
species and
with 50 per
total of 20 t
times at 5 p
of removing
checking "p
a single test
mortality.

No size
on heat-tole
ever, were st
such that th
sample was
was on han

with upper lethal temperatures, it was apparent that at higher acclimations the coho were conforming very closely to the reaction times already determined for the spring salmon. The remaining tests in this series were therefore bypassed to permit more intensive investigation in the relatively unexplored low-lethal range. In retrospect this was perhaps unfortunate as it now appears that a difference between coho and spring salmon, where such occurs, is present in their high-temperature tolerance from low acclimations.

The results for the coho appear in Table IV and Figure 8.

PROLONGED EXPOSURE. An experiment concerning long-term exposure to high temperatures was performed with sockeye and chum salmon acclimated to 5°C. These fish do not feed readily at critical high temperatures, so prolonged exposure would inevitably cause death from malnutrition if from no other cause. In this respect the two species are alike. Since they differ in migratory habits, the sockeye remaining in fresh water usually for at least one year longer than chum salmon, it is quite possible that they differ in ability to tolerate fresh-water conditions (cf. Hoar and Bell, 1950). So far, few experiments with high temperatures have involved exposure times in excess of 10,000 minutes (approximately one week), particularly for fish from acclimation temperatures below 15°C. The results for 30,000 minutes' exposure for these two species are recorded in Figures 9 and 10. The sockeye typically showed no additional mortalities beyond 3,000 minutes. The pattern of mortality in the chum salmon was very similar to that for the sockeye, with no further deaths recorded between the end of the first day (1,440 minutes) and the end of a week (10,080 minutes). Beyond this period a rather unexpected but orderly progress of mortality appeared in the chum salmon samples. Speculation as to the cause of death might be made in the direction of the relative intolerance of this species to fresh water; further experimentation is desirable.

LOWER LIMITS OF TEMPERATURE TOLERANCE

SPRING SALMON. The method of determining the resistance times at low-lethal levels of temperature usually involved two samples of ten fish of the same species and acclimation for each test temperature. The removal of one sample with 50 per cent predicted dead changed each test before completion from a total of 20 to a total of 10 subjects, hence the plotting of the individual resistance times at 5 per cent intervals changed to 10 per cent intervals following the time of removing the first sample. The slight inconvenience involved was a result of checking "predicted" against "actual" dead. In this manner all mortalities from a single test temperature could be plotted in determining the time to 50 per cent mortality.

No size effect in Pacific salmon of the same age was demonstrated from data on heat-tolerance (Table XIII). Results of experiments on cold-tolerance, however, were strongly suggestive of a size influence. The method employed was not such that the individual fish, once dead, could be later identified when the whole sample was removed from a lethal bath. A group of fish, some living, some dead, was on hand. By considering only those cases in which at least three but not




FIGURE 9. Times to death at various high temperatures among young sockeye salmon acclimated to 5°C. and tested for 30,000 minutes (approximately three weeks).




FIGURE 10. Times to death at various high temperatures among young chum salmon acclimated to 5°C. and tested for 30,000 minutes (approximately three weeks).

more than
showed nea
sample as
weights we
were: living
statistically

Eviden
the first, o
and was fo
the method
remainder.
recorded o
size factor
firmatory o
comparativ
stitute the

Prelim
had been i
24° and 20
with upper
temperatures
within the
"expensive
for dead th
tests at 1°C

The m
temperatur
of death. 7
throughout

PINK S
climated p
troughs ap
some intol
standing 5
stopped.

SOCKE
mersion in
salmon. Fo
temperatur
Even 10°C
(three days
those take

The s
nounced a
low tempe

more than seven had succumbed, comparison of the sizes among all specimens showed nearly twice as many living fish above the mean length for the total sample as dead fish. Since the measurements were made some time after death, weights were not taken. The mean lengths and standard errors of the two groups were: living = 5.00 ± 0.046 cm., dead = 4.75 ± 0.043 cm. The data show a statistically significant difference (Table XIV).

Evidence was procured for two separate lethal effects of low temperature: the first, occurring only at the lowest temperatures, was very rapid and decisive, and was followed by a second series of deaths after considerable delay. Owing to the methods employed, the first mortalities could not be sorted out from the remainder. They constituted a relatively small fraction of the total dead. Unrecorded observation at the time of an experiment revealed an apparently distinct size factor in this primary phase of death, the smallest fish dying first. Confirmatory experiments will be required for adequate proof. The presence of these comparatively smaller fish in the samples of dead fish recorded above may constitute the main source of difference between the two major groups.

Preliminary tests with spring salmon, before the system of predicting death had been instituted, were performed at 0.5°C . intervals from 0° to 3°C . (Figure 11, 24° and 20°C . acclimations; Table V.) The reasoning, derived from experience with upper lethal experiments, was fallacious in part since the range of low temperatures causing death for 24°C . acclimated salmon was almost twice as great, within the same time limits (covering 6 rather than 3 degrees C.). The doubly "expensive" technique of closely spaced temperature tests coupled with sampling for dead fish throughout the experiment was replaced by the predicting system for tests at 1°C . interval in the remaining cases reported for the spring salmon.

The median resistance times follow a rather varied relation with increasing temperature (Figure 11), and are apparently complicated by more than one cause of death. The sigmoid shape of the curves is one characteristic which persists throughout the remaining species, particularly at higher acclimations.

PINK SALMON. Although a few preliminary tests were made with 10°C . acclimated pink salmon, $5\frac{1}{2}$ months old, mortality in the fresh-water holding troughs appeared during the last two weeks in sufficient proportions to signify some intolerance to these conditions. Small samples were incapable of withstanding 5°C . Attempts to continue work with this species in fresh water were stopped.

SOCKEYE SALMON. The inability of Pacific salmon to tolerate sudden immersion in low temperatures was well exemplified in the work with young sockeye salmon. For survival in nature the necessity for this species to acclimate to low temperatures is emphasized by the results illustrated in Figure 12 (Table V). Even 10°C . acclimated samples succumbed to 2° and 3°C . within 4,000 minutes' (three days') exposure, and a temperature of 0°C . caused some mortality amongst those taken from holding tanks at 5°C .

The sigmoid pattern of the resistance time-temperature curves is most pronounced among the sockeye. An initial period of rapidly increasing resistance to low temperature (for 20° to 10°C . acclimation) is followed by relatively little




FIGURE 11. Median resistance times to low temperatures among young spring salmon acclimated to temperatures indicated. Arrows signify tests at temperatures which caused less than 50 per cent death for 5,500 minutes' exposure if placed above the line, or greater than 50 per cent by the time indicated if below the line.




FIGURE 12. Median resistance times to low temperatures among young sockeye salmon acclimated to temperatures indicated. Arrows used as stated in Figure 11.

change
mortal
increas
least 4
test ter
5,500 m
CH
exhibit
When
4,000 m
the ver
the sca
salmon
CO
to low
nificant
as low
the free
the oth
MIX
observe
very ra
samples
tempera
observe
and dis
He reco
transfer
cutaneo
penetrat
was mo
satisfact
critical
sockeye
ting the
clearly
1°C. cau
tures);
4° to 6°
mortalit
mortalit
caused d
of morta
layed to
is conseq
the samp

change in resistance for two and three degrees' increase in temperature. No mortality occurs within a degree or so above this latter zone even for a considerably increased exposure time. The presence of this "break" at an exposure time of at least 4,000 minutes for fish acclimated to 10°C. and higher (usually occurring at test temperatures of 2°C. and above) supports the overall experimental time of 5,500 minutes for lower lethal-temperature determinations.

CHUM SALMON. The greatest difference in response to low temperatures was exhibited by the chum salmon from different acclimations (Table V, Figure 13). When taken from 23°C. and put at 7°C., 50 per cent mortality occurred by 4,000 minutes. From 10°C., however, mortality at 0°C. was observed only toward the very end of the imposed test time. This was somewhat surprising in view of the scant but indicative findings on the intolerance discovered for the pink salmon, to which the chums appear to be most closely related (Milne, 1948).

COHO SALMON. The data and curves illustrating the median resistance times to low temperatures for coho salmon appear in Table V and Figure 14. The significant mortality among members of this species at 0°C. when acclimated to as low as 5°C. demonstrates how confined the coho are to temperatures above the freezing point of water. In general their reactions show the same trends as in the other Pacific salmon.

MIXED LETHAL EFFECTS. Brief mention was made of two distinct responses observed while studying the lethal effects of low temperatures. The presence of very rapid mortalities in contrast with delayed lethal effects, either between samples at slightly different temperatures, or within samples at given critical temperatures, was apparent from even superficial examination. Doudoroff (1942) observed somewhat similar phenomena among young greenfish, *Girella nigricans*, and distinguished between "primary chill-coma" and "secondary chill-coma". He records that "The initial shock was not manifest until several seconds after transfer to the low temperature, and apparently was not due to stimulation of the cutaneous sense organs, but was produced only when the low temperature had penetrated internally, probably to the central nervous system. Accordingly, it was more delayed in large specimens than in small ones". The discovery of a satisfactory criterion of cold-death among the Pacific salmon permitted a more critical study of the time-temperature-acclimation relations for death. Results for sockeye salmon will be presented, being more extensive but not unique. By plotting the data on probability \times logarithmic paper it was possible to discriminate clearly between the two trends of death. From 23°C. acclimation (Figure 15a), 1°C. caused rapid death for all individuals (as presumably would lower temperatures); 2° to 3°C. split the samples into rapid and delayed deaths as shown; 4° to 6°C. resulted in only delayed deaths; above 6°C., less than 50 per cent mortality was obtained. From 20°C. acclimation (Figure 15b) 0°C. caused rapid mortality; 0.5°C. divided the sample into the two types of death; 1°C. and above caused delayed to no mortality. From 15°C. acclimation (Figure 15c) one case of mortality was observed for a lethal-test temperature of 0°C. followed by delayed to no mortality at higher temperatures. The time to 50 per cent. mortality is consequently affected by per cent occurrence of "primary" cold-deaths within the sample and by the size of the fish which chance to be present in that sample.




FIGURE 13. Median resistance times to low temperatures among young chum salmon acclimated to temperatures indicated. Arrows used as stated in Figure 11.




FIGURE 14. Median resistance times to low temperatures among young coho salmon acclimated to temperatures indicated. Arrows used as stated in Figure 11.




FIGURE 15a. Time




FIGURE 15b. Time




FIGURE 15a. Times to death at various low temperatures among young sockeye salmon acclimated to 23°C. Plotted on probit \times logarithmic axes.




FIGURE 15b. Times to death at various low temperatures among young sockeye salmon acclimated to 20°C. Plotted on probit \times logarithmic axes.




FIGURE 15c. Times to death at various low temperatures among young sockeye salmon acclimated to 15°C. Plotted on probit X logarithmic axes.

A reason for greater variability in cold-temperature resistance and the variety of inflections making up the median resistance time-temperature curves can be attributed partly to these causes.

The work of Doudoroff (1942, 1945) was suggestive of an osmoregulatory problem in the delayed cold-deaths, possibly acting as an accessory lethal factor. To test the hypothesis, a preliminary experiment using two groups of 20°C. acclimated sockeye, one in fresh water and the other in Atlantic sea water diluted to 9.9 parts per thousand, was performed at comparable low temperatures (Figure 16). Both rapid and delayed cold-deaths were observed at 0.2°C without significant difference in response within the two media. At 0.7°C. an indication of greater resistance among the last surviving members in the saline solution was present. At 3.2°C. a decided increase in the resistance was observed but not to the point of eliminating death from low temperature in part of the sample.

Two conclusions can be drawn from this experiment. At the lowest lethal levels of temperature a medium of salt water (slightly hypertonic) does not alter the course of death from that observed in fresh water. Such a medium does, however, reduce the lethal effects of low temperature for delayed cold-death when resistance times exceed 1,000 minutes (about 17 hours). From these it would seem that three causes for death are involved: one, a very rapid agent usually effective within 60 minutes of exposure, a second, not so rapid in action, and a third which is related to osmotic balance.

FIGURE 16. acclimated (slightly hypotonic)

The osmotic and Hoar Atlantics. These find lethal temperature agent at from high involved

PREFERR

SPRING of preference lined under surfaces, demonstration of the identity gradient demonstration




FIGURE 16. Times to death at various low temperatures for samples of young sockeye salmon acclimated to 20°C. and tested in fresh water and in sea water diluted to 9.9 parts per thousand (slightly hypertonic). Where a difference in response was apparent a broken line has been used for the salt-water data.

The observation that size appears to have some bearing on resistance to osmotic stress was reported by Wilder (1944) for sea-run speckled trout. Huntsman and Hoar (1940) using salinities of 20 and 28 parts per thousand concluded that Atlantic salmon parr "as they increase in size become more resistant to sea water". These findings in conjunction with the size effect among Pacific salmon at low lethal temperatures support the possibility of an osmotic factor acting as a lethal agent at low temperatures. Conversely the lack of any size relation in deaths from high temperatures might be taken to indicate other than osmotic factors involved in these mortalities.

PREFERRED TEMPERATURES

SPRING SALMON. Some of the inherent problems concerning the determination of preferred temperatures for fish in a vertical temperature gradient were outlined under "Methods". The presence of other gradients (gravity, distance from surfaces, etc.), if impossible to remove or control, must be accounted for by demonstrating the suppression of these in relation to preference for some level of the identity under investigation. By changing the position of the temperature gradient within the tank, without altering other relations, it was possible to demonstrate the selective aggregation of spring salmon in the region of 12° to

13°C. (acclimation of 20°C.) despite a highly significant difference in their position within the tank. This distribution in space, varied yet remaining relatively constant with respect to temperature, has been depicted for this species in Figure 17. An attempt has been made to conform to the relative dimensions of the tank and distribution of the fish, such that the figure presents a "graphical picture" in every sense of the word.




FIGURE 17. Frequency distribution of young spring salmon acclimated to 20°C. in three successive temperature gradients, using the same tank. The calculated preferred temperatures are noted for each distribution.

A few observations which apply to all the species are of significance. No matter what the gradient, even reaching the lethal level, the fish would make feeding excursions to the surface if the appropriate movements of the investigator, which always accompanied feeding, were made. On some, although infrequent, occasions, under the routine procedure of low illumination and night-time recording, no selected temperatures could be determined. Activity of the fish was too great, being more limited by the surfaces of the tank than by any other recognizable feature. No suggestion to account for the increased "excitability" on these particular occasions can be advanced. Adequate feeding and uniformly quiet conditions were always observed prior to introducing a temperature gradient. A natural "exploring" activity is a characteristic of the group.

peratures for acclimation of species tolerant species observed

D. 15°C., experiment was distributed 12° to only can generalization to greater what is

On final and

COMPARISON

RESISTANCE affects levels of and in posed:

(1) acclimation

(2) of resistance

resistance

(3) significance

(4) median

Each variance

different of the

basic conditions

acclimation considered

Species analysis

References

PINK, SOCKEYE, CHUM AND COHO SALMON. The distribution in various temperature gradients as indicated by the preferred temperatures (mean and mode) for acclimations, mostly including 10°, 15° and 20°C. in the Pacific salmon species, are presented in the graphs used to display the zones of temperature tolerance (Figures 20 to 24). The variation in response between the different species was not sufficient to warrant dealing with them separately. Some general observations can be made.

Despite considerable difference in temperature-acclimation, amounting to 15°C., comparatively little difference in preferred temperature was observed experimentally. On the average no greater difference than 3°C. (11° to 14°C.) was displayed between means, and the region of greatest preference lay in the 12° to 14°C. stratum. The pink salmon from 20°C. acclimation constituted the only case in which a preferred temperature as high as 17.7°C. was observed. The general avoidance of temperatures above 15°C. for all species, in spite of acclimation to this level and to 20° and 24°C., was very marked. A tendency to show greater dispersal in the fish from higher acclimations is suggested by the somewhat larger standard deviations for these samples.

Only a record of the mode was available for 5°C. acclimated sockeye on final analysis. Unfortunately no more supporting data are on hand.

COMPARISON OF TEMPERATURE RESISTANCE

RESISTANCE TIMES TO HIGH TEMPERATURES. The extent to which acclimation affects the resistance times to high temperatures within each species and the levels of tolerance characteristic of each species have been presented graphically and in tabular form. If comparison is to be made a variety of questions may be posed:

- (1) How do these species differ in their resistance times at different levels of acclimation?
- (2) If a difference exists, is it a matter of a difference in slope (rate of change of resistance time with temperature), a difference in overall level of temperature resistance, or a difference in both slope and level?
- (3) What measure of difference can be quoted and with what statistical significance?
- (4) Can the postulated straight-line relation between the logarithm of the median resistance time and temperature be justified?

Each of these questions can be answered by the method of analysis of variance (Appendix I) which permits a sorting out of the sources of specific difference from those resulting from sample variability coupled with interaction of the factors of acclimation, lethal test temperature and resistance time. Three basic comparisons can be made involving two of the variates—species, lethals¹ or acclimations²—in as many cases as exist in terms of the third. These are now considered:

Species × *lethals*, for three levels of acclimation. The results from three analysis-of-variance tables (Table VI) demonstrate a highly significant difference

¹ Refers to lethal test temperatures.

² Refers to acclimation temperatures.

between species ($P < .01$) which increases with lower levels of acclimation, that is, at a lower level of acclimation temperature for a comparable treatment of lethal temperature the difference is more distinct. This is the crucial test. Figures 18a and b illustrate this relation. A further analysis to determine which species are contributing to the difference is considered later. As might be expected, the




FIGURE 18a. Comparative median resistance times to high temperatures among young Pacific salmon acclimated to 20°C. Lines have been drawn for the most and least tolerant species.




FIGURE 18b. Comparative median resistance times to high temperatures among young Pacific salmon acclimated to 15°C.

differences for are also highly

Species \times is a case in wh lethal levels is results in Tab crease at lowe as that for a lower levels of If a lethal ten have little co contrast, then that will best

Lethals \times of-variance ta comparisons. temperatures in each of the species, the so the others (P apparent from

It is poss *species* \times *letha* response amon levels of temp lethality, redu of the three fi the interaction 3.0, and is su *species* \times *letha*

A further may be attrib in the first and the total sum extract each (Fisher and Ya for single degr and can be co Such a searchi findings from made by use o cation of the f

(1) No sig exists between

(2) Spring from that of e

differences for all species in their response to different lethal test temperatures are also highly significant.

Species \times *acclimations*, for four levels of lethal test temperature. This is a case in which the lumping of resistance times of three acclimations for single lethal levels is used to distinguish between species. It can be concluded from the results in Table VII that between species at each lethal level, and with some increase at lower levels, there is a consistent significant difference, yet not as great as that for a series of lethals at one acclimation. The increasing significance at lower levels of lethal temperature is inherent in the study of temperature effects. If a lethal temperature approaching that for boiling water were used, it would have little consequence what species or acclimation characterized the fish. In contrast, therefore, it is the temperatures which just cause distinct mortality that will best demonstrate specific and acclimation differences.

Lethals \times *acclimations*, for different species. These five "4 \times 3" analysis-of-variance tables (computed in Table VIII) complete the series of first-order comparisons. In essence they signify what has already been stated, namely, that temperatures of both acclimation and lethal level have highly significant effects in each of the species (former tables only applied to the group as a whole). One species, the sockeye ($P < .05$), does not show the same degree of significance as the others ($P < .01$). There is a greater variability in this species which is apparent from the diversity in the points plotted in Figure 6.

It is possible to draw up a table of second-order comparison, for example, *species* \times *lethals* \times *acclimations*. The very fact of a difference in temperature-response among the species which resulted in the use of somewhat different lethal levels of temperature, to bridge the cases from non-lethality to fairly rapid lethality, reduces this table for like comparisons below a feasible size. A survey of the three first-order tables, however, reveals that the error term, signified by the interaction component in each table, lies mainly between values of 1.0 and 3.0, and is sufficiently consistent to support the view that the interaction of *species* \times *lethals* \times *acclimations* would also be consistent throughout the relation.

A further step involves assigning the particular amount of variation which may be attributed to each independent comparison (degrees of freedom). Thus, in the first analysis concerning *species* \times *lethals* at 20°C. acclimation (Table VI), the total sum of squares was 1249.02 with 29 degrees of freedom. It is possible to extract each independent comparison by the use of appropriate multipliers (Fisher and Yates, 1948; see Appendix). Since these are then the sums of squares for single degrees of freedom they also equal the mean square or variance (s^2), and can be compared for significance directly with the "error" by an F test. Such a searching analysis has been carried out in this particular case only. The findings from it concerning the *species* \times *lethals*, and the statements already made by use of the more generalized analysis tables submitted, permit the application of the following conclusions to the whole problem (consult Table IX):

- (1) No significant difference in response to upper lethal levels of temperature exists between spring and coho salmon (acclimated to 10°C. and above).
- (2) Spring and coho salmon show a highly significant difference in response from that of either sockeye, pink or chum.

(3) Pink and chum salmon show a barely significant difference from each other, but not from sockeye in either case.

(4) There is a very highly significant "linear fitness" of the logarithmic relation for these data which accounts for 98 per cent of the variance; the balance can be attributed to curvature.

Of the total 29 degrees of freedom, 20 were assigned to the "error" component (Table VI). This component is constituted of the variability of the organisms resulting in slight deviations from the postulated relation and their interaction (*lethals* \times *species*), plus the variations due to experimental procedure which can never be entirely eliminated. By the same method of using appropriate multipliers, 20 separate components each representing one degree of freedom were obtained. A study of these provided no value which in itself was significantly different from that to be expected from the variability of the material. Consequently it can be concluded that:

(5) The *slopes* of the lines relating resistance time to temperature for the five species are not significantly different (acclimations 10° to 20°C.) and

(6) The same relation can be applied to each species with equal confidence, differing only in the temperature *level* at which this relation exists.

RESISTANCE TIMES TO LOW TEMPERATURES. The constancy of relations in the upper temperature levels is in contrast to that for lower temperatures. No systematic testing of the latter data is possible from present knowledge of the subject. A graphical presentation of some of the responses among the young Pacific salmon taken from the same acclimation temperatures (Figure 19a and b) serves to illustrate this phenomenon. A fair similarity in pattern for 23°C. acclimated species becomes progressively more variable with decreasing acclimation. The chum salmon, which at first seemed to be among the most sensitive judging from high acclimations, were the least sensitive from an acclimation of 10°C.

The spring salmon were consistently the most sensitive to the lowest temperatures (0°C.), but showed a rapid increase in resistance at slightly higher temperatures.

A striking intolerance to low temperatures characterizes all five species. Further study in this field of temperature-responses will be necessary to clarify some of the complexity which has appeared in these findings. The possible influence of size has been pointed out. The mean fork-lengths of the fish used are reported in Table III.

ZONES OF THERMAL TOLERANCE

The concept of a zone of thermal tolerance bounded by upper and lower incipient lethal temperatures for the greatest possible range of temperature-acclimation, and terminated by ultimate lethal temperatures, was advanced by Fry *et al.* (1942) for the goldfish. The freezing point of water limited the minimum acclimation to 0°C. for fresh-water fish. By construction of a trapezium relating these confining temperatures, calculation of the area of the zone of tolerance in "degrees Centigrade squared" gave quantitative expression for an otherwise qualitative description. Various species of fish have since been described in this




FIGURE 19a. Co salmon ac




FIGURE 19b. Co salmon




FIGURE 19a. Comparative median resistance times to low temperatures among young Pacific salmon acclimated to 23°C. Arrows used as stated in Figure 11.




FIGURE 19b. Comparative median resistance times to low temperatures among young Pacific salmon acclimated to 15°C. Arrows used as stated in Figure 11.

fashion (Brett, 1944; Fry *et al.*, 1946; Hart, 1947, 1949), all conforming within slight variations to the original pattern for the goldfish. An increasing acclimation temperature has always resulted in an increasing lethal temperature in some fractional proportion. The construction of a diagonal line at an angle of 45° to the axes (Figure 20) has provided a ready means of determining the ultimate upper lethal temperature, since this line represents the locus of all points for which lethal temperature equals acclimation temperature. Characteristically, for upper incipient lethal temperatures, no species has been found to exploit the full possibility suggested by extrapolation of this linear relation (between upper lethal temperature and acclimation temperature), always dropping short as a result of uniform intolerance at the highest acclimations, providing a "plateau" in a graphical presentation. Although the resistance time to a high temperature may be lengthened through higher acclimation, it was pointed out that this resistance is finite and the lethal temperature remains unaltered for the relation designated by the "plateau". These relations are illustrated in Figures 20 to 24 for the Pacific salmon. Some new aspects, differing from the normal trapezium, are apparent. The lethal-temperature points are not always best represented by a straight line, particularly in the lower lethal bracket. A high degree of sensitivity to low temperatures among the Pacific salmon almost confines these species to acclimation temperatures above 0°C . Some death at this level was observed among 5°C . acclimated samples (Table XII). In a preliminary acclimation culture of spring salmon at 3°C ., high mortality occurred in the presence of healthy sockeye, chum and coho salmon of the same temperature history. This intolerance of the springs might account for the rapid falling away of the upper lethal temperature for low acclimations, which is not apparent for the coho salmon, so similar in other respects.

The spring and coho salmon had the greatest tolerance and were practically identical in area (529 and 528 degrees C. squared respectively). The sockeye were intermediate (505 degrees C. squared) and the chum salmon least (468 degrees C. squared). The line relating upper lethal temperature to acclimation temperature for the pink salmon was very similar to that for the chum salmon. The apparent intolerance of the pinks to low temperatures would restrict their zone of tolerance even more than in the case of the chums, placing them lowest in order of eurythermal relations. With the possible exception of the pink salmon, and notwithstanding the variable nature of the lines relating lower lethal temperature to acclimation temperature, most of the difference in areas is a result of difference in upper lethal temperature.

DISCUSSION AND CONCLUSIONS

TIME AND TEMPERATURE

Division of response to extremes of temperature into zones of tolerance and resistance, previously set forth for other fishes, has been appropriate for similar distinctions among the Pacific salmon. Although the pattern of resistance to low temperatures was quite different from that for high temperatures, the same factors of tolerance and resistance were equally applicable.

Heilbr
the temper
These have
and Belehr
"there are
due to the

Fig
pre
wi
Ce
W

extremes of
which an or
temperature
characterist
influenced w
when the ra
sample at t
which no m
has been ex
tures (high
can be deter

Heilbrunn (1943) has compiled a considerable number of records concerning the temperature at which thermal death occurs in a wide variety of organisms. These have been taken largely from reviews by Kanitz (1915), Uvarov (1931), and Belehradek (1935). In criticism of these data Heilbrunn (p. 420) comments "there are very few useful records of heat (or cold) death temperatures. This is due to the fact that many authors have neglected the time factor". Death from




FIGURE 20. Thermal tolerance for young spring salmon in fresh water. The preferred temperatures are represented by a central point for the mean, with limits for one standard deviation dotted above and below. The degree Centigrade in which the mode occurred is represented by a solid vertical line. Where an experiment was performed but resulted in less than 50 per cent mortality a "V" has been inserted.

extremes of temperature is not just dependent on a threshold level below or above which an organism either lives or dies, but may be considered a resultant of both temperature and exposure time. At each lethal level of temperature there is a characteristic rate of dying (rate of mortification, Fry *et al.*, 1946) which may be influenced within limits by acclimation. A threshold level is approached, however, when the rate of dying approaches zero, as in the case calculated for half the sample at the incipient lethal temperature. Consideration of the time beyond which no more mortality may be expected from temperature as a primary cause has been extended in the present study for low lethal temperatures. Temperatures (high or low) which would not cause mortality, regardless of acclimation, can be determined from the zone of tolerance. Very similar limits can also be set




FIGURE 21. Thermal tolerance for young pink salmon. No lower lethal experiments were performed. Preferred temperatures plotted as in Figure 20.




FIGURE 22. Thermal tolerance for young sockeye salmon. Preferred temperatures plotted as in Figure 20.




FIGURE 23. Thermal tolerance for young chum salmon. Preferred temperatures and "V" plotted as in Figure 20.




FIGURE 24. Thermal tolerance for young coho salmon. Preferred temperature plotted as in Figure 20.

by examining the resistance times for various acclimation temperatures as in Figure 25 for sockeye salmon. The points beyond which no more deaths are likely to occur have been indicated by a dotted line. Thus, temperatures between 8° and 21°C. are unlikely to cause death among sockeye no matter what the acclimation, nor what the exposure time. The higher or lower the temperature is above or below these limits, the longer the test time necessary to determine the lethal temperature. A somewhat different series of lethal temperatures would have been quoted if no experiments had been continued beyond 24 hours (1,440 min.) for both upper and lower lethal levels.


Future experiments on temperature-tolerance will continue to add to our knowledge of the factors affecting temperature relations in fish. In the investigation of Pacific salmon, emphasis has been placed on uniform treatment of all samples such that, although the methods and later analyses may change, the differences set forth should remain unaltered. One limitation should be pointed out. The results apply to the stocks of salmon from which the eggs were collected (Table I) and may therefore be traced back to comparatively few females. The possibility of variation both within and between local stocks cannot be disregarded.

COMPARISON WITH SOME OTHER SALMONIDS

The resistance times for six species of salmonids determined for samples acclimated to 20°C. have been plotted in Figure 26 (from Fry, 1947b; present paper). Data for only two of the species of *Oncorhynchus* are included. These two, the spring salmon (*O. tshawytscha*) and the chum salmon (*O. keta*) were respectively the most and least resistant to the same test temperatures; the remaining three species occur at intermediate positions (Figure 18). It will be seen that the two species of *Salmo* occupy positions distinct both from each other and from the remaining species. *Salmo salar*, the Atlantic salmon, is the most resistant of the salmonoid group. The members of the Pacific salmon species lie in a compact series intermediate between *Salmo* and *Cristivomer*, while *Salvelinus fontinalis* approaches the resistance of *Salmo trutta* at the highest test temperatures (28.0° and 28.5°C.) but drops below it for lower temperatures, falling within the range for *Oncorhynchus*. Except for the speckled trout (*Salvelinus fontinalis*), no further comparisons are warranted until more experimental data are obtained. A marked difference in the lower lethal relations between the speckled trout and the Pacific salmon exists, sufficient to suggest a qualitative difference rather than a purely quantitative one. The former was found to be resistant to low temperatures by Fry *et al.*, (1946) who report that the lower lethal temperature "was only just above 0°C. when the acclimation temperature was 24°C". This results in a comparatively large zone of tolerance (625 units) exceeding that for Pacific salmon which ranged from 468 to 529 units.

SOME ECOLOGICAL RELATIONS

Limits of tolerance to extremes of temperature among young Pacific salmon have been determined. Significant differences exist. Whether these differences are sufficient to account for some of the distinctive habits which characterize the different species is a matter of conjecture.






FIGURE 25. Median times to death at high and low temperatures for young sockeye. Broken lines indicate levels of temperature causing little or no mortality for continued exposure. Dotted lines join the approximate points of inflection.

Young sockeye are usually lake-dwellers, frequenting the open water and subsurface regions, probably in the cooler temperatures in the vicinity of the thermocline (Ricker, 1937). During the summer they are rarely seen or caught in the shallow, littoral zones of the lake (Brett and McConnell, 1950) where young coho, shown to have a higher temperature-resistance, may be found on occasion in abundance. The sockeye migrate in the spring of the year shortly after the ice has left more northern lakes, or following rising spring temperatures




FIGURE 26. Median resistance times for different salmonoids acclimated to 20°C. Data for other than *Oncorhynchus* from Fry (1947).

in such a lake as Cultus which may remain "open" throughout the year (Foerster, 1937). Termination of the migration is closely correlated with ascending surface temperatures and the presence of a well established epilimnion which Foerster (1937) points out may form a temperature block. Coho continue to migrate downstream after sockeye have ceased to move.

Young pink and chum usually do not experience warm lake or river waters. Their migration to sea almost immediately after hatching in the spring of the year eliminates any such high-temperature experience at this stage of life. Their intolerance to fresh water would probably be more pronounced at higher temperatures. The limited success of Chin and Kuroda (1935) in holding chum salmon in fresh water beyond one year depended in part on low temperatures. The interaction of thermal and osmotic stresses requires investigation.

The spring salmon, although a first year migrant to sea, may be found during the summer in streams and rivers, and is more like the coho in this res-

pect. These of their re- remain un- indicate a waters and

On the siderations be the mos This group made on t

Five s- spring (*O. nerka*), the averaging periments

Two r- for acclima- tion and i- four month- temperatur- temperatur- lower leth-

RESIST- of analysis- acclimation

(1) No exists betw-

(2) Sp- from that

(3) Pi- other, but

(4) T- relation for temperatur-

(5) T- the five spe-

(6) T- differing or

The u- 25.1°C., co-

CRITER- the approx- perceptibly

pect. These two species were the most tolerant to high temperatures. The problem of their relative abilities to remain active at high and low temperatures must remain unanswered for the present. Observations, both field and laboratory, indicate a greater ability to feed and sport (jumping, darting, etc.) in warm waters among the spring and coho than among the remaining three species.

On the basis of morphological studies, rates of growth and life history considerations, Milne (1948) concluded that the pink and chum salmon appeared to be the most specialized, and the spring and coho probably the most primitive. This grouping of the species within the genus is in accord with distinctions made on the basis of temperature-tolerance.

SUMMARY

Five species of Pacific salmon are found in North American waters, the spring (*Oncorhynchus tshawytscha*), the pink (*O. gorbuscha*), the sockeye (*O. nerka*), the chum (*O. keta*) and the coho (*O. kisutch*). Young of these species, averaging 4 to 5 cm. in length and 1 gm. in weight, were used in a series of experiments concerning tolerance to high and to low temperatures.

Two months after hatching, each stock of fish was divided into five groups for acclimation to 5°, 10°, 15°, 20° and 23°C. (24°C. in spring and pink); acclimation and lethal-temperature experiments continued throughout the following four months. Resistance times were determined at intervals of 0.5°C. for high temperatures and 1.0°C. for low temperatures (0.0°C. and above). Upper lethal temperatures were calculated for exposures of 10,000 minutes (one week) and lower lethals for exposures of 5,500 minutes (approximately four days).

RESISTANCE TO HIGH TEMPERATURES. A statistical analysis, using the methods of analysis of variance for test temperatures ranging from 24.5° to 27.5°C. and acclimations of 10° to 20°C., for all species, led to the following conclusions:

- (1) No significant difference in response to upper lethal levels of temperature exists between spring and coho salmon.
- (2) Spring and coho salmon show a highly significant difference in response from that of either sockeye, pink or chum ($P < .01$).
- (3) Pink and chum salmon show a barely significant difference from each other, but not from sockeye in either case ($P = .05$).
- (4) There is a very highly significant ($P < .01$) linear fit of the logarithmic relation for these data (logarithm of the median time to death in relation to the temperature causing that death).
- (5) The slopes of the lines relating log resistance time to temperature for the five species are not significantly different.
- (6) The same relation can be applied to each species with equal confidence, differing only in the temperature level at which this relation exists.

The ultimate upper lethal temperatures for each species were: spring—25.1°C., coho—25.0°C., sockeye—24.4°C., pink—23.9°C., chum—23.8°C.

CRITERION OF DEATH AT LOW TEMPERATURES. It was discovered that with the approach of death from a low temperature the opercula commence to fan out perceptibly. This criterion was shown to be significant when compared with data

from recovery tests at warmer temperatures. Thus, the individual times to death at given low temperatures could be recorded and median resistance times determined for each sample.

RESISTANCE TO LOW TEMPERATURES. The relation between resistance time to low temperature and the level of that temperature has not been resolved into a simple equation as has been the case for heat-tolerance relations. A variable but usually sigmoid to double sigmoid pattern characterized the curves when plotted on normal axes. An initial period of rapidly increasing resistance to low temperature was followed by relatively little change in resistance for two and three degrees increase in temperature (15° to 23°C . acclimation). No mortality usually occurred within a degree or so above this latter zone for a considerably increased exposure time up to 5,500 minutes.

The young salmon were very sensitive to low temperatures. Among the four species tested, the coho and sockeye salmon could not tolerate long exposure (four days) to 0°C . even when taken from holding temperatures as low as 5°C .

The lower lethal temperatures for the highest acclimation, 23°C ., were: spring— 7.4°C ., coho— 6.4°C ., sockeye— 6.7°C ., chum— 7.3°C .

MIXED LETHAL EFFECT OF LOW TEMPERATURES. From acclimation temperatures of 20°C . and above, mixed responses were noted in the lethal baths. A rapid death of all fish occurred at the lowest temperatures. Temperatures slightly above this level caused rapid death in part of the sample followed by a long delay and then death of the remainder. Temperatures somewhat higher, yet still low enough to cause death, did so only after prolonged exposure. By plotting the data on probability \times logarithmic paper it was possible to discriminate clearly between the two trends of death. Exposure to the same low temperatures in lethal baths containing 9.9‰ sea water instead of fresh water (slightly hypertonic) resulted in partially increased tolerance among sockeye salmon.

It therefore appears that three causes for death may be involved: one, a very rapid agent usually effective within 60 minutes of exposure, a second not so rapid in action, and a third which is related to osmotic balance.

SIZE EFFECT. No significant difference in the size of the first and last fish to die from high temperatures was present. For death from low temperatures, however, the size distribution of the dead fish from samples in which 30 to 50 per cent of the fish died showed a significantly lower mean length than in the balance of living fish. The earlier death among smaller fish appeared to be partly the result of greater susceptibility to "rapid" cold-death.

ZONES OF THERMAL TOLERANCE. The zones of thermal tolerance were constructed graphically and the areas calculated in units of degrees Centigrade squared. The high degree of sensitivity to low temperatures, almost confining these species to acclimation temperatures above 0°C ., results in a comparatively low thermal tolerance rating. The spring and coho were almost identical, with 529 and 528 units respectively; the sockeye was next with 505 units; followed by the chum with 468 units. Lacking low lethal-temperature data, the pink salmon cannot be included, but one preliminary experiment suggested a lower resistance than in the chum salmon, and consequently a smaller zone of tolerance. Even

the me
trout S
PR
temper
in accli
differen
acclima
CO
emphas
by pres
resistan
than in
is more
salmon
GE
stenoth
biokine
the spr
the pink
taxonom

BELEHRA
1935

BINET, L
Gén.

BLISS, C
The

Calc

BRETT, J
Soc.

Ser.

Ser.

BURROWS
Cult

CHIN, Z.
long

DAVENPORT
orga

DAY, F. E
DOUDGERO

with

ment

DYMOND,
1936

the most tolerant of the Pacific salmon was considerably below the speckled trout *Salvelinus fontinalis*, with a calculated area of 625 units.

PREFERRED TEMPERATURES. Comparatively little difference in preferred temperature was recorded experimentally, either between species or for differences in acclimation amounting to 15°C. in some instances. On the average, no greater difference than 3°C. (11° to 14°C.) was displayed between means from different acclimations. The region of greatest preference lay in the 12° to 14°C. stratum.

COMPARISON OF TEMPERATURE TOLERANCE. In making comparisons, most emphasis was placed on the resistance times to high temperatures. Experiments by previous investigators with other salmonoids demonstrate a greater heat resistance in two species of *Salmo* and lesser resistance in *Cristivomer namaycush* than in the *Oncorhynchus* group (all acclimated to 20°C.). *Salvelinus fontinalis* is more resistant to temperatures of 27°C. and above than any of the Pacific salmon; however, below this level, the generic distinction does not apply.

GENERAL CONCLUSIONS. The species of Pacific salmon are comparatively stenothermal. An intolerance to low temperatures particularly restricts their biokinetic range. For prolonged exposure (up to one week) to high temperatures the spring and coho salmon were most resistant, the sockeye intermediate and the pink and chum salmon least resistant. These differences are in keeping with taxonomic conclusions and certain ecological distinctions.

REFERENCES

- BELEHRADEK, J. Temperature and living matter. *Protoplasma Monographien*, 8, 1-220, Berlin, 1935.
- BINET, L., AND G. MORIN. Action de la chaleur sur les poissons. *Jour. de Physiol. et de Pathol. Gén.*, 32, 272-379, 1934.
- BLISS, C. I. The calculation of the dosage-mortality curve. *Ann. Appl. Biol.*, 22, 134-167, 1935a. The comparison of dosage-mortality data. *Ann. Appl. Biol.*, 22, 307-333, 1935b. Calculation of the time-mortality curve. *Ann. Appl. Biol.*, 24, 815-852, 1937.
- BRETT, J. R. Tempering versus acclimation in the planting of speckled trout. *Trans. Am. Fish. Soc.*, 70, 397-403, 1941.
- Some lethal temperature relations of Algonquin Park fishes. *Univ. Toronto Stud., Biol. Ser.*, No. 53 (*Pub. Ont. Fish. Res. Lab.*, No. 63), 49 pp., 1944.
- Rate of gain of heat-tolerance in goldfish (*Carassius auratus*). *Univ. Toronto Stud., Biol. Ser.*, No. 52 (*Pub. Ont. Fish. Res. Lab.*, No. 64), 9-28, 1946.
- BURROWS, R. E. Fish-cultural procedures for use at the Grand Coulee stations. *Western Fish-Cultural Investigations, etc.*, 46 pp., Portland, Oregon (no date, mimeographed).
- CHIN, Z., AND K. KURODA. Sur l'élevage du saumon (*Oncorhynchus keta* Walbaum) pendant une longue durée dans l'eau douce. *Keijo Jour. Med.*, 6, 30-40, Keijo, Japan, 1935.
- DAVENPORT, C. B., AND W. E. CASTLE. Studies on morphogenesis. III. On the acclimatization of organisms to high temperature. *Arch. Entwicklungsmech.*, 2, 227-249, 1895.
- DAY, F. Effects of elevated temperature on fishes. *U.S. Fish. Comm. Bull.*, 5, 142-144, 1885.
- DOUDOROFF, P. Reactions of marine fishes to temperature gradients. *Biol. Bull.*, 75, 494-509, 1938.
- The resistance and acclimation of marine fishes to temperature changes. I. Experiments with *Girella nigricans* (Ayres). *Biol. Bull.*, 83, 219-244, 1942.
- The resistance and acclimatization of marine fishes to temperature changes. II. Experiments with *Fundulus* and *Atherinops*. *Biol. Bull.*, 88(2), 194-206, 1945.
- DYMOND, J. R. Some fresh-water fishes of British Columbia. *Rep. B.C. Comm. Fish.*, 1935, 60-73, 1936.

- ELLIS, C. H. Salmon hatcheries. *Ann. Rep. Washington, Dept. Fish.*, 1948, 59-82, 1948.
- FISHER, R. A. The design of experiments. 4th ed., Edinburgh: Oliver and Boyd, 1945.
- FOERSTER, R. E. The relation of temperature to the seaward migration of young sockeye salmon (*Oncorhynchus nerka*). *J. Biol. Bd. Can.*, 3(5), 421-438, 1937.
- FRY, F. E. J. Effects of the environment on animal activity. *Univ. Toronto Stud., Biol. Ser.*, No. 55 (*Pub. Ont. Fish. Res. Lab.*, No. 68), 62 pp., 1947a.
- Temperature relations of salmonoids. *Proc. Nat. Com. Fish. Cult.*, 10th meet., 1947, Appendix "D", 1947b.
- FRY, F. E. J., J. R. BRETT AND G. H. CLAWSON. Lethal limits of temperature for young goldfish. *Rev. Can. de Biol.*, 1, 50-56, 1942.
- FRY, F. E. J., J. S. HART AND K. F. WALKER. Lethal temperature relations for a sample of young speckled trout, *Salvelinus fontinalis*. *Univ. Toronto Stud., Biol. Ser.*, No. 54 (*Pub. Ont. Fish. Res. Lab.*, No. 66), 9-35, 1946.
- GRAHAM, J. M. Some effects of temperature and oxygen pressure on the metabolism and activity of the speckled trout, *Salvelinus fontinalis*. *Can. Jour. Res.*, D, 27, 270-288, 1949.
- HART, J. S. Lethal temperature relations of certain fish of the Toronto region. *Trans. Roy. Soc. Can.*, 41(3), 57-71, 1947.
- Geographic variations of some physiological and morphological characteristics in fish. *Ph.D. thesis, Dept. of Zool., Univ. of Toronto*, 175 pp., 1949.
- HATHAWAY, E. S. Quantitative study of the changes produced by acclimation on the tolerance of high temperatures by fishes and amphibians. *Bull. U.S. Bur. Fish.*, 43(2), 169-192, 1927.
- HEILBRUNN, L. V. An outline of general physiology, 2nd ed., Philadelphia: W. B. Saunders, 748 pp., 1943.
- HOAR, W. S., AND G. M. BELL. The thyroid gland in relation to seaward migration of Pacific salmon. *Can. Jour. Res.*, D, 28, 126-136, 1950.
- HUNTSMAN, A. G. Death of salmon and trout with high temperature. *J. Fish. Res. Bd. Can.*, 5, 485-501, 1942.
- HUNTSMAN, A. G., AND W. S. HOAR. Resistance of Atlantic salmon to sea water. *Jour. Fish. Res. Bd. Can.*, 4, 409-411, 1940.
- HUNTSMAN, A. G., AND M. I. SPARKS. Limiting factors for marine animals. III. Relative resistance to high temperatures. *Contr. Can. Biol.*, 2, 95-114, 1924.
- KANITZ, A. Temperatur und Lebensvorgänge. Berlin, 1915.
- LITCHFIELD, J. T., JR., AND F. WILCOXON. A simplified method of evaluating dose-effect experiments. *Jour. Pharm. and Exp. Therapeut.*, 95(2), 99-113, 1949.
- LOEB, J., AND H. WASTENEYS. On the adaptation of fish (*Fundulus*) to higher temperatures. *Jour. Exp. Zool.*, 12, 543-557, 1912.
- MAUREL, E., AND LAGRIFFE. Action comparée de la chaleur et du froid sur certains poissons. *C. R. Soc. Biol.*, 51, 915-918, 1899.
- MILNE, D. J. The growth, morphology and relationship of the species of Pacific salmon and the steelhead trout. *Ph.D. Thesis, Dept. Zool., McGill Univ.*, 101 pp., 1948.
- OGLE, C., AND C. A. MILLS. Animal adaptation to environmental temperature conditions. *Amer. Jour. Physiol.*, 103, 606-612, 1933.
- REGNARD, P. Action des très basses températures sur les animaux aquatiques. *C. R. Soc. Biol.*, 2, 652-653, 1895.
- RICKER, W. E. Physical and chemical characteristics of Cultus lake, British Columbia. *J. Biol. Bd. Can.*, 3, 363-402, 1937.
- SNEDECOR, G. W. Statistical methods. 4th ed., 1-485, Iowa State College Press, 1946.
- SUMNER, F. B., AND P. DOUDOROFF. Some experiments upon temperature acclimatization and respiratory metabolism in fishes. *Biol. Bull.*, 74, 403-429, 1938.
- SULLIVAN, C. M. Aspects of the physiology of temperature selection in speckled trout (*Salvelinus fontinalis*). *Ph.D. thesis, Dept. of Zool., Univ. of Toronto*, 1-177, 1949.
- UVAROV, B. P. Insects and climate. *Trans. Entomol. Soc. London*, 79, Part 1, 1931.
- WEIGMANN, R. Die Wirkung starker Abkühlung auf Amphibien und Reptilien. *Zeitschr. f. wissenschaft. zool.*, 134, 641-692, 1929.

Zell
Biol
WILDER
spec
YATES,
Scie

Weitere Untersuchungen über die Kältebeständigkeit poikilothermer Wirbeltiere.
Zeitschr. f. wissenschaft. Zool., 136, 195-209, 1930.

Zur Kältebeständigkeit poikilothermer Tiere, Untersuchung an Schnecken und Fischen.
Biol. Zentralblatt, 56, 301-322, 1936.

WILDER, D. G. A comparative study of anadromous and fresh water populations of the eastern speckled trout (*Salvelinus fontinalis*). *Ph.D. thesis, Univ. of Toronto*, 1944.

YATES, F. The design and analysis of factorial experiments. Harpenden: Imperial Bureau of Soil Science, 1937.