

## Susitna-Watana Hydroelectric Project Document ARLIS Uniform Cover Page

|  |   |
|--|---|
| <p><b>Title:</b></p> <p>Study of fish distribution and abundance in the upper Susitna River (Study 9.5), Evaluation of 2014 study modifications in the Black River technical memorandum</p>  | <h1 style="margin: 0;">SuWa 270</h1>  |
| <p><b>Author(s) – Personal:</b></p>  |   |
| <p><b>Author(s) – Corporate:</b></p> <p>R2 Resource Consultants, Inc.</p>  |   |
| <p><b>AEA-identified category, if specified:</b></p> <p>December 17, 2014 technical memorandum filings</p>   |   |
| <p><b>AEA-identified series, if specified:</b></p>   |   |
| <p><b>Series (ARLIS-assigned report number):</b></p> <p>Susitna-Watana Hydroelectric Project document number 270</p>   | <p><b>Existing numbers on document:</b></p>                                     |
| <p><b>Published by:</b></p> <p>[Anchorage, Alaska : Alaska Energy Authority, 2014]</p>   | <p><b>Date published:</b></p> <p>December 2014</p>                              |
| <p><b>Published for:</b></p> <p>Prepared for Alaska Energy Authority</p>   | <p><b>Date or date range of report:</b></p>                                     |
| <p><b>Volume and/or Part numbers:</b></p> <p>Attachment A</p>  | <p><b>Final or Draft status, as indicated:</b></p>                              |
| <p><b>Document type:</b></p> <p>Technical memorandum</p>   | <p><b>Pagination:</b></p> <p>v, 28 p.</p>                                       |
| <p><b>Related work(s):</b></p> <p>Cover letter to this report: Susitna-Watana Hydroelectric Project, FERC Project no. 14241-000; Filing of additional information in response to October 2014 Initial Study Plan Meetings. (SuWa 269)</p> <p>Attachment B (SuWa 271)</p> | <p><b>Pages added/changed by ARLIS:</b></p> <p>Added cover letter (2 pages)</p> |
| <p><b>Notes:</b></p>   |   |

All reports in the Susitna-Watana Hydroelectric Project Document series include an ARLIS-produced cover page and an ARLIS-assigned number for uniformity and citability. All reports are posted online at <http://www.arlis.org/resources/susitna-watana/>





December 17, 2014

Ms. Kimberly D. Bose  
Secretary  
Federal Energy Regulatory Commission  
888 First Street, N.E.  
Washington, D.C. 20426

**Re: Susitna-Watana Hydroelectric Project, Project No. 14241-000**

**Filing of Additional Information in Response to  
October 2014 Initial Study Plan Meetings**

Dear Secretary Bose:

By letters dated January 28, 2014 and October 3, 2014, the Federal Energy Regulatory Commission (Commission or FERC) modified the procedural schedule for the preparation and review of the Initial Study Report (ISR) for the proposed Susitna-Watana Hydroelectric Project, FERC Project No. 14241 (Project).<sup>1</sup> As required by the Commission's January 28 letter, the Alaska Energy Authority (AEA) filed the ISR with the Commission on June 3, 2014 and conducted ISR meetings on October 15, 16, 17, 21, 22, and 23, 2014. As required by the Commission's October 3 letter, AEA will be conducting additional ISR meetings on January 7 and 8, 2015.

In response to the October ISR meetings, AEA and licensing participants identified two technical memoranda that AEA would file with the Commission in December 2014, ahead of the January ISR meetings. In accordance, AEA is filing and distributing the following technical memoranda:

- **Attachment A: *Study of Fish Distribution and Abundance in the Upper Susitna River (Study 9.5) - Evaluation of 2014 Study Modifications in the Black River Technical Memorandum.*** This technical memorandum describes how the modifications to the Study of Fish Distribution and Abundance in the Upper Susitna River (Study 9.5) outlined in the ISR were applied to the Black River during the 2014 study year.
- **Attachment B: *River Productivity Study (Study 9.8) - Fish Diet Sample Size Sufficiency Analysis Technical Memorandum.*** This technical memorandum describes an analysis of stomach contents samples conducted after field

---

<sup>1</sup> Letter from Jeff Wright, FERC Office of Energy Projects, to Wayne Dyok, Alaska Energy Authority, Project No. 14241-000 (issued Jan. 28, 2014); Letter from Jeff Wright, FERC Office of Energy Projects, to Wayne Dyok, Alaska Energy Authority, Project No. 14241-000 (issued Oct. 3, 2014).

sampling was completed to determine whether the sample size targets and the actual sample sizes were sufficient to meet the Study Plan objectives.

Additionally, AEA notes that data collected during the Study Plan implementation, to the extent they have been verified through AEA's quality assurance and quality control (QAQC) procedures and are publicly available, can be accessed at [http://gis.suhydro.org/isr\\_mtg](http://gis.suhydro.org/isr_mtg). On December 17, 2014, AEA posted the following data to this website:

- *Baseline Water Quality Data (Study 5.5)*, 2014 QAQC water quality data and DVRs per the Quality Assurance Project Plan.

Finally, AEA notes that it has posted the agenda and PowerPoint presentations for the upcoming January ISR meetings to the Project website (<http://www.susitna-watanahydro.org/meetings/>).

AEA appreciates the opportunity to provide this additional information to the Commission and licensing participants, which it believes will be helpful in determining the appropriate development of the 2015 study plan as set forth in the ISR. If you have questions concerning this submission please contact me at [wdyok@aidea.org](mailto:wdyok@aidea.org) or (907) 771-3955.

Sincerely,



Wayne Dyok  
Project Manager  
Alaska Energy Authority

Attachments

cc: Distribution List (w/o Attachments)

**Susitna-Watana Hydroelectric Project**  
**(FERC No. 14241)**

**Study of Fish Distribution and Abundance**  
**in the Upper Susitna River (Study 9.5)**

**Evaluation of 2014 Study Modifications**  
**in the Black River**  
**Technical Memorandum**

Prepared for  
Alaska Energy Authority



Prepared by  
R2 Resource Consultants, Inc.

December 2014

## TABLE OF CONTENTS

|           |   |           |
|-----------|---|-----------|
| <b>1.</b> | <b>Introduction.....</b>  | <b>1</b>  |
| <b>2.</b> | <b>Sampling Decision: Increased Sampling Effort in Select Upper River Tributaries....</b> | <b>1</b>  |
| 2.1.      | Sampling in 2013 .....  | 1         |
| 2.2.      | Measures of Sampling Sufficiency .....  | 1         |
| 2.3.      | Increased Sampling Effort .....   | 2         |
| 2.4.      | Implementation of Increased Tributary Sampling Effort in 2014 .....                       | 2         |
| <b>3.</b> | <b>Results of Comparison of sampling protocols in the Black River .....</b>               | <b>3</b>  |
| 3.1.      | Fish Distribution .....   | 3         |
| 3.1.1.    | Species Richness .....  | 3         |
| 3.2.      | Relative Fish Abundance .....   | 4         |
| 3.2.1.    | CPUE .....  | 4         |
| 3.3.      | Fish-Habitat Associations .....   | 6         |
| <b>4.</b> | <b>Discussion.....</b>  | <b>6</b>  |
| <b>5.</b> | <b>Literature Cited .....</b>   | <b>7</b>  |
| <b>6.</b> | <b>Tables .....</b>   | <b>9</b>  |
| <b>7.</b> | <b>Figures.....</b>   | <b>21</b> |

**LIST OF TABLES**

Table 2.3-1. 2013-2014 tributary sampling summary and proposed future Upper River tributary sampling length targets. .... 9

Table 2.4-1. Black River sample unit length (meters) by tributary channel /macrohabitat and mesohabitat type for GRTS sampling approach 2013 and 2014. .... 10

Table 2.4-2. Black River mesohabitat unit count (number of replicate mesohabitat units) by tributary macrohabitat and mesohabitat type for GRTS sampling approach 2013 and 2014. .... 11

Table 3.1-1. Total observations of fish species in the Black River by habitat category using full and subsampling approaches during 2014. .... 12

Table 3.1-2. Summary of sampling sufficiency measures for the Black River in 2013 and 2014. .... 13

Table 3.2-1. Summary of juvenile Arctic grayling CPUE for backpack electrofishing in fish/hour by habitat type in the Black River for three sampling periods in 2014. .... 13

Table 3.2-2. Summary of subadult/adult Arctic grayling CPUE for backpack electrofishing in fish/hour by habitat type in the Black River for three sampling periods in 2014. .... 14

Table 3.2-3. Summary of total Arctic grayling CPUE for backpack electrofishing in fish/hour by habitat type in the Black River for three sampling periods in 2014. .... 15

Table 3.2-4. Summary of juvenile burbot CPUE for backpack electrofishing in fish/hour by habitat type in the Black River for three sampling periods in 2014. .... 16

Table 3.2-5. Summary of subadult/adult burbot CPUE for backpack electrofishing in fish/hour by habitat type in the Black River for three sampling periods in 2014. .... 17

Table 3.2-6. Summary of total burbot CPUE for backpack electrofishing in fish/hour by habitat type in the Black River for three sampling periods in 2014. .... 18

Table 3.2-7. Summary of total sculpin CPUE for backpack electrofishing in fish/hour by habitat type in the Black River for three sampling periods in 2014. .... 19

Table 3.3-1. Total observations of fish species in the Black River by season and mesohabitat type using full and subsampling approaches during 2014. .... 20

**LIST OF FIGURES**

Figure 3.1-1. Species accumulation curves from the Black River GRTS sampling sites during full and subsampling in 2014. .... 21

Figure 3.2-1. Boxplots comparing 2014 subsampling to 2014 full sample in the Black River based on CPUE for backpack electrofishing in fish/hour for juvenile Arctic grayling during three sampling events. .... 22

Figure 3.2-2. Boxplots comparing 2014 subsampling to 2014 full sample in the Black River based on CPUE for backpack electrofishing in fish/hour for subadult/adult Arctic grayling during three sampling events. .... 23

Figure 3.2-3. Boxplots comparing 2014 subsampling to 2014 full sample in the Black River based on CPUE for backpack electrofishing in fish/hour for total Arctic grayling during three sampling events..... 24

Figure 3.2-4. Boxplots comparing 2014 subsampling to 2014 full sample in the Black River based on CPUE for backpack electrofishing in fish/hour for juvenile burbot during three sampling events..... 25

Figure 3.2-5. Boxplots comparing 2014 subsampling to 2014 full sample in the Black River based on CPUE for backpack electrofishing in fish/hour for subadult/adult burbot during three sampling events..... 26

Figure 3.2-6. Boxplots comparing 2014 subsampling to 2014 full sample in the Black River based on CPUE for backpack electrofishing in fish/hour for total burbot during three sampling events..... 27

Figure 3.2-7. Boxplots comparing 2014 subsampling to 2014 full sample in the Black River based on CPUE for backpack electrofishing in fish/hour for total sculpin during three sampling events..... 28

## LIST OF ACRONYMS AND SCIENTIFIC LABELS

| <b>Abbreviation</b> | <b>Definition</b>   |
|---------------------|---|
| ADF&G               | Alaska Department of Fish and Game                                  |
| AEA                 | Alaska Energy Authority   |
| CPUE                | Catch per unit effort   |
| CW                  | Channel width   |
| FDA UP              | Study of Fish Distribution and Abundance in the Upper Susitna River |
| FERC                | Federal Energy Regulatory Commission                                |
| GRTS                | Generalized random tessellation stratified sampling                 |
| IP                  | Implementation Plan   |
| ISR                 | Initial Study Report  |
| km                  | Kilometer   |
| km <sup>2</sup>     | Square kilometers   |
| m                   | Meter   |
| MC                  | Main Channel  |
| Mi                  | Mile  |
| mi <sup>2</sup>     | Square miles  |
| OC                  | Off-Channel   |
| OCH                 | Off-Channel Habitat   |
| PRM                 | Project river mile  |
| RSP                 | Revised Study Plan  |
| SE                  | Standard Error  |
| SPD                 | Study Plan Determination  |
| SR                  | Species richness  |
| TSR                 | True species richness   |



## 1. INTRODUCTION

In 2013, AEA's study teams conducted the first year of data collection for the Fish Distribution and Abundance in the Upper Susitna River Study (Study 9.5). Objective 1 of the Study of Fish Distribution and Abundance in the Upper Susitna River was to describe the seasonal distribution, relative abundance (as determined by catch per unit effort [CPUE], fish density, and counts), and fish-habitat associations of resident fishes, juvenile anadromous salmonids, and the freshwater life stages of non-salmon anadromous species (RSP Section 9.5.1; AEA 2012). Sampling in 2013 was effective at documenting fish distribution (Task A). Relative abundance estimates were effectively generated for all sampled habitats (Task B). However, analysis of habitat associations (Task C) was limited by the low number of off-channel habitats in the mainstem Susitna River and the low number of rare habitat types in Upper River tributaries.

Modifications to the Study Plan were presented in Part C, Section 7 of the Initial Study Report (ISR) filed with FERC June 3, 2014 (AEA 2014a). AEA implemented the following proposed modifications in 2014 to gather additional information; to meet study plan objectives; and better inform the second study year. This technical memorandum describes how the modifications to the Study of Fish Distribution and Abundance in the Upper Susitna River (Study 9.5) outlined in the ISR were applied to the Black River during the 2014 study year.

In response to the October 2014 ISR meetings, AEA informed the licensing participants that AEA would be filing this TM with the Commission ahead of the January 2015 ISR meetings.

## 2. SAMPLING DECISION: INCREASED SAMPLING EFFORT IN SELECT UPPER RIVER TRIBUTARIES

### 2.1. Sampling in 2013

The April 2013 FERC Study Plan Determination (SPD) recommended scaling sampling in proportion to stream size (FERC 2013, p. B-124). To achieve a spatially-balanced and random sample of fish habitats within Upper River tributaries, the length of the tributaries were divided into sampling panels that were 200, 400, or 800 m long, depending on the tributary drainage area, and the required percentage of stream length was sampled using a generalized random tessellation stratified (GRTS) sampling methodology. The original sampling plan (ISR 9.5 Section 4.1.2.1) was to survey the GRTS panel for mesohabitat types, and to select one unit of each mesohabitat type and sample 40 m (131 ft) of each selected unit. The FERC SPD recommended all the classified mesohabitat units be sampled. However, logistical constraints in 2013 required sub-sampling 100 m (328 ft)-long units within GRTS panels. Specifically, within a selected GRTS panel, fish sampling occurred in either a complete mesohabitat unit or up to 00 m (656328 ft) per mesohabitat for each mesohabitat type present (ISR 9.5 Section 7.1.2.4).

### 2.2. Measures of Sampling Sufficiency

Post-season analysis indicated that the 2013 tributary sampling program was effective at documenting the fish species present within Upper River tributaries. The analysis consisted of

comparing the total number of species found in a tributary, referred as observed species richness (SR), and an estimate of true species richness (TSR) in a tributary (Cochran 1977).

However, as indicated in Section 7.1.2.4 of Study 9.5 Initial Study Report (AEA 2014b), the 2013 sub-sampling may have been inconsistent with the intent of the April 2013 FERC SPD, with smaller basins receiving proportionally more effort, and larger basins receiving proportionally less. In addition, a post-2013 field season review of the remote video within each GRTS panel indicated that there were some habitat types that were under-represented in 2013 fish sampling and the fish-habitat association analysis likely would benefit from additional replicates.

Sampling sufficiency for characterizing fish distribution is often evaluated in relation to channel width (Paller 1995, Patton et al. 2000, Hughes et al. 2002, Maret and Ott 2003, Reynolds et al. 2003, Kirsch et al. 2014). Fish sampling and habitat surveys completed in 2013 provided channel width information that was not available to incorporate into the Implementation Plan (AEA 2013). The AEA study team reviewed the 2013 sampling effort in the context of field measurements of channel width to prioritize additional sampling. Kirsch et al. (2014) recommended sampling lengths of 40 wetted channel widths for wadeable streams, 120 channel widths for nonwadeable streams in basins with a watershed area of 100-300 km<sup>2</sup> (38.6 – 115.8 mi<sup>2</sup>), and more than 140 channel widths in nonwadeable streams in larger drainage basins.

### **2.3. Increased Sampling Effort**

AEA proposed to apply the recommendation from Kirsch et al. 2014 for determining the length of Upper River tributaries to sample during the next year of sampling as described in Upper River technical memorandum filed September 17, 2014 (R2 Resource Consultants 2014). The stream-specific sample length changes for all Upper River tributary waters were presented in the September 2014 technical memorandum and are included in Table 2.3-1 for ease of access.

AEA proposed to maintain the spatial configuration of the original GRTS panel sampling and apportioned the additional sampling length within the existing panels by selecting the number of fully-sampled panels necessary to achieve the sampling length target as described in ISR Section 9.5.7.1.2.4. However, in the Black River the total length of main channel habitat within GRTS panels selected for sampling in 2013 was short target sampling lengths, so two more panels were added to the 2014 fish surveys.

### **2.4. Implementation of Increased Tributary Sampling Effort in 2014**

As described in ISR Section 9.5.7.1.2.4, AEA implemented the recommended increase in sampling in the Black River in 2014. The proposed increase in sampling length for the Black River was more than triple the effort expended in 2013 (Table 2.4-1). In 2013, the 100 m (328 ft) sub-sampling approach occurred in six GRTS panels (Panels 1, 2, 4, 6, 7, 9) and resulted in sampling of 11 mesohabitat units within 1,050 m (0.65 mi) of sample unit length (Table 2.4-2). In 2014, sampling the full length of all available main channel mesohabitats and 20x wetted-widths of off-channel habitats present within in the same six panels resulted in 19 mesohabitat units for a total length of 2,724 m (1.69 mi) sampled (Table 2.4-2). In order to achieve the target length of 3,178 m (1.97 mi) of recommended sampling (Table 2.3-1), two additional 400 m

panels were added using the GRTS methodology (Panels 3 and 5). These eight panels also included 402 m of off-channel habitats so the total effort in 2014 completed surveys in 28 mesohabitat units for a total of 3,619 m (2.25 mi).

### **3. RESULTS OF COMPARISON OF SAMPLING PROTOCOLS IN THE BLACK RIVER**

The 2014 sampling was conducted so as to first replicate the 2013 survey length and then extend the sampling to the full targeted sample length. Breaking the data in this way facilitated comparison between the subsample and full sample approaches and avoided concerns about interannual variability of data that could result from comparing 2013 and 2014 data sets. For the remainder of the memorandum, the replicated data set is referred to as the 2014 subsample while the data collected from the fully expanded effort is referred to as the ‘full sample’. The purpose of such a comparison was to determine if completing the full sampling approach improved AEA’s ability to meet study objectives. The evaluation that follows is based on various species metrics including fish distribution, species richness, relative abundance, and fish-habitat associations.

#### **3.1. Fish Distribution**

Five fish species were identified in the Black River system during the full 2014 sampling effort: Arctic grayling (*Thymallus arcticus*), burbot (*Lota lota*), longnose sucker (*Catostomus catostomus*), sculpin (Cottid sp.), and round whitefish (*Prosopium cylindraceum*) (Table 3.1-1). These species represent resident salmonid or non-salmonid functional groups and all were observed previously in the Black River. Notably absent in 2014, were anadromous juvenile Chinook salmon (*Oncorhynchus tshawytscha*), observed in Black River during subsampling in 2013.

In replicating the 2013 survey effort, the 2014 subsample resulted in the collection of three of the five species documented: Arctic grayling, burbot, and sculpin. Longnose sucker and round whitefish were found in the extended survey length in low numbers; only 1 longnose sucker and 5 whitefish were observed out of 3,193 total fish observations (Table 3.1-1). Due to their relative rarity in the Black River habitats, the subsampling approach was insufficient to reliably detect these species.

##### **3.1.1. Species Richness**

The observed species richness (SR), or the total number of species found in the Black River in 2014, and an estimate of true species richness (TSR) following the concepts of Cochran (1977) were used as measures of sampling sufficiency, indicative of the success of the full and subsampling approaches in detecting species’ presence. The SR and TSR values, and when these metrics were first achieved within the GRTS panel matrix, are shown in Table 3.1-2 and depicted in Figure 3.1-1. As discussed above, the observed SR from the full sample was five species collected from eight GRTS panels. The five species were detected within the first two GRTS panels. The estimated TSR for the Black River was 5.6 species. The subsample returned only three species from six GRTS panels and all three were detected in the first site. The SR and TSR

were both calculated at 3.0 fish for the subsampling effort. These calculations highlight the limitations of this analysis when all species are either very common (observed at all six sites) or very rare (never observed).

## 3.2. Relative Fish Abundance

Fish counts across all sites, mesohabitats, sampling methods, and seasons are shown in Table 3.1-1 for the full and subsamples. Sculpin dominated the catches followed by Arctic grayling, and burbot using both the full and sub-sampling methods. Since the full sample effort totaled 3.4 times the stream distance of the subsampling ( $3,619/1,050\text{m} = 3.4\text{x}$ ), it was expected that the numeric factor between fish counts of fish from the two approaches should be in the range of 3.4x (or conversely, the subsample ran slightly less than 30 percent of the full sample). As expected, the count expansion factors for Arctic grayling, burbot, and sculpin ranged between 2.7 and 3.9x (Table 3.1-1). This finding suggests the subsampling approach provided consistent information compared to the full sampling regarding the most abundant species.

### 3.2.1. CPUE

Fish abundance information can be somewhat biased as a function of sampling gear type and the level of effort expended during the surveys. For this reason fish counts are often reported in terms of relative fish abundance and the counts are normalized with respect to effort. For comparative purposes, catch per unit of effort (CPUE) was calculated for the most frequently used sampling technique performed during the 2014 surveys, backpack electrofishing.

Analyses were conducted for the three species most often observed, with several lifestage groupings:

- Arctic grayling juveniles, subadult/adults, and total of all lifestages,
- Burbot juveniles, subadults/adults, and total of all lifestages, and
- Sculpin total of all lifestages.

CPUE was estimated as catch per hour of shocking time for each species/lifestage combination within each mesohabitat unit sampled using backpack electrofishing. Mesohabitats were the primary mesohabitat sampling units and were sampled as clusters defined by GRTS panels. Average CPUE and the associated standard error (SE) for each mesohabitat type was estimated using a combined ratio estimate (Cochran 1977). Calculations were performed with package *survey* (Lumley 2004, Lumley 2014) in the statistical software *R* (version 3.1.1; R Core Team 2014).

CPUE for backpack electrofishing results by mesohabitat type and sampling events (seasons) are shown in Tables 3.2-1 through 3.2-7. The mean, standard error, and number of replicate mesohabitat units sampled for the full and subsampling approaches are shown in each of the tables. Boxplots comparing the subsample and the full sample CPUE results for the three species and three sampling events, early summer, late summer, and fall are presented by habitat type and life history stage in Figures 3.2-1 to 3.2-7.

### 3.2.1.1. *Arctic Grayling*

The CPUE for juvenile Arctic grayling ranged between 0 and 10 fish/hr during the full sample survey depending on habitat type. The same metric for the subsample ranged from 0 to 11 fish/hr. The CPUE results were nearly identical for all habitat types and all seasons during the full and subsample surveys (Table 3.2-1, Figure 3.2-1). The standard errors were generally lower for the full sample, as expected due to increased sampling. This finding suggests that the main benefit for an expanded sampling effort for juvenile Arctic grayling in terms of relative abundance was an increase in precision.

Given the lower abundances of subadult and adult life stages, it appears the grayling capture rate was slightly greater in the full sample compared to the subsample for boulder riffle, rapid, and upland slough habitats (Table 3.2-2, Figure 3.2-2). A small improvement in accuracy of relative abundance for the full sample was apparent for subadult/adult Arctic grayling.

The total CPUE for all Arctic grayling, regardless of life history stage, indicated similar findings as for the juvenile grayling. Since juveniles comprise at least 63 percent of this total, and since the added benefit ascribed to the subadult/adult class was small, the full sampling effort did not provide considerable improvement in CPUE accuracy for this species compared to what could be determined from the subsample (Table 3.2-3, Figure 3.2-3). However, there were improvements in precision of the estimates, which can be helpful in comparing abundance among habitats.

### 3.2.1.2. *Burbot*

The CPUE for juvenile burbot ranged between 0 and 4 fish/hr during the full sample survey and from 0 to 8 fish/hr for the subsample. The mean CPUE results differed between the full and subsampling surveys (Table 3.2-4, Figure 3.2-4). This finding suggests there may have been added benefit in accuracy as well as precision (reduced SE estimates) for the full sample effort for juvenile burbot.

There were no adult and very few subadult burbot captured during 2014 by any of the sampling approaches. Given the rare occurrences of this life history stage, it appears the burbot capture rate was slightly greater during the full sample compared to the subsample for boulder riffle, and rapid habitats (Table 3.2-5; Figure 3.2-5). A small added benefit for the full sample effort was apparent for subadult burbot.

The total CPUE for all burbot, regardless of life history stage, indicated similar findings as the juvenile burbot. Small gains in accuracy and precision during the full sample were observed, particularly in riffle habitats and in the rapid habitat that was not sampled in the subsample approach (Table 3.2-6, Figure 3.2-6).

### 3.2.1.3. *Sculpin*

Sculpin were the most abundant species observed during the surveys. Given the small overall size of the sculpin, the total of all life histories was evaluated for this species.

The mean CPUE for sculpin ranged between 23 and 82 fish/hr during the full sample survey and from 37 to 91 fish/hr for the subsample. In some habitats, the CPUE data show large differences

in mean estimates between the full and subsample methods, but the mean estimates using the full sample methodology were not always higher or lower (Table 3.2-7, Figure 3.2-7). Thus, the subsample did not result in consistently biased estimates of average CPUE. Precision, however, was substantially better (lower SE) with the full compared to the subsampling approach, with a few exceptions. In these few cases when the subsampling method offered lower standard errors than the full survey, the sample size was very small and likely resulted in an underestimate in the overall variability in CPUE. The full sample resulted in more realistic variability measurements and better precision in estimating CPUE for sculping. This finding suggests that the only added benefit for an expanded sampling effort for sculpin is in small increases in estimating error.

### 3.3. Fish-Habitat Associations

For consistency with the ISR (AEA 2014), this section documents the total observations (counts) of fish species and life history types among mesohabitats during the 2014 sub- and full sampling. The total observations of fish species in the Black River system by season and macrohabitat type are presented in Table 3.3-1. When these count data are reviewed simultaneously with the increased sampling of mesohabitats depicted in Tables 3.2-1 and 3.2-2 (taken from R2 Resource Consultants 2014), it is clear that the 2014 sampling effort resulted in increased replicates of fish counts across habitats, including rarer habitats. This increased replication will better support a full evaluation of fish-habitat association for the USR once the study modification is implemented in Upper River tributaries during the next year of study.

In addition, some general observations based on fish counts by habitat, including seasonal shifts in habitat associations as fish grew and matured and as water temperature declined are presented below.

- Highest counts of Arctic grayling, sculpin, and burbot were in boulder riffle habitat, followed closely by counts in run/glide habitat
- Arctic grayling counts lowest in upland slough habitat
- Overall trend for Arctic grayling, sculpin, and burbot was for reduced counts from summer to fall
- Patterns in habitat associations were similar across life stages for Arctic grayling
- Sculpin found in all habitat types sampled
- Round whitefish and longnose sucker were rare in all Black River mesohabitats

## 4. DISCUSSION

This technical memorandum was prepared to assess whether additional sampling effort improved AEA's ability to meet study objectives including fish distribution, relative abundance, or habitat associations in the Black River. The subsampling approach performed adequately where species and habitats were abundant. The expanded, full sampling approach provided the greatest return with respect to rare habitats and rare species and, thus, confirms the adequacy of the ISR proposed modification. Thus, AEA recommends continuing future surveys using the full sampling approach. After successfully implementing the full sampling approach in the Black River, AEA recommends adopting the tributary sampling modifications and targets from the Initial Study Report 7.1.2.4, as summarized in Section 2.3 above.

In applying this modified approach, the sampling length in all but one tributary would be maintained or increased beyond that accomplished in 2013 (R2 Resource Consultants 2014). The sample length for each tributary will be developed for the length of main-channel to be sampled and will be accomplished by sampling the fewest number of GRTS panels possible to accommodate the target length. The use of the GRTS panel process for selection will ensure that survey sites are spatially balanced throughout the mainstem. In addition, because the target lengths are based on main channel panels, the length of off-channel habitat surveyed will be in addition to the length of sample targets, as was evident for the Black River in 2014. As the application of this modification in the Black River has shown, this modified approach will allow for inclusion of additional mesohabitat replicates and will improve AEA's ability to discuss fish use of habitats for rare species and habitats in Upper River tributaries.

## 5. LITERATURE CITED

- Alaska Energy Authority (AEA). 2012. Revised Study Plan: Susitna-Watana Hydroelectric Project FERC Project No. 14241. December 2012. Prepared for the Federal Energy Regulatory Commission by the Alaska Energy Authority, Anchorage, Alaska. <http://www.susitna-watanahydro.org/study-plan>.
- Alaska Energy Authority (AEA). 2013. Final Susitna River Fish Distribution and Abundance Implementation Plan: Susitna-Watana Hydroelectric Project FERC Project No. 14241. March 2013. Prepared for the Federal Energy Regulatory Commission by the Alaska Energy Authority, Anchorage, Alaska. <http://www.susitna-watanahydro.org/wp-content/uploads/2013/03/Attachment-A.pdf>.
- Alaska Energy Authority. 2014a. Initial Study Report, Salmon Escapement Study, Study Plan Section 9.5, Part C Susitna-Watana Hydroelectric Project, FERC Project No. 14241. June 2014. Anchorage, Alaska. <http://www.susitna-watanahydro.org/type/documents>.
- Alaska Energy Authority (AEA). 2014b. Initial Study Report: Susitna-Watana Hydroelectric Project FERC Project No. 14241. June 2014. Prepared for the Federal Energy Regulatory Commission by the Alaska Energy Authority, Anchorage, Alaska. <http://www.susitna-watanahydro.org/type/documents>.
- Cochran, W.G. 1977. Sampling Techniques. Third Edition. John Wiley & Sons, New York.
- Federal Energy Regulatory Commission (FERC), Office of Energy Projects. 2013. Study Plan Determination for the Susitna-Watana Hydroelectric Project. Federal Energy Regulatory Commission, Washington, D.C. Issuance: 20130201-3041.
- Hughes, R.M., P.R. Kaufmann, A.T. Herlihy, S.S. Intelmann, S.C. Corbett, M.C. Arbogast and R.C. Hjort. 2002. Electrofishing Distance Needed to Estimate Fish Species Richness in Raftable Oregon Rivers. North American Journal of Fisheries Management. 22(4).
- Kirsch, J.M., J.D. Buckwalter, and D.J. Reed. 2014. Fish inventory and anadromous cataloging in the Susitna River, Matanuska River, and Knik River basins, 2003 and 2011. Alaska Department of Fish and Game, Fishery Data Series No. 14-04, Anchorage.

- Lumley, T. (2004). Analysis of complex survey samples. *Journal of Statistical Software* 9(1): 1-19.
- Lumley, T. (2014). “survey: analysis of complex survey samples”. R package version 3.30.
- Maret, T.R. and D.S. Ott. 2003. Assessment of fish assemblages and minimum sampling effort required to determine biotic integrity of large river in southern Idaho, 2002: U.S. Geological Survey Water-Resources Investigations Report 2003-4274, 16 p.
- Paller, M.H. 1995. Relationships among number of fish species sampled, reach length surveyed, and sampling effort in South Carolina Coastal Plain streams. *North American Journal of Fisheries Management*. 15: 110–120.
- Patton, T.M., Hubert, W.A., Rahel, F.J., and Gerow, K.G. 2000. Effort needed to estimate species richness in small streams on the Great Plains in Wyoming. *North American Journal of Fisheries Management* 20: 394–398.
- R Core Team (2014). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- R2 Resource Consultants, Inc. 2014. Proposed 2015 Modifications to Fish Distribution and Abundance Study Plan Implementation Technical Memorandum. Susitna-Watana Hydroelectric Project. September 2014.
- Reynolds, L., A.T. Herlihy, P.R. Kaufmann, S.V. Gregory, and R.M. Hughes. 2003. Electrofishing effort requirements for assessing species richness and biotic integrity in western Oregon streams. *North American Journal of Fisheries Management* 23(2): 450-461.



## 6. TABLES

Table 2.3-1. 2013-2014 tributary sampling summary and proposed future Upper River tributary sampling length targets.

| GRTS Sampled Tributaries         | Drainage Basin Area (km <sup>2</sup> ) | Chinook salmon presence | GRTS Sampling Unit Size (m) | Number of GRTS Population Sample Units | Number of 2013 Sample Sites | Number of mesohabitats sampled 2013 | Meters Sampled 2013 | % Sampled 2013 | Number of mesohabitats sampled 2014 | Meters sampled 2014 | Average Wetted width (m) | Channel Widths Sampled 2013 | Kirsch et al. 2014 target (CW) | Kirsch et al. 2014 target (m) | Kirsch et al. 2014 target (%) | Proposed Change (m) |
|----------------------------------|--|-------------------------|-----------------------------|--|-----------------------------|-------------------------------------|---------------------|----------------|-------------------------------------|---------------------|--------------------------|-----------------------------|--------------------------------|-------------------------------|-------------------------------|---------------------|
| Oshetna River (PRM 235.1)        | 1424.5                                 | yes                     | 800                         | 52                                     | 13                          | 28                                  | 2,604               | 6%             | --                                  | --                  | 36                       | 73                          | 140                            | 5,026                         | 12%                           | 2,422               |
| Black River                      | NA                                     | no                      | 400                         | 24                                     | 6                           | 11                                  | 1,050               | 11%            | 28                                  | 3619                | 23                       | 46                          | 140                            | 3,178                         | 33%                           | 2,128               |
| Goose Creek (PRM 232.8)          | 269.1                                  | no                      | 200                         | 81                                     | 20                          | 38                                  | 3,107               | 19%            | --                                  | --                  | 14                       | 219                         | 120                            | 1,704                         | 11%                           | -1,403              |
| Kosina Creek (PRM 209.1)         | 1036.5                                 | yes                     | 800                         | 24                                     | 6                           | 10                                  | 1,000               | 5%             | --                                  | --                  | 32                       | 31                          | 120                            | 4,522                         | 24%                           | 3,522               |
| Tsisi Creek                      | NA                                     | no                      | 400                         | 23                                     | 6                           | 10                                  | 980                 | 11%            | --                                  | --                  | 14                       | 69                          | 140                            | 1,988                         | 22%                           | 1,008               |
| Watana Creek (PRM 196.9)         | 452.7                                  | yes                     | 400                         | 60                                     | 15                          | 30                                  | 2,561               | 11%            | --                                  | --                  | 11                       | 231                         | 140                            | 1,554                         | 6%                            | --                  |
| Watana Creek Tributary           | NA                                     | no                      | 200                         | 67                                     | 13                          | 18                                  | 1,459               | 11%            | --                                  | --                  | 10                       | 154                         | 140                            | 1,330                         | 10%                           | --                  |
| Unnamed Tributary (PRM 194.8)    | 321.2                                  | no                      | 400                         | 32                                     | 2                           | 4                                   | 300                 | 2%             | --                                  | --                  | 3                        | 88                          | 140                            | 476                           | 4%                            | 176                 |
| <b>GRTS Total</b>                | --                                     | --                      | --                          | <b>454</b>                             | <b>81</b>                   | <b>149</b>                          | <b>13,061</b>       | <b>8%</b>      | --                                  | --                  | --                       | --                          |                                | <b>19,778</b>                 | <b>12%</b>                    | <b>7,853</b>        |
| <b>Direct sample Tributaries</b> |  |                         |                             |  |                             |                                     |                     |                |                                     |                     |                          |                             |                                |                               |                               |                     |
| Jay Creek (PRM 211)              | 160.1                                  | no                      | NA                          | --                                     | NA                          | 8                                   | 324                 | --             |                                     |                     | 14                       | --                          | --                             | --                            | --                            | --                  |
| Unnamed Tributary (PRM 206.3)    | <80.3                                  | no                      | NA                          | --                                     | NA                          | --                                  | --                  | --             | 3                                   | 263                 | 6.9                      | --                          | --                             | --                            | --                            | Direct              |
| Unnamed Tributary (PRM 204.5)    | <80.3                                  | no                      | NA                          | --                                     | NA                          | --                                  | --                  | --             | 2                                   | 330                 | 4.5                      | --                          | --                             | --                            | --                            | Direct              |
| Unnamed Tributary (PRM 197.7)    | <80.3                                  | no                      | NA                          | --                                     | NA                          | --                                  | --                  | --             | 5                                   | 358                 | 7.1                      | --                          | --                             | --                            | --                            | Direct              |
| Deadman Creek (PRM 189.4)        | 453.5                                  | no                      | NA                          | --                                     | NA                          | --                                  | --                  | --             | 5                                   | 357                 | 28.4                     | --                          | --                             | --                            | --                            | --                  |
| <b>Direct Sample Total</b>       | --                                     | --                      | --                          | --                                     | --                          | <b>8</b>                            | <b>324</b>          | --             | <b>15</b>                           | <b>1,308</b>        | --                       | --                          | --                             | --                            | --                            | --                  |

Table 2.4-1. Black River sample unit length (meters) by tributary channel /macrohabitat and mesohabitat type for GRTS sampling approach 2013 and 2014.

| Year              | Trib Hab Type         | Single Channel |            |            | Split Channel |                |            |            | Complex Channel |             |                |            | Off-Channel Habitat |               |               |                | OCH Total | Grand Total |            |            |              |
|-------------------|-----------------------|----------------|------------|------------|---------------|----------------|------------|------------|-----------------|-------------|----------------|------------|---------------------|---------------|---------------|----------------|-----------|-------------|------------|------------|--------------|
|                   | Trib MC/OC HabType    | Main Channel   |            |            | Primary       |                | Secondary  |            | Primary         |             | Secondary      | Tertiary   | Tributary           |               | Upland Slough |                |           |             |            |            |              |
|                   | Mesohabitat           | Boulder riffle | Rapid      | Run/Glide  | Single Total  | Boulder riffle | Riffle     | Run/Glide  | Run/Glide       | Split Total | Boulder riffle | Run/Glide  | Riffle              | Complex Total | Run/Glide     | Boulder Riffle |           |             | Pool       | Run/Glide  |              |
|                   |                       |                |            |            |               |                |            |            |                 |             |                |            |                     |               |               |                |           |             |            |            |              |
| 2013              | Black River: Panel 01 |                |            | 100        | 100           |                |            |            |                 |             |                |            |                     |               |               |                | 100       | 100         | 200        |            |              |
|                   | Black River: Panel 02 |                |            |            |               |                |            |            |                 |             | 100            | 100        |                     | 200           |               |                |           | 100         | 100        | 300        |              |
|                   | Black River: Panel 04 | 100            |            |            | 100           |                |            |            |                 |             |                |            |                     |               |               |                |           |             |            | 100        |              |
|                   | Black River: Panel 06 |                |            |            |               | 100            |            | 100        | 50              | 250         |                |            |                     |               |               |                |           |             |            |            | 250          |
|                   | Black River: Panel 07 | 100            |            |            | 100           |                |            |            |                 |             |                |            |                     |               |               |                |           |             |            |            | 100          |
|                   | Black River: Panel 09 | 100            |            |            | 100           |                |            |            |                 |             |                |            |                     |               |               |                |           |             |            |            | 100          |
| <b>2013 Total</b> |                       | <b>300</b>     | <b>-</b>   | <b>100</b> | <b>400</b>    | <b>100</b>     | <b>-</b>   | <b>100</b> | <b>50</b>       | <b>250</b>  | <b>-</b>       | <b>100</b> | <b>100</b>          | <b>-</b>      | <b>200</b>    | <b>-</b>       | <b>-</b>  | <b>100</b>  | <b>100</b> | <b>200</b> | <b>1,050</b> |
| 2014              | Black River: Panel 01 | 104            |            | 296        | 400           |                |            |            |                 |             |                |            |                     |               |               |                | 127       | 127         | 127        | 527        |              |
|                   | Black River: Panel 02 |                |            |            |               |                | 100        | 245        |                 | 345         |                |            | 55                  |               | 55            |                |           |             | 140        | 140        | 540          |
|                   | Black River: Panel 03 | 43             | 282        | 75         | 400           |                |            |            |                 |             |                |            |                     | 51            | 44            |                |           |             | 95         | 495        |              |
|                   | Black River: Panel 04 |                | 90         |            | 90            |                |            | 310        |                 | 310         |                |            |                     |               |               |                |           |             | 40         | 40         | 440          |
|                   | Black River: Panel 05 |                |            |            |               |                |            |            |                 |             | 400            |            |                     |               | 400           |                |           |             |            |            | 400          |
|                   | Black River: Panel 06 | 150            |            | 100        | 250           | 100            |            |            |                 | 100         |                | 50         |                     |               | 50            |                |           |             |            |            | 400          |
|                   | Black River: Panel 07 | 210            | 190        |            | 400           |                |            |            | 17              | 17          |                |            |                     |               |               |                |           |             |            |            | 417          |
|                   | Black River: Panel 09 | 280            |            |            | 280           |                |            |            |                 |             | 120            |            |                     |               | 120           |                |           |             |            |            | 400          |
| <b>2014 Total</b> |                       | <b>787</b>     | <b>562</b> | <b>471</b> | <b>1,820</b>  | <b>100</b>     | <b>100</b> | <b>555</b> | <b>17</b>       | <b>772</b>  | <b>520</b>     | <b>50</b>  | <b>55</b>           | <b>-</b>      | <b>625</b>    | <b>51</b>      | <b>44</b> | <b>127</b>  | <b>180</b> | <b>402</b> | <b>3,619</b> |

Table 2.4-2. Black River mesohabitat unit count (number of replicate mesohabitat units) by tributary macrohabitat and mesohabitat type for GRTS sampling approach 2013 and 2014.

| Year              | Tributary Habitat Type | Single Channel |          |           | Split Channel  |          |           |           | Split Total | Complex Channel |           |           | Complex Total | Off-Channel Habitat |                |          |           | OCH Total | Grand Total |          |           |
|-------------------|------------------------|----------------|----------|-----------|----------------|----------|-----------|-----------|-------------|-----------------|-----------|-----------|---------------|---------------------|----------------|----------|-----------|-----------|-------------|----------|-----------|
|                   | Trib MC/OC HabType     | Main Channel   |          |           | Primary        |          | Secondary | Primary   |             | Secondary       | Tertiary  | Tributary |               | Upland Slough       |                |          |           |           |             |          |           |
|                   | Mesohabitat            | Boulder riffle | Rapid    | Run/Glide | Boulder riffle | Riffle   | Run/Glide | Run/Glide |             | Boulder riffle  | Run/Glide | Riffle    |               | Run/Glide           | Boulder Riffle | Pool     | Run/Glide |           |             |          |           |
| 2013              | Black River: Panel 01  |                |          | 1         | 1              |          |           |           |             |                 |           |           |               |                     |                | 1        |           | 1         | 2           |          |           |
|                   | Black River: Panel 02  |                |          |           |                |          |           |           |             | 1               | 1         |           |               |                     |                |          | 1         | 1         | 3           |          |           |
|                   | Black River: Panel 04  | 1              |          |           | 1              |          |           |           |             |                 |           |           |               |                     |                |          |           |           | 1           |          |           |
|                   | Black River: Panel 06  |                |          |           |                | 1        | 1         | 1         | 3           |                 |           |           |               |                     |                |          |           |           | 3           |          |           |
|                   | Black River: Panel 07  | 1              |          |           | 1              |          |           |           |             |                 |           |           |               |                     |                |          |           |           | 1           |          |           |
|                   | Black River: Panel 09  | 1              |          |           | 1              |          |           |           |             |                 |           |           |               |                     |                |          |           |           | 1           |          |           |
| <b>2013 Total</b> |                        | <b>3</b>       | <b>-</b> | <b>1</b>  | <b>4</b>       | <b>1</b> | <b>-</b>  | <b>1</b>  | <b>1</b>    | <b>3</b>        | <b>-</b>  | <b>1</b>  | <b>1</b>      | <b>-</b>            | <b>2</b>       | <b>-</b> | <b>-</b>  | <b>1</b>  | <b>1</b>    | <b>2</b> | <b>11</b> |
| 2014              | Black River: Panel 01  | 1              |          | 1         | 2              |          |           |           |             |                 |           |           |               |                     |                |          | 1         |           | 1           | 3        |           |
|                   | Black River: Panel 02  |                |          |           |                | 1        | 1         |           |             | 2               |           | 1         |               |                     |                |          |           | 1         | 1           | 4        |           |
|                   | Black River: Panel 03  | 1              | 2        | 2         | 5              |          |           |           |             |                 |           |           |               | 2                   | 1              |          |           |           | 3           | 8        |           |
|                   | Black River: Panel 04  |                | 1        |           | 1              |          | 1         |           |             | 1               |           |           |               |                     |                |          |           | 1         | 1           | 3        |           |
|                   | Black River: Panel 05  |                |          |           |                |          |           |           |             | 1               |           |           |               |                     | 1              |          |           |           |             | 1        |           |
|                   | Black River: Panel 06  | 1              |          | 1         | 2              | 1        |           |           |             | 1               | 1         |           |               |                     | 1              |          |           |           |             | 4        |           |
|                   | Black River: Panel 07  | 1              | 1        |           | 2              |          |           | 1         |             | 1               |           |           |               |                     |                |          |           |           |             | 3        |           |
|                   | Black River: Panel 09  | 1              |          |           | 1              |          |           |           |             | 1               |           |           |               |                     | 1              |          |           |           |             | 2        |           |
| <b>2014 Total</b> |                        | <b>5</b>       | <b>4</b> | <b>4</b>  | <b>13</b>      | <b>1</b> | <b>1</b>  | <b>2</b>  | <b>1</b>    | <b>5</b>        | <b>2</b>  | <b>1</b>  | <b>1</b>      | <b>-</b>            | <b>4</b>       | <b>2</b> | <b>1</b>  | <b>1</b>  | <b>2</b>    | <b>6</b> | <b>28</b> |

Table 3.1-1. Total observations of fish species in the Black River by habitat category using full and subsampling approaches during 2014.

| Habitat Category        | 2014 Full Sample <sup>a</sup> |                 |                 |             |                              | 2014 Subsample <sup>a</sup> |                 |                 |             |                              |
|-------------------------|-------------------------------|-----------------|-----------------|-------------|------------------------------|-----------------------------|-----------------|-----------------|-------------|------------------------------|
|                         | Burbot                        | Arctic Grayling | Longnose Sucker | Sculpin Sp. | Round Whitefish <sup>b</sup> | Burbot                      | Arctic Grayling | Longnose Sucker | Sculpin Sp. | Round Whitefish <sup>b</sup> |
| Black River Mainstem    | 101                           | 422             | 0               | 2,147       | 5                            | 37                          | 122             | 0               | 736         | 0                            |
| Unnamed Tributary       | 3                             | 52              | 0               | 206         | 0                            | NS                          | NS              | NS              | NS          | NS                           |
| Upland Slough           | 9                             | 10              | 1               | 237         | 0                            | 5                           | 2               | 0               | 207         | 0                            |
| Total Observations      | 113                           | 484             | 1               | 2,590       | 5                            | 42                          | 124             | 0               | 943         | 0                            |
| Factor (Full/subsample) | 2.7                           | 3.9             | -               | 2.7         | -                            | 0.37                        | 0.26            | -               | 0.36        | -                            |

<sup>a</sup> Counts from all sampling methods

<sup>b</sup> Whitefish total includes unidentified species

NS = Not surveyed during 2014 under the subsampling approach

0 = Surveyed in 2014 without any recorded fish observations by any of the collection methods

Table 3.1-2. Summary of sampling sufficiency measures for the Black River in 2013 and 2014.

| Upper River Tributary        | Number of Sample Sites | SR <sup>a</sup> | Site when SR first observed | TSR <sub>H-T</sub> <sup>b</sup> | Site when TSR <sub>H-T</sub> -1 first observed | TSR <sub>H-T</sub> minus SR | Percent of TSR observed |
|------------------------------|------------------------|-----------------|-----------------------------|---------------------------------|--|-----------------------------|-------------------------|
| Black River Subsample 2013   | 6                      | 6               | 3                           | 6.6                             | 3  | 0.6                         | 91%                     |
| Black River Subsample 2014   | 6                      | 3               | 1                           | 3.0                             | 1  | 0.0                         | 100%                    |
| Black River Full Sample 2014 | 8                      | 5               | 2                           | 5.6                             | 2  | 0.6                         | 89%                     |

<sup>a</sup> Observed species richness - the total number of species found in a tributary

<sup>b</sup> Horvitz-Thompson estimate (Cochran 1977) of the true species richness in a tributary

Table 3.2-1. Summary of juvenile Arctic grayling CPUE for backpack electrofishing in fish/hour by habitat type in the Black River for three sampling periods in 2014.

|                                    |              | Arctic Grayling Juveniles    |      |     |                              |      |     |
|------------------------------------|--------------|------------------------------|------|-----|------------------------------|------|-----|
|                                    |              | 2014 Full Sample             |      |     | 2014 Subsample               |      |     |
|                                    |              | Number of Units <sup>1</sup> | Mean | SE  | Number of Units <sup>1</sup> | Mean | SE  |
| Boulder Riffle                     | Early Summer | 12                           | 5.9  | 1.3 | 4                            | 5.0  | 2.9 |
|                                    | Late Summer  | 11                           | 9.9  | 1.9 | 4                            | 11.0 | 0.4 |
|                                    | Fall         | 14                           | 5.2  | 1.2 | 4                            | 3.8  | 1.2 |
| Run/Glide                          | Early Summer | 9                            | 9.1  | 3.0 | 4                            | 7.6  | 3.2 |
|                                    | Late Summer  | 11                           | 4.9  | 1.1 | 6                            | 5.9  | 2.9 |
|                                    | Fall         | 9                            | 3.4  | 1.1 | 4                            | 5.1  | 1.7 |
| Rapids                             | Early Summer | 4                            | 6.6  | 2.9 | 0                            | n/a  | n/a |
|                                    | Late Summer  | 4                            | 1.4  | 1.7 | 0                            | n/a  | n/a |
|                                    | Fall         | 2                            | 7.8  | 1.1 | 0                            | n/a  | n/a |
| Riffles                            | Early Summer | 2                            | 2.3  | n/a | 1                            | 4.6  | n/a |
|                                    | Late Summer  | 1                            | 0.0  | n/a | 0                            | n/a  | n/a |
|                                    | Fall         | 2                            | 9.3  | n/a | 1                            | 7.0  | n/a |
| Upland Sloughs (Pools + Run/Glide) | Early Summer | 5                            | 0.7  | 0.5 | 2                            | 0.0  | n/a |
|                                    | Late Summer  | 4                            | 0.0  | 0.0 | 2                            | 0.0  | n/a |
|                                    | Fall         | 4                            | 1.4  | 0.9 | 2                            | 2.9  | 2.5 |

<sup>1</sup>Replicate mesohabitat units

Table 3.2-2. Summary of subadult/adult Arctic grayling CPUE for backpack electrofishing in fish/hour by habitat type in the Black River for three sampling periods in 2014.

|                                       |              | Arctic Grayling Subadults/Adults |      |      |                              |      |      |
|---------------------------------------|--------------|----------------------------------|------|------|------------------------------|------|------|
|                                       |              | 2014 Full Sample                 |      |      | 2014 Subsample               |      |      |
|                                       |              | Number of Units <sup>1</sup>     | Mean | SE   | Number of Units <sup>1</sup> | Mean | SE   |
| Boulder Riffle                        | Early Summer | 12                               | 1.6  | 0.84 | 4                            | 0.4  | 0.36 |
|                                       | Late Summer  | 11                               | 1.0  | 0.38 | 4                            | 0.0  | n/a  |
|                                       | Fall         | 14                               | 0.6  | 0.23 | 4                            | 0.6  | 0.50 |
| Run/Glide                             | Early Summer | 9                                | 1.1  | 0.51 | 4                            | 1.8  | 0.63 |
|                                       | Late Summer  | 11                               | 0.0  | n/a  | 6                            | 0.0  | n/a  |
|                                       | Fall         | 9                                | 0.4  | 0.30 | 4                            | 0.0  | n/a  |
| Rapids                                | Early Summer | 4                                | 5.2  | 2.00 | 0                            | n/a  | n/a  |
|                                       | Late Summer  | 4                                | 0.0  | n/a  | 0                            | n/a  | n/a  |
|                                       | Fall         | 2                                | 0.0  | n/a  | 0                            | n/a  | n/a  |
| Riffles                               | Early Summer | 2                                | 0.7  | n/a  | 1                            | 0.0  | n/a  |
|                                       | Late Summer  | 1                                | 0.0  | n/a  | 0                            | n/a  | n/a  |
|                                       | Fall         | 2                                | 0.0  | n/a  | 1                            | 0.0  | n/a  |
| Upland Sloughs<br>(Pools + Run/Glide) | Early Summer | 5                                | 0.7  | 0.54 | 2                            | 0.0  | n/a  |
|                                       | Late Summer  | 4                                | 0.0  | n/a  | 2                            | 0.0  | n/a  |
|                                       | Fall         | 4                                | 0.0  | n/a  | 2                            | 0.0  | n/a  |

<sup>1</sup>Replicate mesohabitat units

Table 3.2-3. Summary of total Arctic grayling CPUE for backpack electrofishing in fish/hour by habitat type in the Black River for three sampling periods in 2014.

|                                       |              | Arctic Grayling Total        |      |     |                              |      |     |
|---------------------------------------|--------------|------------------------------|------|-----|------------------------------|------|-----|
|                                       |              | 2014 Full Sample             |      |     | 2014 Subsample               |      |     |
|                                       |              | Number of Units <sup>1</sup> | Mean | SE  | Number of Units <sup>1</sup> | Mean | SE  |
| Boulder Riffle                        | Early Summer | 12                           | 7.5  | 1.7 | 4                            | 5.4  | 3.0 |
|                                       | Late Summer  | 11                           | 11.0 | 1.8 | 4                            | 11.0 | 0.4 |
|                                       | Fall         | 14                           | 5.8  | 1.3 | 4                            | 4.4  | 1.4 |
| Run/Glide                             | Early Summer | 9                            | 10.0 | 2.9 | 4                            | 9.4  | 3.1 |
|                                       | Late Summer  | 11                           | 4.9  | 1.1 | 6                            | 5.9  | 2.9 |
|                                       | Fall         | 9                            | 3.8  | 1.3 | 4                            | 5.1  | 1.7 |
| Rapids                                | Early Summer | 4                            | 12.0 | 3.7 | 0                            | n/a  | n/a |
|                                       | Late Summer  | 4                            | 1.4  | 1.7 | 0                            | n/a  | n/a |
|                                       | Fall         | 2                            | 7.8  | 1.1 | 0                            | n/a  | n/a |
| Riffles                               | Early Summer | 2                            | 3.0  | n/a | 1                            | 4.6  | n/a |
|                                       | Late Summer  | 1                            | 0.0  | n/a | 0                            | n/a  | n/a |
|                                       | Fall         | 2                            | 9.3  | n/a | 1                            | 7.0  | n/a |
| Upland Sloughs<br>(Pools + Run/Glide) | Early Summer | 5                            | 1.5  | 1.1 | 2                            | 0.0  | n/a |
|                                       | Late Summer  | 4                            | 0.0  | n/a | 2                            | 0.0  | n/a |
|                                       | Fall         | 4                            | 1.4  | 0.9 | 2                            | 2.9  | 2.5 |

<sup>1</sup>Replicate mesohabitat units

Table 3.2-4. Summary of juvenile burbot CPUE for backpack electrofishing in fish/hour by habitat type in the Black River for three sampling periods in 2014.

|                                       |              | Burbot Juveniles             |      |      |                              |      |     |
|---------------------------------------|--------------|------------------------------|------|------|------------------------------|------|-----|
|                                       |              | 2014 Full Sample             |      |      | 2014 Subsample               |      |     |
|                                       |              | Number of Units <sup>1</sup> | Mean | SE   | Number of Units <sup>1</sup> | Mean | SE  |
| Boulder Riffle                        | Early Summer | 12                           | 2.1  | 0.52 | 4                            | 2.9  | 1.0 |
|                                       | Late Summer  | 11                           | 4.3  | 1.20 | 4                            | 8.3  | 2.7 |
|                                       | Fall         | 14                           | 1.2  | 0.29 | 4                            | 2.4  | 0.7 |
| Run/Glide                             | Early Summer | 9                            | 2.4  | 0.92 | 4                            | 3.0  | 2.0 |
|                                       | Late Summer  | 11                           | 0.7  | 0.35 | 6                            | 0.4  | 0.2 |
|                                       | Fall         | 9                            | 0.0  | n/a  | 4                            | 0.0  | n/a |
| Rapids                                | Early Summer | 4                            | 4.4  | 3.30 | 0                            | n/a  | n/a |
|                                       | Late Summer  | 4                            | 1.4  | 1.70 | 0                            | n/a  | n/a |
|                                       | Fall         | 2                            | 0.0  | n/a  | 0                            | n/a  | n/a |
| Riffles                               | Early Summer | 2                            | 0.7  | n/a  | 1                            | 0.0  | n/a |
|                                       | Late Summer  | 1                            | 3.3  | n/a  | 0                            | n/a  | n/a |
|                                       | Fall         | 2                            | 0.0  | n/a  | 1                            | 0.0  | n/a |
| Upland Sloughs<br>(Pools + Run/Glide) | Early Summer | 5                            | 0.9  | 0.69 | 2                            | 2.3  | 2.0 |
|                                       | Late Summer  | 4                            | 0.0  | n/a  | 2                            | 0.0  | n/a |
|                                       | Fall         | 4                            | 1.1  | 0.99 | 2                            | 2.1  | 1.9 |

<sup>1</sup>Replicate mesohabitat units



Table 3.2-5. Summary of subadult/adult burbot CPUE for backpack electrofishing in fish/hour by habitat type in the Black River for three sampling periods in 2014.

|                                       |              | Burbot Subadults/adults <sup>1</sup> |      |      |                              |      |     |
|---------------------------------------|--------------|--------------------------------------|------|------|------------------------------|------|-----|
|                                       |              | 2014 Full Sample                     |      |      | 2014 Subsample               |      |     |
|                                       |              | Number of Units <sup>1</sup>         | Mean | SE   | Number of Units <sup>2</sup> | Mean | SE  |
| Boulder Riffle                        | Early Summer | 12                                   | 0.4  | 0.32 | 4                            | 0.0  | n/a |
|                                       | Late Summer  | 11                                   | 0.1  | 0.08 | 4                            | 0.0  | n/a |
|                                       | Fall         | 14                                   | 0.0  | n/a  | 4                            | 0.0  | n/a |
| Run/Glide                             | Early Summer | 9                                    | 0.0  | n/a  | 4                            | 0.0  | n/a |
|                                       | Late Summer  | 11                                   | 0.0  | n/a  | 6                            | 0.0  | n/a |
|                                       | Fall         | 9                                    | 0.0  | n/a  | 4                            | 0.0  | n/a |
| Rapids                                | Early Summer | 4                                    | 1.1  | 0.66 | 0                            | n/a  | n/a |
|                                       | Late Summer  | 4                                    | 0.0  | n/a  | 0                            | n/a  | n/a |
|                                       | Fall         | 2                                    | 0.0  | n/a  | 0                            | n/a  | n/a |
| Riffles                               | Early Summer | 2                                    | 0.0  | n/a  | 1                            | 0.0  | n/a |
|                                       | Late Summer  | 1                                    | 0.0  | n/a  | 0                            | n/a  | n/a |
|                                       | Fall         | 2                                    | 0.0  | n/a  | 1                            | 0.0  | n/a |
| Upland Sloughs<br>(Pools + Run/Glide) | Early Summer | 5                                    | 0.0  | n/a  | 2                            | 0.0  | n/a |
|                                       | Late Summer  | 4                                    | 0.0  | n/a  | 2                            | 0.0  | n/a |
|                                       | Fall         | 4                                    | 0.0  | n/a  | 2                            | 0.0  | n/a |

<sup>1</sup>No adult burbot were collected. <sup>2</sup>Replicate mesohabitat units

Table 3.2-6. Summary of total burbot CPUE for backpack electrofishing in fish/hour by habitat type in the Black River for three sampling periods in 2014.

|                                       |              | Burbot Total                 |      |      |                              |      |     |
|---------------------------------------|--------------|------------------------------|------|------|------------------------------|------|-----|
|                                       |              | 2014 Full Sample             |      |      | 2014 Subsample               |      |     |
|                                       |              | Number of Units <sup>1</sup> | Mean | SE   | Number of Units <sup>1</sup> | Mean | SE  |
| Boulder Riffle                        | Early Summer | 12                           | 3.0  | 0.96 | 4                            | 2.9  | 1.0 |
|                                       | Late Summer  | 11                           | 4.5  | 1.10 | 4                            | 8.3  | 2.7 |
|                                       | Fall         | 14                           | 1.2  | 0.29 | 4                            | 2.4  | 0.8 |
| Run/Glide                             | Early Summer | 9                            | 2.4  | 0.92 | 4                            | 3.0  | 2.0 |
|                                       | Late Summer  | 11                           | 0.7  | 0.35 | 6                            | 0.4  | 0.2 |
|                                       | Fall         | 9                            | 0.0  | n/a  | 4                            | 0.0  | n/a |
| Rapids                                | Early Summer | 4                            | 6.6  | 2.70 | 0                            | n/a  | n/a |
|                                       | Late Summer  | 4                            | 1.4  | 1.70 | 0                            | n/a  | n/a |
|                                       | Fall         | 2                            | 0.0  | n/a  | 0                            | n/a  | n/a |
| Riffles                               | Early Summer | 2                            | 0.7  | n/a  | 1                            | 0.0  | n/a |
|                                       | Late Summer  | 1                            | 3.3  | n/a  | 0                            | n/a  | n/a |
|                                       | Fall         | 2                            | 0.0  | n/a  | 1                            | 0.0  | n/a |
| Upland Sloughs<br>(Pools + Run/Glide) | Early Summer | 5                            | 0.9  | 0.69 | 2                            | 2.3  | 2.0 |
|                                       | Late Summer  | 4                            | 0.0  | n/a  | 2                            | 0.0  | 0.0 |
|                                       | Fall         | 4                            | 1.1  | 0.99 | 2                            | 2.1  | 1.9 |

<sup>1</sup>Replicate mesohabitat units

Table 3.2-7. Summary of total sculpin CPUE for backpack electrofishing in fish/hour by habitat type in the Black River for three sampling periods in 2014.

|                                       |              | Sculpin Total                |      |     |                              |      |     |
|---------------------------------------|--------------|------------------------------|------|-----|------------------------------|------|-----|
|                                       |              | 2014 Full Sample             |      |     | 2014 Subsample               |      |     |
|                                       |              | Number of Units <sup>1</sup> | Mean | SE  | Number of Units <sup>1</sup> | Mean | SE  |
| Boulder Riffle                        | Early Summer | 12                           | 68   | 11  | 4                            | 39   | 5.3 |
|                                       | Late Summer  | 11                           | 53   | 9.9 | 4                            | 77   | 12  |
|                                       | Fall         | 14                           | 30   | 5.3 | 4                            | 37   | 6.8 |
| Run/Glide                             | Early Summer | 9                            | 69   | 9.1 | 4                            | 91   | 11  |
|                                       | Late Summer  | 11                           | 82   | 15  | 6                            | 75   | 15  |
|                                       | Fall         | 9                            | 51   | 9.0 | 4                            | 68   | 20  |
| Rapids                                | Early Summer | 4                            | 46   | 12  | 0                            | n/a  | n/a |
|                                       | Late Summer  | 4                            | 77   | 1.9 | 0                            | n/a  | n/a |
|                                       | Fall         | 2                            | 43   | 15  | 0                            | n/a  | n/a |
| Riffles                               | Early Summer | 2                            | 52   | n/a | 1                            | 54   | n/a |
|                                       | Late Summer  | 1                            | 23   | n/a | 0                            | n/a  | n/a |
|                                       | Fall         | 2                            | 47   | n/a | 1                            | 52   | n/a |
| Upland Sloughs<br>(Pools + Run/Glide) | Early Summer | 5                            | 35   | 11  | 2                            | 63   | 26  |
|                                       | Late Summer  | 4                            | 35   | 19  | 2                            | 70   | 24  |
|                                       | Fall         | 4                            | 45   | 15  | 2                            | 91   | 5.5 |

<sup>1</sup>Replicate mesohabitat units

Table 3.3-1. Total observations of fish species in the Black River by season and mesohabitat type using full and subsampling approaches during 2014.

| Mesohabitat Type            | Study Period | Full 2014 Sampling Approach <sup>a</sup> |                 |                 |                 |                 |                 |                              | 2014 Subsampling Approach <sup>a</sup> |                 |                 |                 |                 |                 |                              |
|-----------------------------|--------------|--|-----------------|-----------------|-----------------|-----------------|-----------------|------------------------------|--|-----------------|-----------------|-----------------|-----------------|-----------------|------------------------------|
|                             |              | Burbot                                   |                 | Arctic Grayling |                 | Longnose Sucker | Sculpin Sp.     | Round Whitefish <sup>b</sup> | Burbot                                 |                 | Arctic Grayling |                 | Longnose Sucker | Sculpin Sp.     | Round Whitefish <sup>b</sup> |
|                             |              | Juvenile                                 | All Life Stages | Juvenile        | All Life Stages | All Life Stages | All Life Stages | All Life Stages              | Juvenile                               | All Life Stages | Juvenile        | All Life Stages | All Life Stages | All Life Stages | All Life Stages              |
| <b>Black River Mainstem</b> |              |  |                 |                 |                 |                 |                 |                              |  |                 |                 |                 |                 |                 |                              |
| Boulder Riffle              | Early Summer | 15                                       | 27              | 66              | 112             | 0               | 504             | 4                            | 6                                      | 11              | 16              | 21              | 0               | 116             | 0                            |
|                             | Late Summer  | 19                                       | 22              | 62              | 92              | 0               | 331             | 0                            | 9                                      | 9               | 17              | 22              | 0               | 120             | 0                            |
|                             | Fall         | 11                                       | 14              | 40              | 58              | 0               | 273             | 0                            | 4                                      | 5               | 7               | 11              | 0               | 67              | 0                            |
| Rapid                       | Early Summer | 5  | 9               | 10              | 26              | 0               | 65              | 0                            | NS                                     | NS              | NS              | NS              | NS              | NS              | NS                           |
|                             | Late Summer  | 4  | 4               | 3               | 6               | 0               | 101             | 0                            | NS                                     | NS              | NS              | NS              | NS              | NS              | NS                           |
|                             | Fall         | 0  | 0               | 6               | 7               | 0               | 35              | 0                            | NS                                     | NS              | NS              | NS              | NS              | NS              | NS                           |
| Riffle                      | Early Summer | 2  | 3               | 3               | 5               | 0               | 96              | 1                            | 1                                      | 2               | 3               | 3               | 0               | 48              | 0                            |
|                             | Late Summer  | 2  | 2               | 0               | 0               | 0               | 11              | 0                            | NS                                     | NS              | NS              | NS              | NS              | NS              | NS                           |
|                             | Fall         | 0  | 0               | 4               | 8               | 0               | 29              | 0                            | 0                                      | 0               | 2               | 6               | 0               | 20              | 0                            |
| Run/Glide                   | Early Summer | 12                                       | 13              | 26              | 39              | 0               | 239             | 0                            | 6                                      | 7               | 18              | 23              | 0               | 141             | 0                            |
|                             | Late Summer  | 6  | 6               | 40              | 50              | 0               | 305             | 0                            | 3                                      | 3               | 19              | 28              | 0               | 142             | 0                            |
|                             | Fall         | 1  | 1               | 14              | 19              | 0               | 158             | 0                            | 0                                      | 0               | 8               | 8               | 0               | 82              | 0                            |
| <b>Unnamed Tributary</b>    |              |  |                 |                 |                 |                 |                 |                              |  |                 |                 |                 |                 |                 |                              |
| Boulder Riffle              | Early Summer | 2  | 2               | 4               | 5               | 0               | 7               | 0                            | NS                                     | NS              | NS              | NS              | NS              | NS              | NS                           |
|                             | Fall         | 0  | 0               | 3               | 5               | 0               | 2               | 0                            | NS                                     | NS              | NS              | NS              | NS              | NS              | NS                           |
| Riffle                      | Late Summer  | 0  | 0               | 3               | 3               | 0               | 8               | 0                            | NS                                     | NS              | NS              | NS              | NS              | NS              | NS                           |
|                             | Fall         | 0  | 0               | 0               | 0               | 0               | 4               | 0                            | NS                                     | NS              | NS              | NS              | NS              | NS              | NS                           |
| Run/Glide                   | Early Summer | 1  | 1               | 11              | 32              | 0               | 2               | 0                            | NS                                     | NS              | NS              | NS              | NS              | NS              | NS                           |
|                             | Late Summer  | 0  | 0               | 0               | 0               | 0               | 24              | 0                            | NS                                     | NS              | NS              | NS              | NS              | NS              | NS                           |
|                             | Fall         | 0  | 0               | 6               | 7               | 0               | 159             | 0                            | NS                                     | NS              | NS              | NS              | NS              | NS              | NS                           |
| <b>Upland Slough</b>        |              |  |                 |                 |                 |                 |                 |                              |  |                 |                 |                 |                 |                 |                              |
| Pool                        | Early Summer | 3  | 4               | 1               | 3               | 1               | 62              | 0                            | 2                                      | 2               | 0               | 0               | 0               | 44              | 0                            |
|                             | Late Summer  | 0  | 0               | 3               | 4               | 0               | 3               | 0                            | NS                                     | NS              | NS              | NS              | NS              | NS              | NS                           |
|                             | Fall         | 1  | 2               | 0               | 1               | 0               | 1               | 0                            | NS                                     | NS              | NS              | NS              | NS              | NS              | NS                           |
| Run/Glide                   | Early Summer | 1  | 1               | 0               | 1               | 0               | 45              | 0                            | 1                                      | 1               | 0               | 1               | 0               | 39              | 0                            |
|                             | Late Summer  | 1  | 1               | 0               | 0               | 0               | 68              | 0                            | 1                                      | 1               | 0               | 0               | 0               | 68              | 0                            |
|                             | Fall         | 1  | 1               | 1               | 1               | 0               | 58              | 0                            | 1                                      | 1               | 1               | 1               | 0               | 56              | 0                            |

<sup>a</sup> Counts from all sampling methods

<sup>b</sup> Whitefish total includes unidentified species

NS = Not surveyed during 2014 under the subsampling approach

0 = Surveyed in 2014 without any recorded fish observations by any of the collection methods

## 7. FIGURES




Figure 3.1-1. Species accumulation curves from the Black River GRTS sampling sites during full and subsampling in 2014. Note: The species accumulation curve generated during subsampling in 2013 is provided for reference.




Figure 3.2-1. Boxplots comparing 2014 subsampling to 2014 full sample in the Black River based on CPUE for backpack electrofishing in fish/hour for juvenile Arctic grayling during three sampling events. The boxes represent the interquartile range (i.e., 1<sup>st</sup> to 3<sup>rd</sup> quartile of data), the black line in the box is the median. The whiskers extend to the full range of the data unless one or more data points are extreme, in which case these points are plotted separately as small circles. Box width is proportional to sample size.




Figure 3.2-2. Boxplots comparing 2014 subsampling to 2014 full sample in the Black River based on CPUE for backpack electrofishing in fish/hour for subadult/adult Arctic grayling during three sampling events. The boxes represent the interquartile range (i.e., 1<sup>st</sup> to 3<sup>rd</sup> quartile of data), the black line in the box is the median. The whiskers extend to the full range of the data unless one or more data points are extreme, in which case these points are plotted separately as small circles. Box width is proportional to sample size.




Figure 3.2-3. Boxplots comparing 2014 subsampling to 2014 full sample in the Black River based on CPUE for backpack electrofishing in fish/hour for total Arctic grayling during three sampling events. The boxes represent the interquartile range (i.e., 1<sup>st</sup> to 3<sup>rd</sup> quartile of data), the black line in the box is the median. The whiskers extend to the full range of the data unless one or more data points are extreme, in which case these points are plotted separately as small circles. Box width is proportional to sample size.






Figure 3.2-4. Boxplots comparing 2014 subsampling to 2014 full sample in the Black River based on CPUE for backpack electrofishing in fish/hour for juvenile burbot during three sampling events. The boxes represent the interquartile range (i.e., 1<sup>st</sup> to 3<sup>rd</sup> quartile of data), the black line in the box is the median. The whiskers extend to the full range of the data unless one or more data points are extreme, in which case these points are plotted separately as small circles. Box width is proportional to sample size.




Figure 3.2-5. Boxplots comparing 2014 subsampling to 2014 full sample in the Black River based on CPUE for backpack electrofishing in fish/hour for subadult/adult burbot during three sampling events. The boxes represent the interquartile range (i.e., 1st to 3rd quartile of data), the black line in the box is the median. The whiskers extend to the full range of the data unless one or more data points are extreme, in which case these points are plotted separately as small circles. Box width is proportional to sample size.




Figure 3.2-6. Boxplots comparing 2014 subsampling to 2014 full sample in the Black River based on CPUE for backpack electrofishing in fish/hour for total burbot during three sampling events. The boxes represent the interquartile range (i.e., 1st to 3rd quartile of data), the black line in the box is the median. The whiskers extend to the full range of the data unless one or more data points are extreme, in which case these points are plotted separately as small circles. Box width is proportional to sample size.




Figure 3.2-7. Boxplots comparing 2014 subsampling to 2014 full sample in the Black River based on CPUE for backpack electrofishing in fish/hour for total sculpin during three sampling events. The boxes represent the interquartile range (i.e., 1st to 3rd quartile of data), the black line in the box is the median. The whiskers extend to the full range of the data unless one or more data points are extreme, in which case these points are plotted separately as small circles. Box width is proportional to sample size.