

07726

PRIVILEGED AND CONFIDENTIAL DOCUMENT
RELEASED FOR SETTLEMENT PURPOSES ONLY;
NO REPRODUCTION OR FURTHER
TRANSMITTAL AUTHORIZED

ALASKA POWER AUTHORITY
SUSITNA HYDROELECTRIC PROJECT

POSITION PAPER
FISHERIES ISSUE F-2.2

EXECUTIVE SUMMARY

Statement of Issue:

Significance of potential changes in pH on salmon and resident fish habitats and populations downstream of the dams.

Position

It is the position of the Alaska Power Authority that potential changes in pH due to inundation of acid bogs and tundra soils by the project reservoirs will not significantly affect the pH of the reservoir or the river downstream of the dams. Therefore, no mitigation measures are needed.

Present Knowledge

The Susitna River watershed above Devil Canyon presently drains large areas of acidic tundra soils and bog or wetland habitats. The mainstem river water, however, contains sufficient alkalinity-producing ions to buffer against biologically harmful pH changes which otherwise might occur due to acidic tributary influents. The wetland mapping recently completed by the USFWS indicates that approximately 2.7 percent of the combined impoundment areas is "bog-like" wetland. The alkalinity buffering system which exists now will continue to minimize pH changes during project construction and operation.

ALASKA POWER AUTHORITY
SUSITNA HYDROELECTRIC PROJECT
POSITION PAPER
FISHERIES ISSUE F-2.2

INTRODUCTION

Statement of Issue:

Significance of potential changes in pH on salmon and resident fish habitats and populations downstream of the dams.

Position

It is the position of the Alaska Power Authority that potential changes in pH due to inundation of acidic bogs and tundra soils by the project reservoirs will not significantly affect the pH of the reservoir or the river downstream of the dams. The amount of such acidic habitat is so small, about three percent of the area to be inundated, and the buffering capacity of the river so large, that potential changes in pH will be neutralized by the natural system.

DISCUSSION

The physical, chemical and biological characteristics of a water body (including its pH and alkalinity) will reflect the basic climatic and hydrologic regimes and biogeochemistry of its entire drainage basin, and not merely the small portion of the drainage which is under water (Hutchinson 1967, 1973, 1975, Wetzel 1975, Vollenweider and Kerekes 1980). The pH of upstream and downstream mainstem Susitna riverine habitats is presently regulated by the carbon dioxide-bicarbonate-carbonate and aluminum silicate dissolution buffering systems (Wetzel 1975, Stumm and Morgan 1970). These two buffering systems maintain pH values between 6.0 and 8.3 in most fresh water ecosystems of North America.

The drainage basin of the Susitna River upstream of the proposed Devil Canyon dam site encompasses approximately 5,810 square miles of unvegetated mountains and highland tundra. The watershed's bedrock, glacial till and glacial outwash materials contain alkalinity-producing carbonate and silicate minerals (APA 1983, R&M Consultants Inc. 1982). Overlying the watershed's lithic material is a substantial area of tundra consisting of saturated, peaty soils, which may be acidic in nature. Sphagnum bogs are frequent on the tundra and they commonly have a pH less than 4.5. Despite substantial inflow from tributary drainages with acidic soils and acidic bogs, the ionic composition of the mainstem river is presently sufficient to buffer tributary acidity and maintain low to moderate alkalinity values. Changes in mainstem pH values are seasonally variable but range between 6.0 and 8.1. The mean annual pH is greater than 7.0 in the mainstem water of the upper river watershed.

A wetlands mapping project has been completed by the USFWS in order to quantify the amount of different wetland types in both Watana and Devil Canyon impoundment zones. The estimated total of vegetated areas which are classifiable as wetlands equals 8,316 acres or 18.8 percent of the combined impoundment areas. The estimated total of all "bog-like" wetlands equals 1,182 acres or approximately 2.7 percent of the combined impoundment zone areas. The pH of the proposed reservoirs and downstream riverine habitats under with-project conditions will be regulated by the same chemical buffering system existing at present. Flooding of the small area of bog habitats is not anticipated to cause a biologically significant change of pH in riverine habitats downstream of the proposed project or in the reservoirs.

A number of references, including review articles, research reports and texts have been reviewed for any discussion of pH changes in reservoirs or their downstream habitats due to bog inundation (Supplementary References). No documentation of such a problem has been located in the open literature dealing with lake or larger reservoir limnology.

REFERENCES

Literature Cited

- Alaska Power Authority 1983. Final application for license for major project, Susitna Hydroelectric Project, before the Federal Energy Regulatory Commission, accepted by FERC July 27, 1983. Vol. 5-A and B, Chpts 1 and 2, Alaska Power Authority. Susitna Hydroelectric Project. 2 vols.
- Hutchinson, G.E. 1967. A Treatise on Limnology. Vol. II. Introduction to Lake Biology and the Limnoplankton. J. Wiley and Sons, New York, 1115 pp.
- Hutchinson, G.E. 1973. Eutrophication. Amer. Scien. 61(3): 269-279.
- Hutchinson, G.E. 1975. A Treatise on Limnology. Vol. I. Geography, Physics and Chemistry. J. Wiley and Sons, New York. (reprint of 1957 edition).
- R&M Consultants, Inc. 1982. Glacial Lake Studies. Interim Report. Prepared for Acres American, Inc., Anchorage, Alaska.
- Stumm, W., and J.J. Morgan. 1970. Aquatic Chemistry. Wiley Interscience, New York, 583 pp.
- Wetzel, R.G. 1975. Limnology. W.B. Saunders, Co., Philadelphia, 743 pp.
- Vollenweider, R.A., and J. Kerekes. 1980. The loading concept as basis for controlling eutrophication. Philosophy and preliminary results of the OECD Programme on eutrophication. Prog. Wat. Tech., Vol. 12, Norway, pp. 5-18. IAWPR/Pergamon Press Ltd.

Supplemental References^{1/}

- Allan, R.J. 1978. Natural controls of dissolved solids in Boundary Reservoir, Saskatchewan. Canadian Wat. Res. J. 3(3): 78-96.
- Baxter, R.H., and P. Glaude. 1980. Environmental effects of dams and impoundments in Canada: experiences and prospects. Can. Bull. Fish Aquat. Sci. 205, 34p
- Bodaly, R.A., D.M. Rosenberg, M.N. Goboury, R.E. Hecky, R.W. Newbury and K. Patalis. 1983. Ecological effects of hydroelectric development in northern Manitoba, Canada: The Churchill-Nelson River Diversion. in, "Effects of Pollutants at the Ecosystem Level," SCOPE, 22, John Wiley and Sons, Toronto. pp. 273-310.
- Campbell, P.G., B. Bobee, A. Caille, M.J. Demalsy, J.L. Sasseville and S.A. Visser. 1975. Pre-impoundment site preparation: a study of the effects of topsoil stripping on reservoir water quality. Verh. Internat. Verein. Limnol., 19, pp.768-1777.
- Denisova, A.I. 1977. Factors governing the biological productivity, regime, and content of biogenic and organic substances in the sequence of Dniepr reservoirs. Water Resources 4(6): 885-895.
- Duffer, W.R., and C.C Harlin. 1971. Changes in water quality resulting from impoundment. National Water Quality Control Research Program, R.S. Kerr Water Research Center USEPA, Ada, Oklahoma, 110 pp.
- Duthie, H.C. 1979. Limnology of subarctic Canadian lakes and some effects of impoundment. Arctic and Alpine Reserach, 11(2): 145-158.

^{1/} Reviewed but not cited in text.

Efford, I.E. 1975. Assessment of the impact of hydro-dams. J. Fish. Res. Board Can. 32: 196-209.

Geen, G.H. 1975. Ecological consequences of the proposed Moran Dam on the Fraser River. J. Fish. Res. Bd. Can. 32(1): 126-135.

Gill, D. 1971. Damming the Mackenzie: A theoretical assessment of the long-term influence of river impoundment on the ecology of the Mackenzie River delta. Proc. Peach-Athabasca Delta Symp. Jan. 14-15, 1971, Univ. of Alberta, Edmonton, pp. 204-234.

Gill, D. and A.D. Cooke. 1974. Controversies over hydroelectric developments in sub-arctic Canada. Polar Record 17(107): 109-127

Grimard, Y. and H.G. Jones 1982. Trophic upsurge in new reservoirs: a model for total phosphorus concentrations. Can J. Fish. Aquat. Sci. 39: 1473-1483.

Grimas, U. 1961. The bottom fauna of natural and impounded lakes in northern Sweden (Amkarvattnet and Blasjon). Rep. No. 42: 183-237, Institute of Freshwater Research, Drottningholm, Sweden.

Grimas, U. 1965. Inlet impoundments. An attempt to preserve littoral animals in regulated subarctic lakes. Rep. No. 46: 22-30. Institute for Freshwater Research, Drottningholm, Sweden.

Grimas, U. and N.A. Nilsson. 1965. On the food chain in some north Swedish river reservoirs. Rep. No. 46: 31-48. Institute of Freshwater Research, Drottningholm, Sweden.

- Gunnison, D., R.L. Chen, and J.M. Brannon. 1983. Relationships of materials in flooded soils and sediments to the water quality of reservoirs - I. Oxygen consumption rates. *Water Res.* 17(11): 1609-1617.
- Hanan, H.H. 1979. Chemical modifications in reservoir-regulated streams. In: "The Ecology of Regulated Streams," Ward, J.V. and J.A. Stanford, eds., Plenum Press, New York, pp. 75-94.
- Harvey, H.H. 1976. Aquatic environmental quality: problems and proposals. *J. Fish. Res. Board Can.* 33: 2634-2670.
- Hecky, R.E., and H.A. Ayles, 1974. Summary of fisheries - limnology investigations on Southern Indian Lake. Report for the Lake Winnipeg, Churchill and Nelson Rivers Study Board, Fisheries and Marine Service, Environment Canada, Winnipeg, Manitoba. 26 pp.
- Jackson, K.S., I.R. Jonasson and G.B. Skippen. 1978. The nature of metals-sediments-water interactions in freshwater bodies, with emphasis on the role of organic matter. *Earth-Science Reviews*: pp. 97-146.
- Lehmkuhl, D.M. 1979. Environmental disturbance and life histories: principles and examples. *J. Fish. Res. Board Can.* 36: 329-334.
- Lindstrom, T. 1973. Life in a lake reservoir: Fewer options, decreased production. *AMBIO*, pp. 145-153.
- Marcotte, P.H. and A. Soucy. 1979. Le deboisement dans les reservoirs. *Geos.*, Spring, pp. 15-17.
- Nilsson, Nils-Arvid. 1964. Effects of impoundment on the feeding habits of brown trout and char in Lake Ransaren (Swedish Lappland). *Verh. Internat. Verein. Limnol.*, XV: 444-452.

- Ostrofsky, M.L., and H.C. Duthie. 1975. Primary productivity, phytoplankton and limiting nutrient factors in Labrador Lakes. Int. Rev. ges. Hydrobiol. 60(2): 145-158.
- Ostrofsky, M.L. and H.C. Duthie. 1978. V. Manmade Lakes: An approach to modelling productivity in reservoirs. Verh. Internat. Verein. Limnol. 20: 1562-1567.
- Ostrofsky, M.L., and H.C. Duthie. 1980. Trophic upsurge and the relationships between phytoplankton biomass and productivity in Smallwood Reservoir, Canada. Can J. Bot. 58: 1174-1180.
- Peterson, L.A., G. Nichols, and R&M Consultants, Inc. 1982. Water quality effects resulting from impoundment of the Susitna River. Prepared for Acres American, Inc. and the Alaska Power Authority. 43 pp.
- Rodhe, W. 1964. IV. Production. 2. Post-primary. Effects of impoundments on water chemistry and plankton in Lake Ransaren (Swedish Lapland). Verh. Internat. Verein. Limnol., XV: 437-443.
- Runnstrom, S. 1964. Effects of impoundment on the growth of Salmo trutta and Salvelinus alpinus in Lake Ransaren (Swedish Lapland). Verh. Internat. Verein. Limnol. XV: 453-461.
- Smith, D.W., and S.R. Justice. 1975. Effects of reservoir clearing on water quality in the Arctic and Subarctic. Rep. No. IWR-58, Institute of Water Resources, Univ. of AK., 13 p.
- Soucy, A. 1978. James Bay Hydroelectric Development: Evolution of Environmental Considerations. Can. Wat. Res. J.3(4): 44-88.
- Spence, J.A., and H.B.N. Hynes. 1971. Differences in fish populations upstream and downstream of a mainstream impoundment. J. Fish. Res. Board Can. 28: 45-46.

Therien, N., G. Spiller and B. Coupal. 1982. Simulation de la decomposition de la matiere vegetale et des sols inondes dans les reservoirs de la region de la Baie de James. Canadian Wat. Res. J., 7(1): 375-396.

Welch, P.S. 1952. Limnology. McGraw-Hill Book Co., New York. 538 pp.