

Fisheries and Habitat Investigations  
of the Susitna River--A Preliminary Study of  
Potential Impacts of the Devils  
Canyon and Watana Hydroelectric Projects

by  
James C. Riis  
and  
Nancy V. Friese

Alaska Department of Fish and Game  
Divisions of Sport and Commercial Fish

March 1978

# TABLE OF CONTENTS

|  | <u>Page</u> |
|--|-------------|
| List of Figures . . . . .                                  | 4           |
| List of Tables . . . . .                                   | 5           |
| List of Appendix Tables . . . . .                          | 6           |
| Summary . . . . .  | 7           |
| Background . . . . .                                       | 9           |
| Description of Area . . . . .                              | 10          |
| Procedures . . . . .                                       | 11          |
| Fisheries . . . . .  | 13          |
| Adults . . . . .   | 13          |
| Juvenile salmon migration . . . . .                        | 13          |
| Juvenile studies . . . . .                                 | 14          |
| Water Quantity . . . . .                                   | 16          |
| Water Quality . . . . .                                    | 17          |
| Findings and Discussion . . . . .                          | 17          |
| Fisheries . . . . .  | 17          |
| Adults . . . . .   | 17          |
| Juvenile salmon migration . . . . .                        | 23          |
| Juvenile studies . . . . .                                 | 28          |
| Aquatic insects and juvenile salmon gut contents . . . . . | 30          |
| Impoundment area fisheries investigations . . . . .        | 37          |
| Water Quantity . . . . .                                   | 40          |
| Water Quality . . . . .                                    | 43          |
| Conclusion . . . . .                                       | 47          |
| Recommendations . . . . .                                  | 48          |
| Acknowledgements . . . . .                                 | 50          |
| Literature Cited . . . . .                                 | 51          |

TABLE OF CONTENTS (cont.)

|                        | <u>Page</u> |
|------------------------|-------------|
| Appendix I . . . . .   | 53          |
| Appendix II . . . . .  | 73          |
| Appendix III . . . . . | 96          |

## LIST OF FIGURES

| <u>Figure</u> |   | <u>Page</u> |
|---------------|---|-------------|
| 1             | The Susitna River drainage, Devils Canyon Project, 1977 . . .   | 12          |
| 2             | Upper Susitna River study area, Devils Canyon Project, 1977 .   | 15          |
| 3             | Locations of adult salmon tag recoveries occurring downstream<br>of the Susitna River fishwheel sites, Devils Canyon Project,<br>1977 (RS-sockeye salmon; PS-pink salmon; CS-chum salmon;<br>SS-coho salmon; KS-chinook salmon) . . . . . | 22          |
| 4             | Susitna River discharge at Gold Creek, Devils Canyon Project,<br>1977 . . . . .   | 41          |

# LIST OF TABLES

| <u>Table</u> |   | <u>Page</u> |
|--------------|---|-------------|
| 1            | Relative magnitude of pink, chum, and sockeye salmon moving past the fishwheel sites as determined by Peterson population estimates, Devils Canyon Project, 1974, 1975, and 1977. <sup>1/</sup> . . . | 18          |
| 2            | Peak chum, pink, and sockeye salmon ground escapement survey counts within the upper Susitna River, Devils Canyon Project, 1977 . . . . .   | 20          |
| 3            | Peak chinook salmon counts within the Susitna River drainage, 1977 . . . . .  | 21          |
| 4            | Rabideux Creek salmon fry trapping, Devils Canyon Project, 1977 . . . . .   | 24          |
| 5            | Montana Creek salmon fry trapping, Devils Canyon Project, 1977  | 26          |
| 6            | Willow Creek chinook salmon fry trapping, Devils Canyon Project, 1977 . . . . .   | 27          |
| 7            | Mean percent composition of gut contents per fish of chinook, sockeye, and coho salmon juveniles in sloughs and clearwater tributaries of the Susitna River, Devils Canyon Project, 1977.             | 32          |
| 8            | Limnological data from selected tributaries to the Susitna River, Devils Canyon Project, 1977 . . . . .   | 38          |
| 9            | Susitna River impoundment area lake surveys, Devils Canyon Project, 1977 . . . . .  | 39          |
| 10           | Water flows of Montana, Rabideux, and Willow creeks from May through November, Devils Canyon Project, 1977. <sup>1/</sup> . . . . .   | 44          |
| 11           | Thermograph set in Susitna River downstream of Parks Highway bridge, daily maximum and minimum water temperature, Devils Canyon Project, 1978 . . . . .   | 45          |

# LIST OF APPENDIX TABLES

## Appendix I

| <u>Table</u> |   | <u>Page</u> |
|--------------|---|-------------|
| 1            | Percent age composition of chinook, sockeye, coho, and chum salmon escapement samples, Devils Canyon Project, 1974, 1975, and 1977 . . . . .  | 54          |
| 2            | Age, length, and sex characteristics of chum, chinook, sockeye, and coho salmon escapement samples, Devils Canyon Project, 1974, 1975, and 1977 . . . . .   | 55          |
| 3            | Analyses of age, length, weight, and condition factors of juvenile sockeye salmon samples from Susitna River sloughs and clearwater tributaries, Devils Canyon Project, 1977 . . .                        | 56-57       |
| 4            | Analyses of age, length, weight, and condition factors of juvenile coho salmon samples from Susitna River sloughs and clearwater tributaries, Devils Canyon Project, 1977 . . . . .                       | 58-59       |
| 5            | Analyses of age, length, weight, and condition factors of juvenile chinook salmon samples from Susitna River sloughs and clearwater tributaries, Devils Canyon Project, 1977 . . .                        | 60-64       |
| 6            | Stomach content analysis of juvenile chinook, coho, and sockeye salmon collected in sloughs and clearwater tributaries of the Susitna River during summer and fall, Devils Canyon Project, 1977 . . . . . | 65-72       |

## Appendix II

|   |  |       |
|---|--|-------|
| 1 | Susitna River discharge at Gold Creek (USGS provisional data) 1977 . . . . .   | 74    |
| 2 | Water quality data and juvenile salmon surveys in sloughs and clearwater tributaries of the Susitna River between the Chulitna River and Portage Creek, Devils Canyon Project, 1977. | 75-81 |
| 3 | Thermograph set in Rabideux Creek, upper sub-area; daily maximum and minimum water temperature, Devils Canyon Project, 1977 . . . . .  | 82    |
| 4 | Thermograph set in Montana Creek, upper sub-area; daily maximum and minimum water temperature, Devils Canyon Project, 1977 . . . . .   | 83    |
| 5 | Water chemistry data, Rabideux Creek, Devils Canyon Project, 1977 . . . . .  | 84    |
| 6 | Water chemistry data, Montana Creek, Devils Canyon Project, 1977 . . . . .   | 85    |
| 7 | United States Department of Interior, Geological Survey . . .  | 86-95 |

## SUMMARY

Biological and water quality and quantity investigations were conducted from May 1, 1977 through March 7, 1978 to obtain baseline data on indigenous fish populations and the existing aquatic habitat of the Susitna River drainage. These investigations conclude a four year series of environmental baseline inventories. They were designed to generate sufficient biological information to enable the Alaska Department of Fish and Game (ADF&G) to prepare a comprehensive biological study plan in the event a final environmental impact study is initiated to determine the feasibility of constructing the proposed Watana and Devils Canyon hydroelectric dams on the Susitna River.

The relative abundance, distribution and migrational timing of adult salmon (Oncorhynchus sp.) were determined within the Susitna River drainage through tag and recovery programs during 1977. The salmon escapement from June 29 through August 14 was estimated to be approximately 237,000 sockeye (O. nerka), 50,000 coho (O. kisutch), and 105,000 chum salmon (O. keta) (Friese, in prep.). An escapement estimate in excess of 100,000 fish was determined for chinook salmon (O. tshawytscha) through aerial surveys (Kubik, 1977; Watsjold, 1977). Population estimates of pink salmon utilizing the drainage in the area of the Susitna and Chulitna river confluence were determined as a part of this study.

Documentation of the outmigration of salmon fry from tributary rearing areas into the mainstem Susitna River was accomplished by intensive investigation of two clearwater tributaries. The objective of these studies was to determine utilization of the mainstem river for rearing during winter months. A total of 25,176 chinook salmon fry were marked

in Montana Creek between July 19 and August 4. A gradual downstream movement of fry was noted from the latter part of August to February. A drastic reduction in population density was found in February and was attributed to low flows which prevailed at the time. Chinook fry were documented overwintering in the Susitna River. No distinct movement of fry was observed in Rabideux Creek.

The relative abundance, distribution, age, length, and weight characteristics, and feeding habits of juvenile salmonids were monitored in sloughs and tributaries of the Susitna River from Portage Creek downstream to the Chulitna River confluence from July 1 through October 5, 1977. The predominant rearing species were chinook and coho salmon. Water quality and quantity determinations were made in conjunction with all juvenile salmon surveys.

The Susitna River was floated from its intersection with the Denali Highway to Devils Canyon during the first two weeks of July to inventory fish species present and survey the aquatic habitat in the areas to be inundated. Arctic grayling (Thymallus arcticus) were abundant in all of the clearwater tributaries within the proposed impoundment area. The headwaters of these tributaries and upland lakes were also surveyed by separate crews. It is apparent that the Watana reservoir, which is projected to have substantial seasonal fluctuations, will alter the fisheries habitat.

Measurements of hydrological and limnological parameters associated with the Susitna River and selected tributaries and sloughs were obtained between the Denali Highway and Montana Creek. A cooperative agreement between the United States Geological Survey (USGS) and the ADF&G was initiated to determine discharge, sediment loads, and standard water



quality analysis of the mainstem Susitna River. This data, along with the water quality and quantity data collected in conjunction with the fisheries studies, will be extremely valuable for future comparisons.

Long term ecological changes to the drainage may be significant due to dam construction. The level and flow patterns of the Susitna River will be altered and will affect the fisheries resources. Extensive research is necessary both upstream and downstream of the proposed dams to adequately assess the potential effects of these impacts on fisheries resources.

The effects of impoundments and construction activities which alter natural flow regimes, water chemistry, mass transport of materials, and quantity of wetted habitat areas are of primary concern. These changes may disrupt the trophic structure and habitat composition and reduce or eliminate terrestrial and aquatic populations. These populations and vegetation in and around the free-flowing rivers have evolved to their current levels due to natural flow variations. Some species may be present only because this particular hydrologic regime exists.

#### BACKGROUND

Background knowledge of the Susitna River basin is limited. The proposed hydroelectric development necessitates gaining a thorough knowledge of its natural characteristics and populations prior to final dam design approval and construction authorization to enable protection of the aquatic and terrestrial populations from unnecessary losses.

The Susitna River basin has long been recognized as an area of high recreational and aesthetic appeal. It is also important habitat to a wide variety of fish species, both resident and anadromous. Five species of Pacific salmon (chinook, coho, chum, pink, and sockeye) utilize the

Susitna River drainage for spawning and rearing. The majority of the chinook, coho, chum, and pink salmon production in the Cook Inlet area occurs within this drainage. Grayling, rainbow trout (Salmo gairdneri), Dolly Varden (Salvelinus malma), burbot (Lota lota), lake trout (Salvelinus namaycush), whitefish (Coregonus sp.), and sculpins (Cottus sp.) are some of the more common and important resident fish species.

Baseline environmental fisheries studies have been conducted by ADF&G intermittently since 1974. The projects were financed with federal funding averaging \$29,000 per year for the first three years. An allocation of \$100,000 was received for this study. The National Marine Fisheries Service (NMFS) and U.S. Fish and Wildlife Service (USFWS) contracted ADF&G to conduct a one-year assessment of salmon populations utilizing the Susitna River in the vicinity of the proposed Devils Canyon dam site during 1974. The objectives of these studies were to determine the adult salmon distribution, relative abundance and migrational timing and to determine juvenile rearing areas (Barrett, 1974). Additional funding was received in 1975, 1976, and 1977 from USFWS to continue and expand these studies and to monitor the physical and chemical parameters associated with the mainstem Susitna (USFWS, 1976 and Riis, 1977). Additional baseline studies will not be initiated during 1978 due to lack of funding.

The proposed hydroelectric project is discussed in Barrett (1974), Friese (1975), USFWS (1976), and Riis (1977). The purpose of this data report is to present the findings of the studies conducted from May 1977 through March 1978 and to make recommendations for future investigations and a final environmental impact statement.

#### DESCRIPTION OF AREA

The Susitna River is approximately 275 miles long from its source in the Alaska Mountain Range to its point of discharge into Cook Inlet

(Figure 1). The major tributaries of the Susitna originate in glaciers and carry a heavy load of glacial silt during ice free months. There are also many smaller tributaries which are perennially silt free. The study area included the majority of the Susitna River between the Denali Highway and Cook Inlet. The entire drainage from Devils Canyon downstream was monitored for chinook salmon escapement. Studies of other anadromous species were more restricted to the mainstem Susitna and adjacent areas between Devils Canyon and Susitna Station.

Two clearwater tributaries, Rabideux and Montana creeks, were selected for intensive juvenile salmon studies. These streams are located downstream of the proposed dam site near the Parks Highway Bridge. A total of 26 clearwater sloughs and eight tributaries were surveyed between the Chulitna River confluence and Devils Canyon area. These areas are described in USFWS (1976). Surveys of the Talkeetna River were conducted, but results are not included within this report. Resident fish were inventoried in the impoundment area upstream of Devils Canyon.

Water quality and quantity sampling stations were monitored in the Susitna River and tributaries. Twenty-six of these sites were clearwater sloughs adjunct to the Susitna River. Three sites were in the mainstem Susitna River and the ten remaining locations were clearwater creeks and rivers flowing into the Susitna River. Site selection was based on proximity to the Devils Canyon dam area and previous Susitna River studies documenting fish usage (Barrett, 1974; USFWS, 1976).

#### PROCEDURES

A field camp was established at Gold Creek for studies downstream of Devils Canyon due to its central location to the sample sites and the




Figure 1. The Susitna River drainage, Devils Canyon Project, 1977.

logistical advantages offered by the Alaska Railroad. Travel on the Susitna River to the sites was accomplished by riverboats equipped with jet outboard motors. Access to sloughs and tributaries downstream from Gold Creek was accomplished with a Zodiac raft. A field camp was also established along the Susitna River five miles upstream from Talkeetna to install and operate fishwheels. Fishwheels were deployed commencing July 5 and were operated through August 27. Methods of operation are discussed by Friese (1975). A field station was located in the vicinity of Talkeetna to conduct Rabideux and Montana creek studies. Avon rubber rafts supported with helicopter and fixed wing aircraft were used for the impoundment area studies.

## FISHERIES

### Adults

Adult salmon escapement was generally determined by tag and recovery population estimates utilizing fishwheels and ground escapement surveys. Methods are discussed in Friese (1975). The Peterson population estimate used to determine salmon abundance is presented in Table 1. Chinook salmon counts were conducted with a Bell-47 helicopter and fixed wing aircraft. Variable mesh gillnets were used to determine species composition in the impoundment area lakes. Electroshockers and angling were also employed to collect adult fish for this study. Sloughs and tributaries in the upper study area were surveyed on the ground according to methods described in Friese (1975).

### Juvenile salmon migration

Intensive fry trapping was undertaken in Rabideux Creek on June 16. The creek was sectioned into three study areas: upper, middle, and lower. Coho salmon yearlings were anesthetized with MS-222 and fin

clipped from June 16 through August 31. The following fin clip codes were used: upper caudal lobe for upper sub-area, one-half dorsal for mid sub-area, and lower caudal lobe for lower sub-area. After marking, the salmon were allowed to recover and were released at the location of capture. Recovery of these marked coho salmon was continued until mid-November when extreme cold weather and icing conditions prevented further intensive work.

Montana Creek was also sectioned into three study sub-areas: upper, middle, and lower. The upper area was approximately eight stream miles above its mouth, the middle about three stream miles, and the lower was from the Parks Highway downstream to its junction with the Susitna River. The upper and middle sections were seined from July 19 through August 4. All chinook salmon fry captured were marked with an upper caudal fin clip for the upper area and a lower caudal fin clip for the middle area. Minnow traps baited with salmon roe were utilized from the latter part of August until the end of February to monitor fry movements and population densities throughout the system.

#### Juvenile studies

Twenty-eight clearwater sloughs and nine tributary streams have previously been identified as observed or potential rearing sites for juvenile salmon in the upper Susitna River between Talkeetna and Devils Canyon (Figure 2) (Barrett, 1974; Friese, 1975). Juvenile salmon were collected from these locations during two different sampling periods during this study. Each slough and tributary were also surveyed biweekly for relative abundance of rearing fish and water quality data. Methods are discussed in Friese (1975). Fry samples for analysis of physical characteristics and feeding habits were collected with dip net, minnow traps, or seine and preserved in a 10 percent formalin solution (Brown, 1971).




Figure 2. Upper Susitna River study area, Devils Canyon Project, 1977.

Summer samples were netted by minnow seine between July 11 and August 5. Juvenile salmon were collected by a combination of minnow seine and minnow traps from September 20 to 24. Fork lengths and scale smears were taken in the field for each individual fish. Specimens, together with incidental catches of other resident fish species, were preserved in 10 percent formalin. Species identification, verified by pyloric caecae counts, and weight determinations were made in the Anchorage laboratory. The gut was dissected from each fish and contents from both hind- and foregut removed. All gut contents from one sampling location were pooled by species for each sampling day to facilitate investigation. Individual stomachs were not examined separately. Insects were identified to order and larval and pupal forms of Diptera to family. Other organisms present were identified to the most convenient taxon, usually order. The major keys used were Pennack (1953), Usinger (1968), Ward and Whipple (1959), and Jacques (1947). Volume percentages were estimated according to four gross categories: Crustacea, immature Insecta, adult Insecta, and other organisms. These estimates reflect the interpretations of the investigator, but it is felt that they gave a close approximation of actual volumes.

#### WATER QUANTITY

Discharge data were collected by ADF&G personnel at many of the slough and tributary sites. Flows were measured with Price AA Gurley current meters. Leupold stage gauges were installed in the sloughs and permanent bench marks were established on the river banks adjacent to the gauges for future location reference (Riis, 1977).

Mainstem Susitna River flow was continually monitored by USGS at their Gold Creek site and three times during the summer at Portage Creek and at the Parks Highway Bridge.



Water flows in Rabideux Creek were measured by recording the height of the water passing through culverts at the Parks Highway, approximately one-half mile above its confluence with the Susitna River. Recordings were converted into cubic feet per second. The River Forecast Center of the National Weather Service monitored water stage and computed flow in Montana and Willow Creeks.

#### WATER QUALITY

Dissolved oxygen, temperature, pH, and specific conductance were measured biweekly and on a random basis in clearwater sloughs and tributaries with a Yellow Springs Instrument Model 57 oxygen and temperature meter, Cole Parmer Digi sense pH meter, and Labline Lectro mho meter, respectively. Alkalinity and hardness were determined with a Hach chemical kit (model DR-EL/2 and model AL36B) using methods outlined by the manufacturers.

Temperature data was continually recorded with Ryan thermographs, Model D-30, at one site on the Susitna River and at three sites in both Rabideux and Montana creeks. Analysis of water samples from the mainstem Susitna were analyzed by the USGS laboratory.

Benthic invertebrates were collected with artificial substrates (McCoy, 1974) and Surber samplers for future analysis.

### FINDINGS AND DISCUSSION

#### FISHERIES

##### Adults

Adult salmon abundance above the Chulitna River confluence was determined by tag and subsequent recovery programs during 1974, 1975, and 1977 (Table 1). The relative magnitude of pink salmon moving past

Table 1 . Relative magnitude of pink, chum, and sockeye salmon moving past the fishwheel sites as determined by Peterson population estimates, Devils Canyon Project, 1974, 1975, and 1977.<sup>1/</sup>

|                          | Species       |               |             |
|--------------------------|---------------|---------------|-------------|
|                          | Pink          | Chum          | Sockeye     |
| <u>1974</u>              |               |               |             |
| M                        | 160           | 568           | 39          |
| R                        | 23            | 74            | 13          |
| C                        | 755           | 3,164         | 336         |
| N                        | 5,040         | 23,970        | 939         |
| Confidence Interval      | 3,836-8,359   | 20,081-30,746 | 709-1,764   |
| <u>1975</u>              |               |               |             |
| M                        | 943           | 674           | 370         |
| R                        | 46            | 8             | 22          |
| C                        | 291           | 139           | 103         |
| N                        | 6,129         | 10,549        | 1,760       |
| Confidence Interval      | 4,977-11,895  | 7,122-35,293  | 1,355-2,865 |
| <u>1977<sup>2/</sup></u> |               |               |             |
| M                        | 429           | 46            | 31          |
| R                        | 64            | 3             | 1           |
| C                        | 6,644         | 2,332         | 661         |
| N                        | 43,857        | ---           | ---         |
| Confidence Interval      | 36,375-57,439 | ---           | ---         |

1/ Calculated by the following formulas:

$$N = \frac{M (C+1)}{R+1}$$

$$95\% \text{ confidence interval around } N \quad R/C = R/C \pm t \frac{\frac{R}{C} (1 - \frac{R}{C})}{C} \frac{(N-C)}{N}$$

where:

N = Population size during time of marking

M = Number of fish marked

C = Total of fish observed for presence of mark during sample census.

R = Total number of marked (recaptured) fish found during sample census.

2/ Population estimates were not determined for chum and sockeye salmon since number of tag recoveries were too low to place confidence limits on estimates.

the fishwheel sites above Talkeetna during 1977 was approximately 44,000 fish. Tag recoveries of other salmon species were too low to determine abundance. Abundance of all salmon species within sloughs and tributaries, with the exception of chinook salmon, was determined by ground escapement surveys. Peak survey counts by species from Portage Creek downstream to the Chulitna River confluence was 1,330 chum, 3,429 pink, and 301 sockeye salmon (Table 2). These estimates are considered minimum escapements, since counts were only conducted within index areas (USFWS, 1976). Migrational timing of coho salmon was too late to determine peak abundance.

The chinook salmon escapement within the drainage was about 100,000 fish (Table 3). The 1977 escapement appears to have a high reproduction potential (Kubik, 1977 and Watsjold, 1977). Historic escapement and harvest data indicate a minimum escapement level of at least 60,000 chinook salmon would be required yearly to restore stocks to historic levels.

Numerous tag recoveries downstream of the tagging project were obtained from the sport fish harvest during 1977 (Figure 3). This "drop-out" phenomenon was also observed during 1974 and 1975. The total magnitude of tagged fish moving downstream was not determined since reporting of tag recoveries was on a voluntary basis. This should, however, be thoroughly evaluated during future studies. If the Chulitna, Susitna, and Talkeetna river confluence area serves as a milling area for fish destined to spawning areas downstream, the project impact area would be greatly expanded and numbers of fish affected increased significantly.

Age, length, and sex composition characteristics were determined from fishwheel catch samples for all species except pink salmon. Results are presented in Appendix I, Tables 1 and 2. Data is comparable with

Table 2. Peak chum, pink and sockeye salmon ground escapement survey counts within the upper Susitna River, Devils Canyon Project, 1977.

| CHUM SALMON          |         |            |            |            |
|----------------------|---------|------------|------------|------------|
| Area                 | Date    | Density    |            | Total      |
|                      |         | Live       | Dead       |            |
| Slough 8A            | 9/22/77 | 34         | 17         | 51         |
| Slough 9             | 8/19/77 | 34         | 2          | 36         |
| Slough 10            | 9/9/77  | 0          | 2          | 2          |
| Slough 11            | 9/22/77 | 79         | 37         | 116        |
| Slough 16            | 8/28/77 | 0          | 4          | 4          |
| Slough 20            | 8/16/77 | 27         | 1          | 28         |
| Slough 21            | 9/20/77 | 187        | 117        | 304        |
| Lane Creek           | 8/19/77 | 0          | 2          | 2          |
| Fourth of July Creek | 8/11/77 | 11         | 0          | 11         |
| Indian River         | 8/18/77 | <u>514</u> | <u>262</u> | <u>776</u> |
| TOTAL                |         | 886        | 444        | 1,330      |

| PINK SALMON          |         |              |            |              |
|----------------------|---------|--------------|------------|--------------|
| Area                 | Date    | Density      |            | Total        |
|                      |         | Live         | Dead       |              |
| Slough 16            | 8/28/77 | 0            | 13         | 13           |
| Lane Creek           | 8/11/77 | 1,190        | 3          | 1,193        |
| Fourth of July Creek | 8/11/77 | 611          | 1          | 612          |
| Indian River         | 8/18/77 | <u>1,031</u> | <u>580</u> | <u>1,611</u> |
| TOTAL                |         | 2,832        | 597        | 3,429        |

| SCKEYE SALMON |         |          |          |          |
|---------------|---------|----------|----------|----------|
| Area          | Date    | Density  |          | Total    |
|               |         | Live     | Dead     |          |
| Slough 8A     | 9/9/77  | 64       | 6        | 70       |
| Slough 8B     | 9/9/77  | 2        | 0        | 2        |
| Slough 9      | 9/9/77  | 6        | 0        | 6        |
| Slough 11     | 9/8/77  | 181      | 33       | 214      |
| Slough 19     | 9/7/77  | 7        | 1        | 8        |
| Indian River  | 8/18/77 | <u>1</u> | <u>0</u> | <u>1</u> |
| TOTAL         |         | 261      | 40       | 301      |

Table 3. Peak chinook salmon counts within the Susitna River drainage, 1977.

| Streams (West Side)    | Count  | Streams (East Side)          | Count   |
|------------------------|--------|------------------------------|---------|
| Deshka River           | 39,642 | Willow Creek                 | 1,065   |
| Alexander Creek        | 13,385 | Montana Creek                | 1,443   |
| Talachulitna River     | 1,856  | Moose Creek                  | 153     |
| Lake Creek             | 7,391  | Prairie Creek                | 5,790   |
| Martin Creek           | 1,060  | Chunilna Creek               | 769     |
| Cache Creek            | 100    | Kashwitna River (North Fork) | 336     |
| Bear Creek             | 298    | Little Willow Creek          | 598     |
| Red Creek              | 1,511  | Sheep Creek                  | 630     |
| Peters Creek           | 3,042  | Indian River                 | 393     |
| Donkey Creek           | 159    | Portage Creek                | 374     |
| Fish Creek (Quits)     | 131    | Chulitna River (East Fork)   | 168     |
| Fish Creek (Kroto S.)  | 132    | Chulitna River (Middle Fork) | 1,782   |
| Unnamed-Kichatna River | 120    | Chulitna River (Mainstem)    | 229     |
| Clearwater Creek       | 47     | Goose Creek                  | 133     |
| Quartz Creek           | 8      | Honolulu Creek               | 36      |
| Canyon Creek           | 135    | Byers Creek                  | 69      |
| Dickason Creek         | 4      | Troublesome Creek            | 95      |
| Unnamed-Hayes River    | 2      | Bunco Creek                  | 136     |
| Rabideux Creek         | 99     |                              |         |
| Total Count            | 69,122 | Total Count                  | 14,199  |
| Estimated Total Count  | 93,411 | Estimated Total Count        | 17,028  |
|                        |        | Total Count                  | 83,321  |
|                        |        | Estimated Total Count        | 109,439 |




Figure 3. Locations of adult salmon tag recoveries occurring downstream of the Susitna River fishwheel sites, Devils Canyon Project, 1977 (RS-sockeye salmon; PS-pink salmon; CS-chum salmon; SS-coho salmon; KS-chinook salmon).

escapement samples obtained from other areas within the drainage (Frieese, in prep.). Carcass data collected in the Deshka River and Alexander Creek revealed a high percentage of five- and six-year-old chinook salmon females (Kubik, 1977).

#### Juvenile salmon migration

Intensive studies of juvenile chinook and coho salmon were conducted in Rabideux and Montana creeks to define the life histories of these species as related to the variable conditions of the drainage. The authors believe that the overwintering period during the first year of life is probably the most critical time for survival of these two species.

Rabideux and Montana creeks were selected for this study due to: accessibility, their opposite physical characteristics, and the difference in the ratio of rearing species. Willow Creek and Indian River were also sampled periodically for comparative purposes.

Rabideux Creek was selected to obtain representative data on coho salmon fry densities and yearling movements. A total of 1,041 yearling cohos were marked. Of these, 274 were marked in the upper sub-area, 753 in the middle sub-area, and 14 in the lower sub-area. Catches of rearing coho and chinook salmon captures and recaptures are presented in Table 4. A total of 159 marked fish were recaptured in the original area of marking and 32 in dispersed areas. An increase in catch per hour of coho salmon fry occurred following August 1 because increased growth made them more susceptible to capture in the 1/4" mesh minnow traps. Fourteen marked yearlings moved downstream, five upstream, and thirteen migrated to small lateral tributaries. No distinct pattern was exhibited, which could be attributed to the fact that environmental conditions are more stable throughout the year in this tributary during this particular year.

Table 4. Rabideaux Creek salmon fry trapping, Devils Canyon Project, 1977.

| Date                | Coho Smolt | Coho Yearling     | Marked Coho Yearling  | Coho Fry | Chinook Fry | No. Traps | Trap Hours | Coho Yearling Per Hour | Coho Fry Per Hour | Chinook Fry Per Hour |
|---------------------|------------|-------------------|-----------------------|----------|-------------|-----------|------------|------------------------|-------------------|----------------------|
| UPPER SECTION       |            |                   |                       |          |             |           |            |                        |                   |                      |
| 6/16-6/30           | 33         | 218               | 5u. <sup>1/</sup>     | 60       | 728         | 67        | 1608       | .14                    | .04               | .45                  |
| 7/10-7/15           | 0          | 56                | 12u.                  | 136      | 650         | 70        | 1680       | .04                    | .08               | .39                  |
| 9/16-9/30           | 0          | 36                | 1u.                   | 27       | 48          | 14        | 336        | .11                    | .08               | .14                  |
| 11/1-11/15          | 0          | 274               | 7m.2u.                | 117      | 0           | 35        | 805        | .35                    | .15               | 0                    |
| MIDDLE SECTION      |            |                   |                       |          |             |           |            |                        |                   |                      |
| 6/16-6/30           | 80         | 361               | 12m.                  | 109      | 1120        | 200       | 4800       | .08                    | .02               | .23                  |
| 7/1-7/15            | 0          | 229               | 26m.                  | 243      | 1284        | 135       | 3240       | .08                    | .08               | .40                  |
| 8/1-8/15            | 0          | 38                | 17m.1u. <sup>1/</sup> | 602      | 249         | 104       | 2496       | .02                    | .24               | .04                  |
| 8/16-8/31           | 0          | 125 <sup>2/</sup> | 64m.2u.               | 3764     | 1479        | 207       | 4968       | .04                    | .76               | .30                  |
| 10/1-10/15          | 0          | 116               | 7m.                   | 960      | 1253        | 59        | 1416       | .09                    | .68               | .88                  |
| 10/16-10/31         | 0          | 58                | 6m.                   | 510      | 133         | 5         | 105        | .61                    | 4.86              | 1.27                 |
| 11/1-11/15          | 0          | 57                | 4m.                   | 1952     | 399         | 23        | 522        | .12                    | 3.74              | .76                  |
| LOWER SECTION       |            |                   |                       |          |             |           |            |                        |                   |                      |
| 6/16-6/30           | 0          | 2                 | 0                     | 2        | 45          | 29        | 696        | tr.                    | tr.               | .06                  |
| 7/1-7/15            | 0          | 2                 | 0                     | 2        | 15          | 29        | 696        | tr.                    | tr.               | .02                  |
| 8/1-8/15            | 0          | 6                 | 1 1.                  | 95       | 50          | 50        | 1200       | tr.                    | .08               | .04                  |
| 9/1-9/15            | 0          | 31                | 1 1.                  | 180      | 797         | 20        | 480        | .07                    | .38               | 1.66                 |
| 9/16-9/30           | 0          | 44                | 2u.1 1.               | 221      | 468         | 47        | 1128       | .04                    | .20               | .41                  |
| 10/1-10/15          | 0          | 125               | 1u.                   | 668      | 3832        | 207       | 4944       | .03                    | .14               | .78                  |
| 10/16-10/31         | 0          | 98                | 3m.                   | 198      | 821         | 44        | 964        | .10                    | .21               | .85                  |
| 11/1-11/15          | 0          | 142               | 4m.                   | 621      | 1449        | 93        | 2078       | .07                    | .30               | .70                  |
| LATERAL TRIBUTARIES |            |                   |                       |          |             |           |            |                        |                   |                      |
| 10/1-10/15          | 0          | 270               | 2u.4m.                | 393      | 76          | 31        | 744        | .37                    | .53               | .10                  |
| 10/16-10/31         | 0          | 231               | 4m.                   | 794      | 117         | 55        | 1212       | .19                    | .66               | .10                  |
| 11/1-11/15          | 0          | 181               | 4m.                   | 588      | 72          | 22        | 506        | .37                    | 1.16              | .14                  |

1/ u - Upper Section marked coho; m.-Middle Section marked coho; 1.-Lower Section marked coho.

2/ 8/31 marking of coho yearlings was terminated.



Other species inhabiting the system were chinook salmon, round whitefish (Prosopium cylindraceum), longnose sucker (Catostomus catostomus), arctic grayling, pink salmon, Dolly Varden, rainbow trout, threespine stickleback (Gasterosteus aculeatus), burbot, slimy sculpin (Cottus cognatus), and the western brook lamprey (Lampetra planeri).

Montana Creek was selected to obtain data on juvenile chinook salmon abundance and migration. A total of 25,176 fry were marked from July 19 through August 14. The distribution of marking was 16,039 in the upper area and 9,137 in the middle area. Species composition of other fish was similar to Rabideux Creek. Table 5 illustrates the findings of trapping in biweekly periods until the first of December. After this time, trapping was conducted one to three days per month.

The chinook salmon catch per hour indicated a gradual population density decline until February when a drastic reduction was recorded (Table 5). The gradual reduction is attributed to fry slowly moving downstream to the Susitna River throughout the season. This is also evidenced by marked fry being recovered below their area of release while no evidence of upstream recoveries was recorded.

Willow Creek was also sampled with minnow traps periodically between August 23 and March 2. This data clearly shows a decline in population density between December and February (Table 6).

The drastic reduction in population density found in February is attributed to the extremely low water conditions encountered at that time. The reduced flow was believed to have eliminated required rearing habitat and forced the juvenile salmonids into the mainstem Susitna River. Traps were set in the Susitna River and one of its sloughs to test this theory. Chinook salmon fry were recovered from the Susitna

Table 5. Montana Creek salmon fry trapping, Devils Canyon Project, 1977.

| Date           | Chinook<br>Fry<br>Unmarked | Chinook<br>Fry<br>Upper Mark | Chinook<br>Fry<br>Lower Mark | Coho<br>Fry | Coho<br>Yearling | Number<br>Traps | Total<br>Trap<br>Hours | Chinook<br>Per Trap | Chinook<br>Per Hour |
|----------------|----------------------------|------------------------------|------------------------------|-------------|------------------|-----------------|------------------------|---------------------|---------------------|
| UPPER SECTION  |                            |                              |                              |             |                  |                 |                        |                     |                     |
| 8/16-8/31      | 178                        | 56                           | --                           | --          | --               | 13              | 312                    | 18.0                | .75                 |
| 9/1-9/15       | 336                        | 6                            | --                           | 1           | 5                | 5               | 115                    | 68.4                | 2.97                |
| 9/16-9/30      | 461                        | 2                            | --                           | 11          | --               | 14              | 294                    | 33.1                | 1.57                |
| 10/1-10/15     | 4188                       | 7                            | --                           | --          | 14               | 110             | 2540                   | 38.1                | 1.65                |
| 10/16-10/31    | 2987                       | 16                           | --                           | 6           | 5                | 74              | 1560                   | 40.6                | 1.93                |
| 11/1-11/15     | 1467                       | 3                            | --                           | 2           | 8                | 37              | 888                    | 39.7                | 1.66                |
| 11/16-11/30    | 410                        | 1                            | --                           | --          | 2                | 17              | 402                    | 24.2                | 1.02                |
| 12/22          | 136                        | --                           | --                           | 2           | --               | 5               | 128                    | 27.2                | 1.06                |
| 1/27           | 185                        | --                           | --                           | 4           | --               | 5               | 126                    | 37.0                | 1.47                |
| 2/23-24        | 126                        | --                           | --                           | 1           | --               | 22              | 440                    | 5.7                 | 0.29                |
| MIDDLE SECTION |                            |                              |                              |             |                  |                 |                        |                     |                     |
| 8/16-8/31      | 1206                       | 6                            | 13                           | --          | --               | 15              | 360                    | 81.7                | 3.40                |
| 9/1-9/15       | 1445                       | 6                            | 8                            | 19          | 1                | 17              | 328                    | 85.8                | 4.45                |
| 9/16-9/30      | --                         | --                           | --                           | --          | --               | --              | --                     | --                  | --                  |
| 10/1-10/15     | 1982                       | 4                            | 4                            | --          | 10               | 39              | 936                    | 51.0                | 2.13                |
| 10/16-10/31    | 3218                       | 5                            | 10                           | 24          | 13               | 65              | 1490                   | 49.7                | 2.17                |
| 11/1-11/15     | 1601                       | 3                            | 5                            | 22          | 3                | 52              | 1208                   | 30.9                | 1.33                |
| 11/16-11/30    | 507                        | 3                            | 1                            | 3           | 3                | 17              | 390                    | 30.1                | 1.31                |
| 12/22          | 187                        | --                           | --                           | --          | 3                | 5               | 120                    | 37.4                | 1.56                |
| 1/27           | 40                         | --                           | --                           | 1           | --               | 7               | 130                    | 5.7                 | 0.31                |
| 2/23-24        | 32                         | --                           | --                           | --          | 1                | 20              | 406                    | 1.6                 | 0.08                |
| LOWER SECTION  |                            |                              |                              |             |                  |                 |                        |                     |                     |
| 8/16-8/31      | 1627                       | 6                            | 9                            | --          | --               | 24              | 576                    | 68.4                | 2.85                |
| 9/1-9/15       | 2077                       | --                           | 2                            | 56          | --               | 30              | 142                    | 69.3                | 14.64               |
| 9/16-9/30      | 891                        | 1                            | 3                            | 7           | 39               | 28              | 423                    | 32.0                | 2.12                |
| 10/1-10/15     | 5002                       | 4                            | 1                            | 100         | 162              | 141             | 3292                   | 35.5                | 1.52                |
| 10/16-10/31    | 2221                       | 6                            | 1                            | 75          | 21               | 54              | 1236                   | 41.3                | 1.80                |
| 11/1-11/15     | 647                        | 1                            | --                           | 3           | --               | 40              | 936                    | 16.2                | 0.69                |
| 11/16-11/30    | 456                        | --                           | --                           | 1           | 3                | 10              | 228                    | 45.6                | 2.00                |
| 12/21-23       | 174                        | 1                            | --                           | --          | 4                | 12              | 288                    | 14.6                | 0.61                |
| 1/27           | 116                        | --                           | --                           | 3           | --               | 5               | 108                    | 23.2                | 1.07                |
| 2/23-24        | 108                        | --                           | --                           | --          | 1                | 18              | 372                    | 6.0                 | 0.29                |

Table 6. Willow Creek chinook salmon fry trapping, Devils Canyon Project, 1977.

| Index<br>Area | <u>1977</u> | <u>Catch/Trap Hour</u> |      | <u>1978</u> |      |
|---------------|-------------|------------------------|------|-------------|------|
|               | 8/23        | 10/26                  | 12/1 | 1/18        | 3/2  |
| #1            | 2.8         | 2.6                    | 1.3  | 1.5         | 1.29 |
| #2            | 3.8         | 3.2                    | 3.3  | 1.3         | 0.28 |
| #3            | 4.2         | 4.1                    | 4.8  | 1.3         | 0.67 |

River at a rate of 0.45 per hour. In the slough they were recovered at a rate of 0.12 per hour. These catch rates document that chinook salmon juveniles utilize the mainstem river for rearing during the winter period.

#### Juvenile studies

Juvenile salmonids were present in all sloughs and clearwater tributaries identified within this study, with the exception of Lane Creek. The absence of juveniles in the latter location does not preclude their presence, since survey conditions of this creek were generally poor for juveniles. Pink salmon were the only species observed spawning within this creek and emergent fry would not be expected to be present when surveys were conducted, since this species migrate toward sea after their emergence from the gravel in late May and early June.

The major species utilizing these areas for rearing during summer months were chinook and coho salmon, although sockeye salmon were also collected. Misidentification of salmon fry samples collected in previous studies, particularly between chinook and coho salmon, was noted during 1977. Samples from previous years were reexamined and correct identification was made. Data indicates chinook salmon were the most abundant rearing species collected during 1974 through 1976.

Estimated fry abundance varied throughout the season. Lowest numbers occurred during late September surveys. This data is concurrent with studies conducted in Willow and Montana creeks (see p.25). Attempts were not generally made to establish migration from the upper sloughs and tributaries to the mainstem river. A limited experiment was, however, conducted in Indian River to determine if migrations observed in Montana and Willow creeks also occurred. A total of 579

chinook salmon fry were trapped during a two hour period on August 18. Large numbers of chinook salmon fry were also observed near the confluence area during late August and September. On August 31 the first chinook salmon fry was trapped in the mainstem Susitna River immediately downstream of Indian River. Logistical problems prevented follow-up studies until March 7. Ten traps were fished on this date for 24 hours in areas where high densities of fry had been observed during the summer. Only four chinook salmon were captured. Data is limited, but it does corroborate findings in Montana Creek. Montana Creek and Indian River have comparable gradients, velocities, pool to riffle ratios, and are representative of most of the clearwater tributaries to the Susitna River. It would be reasonable to speculate that life history information of salmon fry from one of these tributaries would be representative of the other.

In addition to the apparent intrasystem migration of juvenile chinook salmon from the lateral tributaries to the Susitna River in the fall, it appears some young-of-the-year chinooks move out of the parent stream in the spring. The majority of the salmon fry observed in sloughs during 1977 were chinook salmon. Adult chinook salmon were not observed spawning in these sloughs during 1976. Observations, therefore, indicate the fry dropped out of spawning areas sometime in the spring into the Susitna River and then moved into the sloughs to rear for the summer.

Definition of the intrasystem migrations for the various life history phases of each species will be important considerations in assessing the potential impacts of this project. It can be assumed that individuals of a species will tend to select areas within a drainage that have the most favorable combinations of hydraulic conditions which

support life history requirements. They will also utilize less favorable conditions, with the probability-of-use decreasing with diminishing favorability of one or several hydraulic conditions (Bovee, 1978). Observations demonstrate that individuals elected to leave an area before conditions became lethal. The movement of rearing salmon fry out of the sloughs in the fall has been documented and is an example of areas where conditions could become lethal.

Data indicates that in early summer salmon rearing conditions are poor in the mainstem Susitna River because of high discharge and sediment loads. The clearwater sloughs and tributary areas are utilized by fry at this time. As the season progresses, discharge and sediment loads of the mainstem Susitna begin to decrease. By fall and winter, the silt load appears to be low enough to transform the mainstem Susitna River into suitable fry rearing habitat to replace slough areas, which are dewatered when mainstem discharge and stage decreases, and tributaries that often freeze in the winter.

Samples for age, length and weight analysis were obtained from each slough during late July and early August and late September. Analysis will not be discussed, but is presented in Appendix I Tables 3, 4, and 5.

#### Aquatic insects and juvenile salmon gut contents

Knowledge of the aquatic insect fauna and its ecology is necessary to assess the potential impacts of the Devils Canyon and Watana dams upon the salmon population downstream. Alterations of currently existing populations would probably have a corollary effect upon rearing fish.

Gut contents of juvenile salmon from sloughs and tributaries between Portage Creek and the Chulitna and Susitna River confluence were

examined to determine feeding habits of rearing fish during 1977.

Studies were considered minimal and further investigations will be required.

Immature members of the Orders Diptera, Plecoptera, Ephemeroptera, Trichoptera, Coleoptera, Hemiptera, and adult forms of Hemiptera and Coleoptera were found in the summer and fall diets of juvenile salmon (Appendix I Table 6). Adult terrestrial insects were estimated to be the largest percentage of the gut contents by volume. Although most of these adult forms were terrestrial, the majority of their life histories were spent in the aquatic environment.

Percent composition of gut contents varied between species of fish examined (Table 7). Feeding habits of chinook and coho salmon were, however, similar during the summer sampling period. Adult Insecta were of primary importance for the latter two species during summer. Sockeye salmon fry fed primarily on Diptera larvae during summer months. Cladocera (Bosminidae) were also found to be important food organisms for sockeye salmon in three sloughs (Appendix I Table 6).

Adult Insecta remained the major food items identified in the fall stomach content samples. Adult Diptera and Hymenoptera comprised approximately 80 percent of the food items in sockeye salmon during the fall as compared to about 18 percent during summer. The importance of immature Insecta and Crustacea apparently decreased appreciably. Change in percent composition of food items per fish was not significant for chinook and coho salmon fry.

Aquatic insects probably play a more important role in the juvenile salmon diet during winter months than in the summer and fall. Many groups of insects (Plecoptera, Ephemeroptera, Trichoptera, and Diptera)

Table 7. Mean percent composition of gut contents per fish of chinook, sockeye, and coho salmon juveniles in sloughs and clearwater tributaries of the Susitna River, Devils Canyon Project, 1977.

| Species | Mean Percent Per Fish |      |           |       |                  |      |               |      |        |      |
|---------|-----------------------|------|-----------|-------|------------------|------|---------------|------|--------|------|
|         | Sample Size           |      | Crustacea |       | Immature Insecta |      | Adult Insecta |      | Other  |      |
|         | Summer                | Fall | Summer    | Fall  | Summer           | Fall | Summer        | Fall | Summer | Fall |
| Chinook | 219                   | 158  | 4         | trace | 24               | 26   | 71            | 62   | 1      | 12   |
| Sockeye | 35                    | 18   | 27        | 2     | 54               | 17   | 18            | 80   | 1      | 1    |
| Coho    | 17                    | 45   | 9         | trace | 17               | 9    | 68            | 69   | 6      | 22   |



are very active during the winter even at water temperatures of 0°C (Hynes, 1970). Conversely, during these cold months terrestrial insects are nonexistent and plankton is either greatly reduced or nil. This would suggest that aquatic insects would probably be a greater proportion of the juvenile salmon diets than in the summer. Additional studies are required to analyze this.

Research and literature in the area of environmental factors affecting aquatic insects is sparse and often times conflicting. There is, apparently, a high degree of variability as to substrate type preference, temperature requirements, and general modes of existence even within the Order level. Evaluating species diversity would probably be the most useful means of monitoring on-going environmental changes in the invertebrate fauna of the river (McCoy, 1974). It would not, however, provide a means to predict whether or how a change will occur. Environmental factors which would probably result in the greatest alterations in the aquatic fauna include: water temperature, flow, substrate types, water clarity, and chemical water quality.

Research in the area of water temperature effects on aquatic fauna are conflicting, but apparently the "environmental clues" for the hatching of eggs, the change from a larval to pupal state, etc., are a combination of threshold temperatures and changing day length (Hynes, 1970). Disruptions in the seasonal pattern of temperature are attributed to have caused extensive alterations in the aquatic insect fauna of the Saskatchewan River (Lehmkuhl, 1972). Hypolimnial water discharge from a dam in the river reduced both diversity and absolute numbers of insects downstream. River temperatures became higher in winter and lower in summer, differing from the norm in such a way that Ephemeroptera eggs

failed to develop into nymphs. Similar temperature effects were thought to have adversely affected other aquatic insect groups at this site, even at a distance of 70 miles downstream. Alteration of natural flow could affect both the respiration of organisms and substrate types.

Most arthropods in still water self-ventilate their gills or respiratory structures. Many immature aquatic insects have lost this function and rely on running water or current to artificially "fan" their gills. A decrease in flow could therefore have an adverse effect upon respiration. The nature of the flow is intimately related to substrate type. A fast current area will generally be clean swept and have a rocky or gravel substrate. The sediment load will drop in slow moving waters and the bottom will become increasingly silty. Each different substrate type supports a completely different benthic fauna. All these current related factors can perhaps best be summarized by Hynes' observation that areas subjected to wide fluctuations in current "are often without much fauna." Neither those organisms adapted to a slow moving area nor those to one of swift water can thrive.

Numerous investigators have established the importance of substrate types upon the nature of the benthic fauna. Each species of aquatic insect seems adapted to a certain substrate type or at least greatly prefers one type to another. Obviously, changes in substrate type will result in altered benthic fauna. This was evidenced when a small beaver dam across a stream in Ontario altered the upstream bottom habitat from swift flowing and stoney to slow moving and silty stones. The total number of aquatic insects were reduced, "especially of Ephemeroptera, Plecoptera, and Trichoptera," while the proportion of Diptera Chironomid larvae was increased (Hynes, 1970). There can be great variations in

substrate preference within each order or even family. Some trends are, however, discernable. In general, rocky or stoney substrates with a swift flow of water will contain both a greater species diversity and a higher biomass than silty substrates with slower moving water. These riffle areas are the most productive regions in running water.

The possible introduction of turbid glacial water by the proposed dam into the clear winter water of the upper Susitna seems to indicate substrate type would be altered to one of increasing silt. This would probably change the aquatic insect fauna and quite possibly reduce its abundance.

Chemical water quality influences upon aquatic insects would be minimal in comparison to the above factors. Lehmkuhl (1972) and Spence and Hynes (1971) discovered no appreciable differences in chemical water quality upstream and downstream from dam impoundments and thus concluded there were no effects from these factors upon benthic invertebrates.

The importance of drift to the relationship between aquatic insects and the diet of juvenile salmon is another factor to consider. Many benthic invertebrates, displaced by crowded conditions and as a means of finding more favorable substrate habitats, leave the substrate and are carried downstream by the water's flow. These are cumulatively called "drift". Investigators have repeatedly found that most of the food items of salmonid fish in flowing water situations consist of drift. Hynes (1970) reports that brown trout feed mostly on drifting organisms. Becker's (1973) food habits study of juvenile chinook salmon on the Columbia River concluded prey items were either drift organisms or adult insects floating on the water's surface. Loftus and Lenon (1977) also believed drift to be an important food source to chinook and chum smolt

on the Salcha River in interior Alaska. A comparison between the gut contents of a limited number of longnose suckers (bottom feeders) collected in our study with that of the juvenile salmon reveals that drift aquatic insects together with floating adult insects were apparently the major food items. The numbers and kinds of organisms in the drift appear to differ substantially when compared to fauna collected strictly on the bottom. As might be expected, heavier organisms such as Trichoptera larvae and their cases, snails, etc., are relatively rare in drift, while Ephemeroptera, Diptera Chironomid larvae, and Plecoptera form a higher percentage than they do on the substrate. Various environmental factors can alter the amount of drift. Investigators have reported varying drift because of ice scouring, water temperature, and daylight changes (Hynes, 1970). The role of drift organisms in both the food habits of rearing salmonid fishes and in the overall ecology of aquatic insects is thus probably of some importance in the Susitna River and should be investigated further.

If a hypolimnial discharge hydroelectric dam is constructed at Devils Canyon, it appears almost certain the downstream benthic fauna will be altered. This will most probably occur because of: 1) changed water temperatures resulting from the hypolimnial discharge which may disrupt the life cycles of certain species; 2) substrate types altered by increased winter turbidity of downstream river water, which will in turn alter the aquatic insects living on the substrate, and 3) discharge flow variations because of varying power demands, which will create areas of the river bottom to which neither swift current associated species nor slow current forms are perfectly adapted for. Which species or group of insects will be most affected, whether they will be major

food items of rearing juvenile salmon or whether the salmon will switch their food preference to the newly abundant forms, and whether the biomass of benthic fauna will decrease, will probably be difficult, if not impossible, to predict. We can only hope to broadly outline what changes may occur.

#### Impoundment area fisheries investigations

Alterations will definitely occur to the fish habitat in the areas to be inundated. The fisheries investigations in the impoundment area during the first two weeks of July revealed that Arctic grayling were abundant in all of the major clearwater tributaries (Table 8). Extreme lake level fluctuations of the Watana reservoir will destroy habitat and affect the high quality fishery which presently exists.

No anadromous species were captured upstream of Devils Canyon during the first two weeks of July. More extensive sampling, however, is necessary throughout the summer to determine if Devils Canyon is a velocity barrier to salmon during different natural flow regimes over a three to five year period.

Lakes in the impoundment area which could be impacted by construction of road or transmission corridors and increased access were also surveyed for species composition (Table 9). Fifteen of the eighteen lakes sampled supported desirable game fish populations.

Construction of the Devils Canyon dam would inundate 7,550 acres and have a surface elevation of 1,450 feet and extend for 28 miles upstream (U.S. Army Corps of Engineers, 1977). Construction of the Watana dam would result in inundation of 43,000 acres with a surface elevation of 2,200 feet extending for 54 miles upstream along the Susitna River. For downstream discharge to remain relatively constant, at least

Table 8. Limmological data from selected tributaries to the Susitna River, Devils Canyon Project, 1977.

| Stream  | Est.<br>Flow<br>(cfs) | Estimated<br>Velocity<br>(fps) | Percent<br>Pools | Bottom<br>Type    | Temp. | pH  | Conduc-<br>tivity | Fish<br>Observed* |
|---------|-----------------------|--------------------------------|------------------|-------------------|-------|-----|-------------------|-------------------|
| Oshetna | 600                   | 3                              | 15               | Rubble<br>Boulder | 13    | 8   | 75                | GR                |
| Goose   | 100                   | 2                              | 40               | Rubble<br>Boulder | 15    | -   | -                 | GR                |
| Jay     | 75                    | 2                              | 40               | Gravel<br>Boulder | 8     | 8.4 | 160               | GR, SK, WF,<br>SC |
| Kosina  | 100                   | 2                              | 30               | Gravel<br>Boulder | 14    | 8   | 65                | GR                |
| Watana  | 300                   | 1.5                            | 20               | Gravel<br>Rubble  | 12    | 7.8 | 110               | GR                |
| Deadman | 900                   | 3                              | 10               | Boulder           | 14    | -   | -                 | GR                |
| Tsusena | 600                   | 2                              | 10               | Gravel<br>Boulder | 6     | 7.8 | 50                | GR                |
| Fog     | 200                   | 1.5                            | 30               | Sand              | 9     | 7.9 | 75                | GR                |

\* GR - Grayling  
SK - Suckers  
SC - Sculpin  
WF - Whitefish

Table 9. Susitna River impoundment area lake surveys, Devils Canyon Project, 1977.

| Lake                 | Location                         | Surface<br>Elevation | Surface<br>Acres | Maximum<br>Depth (Ft) | Fish Species<br>Present*      |
|----------------------|----------------------------------|----------------------|------------------|-----------------------|-------------------------------|
| Clarence             | T30N, R9E, S19, 20               | 2,900                | 299              | 35                    | LT, GR, WF                    |
| Fog 1                | T31N, R5E, S9                    | 2,230                | 147              | 72                    | DV, SC                        |
| 2                    | T31N, R5E, S8                    | 2,230                | 237              | 50                    | DV, SC                        |
| 3                    | T31N, R5E, S15                   | 2,110                | 339              | 81                    | DV, SC                        |
| 4                    | T31N, R5E, S13                   | 2,300                | 358              | 9                     | DV, SC                        |
| 5                    | T31N, R6E, S7                    | 2,300                | 269              | 6                     | -                             |
| George               | T6N, R7W, S20, 29                | 2,400                | 80               | 18                    | GR, LNS                       |
| Louise               | T32N, R6E, S7                    | 2,362                | 155              | 155                   | LT, BB, WF, GR                |
| Connor               | T6N, R7W, S28                    | 2,450                | 18               | 13                    | GR                            |
| Tsusena Butte        | T33N, R5E, S21                   | 2,493                | 190              | 110                   | GR, LT, WF                    |
| Pistol               | T32N, R6E, S7                    | 2,350                | 205              | -                     | -                             |
| Big                  | T32S, R3, 4W, S25,<br>18, 19, 30 | 3,070                | 1,080            | 80                    | LT, WF                        |
| Deadman              | T22S, R4W, S13, 14               | 3,064                | 380              | 70                    | LT, GR, WF                    |
| Watana               | T30N, R7W, S36                   | 3,000                | 300              | 30                    | LT, WF, GR                    |
| Square               | T30N, R3E, S35                   | 1,935                | 230              | 34                    | -                             |
| Little Moose<br>Horn | T30N, R3E, S36                   | 1,850                | 120              | 33                    | GR, LT, LNS                   |
| Stephan              | T30N, R3E, S2,10,16              | 1,862                | 840              | 95                    | LT, RT, RS<br>SS, GR, WF, LNS |

\* Species: GR - Grayling      WF - White Fish      RS - Sockeye Salmon  
RT - Rainbow Trout      SC - Sculpin      LNS - Long Nosed Sucker  
DV - Dolly Varden      SS - Coho Salmon  
LT - Lake Trout      BB - Burbon

one of these reservoirs will have to fluctuate considerably. The Watana reservoir is projected to have the most extreme fluctuations. The majority of the clearwater tributaries to be inundated are found within this section of river and, of the two impoundments, greater impacts will probably occur here since loss of portions of these tributaries is inevitable if the two dams are built. If salmon utilize the area above the Devils Canyon dam site, however, both the Devils Canyon and Watana dams and impoundments could adversely impact migration. Reservoir fluctuations could have a variety of effects on the tributaries. The mouths of these tributaries and stretches of water upstream provide some of the most productive fishery habitat in this area. Some tributaries have steep gradients upstream of the mouth area which act as migration barriers and do not appear to support fish species.

In tributaries where the full pool would extend up to the base of steep tributary gradients or waterfalls, critical lotic habitat would be lost. Periods of lowered pool levels could have a suction effect and result in the erosion and formation of channels with steep gradients which may block intersystem fish migrations and eliminate suitable fishery habitat. Preliminary data on fish species present demonstrates that additional information is required to evaluate the full effects of inundation and regulation in these areas.

#### WATER QUANTITY

Between May 17 and June 14, 1977 the unregulated flow of the Susitna River increased from 13,600 cubic feet per second (cfs) to a peak discharge of 52,600 cfs (Figure 4; Appendix II, Table 1). By July 20, the flow decreased to 22,400 cfs and fluctuated around 20,000 cfs until August 25. On September 6 the flow dropped to 9,520 cfs and then increased to






Figure 4. Susitna River discharge at Gold Creek, Devils Canyon Project, 1977.

16,900 cfs one week later. The flow decreased to 9,840 cfs on September 27 which again was followed by increased flow until the last reading of 12,500 cfs was made on September 30.

Fluctuations in flow during August and September were attributed to heavy rain. Stage fluctuations within the majority of clearwater sloughs of the Susitna River, related directly to mainstem discharge variations (Appendix II, Table 2). Downstream flow is projected to be maintained at a constant rate of approximately 7,000 to 8,000 cfs at Gold Creek after completion of the dams. Slough surveys were terminated near the end of September when the flow was approximately 15,000 cfs. It was not possible to observe the sloughs during this study when the mainstem flow was 8,000 cfs due to freezing conditions. Observations during the 1976 study, however, concluded that 75 percent of the rearing sloughs studied were undesirable habitat when the flow in the mainstem was 7,000 cfs (Riis, 1977).

Habitat requirements for passage, spawning, egg incubation, fry, juvenile, and adult phases of the salmon species studied are quite specific. The USFWS Cooperative Instream Flow Service Group has developed criteria which demonstrate the narrow tolerances of certain salmonid species to hydraulic parameters of velocity, depth, substrate and temperature (Bovee, 1978). The seasonally wide fluctuations of water velocity, depth, temperature, substrate, and sediment of the mainstem Susitna, its sloughs and tributaries determine to some extent the intrasystem migrations of fish seeking more desirable environments. Thus, any alterations to the existing aquatic ecosystem which restrict or reduce the availability of required habitat, will also reduce fish production.

Low flows were encountered in Rabideux Creek from mid-June through the end of August (Table 10). The lowest flow recorded was 24.3 cfs on August 23. The highest flow was 440.7 cfs on September 29 and was apparently due to the heavy rains encountered at that time.

#### WATER QUALITY

Ryan thermographs were installed in the upper sub-areas of Rabideux and Montana creeks. Water temperatures in Rabideux Creek ranged up to five degrees celsius ( $^{\circ}\text{C}$ ) higher than Montana Creek during corresponding time periods. The high recorded in Rabideux Creek was  $18.8^{\circ}\text{C}$  on both July 11 and 12; the low of  $1.7^{\circ}\text{C}$  occurred on October 22 and 23 at which time the thermograph was removed (Appendix II, Table 3). In Montana Creek, a high of  $15.0^{\circ}\text{C}$  was recorded on July 28 and the low of  $0.0^{\circ}\text{C}$  was recorded from November 3 through 6 at which time recording was terminated (Appendix II, Table 4).

A thermograph was also installed in the Susitna River at the Parks Highway bridge. When installed on June 27 the temperature was  $10.5^{\circ}\text{C}$  and the highest water temperature of  $14^{\circ}\text{C}$  was reached on July 12 followed by temperatures fluctuating between  $13.5^{\circ}\text{C}$  and  $10^{\circ}\text{C}$  when a steady decline began on August 25 and continued to the lowest reading of  $2^{\circ}\text{C}$  on October 2 (Table 11).

Temperatures at all other sampling sites were measured with a combined dissolved oxygen and temperature meter and/or a pocket thermometer. Data is presented in Appendix II, Table 2.

Water chemistry of Rabideux and Montana creeks was measured throughout the season. Determinations of dissolved oxygen, pH, hardness, and total alkalinity are presented in Appendix II, Tables 5 and 6.

In Rabideux Creek, dissolved oxygen ranged from a low of 6 ppm in the upper sub-area to a high of 11 ppm recorded in all areas. Hydrogen

Table 10. Water flows of Montana, Rabideux, and Willow creeks from May through November, Devils Canyon Project, 1977.<sup>1/</sup>

MONTANA CREEK

| Date  | Flow (cfs) |
|-------|------------|
| 5/1   | 935        |
| 5/21  | 2,000      |
| 6/5   | 4,800      |
| 6/20  | 1,764      |
| 7/1   | 935        |
| 7/21  | 935        |
| 8/6   | 233        |
| 8/22  | 153        |
| 9/1   | 103        |
| 9/29  | 1,349      |
| 10/15 | 394        |
| 11/9  | 490        |

RABIDEUX CREEK

| Date | Flow (cfs) |
|------|------------|
| 4/13 | 325.4      |
| 5/25 | 128.7      |
| 6/7  | 116.7      |
| 6/17 | 50.2       |
| 6/30 | 33.2       |
| 7/13 | 36.7       |
| 7/26 | 31.4       |
| 8/23 | 24.3       |
| 8/31 | 29.2       |
| 9/21 | 242.9      |
| 9/29 | 440.7      |

WILLOW CREEK

| Date  | Flow (cfs) |
|-------|------------|
| 5/1   | 443        |
| 5/30  | 1,590      |
| 6/15  | 3,320      |
| 6/29  | 1,900      |
| 7/15  | 951        |
| 7/30  | 525        |
| 8/15  | 409        |
| 8/30  | 322        |
| 9/16  | 1,590      |
| 9/29  | 2,070      |
| 10/15 | 525        |
| 10/30 | 348        |
| 11/8  | 676        |

<sup>1/</sup> Montana and Willow creeks data is provisional and was obtained from the National Weather Service.

Table 11. Thermograph set in Susitna River downstream of Parks Highway Bridge, daily maximum and minimum water temperature, Devils Canyon Project, 1978.

| Date | Temp. °C |      | Date | Temp. °C |      | Date  | Temp. °C |      |
|------|----------|------|------|----------|------|-------|----------|------|
|      | Min.     | Max. |      | Min.     | Max. |       | Min.     | Max. |
| 6/27 | 10.5     | 10.5 | 7/30 | 12.5     | 12.5 | 9/12  | 7.5      | 8.0  |
| 6/28 | 10.5     | 10.5 | 7/31 | 11.0     | 12.5 | 9/13  | 7.5      | 7.5  |
| 6/29 | 10       | 10.5 | 8/1  | 10.0     | 10.5 | 9/14  | 7.5      | 7.5  |
| 6/30 | 10       | 10   | 8/2  | 10.0     | 10.0 | 9/15  | 6.0      | 7.5  |
| 7/1  | 10.5     | 10.5 | 8/3  | 10.0     | 11.0 | 9/16  | 6.0      | 6.5  |
| 7/2  | 10.5     | 10.5 | 8/4  | 11.0     | 11.0 | 9/17  | 6.5      | 6.5  |
| 7/3  | 10       | 10.5 | 8/5  | 11.0     | 11.0 | 9/18  | 6.5      | 6.5  |
| 7/4  | 9.5      | 10   | 8/6  | 10.5     | 11.0 | 9/19  | 6.0      | 6.5  |
| 7/5  | 9.5      | 10   | 8/7  | 11.0     | 11.0 | 9/20  | 5.5      | 6.5  |
| 7/6  | 10       | 11   | 8/8  | 10.0     | 10.5 | 9/21  | 5.5      | 5.5  |
| 7/7  | 12       | 12.5 | 8/9  | 10.0     | 11.5 | 9/22  | 5.5      | 6.0  |
| 7/8  | 12       | 13   | 8/10 | 11.0     | 11.5 | 9/23  | 5.5      | 6.0  |
| 7/9  | 12       | 13   | 8/11 | 10.5     | 11.0 | 9/24  | 5.0      | 5.5  |
| 7/10 | 12.5     | 13.5 | 8/12 | 10.5     | 11.0 | 9/25  | 4.5      | 5.0  |
| 7/11 | 13       | 13.5 | 8/13 | 10.5     | 11.0 | 9/26  | 4.5      | 5.0  |
| 7/12 | 13.5     | 14   | 8/14 | 10.5     | 11.0 | 9/27  | 5.0      | 5.0  |
| 7/13 | 13       | 13.5 | 8/15 | 10.5     | 11.0 | 9/28  | 5.0      | 5.0  |
| 7/14 | 11       | 13   | 8/16 | 11.0     | 11.0 | 9/29  | 4.5      | 5.0  |
| 7/15 | 10.5     | 11   | 8/17 | 11.0     | 11.0 | 9/30  | 3.0      | 4.5  |
| 7/16 | 10.5     | 11.5 | 8/18 | 10.0     | 10.5 | 10/1  | 2.5      | 3.0  |
| 7/17 | 11.3     | 12   | 8/19 | 10.5     | 12.0 | 10/2  | 2.0      | 2.5  |
| 7/18 | 12       | 12   | 8/20 | 11.0     | 12.0 | 10/3  | 2.0      | 2.0  |
| 7/19 | 11.5     | 11.5 | 8/21 | 10.5     | 12.0 | 10/4  | 2.0      | 3.0  |
| 7/20 | 11.5     | 11.5 | 8/22 | 11.0     | 11.5 | 10/5  | 2.5      | 3.0  |
| 7/21 | 11       | 11   | 8/23 | 11.0     | 12.0 | 10/6  | 2.0      | 2.5  |
| 7/22 | 11       | 11.5 | 8/24 | 10.5     | 11.5 | 10/7  | 2.5      | 2.5  |
| 7/23 | 11       | 11.5 | 8/25 | 9.5      | 10.5 | 10/8  | 2.5      | 3.0  |
| 7/24 | 11       | 11.5 | 8/26 | 9.0      | --   | 10/9  | 3.0      | 3.5  |
| 7/25 | 11.5     | 11.5 | --   | --       | --   | 10/10 | 3.5      | 3.5  |
| 7/26 | 11.5     | 11.5 | --   | --       | --   | 10/11 | 3.5      | 4.0  |
| 7/27 | 10.5     | 12.0 | 9/9  | 8.0      | --   | 10/12 | --       | 3.5  |
| 7/28 | 11.0     | 12.5 | 9/10 | 7.5      | 8.0  | --    | --       | --   |
| 7/29 | 12.0     | 13.0 | 9/11 | 7.5      | 8.0  | --    | --       | --   |

ion (pH) concentrations were found to be relatively stable ranging from a low of 6.5 to a high of 7.7. Both hardness and total alkalinity were found to range between 17 mg/l to 68 mg/l. The higher readings occurred during the warmer summer months.

Montana Creek exhibited less fluctuation in chemical water characteristics than Rabideux Creek. The dissolved oxygen ranged from 9 to 12 ppm, pH from 6.8 to 7.7, and hardness and total alkalinity from 17 to 34 mg/l.

Water samples were collected jointly by ADF&G and USGS from three sites on the Susitna and the USGS laboratory carried out the complete standard chemical analysis. This data is presented in Appendix II, Table 7 and considerably expands the data base which will be used for future comparisons.

Field determinations of dissolved oxygen, pH, hardness, total alkalinity and specific conductance were collected in clearwater sloughs and tributaries and are tabulated in Appendix II, Table 2. The findings were within acceptable limits for fish life and were in the range of expected results for natural waters in southcentral Alaska.

## CONCLUSION

Baseline inventory studies, to date, emphasize the need to initiate a comprehensive study to properly assess the potential environmental impacts to the aquatic ecosystem of the Susitna drainage by the proposed Watana and Devils Canyon hydroelectric project prior to final design approval and construction authorization.

The Susitna River is a product of its tributaries. All aquatic habitat and populations (within the power transmission corridor site, construction road routes, and above and below the proposed dam sites) which would be directly or indirectly affected during construction and after completion of the project must be carefully evaluated. It is imperative to thoroughly investigate the interrelationships between the aquatic biology and the water quantity and quality of the existing free flowing Susitna River system. Recreational, social, economic, and aesthetic considerations should also be included.

With this information the Alaska Department of Fish and Game will be able to provide the input for preventing unnecessary losses of the fisheries and related resources held in high esteem by the people of Alaska and the Nation as a whole.

## RECOMMENDATIONS

Continued collection of biological data and completion of resource assessment in the area affected by the proposed hydroelectric project is essential to understanding the potential impacts of the proposed action. Appendix III is a summary of ADF&G's recommendations for essential aquatic studies.

Direct studies of aquatic and terrestrial species can delineate a population and indicate their distribution throughout the year and define why species are there to a certain extent. Seasonal life history studies must be accompanied by habitat studies if we are to determine the full significance of habitat alteration to the population.

The studies identified for the pre-authorization environmental assessment are necessary to predict the impacts of hydroelectric development on the ecosystem. The objectives of the biological investigations are based upon the assumption that the Devils Canyon and Watana two dam plan will be selected. It must be realized that as the plan evolves and new information becomes available, the program must be flexible enough to permit adjustment in study direction. If other basin development schemes are proposed, study time and costs will have to be reevaluated. Capital requirements for each year were based upon FY-78 dollars. Inflation will therefore necessitate annual supplemental allocations which represent revised cost estimates. The proposals are closely integrated and demonstrate the need for continuity. The design, timing, manpower requirements, and funding levels of the individual projects have been coordinated.

A team of resource specialists representing various scientific disciplines will be required to carry out field investigations in habitat



assessment. Adequate time will be required to organize study personnel and procure equipment prior to the first field season. An untimely delay could prevent the initiation of the field studies one year.

## ACKNOWLEDGEMENTS

Several biologists with the Alaska Department of Fish and Game gave freely of their time and expertise to assist us during this study. We would like to express our thanks to Paul Janke, Larry Engel, Roger Wadman, Stan Kubik, Larry Heckart, Fred Williams, Mike Mills, Dave Watsjold and Bob Chlupach.

The temporary biologists who worked on this project deserve credit for long hours spent collecting and analyzing data: Christopher Estes, Jeff Hock, Terry Bradley, Pete Probasco, Kevin Delaney, Ward Knous, Steven Sykes, Doug Bue, Dave Clausen and Dave Colwell. Laboratory analysis and literature research on aquatic insect samples were conducted by Dave Clausen. Special appreciation is extended to Christopher Estes for his participation in the preparation of this document.

Funding for this study was provided by the U. S. Fish and Wildlife Service. Individuals from the USFWS also assisted with the Chinook fry marking operation in Montana Creek.

The USGS cooperated in every way possible in collecting and analyzing water samples.

Appreciation is also extended to those individuals not mentioned here who assisted with this project.

## LITERATURE CITED

- Alaska District U.S. Army Corps of Engineers. 1977. Plan of Study for Susitna hydropower feasibility analysis. Prepared under contract agreement for the state of Alaska. September. 297 pp.
- Barrett, B.M. 1974. An assessment study of the anadromous fish population in the upper Susitna River watershed between Devils Canyon and the Chulitna River. Cook Inlet Data Report No. 74-2. Alaska Department of Fish and Game. Division of Commercial Fisheries. 56 pp.
- Becker, C.D. 1973. Food and growth parameters of juvenile chinook salmon, Oncorhynchus tshawytscha, in central Columbia River. Fishery Bulletin. Vol. 71. No. 2. 387-400 pp.
- Bovee, K.D. 1978. Probability-of-use criteria for the family Salmonidae FWS/OBS - 78/07 Cooperative Instream Flow Service Group. Fort Collins, Colorado.
- \_\_\_\_\_, and T. Cochnauer. 1977. Development and evaluation of weighted criteria probability-of-use curves for instream flow assessment: Fisheries FWS/OBS - 77/63 Cooperative Instream Flow Service Group. Fort Collins, Colorado. 39 pp.
- Brown, C.J.D. 1971. Fishes of Montana Big Sky Books. Bozeman, Montana. 207 pp.
- Friese, N.V. 1975. Preauthorization assessment of anadromous fish populations of the upper Susitna River watershed in the vicinity of the proposed Devils Canyon hydroelectric project. Cook Inlet Data Report No 75-2. Alaska Department of Fish and Game. Division of Commercial Fisheries. 121 pp.
- Friese, N.V. (in prep). Susitna River salmon escapement studies, 1977. Cook Inlet Data Report Series. Alaska Department of Fish and Game. Division of Commercial Fisheries.
- Hynes, H.B.N. 1970. The ecology of running water. Univ. of Toronto Press, Toronto.
- Jaques, H.C. 1947. How to know the insects. Second Edition. Wm. C. Brown Co. Dubuque, Iowa. 205 pp.
- Kubik, S.W. and R. Wadman. 1977. Inventory and cataloging of sport fish waters of the lower Susitna River and central Cook Inlet drainages. Annual Report of Progress, 1976-1977. Project F-9-9, Vol. 18, No. G-I-H. 28 pp.
- Lehmkuhl, D.M. 1972. Change in thermal regime as a cause of reduction of benthic fauna downstream of a reservoir. J. Fish. Res. Bd. Canada. Vol. 29, No. 9. 1329-1332 pp.
- Loftus, W.F. and H.L. Lenon. 1977. Food habits of the salmon smolts, Oncorhynchus tshawytscha and O. keta, from the Salcha River, Alaska. Trans. Am. Fish. Soc. Vol. 106, No. 3. 235-240 pp.

# LITERATURE CITED (cont.)

- McCoy, G. 1974. Preconstruction assessment of biological quality of the Chena and Little Chena rivers in the vicinity of Chena Lakes Flood Control Project near Fairbanks, Alaska. U.S. Geological Survey, Water Resources Investigations 29-74. 84 pp.
- Pennak, R.W. 1953. Freshwater invertebrates of the United States. The Ronald Press Co. New York.
- Riis, J.C. 1977. Pre-authorization assessment of the proposed Susitna River hydroelectric projects: preliminary investigations of water quality and aquatic species composition. Alaska Department of Fish and Game. Division of Sport Fish. 91 pp.
- Spence, J.A. and H.B.N. Hynes. 1971a. Differences in benthos upstream and downstream of an impoundment. J. Fish. Res. Bd. Canada. Vol. 28, No. 1. 35-43 pp.
- . 1971b. Differences in fish populations upstream and downstream of a mainstem impoundment. J. Fish. Res. Bd. Canada. Vol. 28, No. 1. 45-46 pp.
- U.S. Fish and Wildlife Service. 1976. Southcentral railbelt area - Susitna River basin. Fish and wildlife studies related to the Corps of Engineers Devils Canyon - Watana Reservoir hydroelectric project. Studies conducted by the Alaska Dept. of Fish and Game under contract agreement with the U.S. Fish and Wildlife Service. February. 185 pp and appendices.
- Usinger, R.L. 1968. Aquatic insects of California. Univ. of Calif. Press. Berkeley.
- Ward, H.B. and G.C. Whipple. 1959. Freshwater Biology. Second Edition. John Wiley and Sons, Inc. New York.
- Watsjold, D.A. 1977. Inventory, cataloging, and population sampling of the sport fish and sport fish waters in upper Cook Inlet. Annual Report of Progress, 1976-1977. Project F-9-9, Vol. 18, No. G-I-D. 48 pp.

## APPENDIX I

Tables in the following appendix include data on adult and juvenile salmonids and stomach content analysis.

## Appendix I

Table 1. Percent age composition of chinook, sockeye, coho, and chum salmon escapement samples, Devils Canyon Project, 1974, 1975, and 1977.

| Year of<br>Return | Age Class |      |      |      |      | Brood Year |      |      |      |      | Sample<br>Size |
|-------------------|-----------|------|------|------|------|------------|------|------|------|------|----------------|
| <u>CHINOOK:</u>   |           |      |      |      |      |            |      |      |      |      |                |
| 1977              | 1.1       | 1.2  | 1.3  | 1.4  | 1.5  | 1970       | 1971 | 1972 | 1973 | 1974 |                |
| Percent           | 9.5       | 9.5  | 52.4 | 28.6 | 0.0  | 0.0        | 28.6 | 52.4 | 9.5  | 9.5  | 100.0          |
| Number            | 2         | 2    | 11   | 6    | 0    | 0          | 6    | 11   | 2    | 2    | 21             |
| 1975              | 1.1       | 1.2  | 1.3  | 1.4  | 1.5  | 1968       | 1969 | 1970 | 1971 | 1972 |                |
| Percent           | 9.3       | 4.6  | 34.9 | 44.2 | 7.0  | 7.0        | 44.2 | 34.9 | 4.6  | 9.3  | 100.0          |
| Number            | 4         | 2    | 15   | 19   | 3    | 3          | 19   | 15   | 2    | 4    | 43             |
| <hr/>             |           |      |      |      |      |            |      |      |      |      |                |
| <u>SOCKEYE:</u>   |           |      |      |      |      |            |      |      |      |      |                |
| 1977              | 1.1       | 1.2  | 1.3  | 2.1  | 2.2  | 1972       | 1973 | 1974 |      |      |                |
| Percent           | 3.3       | 16.7 | 76.7 | 3.3  | 0.0  | 76.7       | 20.0 | 3.3  |      |      | 100.0          |
| Number            | 1         | 5    | 23   | 1    | 0    | 23         | 6    | 1    |      |      | 30             |
| 1975              | 1.1       | 1.2  | 1.3  | 2.1  | 2.2  | 1970       | 1971 | 1972 |      |      |                |
| Percent           | 6.3       | 41.8 | 37.9 | 0.0  | 14.0 | 51.9       | 41.8 | 6.3  |      |      | 100.0          |
| Number            | 5         | 33   | 30   | 0    | 11   | 41         | 30   | 5    |      |      | 79             |
| 1974              | 1.1       | 1.2  | 1.3  | 2.1  | 2.2  | 1969       | 1970 | 1971 |      |      |                |
| Percent           | 27.9      | 46.5 | 4.7  | 11.6 | 9.3  | 14.0       | 58.1 | 27.9 |      |      | 100.0          |
| Number            | 12        | 20   | 2    | 5    | 4    | 6          | 25   | 12   |      |      | 43             |
| <hr/>             |           |      |      |      |      |            |      |      |      |      |                |
| <u>COHO:</u>      |           |      |      |      |      |            |      |      |      |      |                |
| 1977              | 1.1       | 1.2  | 2.0  | 2.1  |      | 1973       | 1974 |      |      |      |                |
| Percent           | 14.3      | 0.0  | 0.0  | 85.7 |      | 85.7       | 14.3 |      |      |      | 100.0          |
| Number            | 1         | 0    | 0    | 6    |      | 6          | 1    |      |      |      | 7              |
| 1975              | 1.1       | 1.2  | 2.0  | 2.1  |      | 1971       | 1972 |      |      |      |                |
| Percent           | 11.8      | 5.9  | 0.0  | 82.3 |      | 88.2       | 11.8 |      |      |      | 100.0          |
| Number            | 2         | 1    | 0    | 14   |      | 15         | 2    |      |      |      | 17             |
| 1974              | 1.1       | 1.2  | 2.0  | 2.1  |      | 1970       | 1971 |      |      |      |                |
| Percent           | 15.9      | 0.0  | 0.9  | 83.2 |      | 84.1       | 15.9 |      |      |      | 100.0          |
| Number            | 18        | 0    | 1    | 94   |      | 95         | 18   |      |      |      | 113            |
| <hr/>             |           |      |      |      |      |            |      |      |      |      |                |
| <u>CHUM:</u>      |           |      |      |      |      |            |      |      |      |      |                |
| 1977              | 0.2       | 0.3  | 0.4  |      |      | 1972       | 1973 | 1974 |      |      |                |
| Percent           | 4.8       | 88.1 | 7.1  |      |      | 7.1        | 88.1 | 4.8  |      |      | 100.0          |
| Number            | 2         | 37   | 3    |      |      | 2          | 37   | 2    |      |      | 42             |
| 1975              | 0.2       | 0.3  | 0.4  |      |      | 1970       | 1971 | 1972 |      |      |                |
| Percent           | 16.4      | 82.0 | 1.6  |      |      | 1.6        | 82.0 | 16.4 |      |      | 100.0          |
| Number            | 21        | 105  | 2    |      |      | 2          | 105  | 21   |      |      | 128            |
| 1974              | 0.2       | 0.3  | 0.4  |      |      | 1969       | 1970 | 1971 |      |      |                |
| Percent           | 48.1      | 33.4 | 18.5 |      |      | 18.5       | 33.4 | 48.1 |      |      | 100.0          |
| Number            | 229       | 159  | 88   |      |      | 88         | 159  | 229  |      |      | 476            |

## Appendix I

Table 2. Age, length, and sex characteristics of chum, chinook, sockeye, and coho salmon escapement samples, Devils Canyon Project, 1974, 1975, and 1977.

| Year of Return  | Age Class | Mean Length (mm) | Standard Deviation (s.) | Range of Lengths | Number Males | Number Females | n   |
|-----------------|-----------|------------------|-------------------------|------------------|--------------|----------------|-----|
| <u>CHUM:</u>    |           |                  |                         |                  |              |                |     |
| 1974            | 0.2       | 545.0            | 32.05                   | 410-650          | 155          | 74             | 229 |
|                 | 0.3       | 614.8            | 33.61                   | 510-695          | 88           | 71             | 159 |
|                 | 0.4       | 627.6            | 30.71                   | 520-695          | 47           | 41             | 88  |
| 1975            | 0.2       | 552.7            | 13.58                   | 530-578          | 11           | 10             | 21  |
|                 | 0.3       | 587.6            | 20.62                   | 532-628          | 55           | 50             | 105 |
|                 | 0.4       | 620.5            | 2.50                    | 618-623          | 0            | 2              | 2   |
| 1977            | 0.2       | 568.5            | 3.50                    | 565-572          | 0            | 2              | 2   |
|                 | 0.3       | 618.3            | 29.05                   | 545-667          | 28           | 9              | 37  |
|                 | 0.4       | 656.7            | 9.43                    | 650-670          | 2            | 1              | 3   |
| <u>CHINOOK:</u> |           |                  |                         |                  |              |                |     |
| 1975            | 1.1       | 389.3            | 31.69                   | 341-421          | 4            | 0              | 4   |
|                 | 1.2       | 483.5            | 6.50                    | 477-490          | 1            | 1              | 2   |
|                 | 1.3       | 710.6            | 84.25                   | 569-812          | 12           | 3              | 15  |
|                 | 1.4       | 856.2            | 62.63                   | 778-990          | 7            | 12             | 19  |
|                 | 1.5       | 937.0            | 45.08                   | 897-1000         | 0            | 3              | 3   |
| 1977            | 1.1       | 371.5            | 28.50                   | 343-400          | 2            | 0              | 2   |
|                 | 1.2       | 580.0            | 5.00                    | 575-580          | 2            | 0              | 2   |
|                 | 1.3       | 816.3            | 59.10                   | 725-920          | 8            | 3              | 11  |
|                 | 1.4       | 994.8            | 52.02                   | 950-1103         | 4            | 2              | 6   |
| <u>SOCKEYE:</u> |           |                  |                         |                  |              |                |     |
| 1974            | 1.1       | 395.5            | 69.14                   | 315-485          | 12           | 0              | 12  |
|                 | 1.2       | 527.8            | 48.99                   | 417-595          | 10           | 10             | 20  |
|                 | 1.3       | 572.5            | 12.50                   | 560-585          | 0            | 2              | 2   |
|                 | 2.1       | 376.6            | 56.94                   | 318-485          | 5            | 0              | 5   |
|                 | 2.2       | 536.3            | 20.12                   | 515-565          | 3            | 1              | 4   |
|                 | 2.2       | 536.3            | 20.12                   | 515-565          | 3            | 1              | 4   |
| 1975            | 1.1       | 352.4            | 37.15                   | 313-423          | 5            | 0              | 5   |
|                 | 1.2       | 471.8            | 42.36                   | 398-548          | 15           | 18             | 33  |
|                 | 1.3       | 576.1            | 26.65                   | 514-638          | 12           | 18             | 30  |
|                 | 2.1       | —                | —                       | —                | 0            | 0              | 0   |
|                 | 2.2       | 532.3            | 39.54                   | 460-576          | 4            | 7              | 11  |
| 1977            | 1.1       | 347.0            | —                       | 347              | 1            | 0              | 1   |
|                 | 1.2       | 451.8            | 27.09                   | 433-505          | 4            | 1              | 5   |
|                 | 1.3       | 596.4            | 30.24                   | 509-639          | 11           | 12             | 23  |
|                 | 2.1       | 371.0            | —                       | 371              | 1            | 0              | 1   |
|                 | 2.2       | —                | —                       | —                | 0            | 0              | 0   |
| <u>COHO:</u>    |           |                  |                         |                  |              |                |     |
| 1974            | 1.1       | 487.9            | 42.92                   | 410-575          | 11           | 7              | 18  |
|                 | 2.0       | 375.0            | —                       | 375              | 1            | 0              | 1   |
|                 | 2.1       | 527.7            | 48.00                   | 376-605          | 49           | 45             | 94  |
| 1975            | 1.1       | 495.5            | 4.50                    | 491-500          | 1            | 1              | 2   |
|                 | 1.2       | 540.0            | —                       | 540              | 1            | 0              | 1   |
|                 | 2.1       | 531.1            | 38.53                   | 454-608          | 5            | 9              | 14  |
| 1977            | 1.1       | 337.0            | —                       | 337              | 1            | 0              | 1   |
|                 | 2.1       | 473.0            | 54.54                   | 400-549          | 5            | 1              | 6   |

## Appendix I

Table 3. Analyses of age, length, weight and condition factors of juvenile sockeye salmon samples from Susitna River sloughs and clearwater tributaries, Devils Canyon Project, 1977.

| Location  | Age Class | Date | Length (mm) |                    |       | Weight (g) |                    |         | Condition Factor |                    |             | Percent 1/<br>Composition | n  |
|-----------|-----------|------|-------------|--------------------|-------|------------|--------------------|---------|------------------|--------------------|-------------|---------------------------|----|
|           |           |      | Mean        | Standard Deviation | Range | Mean       | Standard Deviation | Range   | Mean             | Standard Deviation | Range       |                           |    |
| Slough 1  | 0.0       | 8/5  | 48.0        | 1.0                | 47-49 | 1.5        | 0.2                | 1.3-1.8 | 1.391            | 0.139              | 1.252-1.530 | 100                       | 2  |
| Slough 3  | 0.0       | 9/24 | 53.0        | -                  | -     | 1.7        | -                  | -       | 1.142            | -                  | -           | 100                       | 1  |
| Slough 5  | 0.0       | 7/27 | 44.0        | -                  | -     | 1.7        | -                  | -       | 1.996            | -                  | -           | 100                       | 1  |
| Slough 6  | 0.0       | 7/27 | 34.3        | 2.9                | 31-38 | 0.4        | 0.0                | 0.4     | 1.029            | 0.250              | 0.729-1.342 | 100                       | 3  |
| Slough 8  | 0.0       | 7/27 | 42.7        | 1.7                | 40-45 | 0.7        | 0.1                | 0.5-0.9 | 0.945            | 0.095              | 0.781-1.006 | 100                       | 4  |
|           | 0.0       | 9/23 | 44.5        | 5.5                | 39-50 | 1.1        | 0.4                | 0.6-1.5 | 1.105            | 0.094              | 1.011-1.200 | 100                       | 2  |
| Slough 8B | 0.0       | 9/23 | 48.9        | 5.3                | 44-60 | 1.3        | 0.4                | 0.9-2.4 | 1.091            | 0.064              | 0.986-1.207 | 100                       | 12 |
| Slough 11 | 0.0       | 9/21 | 51.5        | 0.5                | 51-52 | 1.7        | 0.1                | 1.6-1.7 | 1.207            | 0.001              | 1.206-1.209 | 100                       | 2  |
| Slough 12 | 0.0       | 8/2  | 36.9        | 4.6                | 29-45 | 0.7        | 0.3                | 0.3-1.2 | 1.300            | 0.277              | 0.932-1.399 | 100                       | 10 |
| Slough 17 | 0.0       | 7/26 | 44.0        | -                  | -     | 0.5        | -                  | -       | 0.988            | -                  | -           | 100                       | 1  |



Appendix I

Table 3. Analyses of age, length, weight and condition factors of juvenile sockeye salmon samples from Susitna River sloughs and clearwater tributaries, Devils Canyon Project, 1977. (continued)

| Location  | Age Class | Date | Length (mm) |                    |       | Weight (g) |                    |         | Condition Factor |                    |             | Percent <sup>1/</sup> Composition | n  |
|-----------|-----------|------|-------------|--------------------|-------|------------|--------------------|---------|------------------|--------------------|-------------|-----------------------------------|----|
|           |           |      | Mean        | Standard Deviation | Range | Mean       | Standard Deviation | Range   | Mean             | Standard Deviation | Range       |                                   |    |
| Slough 19 | 0.0       | 7/26 | 32.4        | 7.9                | 25-51 | 0.4        | 0.5                | 0.1-1.9 | 0.803            | 0.335              | 0.370-1.432 | 100                               | 12 |
|           | 0.0       | 8/2  | 53.5        | 1.5                | 52-55 | 1.5        | 0.1                | 1.4-1.7 | 1.009            | 0.013              | 0.996-1.022 | 100                               | 2  |
|           | 0.0       | 9/21 | 50.0        | -                  | -     | 1.5        | -                  | -       | 1.200            | -                  | -           | 100                               | 1  |

<sup>1/</sup> Percent composition of each age class within sampling period.

## Appendix I

Table 4. Analyses of age, length, weight and condition factors of juvenile coho salmon samples from Susitna River sloughs and clearwater tributaries, Devils Canyon Project, 1977.

| Location        | Age Class | Date | Length (mm) |                       |        | Weight (g) |                       |           | Condition Factor |                       |             | Percent 1/<br>Composition | n  |
|-----------------|-----------|------|-------------|-----------------------|--------|------------|-----------------------|-----------|------------------|-----------------------|-------------|---------------------------|----|
|                 |           |      | Mean        | Standard<br>Deviation | Range  | Mean       | Standard<br>Deviation | Range     | Mean             | Standard<br>Deviation | Range       |                           |    |
| Slough 1        | 0.0       | 9/24 | 54.6        | 3.9                   | 49-61  | 2.0        | 0.5                   | 1.3-2.9   | 1.199            | 0.084                 | 1.022-1.315 | 93                        | 13 |
|                 | 1.0       | 9/24 | 80.0        | -                     | -      | 6.0        | -                     | -         | 1.172            | -                     | -           | 7                         | 1  |
| Slough 4        | 0.0       | 9/24 | 62.7        | 3.3                   | 59-67  | 2.9        | 0.4                   | 2.5-3.6   | 1.196            | 0.017                 | 1.175-1.217 | 23                        | 3  |
|                 | 1.0       | 9/24 | 75.4        | 8.3                   | 68-99  | 5.9        | 3.3                   | 3.7-15.8  | 1.268            | 0.126                 | 1.152-1.628 | 77                        | 10 |
| Slough 5        | 0.0       | 9/23 | 77.0        | -                     | -      | 6.2        | -                     | -         | 1.358            | -                     | -           | 25                        | 1  |
|                 | 1.0       | 9/23 | 105.3       | 8.9                   | 93-114 | 14.9       | 3.0                   | 10.7-17.5 | 1.267            | 0.104                 | 1.120-1.351 | 75                        | 3  |
| Slough 6        | 0.0       | 7/27 | 57.0        | -                     | -      | 1.9        | -                     | -         | 1.026            | -                     | -           | 100                       | 1  |
| Slough 6A       | 0.0       | 7/27 | 49.5        | 1.5                   | 48-51  | 1.5        | 0.0                   | 1.5       | 1.243            | 0.113                 | 1.113-1.356 | 100                       | 2  |
| Slough<br>or 8A | 0.0       | 9/23 | 63.0        | -                     | -      | 3.0        | -                     | -         | 1.216            | -                     | -           | 100                       | 1  |
| Slough 8C       | 0.0       | 9/23 | 47.0        | -                     | -      | 1.2        | -                     | -         | 1.156            | -                     | -           | 100                       | 1  |
| Slough 10       | 0.0       | 7/27 | 57.0        | -                     | -      | 2.1        | -                     | -         | 1.134            | -                     | -           | 100                       | 1  |

## Appendix I

Table 4. Analyses of age, length, weight and condition factors of juvenile coho salmon samples from Susitna River sloughs and clearwater tributaries, Devils Canyon Project, 1977. (continued)

| Location       | Age Class | Date | Length (mm) |                    |       | Weight (g) |                    |         | Condition Factor |                    |             | Percent <sup>1</sup> /<br>Composition | n |
|----------------|-----------|------|-------------|--------------------|-------|------------|--------------------|---------|------------------|--------------------|-------------|---------------------------------------|---|
|                |           |      | Mean        | Standard Deviation | Range | Mean       | Standard Deviation | Range   | Mean             | Standard Deviation | Range       |                                       |   |
| Slough 13      | 0.0       | 9/22 | 59.0        | -                  | -     | 2.2        | -                  | -       | 1.071            | -                  | -           | 100                                   | 1 |
| Slough 16      | 0.0       | 9/21 | 63.0        | 2.0                | 61-65 | 3.2        | 0.3                | 2.9-3.5 | 1.276            | 0.002              | 1.274-1.278 | 100                                   | 2 |
| Slough 19      | 0.0       | 9/21 | 71.0        | 2.0                | 69-73 | 4.7        | 0.7                | 4.0-5.3 | 1.290            | 0.072              | 1.218-1.362 | 100                                   | 2 |
| Slough 21      | 0.0       | 9/20 | 56.0        | -                  | -     | 1.5        | -                  | -       | 0.854            | -                  | -           | 100                                   | 1 |
| Chase Creek    | 0.0       | 8/6  | 43.0        | 2.0                | 41-45 | 0.9        | 0.1                | 0.8-1.1 | 1.184            | 0.023              | 1.161-1.207 | 100                                   | 2 |
| Whiskers Creek | 0.0       | 8/5  | 43.0        | 5.0                | 38-48 | 0.9        | 0.3                | 0.6-1.2 | 1.089            | 0.004              | 1.085-1.093 | 100                                   | 2 |
|                | 0.0       | 9/24 | 50.7        | 4.3                | 46-57 | 1.7        | 0.4                | 1.1-2.2 | 1.243            | 0.837              | 1.130-1.356 | 100                                   | 6 |

<sup>1</sup>/ Percent composition of each age class within sampling period.

## Appendix I

Table 5. Analyses of age, length, weight and condition factors of juvenile chinook salmon samples from Susitna River sloughs and clearwater tributaries, Devils Canyon Project, 1977.

| Location  | Age Class | Date | Length (mm) |                    |       | Weight (g) |                    |         | Condition Factor |                    |             | Percent/Composition | n  |
|-----------|-----------|------|-------------|--------------------|-------|------------|--------------------|---------|------------------|--------------------|-------------|---------------------|----|
|           |           |      | Mean        | Standard Deviation | Range | Mean       | Standard Deviation | Range   | Mean             | Standard Deviation | Range       |                     |    |
| Slough 1  | 0.0       | -    | -           | -                  | -     | -          | -                  | -       | -                | -                  | -           | -                   | -  |
| Slough 2  | 0.0       | 8/5  | 49.1        | 4.5                | 42-55 | 1.5        | 0.4                | 0.7-1.7 | 1.152            | 0.127              | 0.945-1.282 | 100                 | 6  |
| Slough 3  | 0.0       | 8/5  | 51.0        | 7.1                | 45-68 | 1.7        | 0.7                | 1.1-3.4 | 1.233            | 0.074              | 1.081-1.348 | 100                 | 10 |
|           | 0.0       | 9/24 | 62.8        | 4.3                | 58-69 | 2.8        | 0.7                | 1.8-3.9 | 1.107            | 0.118              | 0.916-1.230 | 100                 | 6  |
| Slough 4  | 0.0       | 9/24 | 59.0        | -                  | -     | 2.4        | -                  | -       | 1.169            | -                  | -           | 100                 | 1  |
| Slough 5  | 0.0       | 7/27 | 55.3        | 3.7                | 51-60 | 1.7        | 0.3                | 1.2-2.0 | 1.011            | 0.135              | 0.904-1.202 | 100                 | 3  |
| Slough 6  | 0.0       | 7/27 | 48.3        | 6.6                | 42-63 | 1.3        | 0.6                | 0.7-2.7 | 1.049            | 0.108              | 0.904-1.026 | 100                 | 7  |
| Slough 6A | 0.0       | 7/27 | 50.7        | 1.7                | 49-54 | 1.2        | 0.1                | 1.0-1.5 | 0.935            | 0.065              | 0.850-1.067 | 88                  | 7  |
|           | 1.0       | 7/27 | 76.0        | -                  | -     | 4.3        | -                  | -       | 0.980            | -                  | -           | 12                  | 1  |
| Slough 8  | 0.0       | 7/27 | 46.2        | 3.9                | 43-53 | 0.9        | 0.2                | 0.7-1.3 | 0.854            | 0.029              | 0.814-0.880 | 100                 | 5  |
|           | 0.0       | 9/23 | 52.0        | 5.7                | 45-60 | 1.7        | 0.6                | 1.0-2.7 | 1.183            | 0.102              | 1.027-1.317 | 100                 | 10 |

## Appendix I

Table 5.

Analyses of age, length, weight and condition factors of juvenile chinook salmon samples from Susitna River sloughs and clearwater tributaries, Devils Canyon Project, 1977. (continued)

| Location  | Age Class | Date | Length (mm) |                    |       | Weight (g) |                    |           | Condition Factor |                    |             | Percent 1/<br>Composition | n  |
|-----------|-----------|------|-------------|--------------------|-------|------------|--------------------|-----------|------------------|--------------------|-------------|---------------------------|----|
|           |           |      | Mean        | Standard Deviation | Range | Mean       | Standard Deviation | Range     | Mean             | Standard Deviation | Range       |                           |    |
| Slough 8A | 0.0       | 8/3  | 54.5        | 1.5                | 53-56 | 2.1        | 0.1                | 1.9-2.2   | 1.264            | 0.011              | 1.253-1.276 | 100                       | 2  |
|           | 0.0       | 9/23 | 64.0        | 2.0                | 62-66 | 3.3        | 0.3                | 3.05-3.70 | 1.283            | 0.003              | 1.280-1.287 | 100                       | 2  |
| Slough 8B | 0.0       | 8/4  | 49.2        | 5.4                | 44-59 | 1.3        | 0.5                | 0.8-2.4   | 1.008            | 0.088              | 0.924-1.169 | 100                       | 5  |
|           | 0.0       | 9/23 | 61.0        | 4.2                | 55-64 | 3.0        | 0.8                | 1.9-3.8   | 1.284            | 0.127              | 1.142-1.450 | 100                       | 3  |
| Slough 8C | 0.0       | 8/4  | 47.6        | 2.2                | 44-51 | 1.1        | 0.1                | 0.8-1.4   | 1.026            | 0.053              | 0.939-1.085 | 100                       | 5  |
|           | 0.0       | 9/23 | 52.7        | 4.0                | 47-57 | 1.8        | 0.3                | 1.2-2.3   | 1.181            | 0.059              | 1.080-1.242 | 100                       | 6  |
| Slough 8D | 0.0       | 9/23 | 56.0        | -                  | -     | 2.1        | -                  | -         | 1.196            | -                  | -           | 100                       | 1  |
| Slough 9  | 0.0       | 7/12 | 49.8        | 4.9                | 44-59 | 1.5        | 0.5                | 0.9-2.5   | 1.185            | 0.082              | 1.056-1.336 | 100                       | 10 |
|           | 0.0       | 7/27 | 50.8        | 3.1                | 46-57 | 1.0        | 0.3                | 0.5-1.7   | 0.725            | 0.199              | 0.427-1.020 | 100                       | 11 |
|           | 0.0       | 9/22 | 56.3        | 6.3                | 46-64 | 2.1        | 0.7                | 1.1-3.1   | 1.154            | 0.400              | 1.080-1.209 | 100                       | 10 |
| Slough 10 | 0.0       | 7/27 | 48.5        | 3.4                | 44-51 | 1.0        | 0.3                | 0.5-1.6   | 0.862            | 0.128              | 0.587-1.055 | 100                       | 9  |

## Appendix I

Table 5. Analyses of age, length, weight and condition factors of juvenile chinook salmon samples from Susitna River sloughs and clearwater tributaries, Devils Canyon Project, 1977. (continued)

| Location  | Age Class | Date | Length (mm) |                    |       | Weight (g) |                    |         | Condition Factor |                    |             | Percent1/<br>Composition | n  |
|-----------|-----------|------|-------------|--------------------|-------|------------|--------------------|---------|------------------|--------------------|-------------|--------------------------|----|
|           |           |      | Mean        | Standard Deviation | Range | Mean       | Standard Deviation | Range   | Mean             | Standard Deviation | Range       |                          |    |
| Slough 11 | 0.0       | 9/21 | 60.3        | 3.5                | 54-69 | 2.5        | 0.5                | 1.3-4.0 | 1.166            | 0.104              | 0.972-1.366 | 100                      | 40 |
| Slough 13 | 0.0       | 9/22 | 56.0        | 3.5                | 53-62 | 2.1        | 0.5                | 1.9-3.0 | 1.221            | 0.052              | 1.142-1.276 | 100                      | 4  |
| Slough 14 | 0.0       | 9/22 | 60.7        | 4.1                | 54-68 | 2.8        | 0.5                | 2.0-3.8 | 1.233            | 0.040              | 1.165-1.296 | 90                       | 9  |
|           | 1.0       | 9/22 | 74.0        | -                  | -     | 5.1        | -                  | -       | 1.259            | -                  | -           | 10                       | 1  |
| Slough 15 | 0.0       | 7/26 | 48.5        | 2.1                | 45-52 | 1.2        | 0.1                | 1.0-1.6 | 1.048            | 0.080              | 0.924-1.175 | 100                      | 10 |
|           | 0.0       | 9/21 | 60.8        | 6.3                | 48-74 | 2.9        | 0.9                | 1.8-4.8 | 1.260            | 0.137              | 0.926-1.628 | 100                      | 19 |
| Slough 16 | 0.0       | 7/26 | 51.7        | 3.1                | 46-58 | 1.5        | 0.3                | 1.0-2.3 | 1.092            | 0.080              | 0.962-1.242 | 100                      | 20 |
|           | 0.0       | 9/21 | 54.8        | 4.4                | 47-63 | 2.1        | 0.5                | 1.4-2.8 | 1.268            | 0.102              | 1.075-1.461 | 93                       | 13 |
|           | 1.0       | 9/21 | 73.0        | -                  | -     | 4.6        | -                  | -       | 1.182            | -                  | -           | 7                        | 1  |
| Slough 17 | 0.0       | 7/11 | 47.9        | 1.0                | 46-50 | 1.3        | 0.1                | 1.2-1.4 | 1.208            | 0.069              | 1.085-1.266 | 100                      | 10 |
|           | 0.0       | 7/26 | 46.1        | 2.6                | 40-50 | 0.9        | 0.1                | 0.7-1.1 | 0.916            | 0.239              | 0.719-1.563 | 100                      | 9  |

## Appendix I

Table 5. Analyses of age, length, weight and condition factors of juvenile chinook salmon samples from Susitna River sloughs and clearwater tributaries, Devils Canyon Project, 1977. (continued)

| Location             | Age Class | Date | Length (mm) |                    |       | Weight (g) |                    |         | Condition Factor |                    |             | Percent 1/<br>Composition | n  |
|----------------------|-----------|------|-------------|--------------------|-------|------------|--------------------|---------|------------------|--------------------|-------------|---------------------------|----|
|                      |           |      | Mean        | Standard Deviation | Range | Mean       | Standard Deviation | Range   | Mean             | Standard Deviation | Range       |                           |    |
| Slough 18            | 0.0       | 7/26 | 50.0        | 3.5                | 46-52 | 1.3        | 0.3                | 1.0-2.2 | 1.079            | 0.065              | 0.963-1.175 | 100                       | 10 |
|                      | 0.0       | 9/21 | 61.7        | 4.5                | 58-69 | 3.1        | 1.0                | 2.3-4.6 | 1.286            | 0.126              | 1.179-1.463 | 100                       | 3  |
| Slough 19            | 0.0       | 8/2  | 60.5        | 3.5                | 57-64 | 2.1        | 0.2                | 1.9-2.4 | 0.970            | 0.055              | 0.915-1.026 | 100                       | 2  |
|                      | 0.0       | 9/21 | 60.3        | 3.7                | 52-65 | 2.7        | 0.4                | 1.7-3.2 | 1.206            | 0.084              | 1.111-1.412 | 100                       | 8  |
| Slough 20            | 0.0       | 7/25 | 54.2        | 4.4                | 46-64 | 1.7        | 0.5                | 0.8-2.8 | 1.048            | 0.128              | 0.822-1.207 | 100                       | 20 |
|                      | 0.0       | 9/20 | 60.7        | 3.9                | 51-68 | 2.7        | 0.5                | 1.5-3.2 | 1.211            | 0.063              | 1.080-1.343 | 100                       | 19 |
| Slough 21            | 0.0       | 7/13 | 45.0        | 1.6                | 43-47 | 1.2        | 0.2                | 1.0-1.5 | 1.340            | 0.078              | 1.258-1.445 | 100                       | 3  |
|                      | 0.0       | 9/20 | 58.9        | 2.5                | 57-63 | 2.3        | 0.3                | 1.9-3.0 | 1.139            | 0.075              | 1.019-1.296 | 100                       | 14 |
| Chase Creek          | 0.0       | 8/6  | 48.7        | 4.1                | 42-54 | 1.3        | 0.3                | 0.8-2.2 | 1.174            | 0.069              | 1.080-1.266 | 100                       | 6  |
| Fourth of July Creek | 0.0       | 8/3  | 49.7        | 4.3                | 40-57 | 1.3        | 0.3                | 0.7-1.8 | 1.009            | 0.076              | 0.873-1.138 | 100                       | 13 |
|                      | 0.0       | 9/22 | 63.0        | 3.0                | 59-68 | 3.2        | 0.3                | 2.9-3.6 | 1.297            | 0.061              | 1.240-1.412 | 100                       | 6  |

## Appendix I

Table 5. Analyses of age, length, weight and condition factors of juvenile chinook salmon samples from Susitna River sloughs and clearwater tributaries, Devils Canyon Project, 1977. (continued)

| Location       | Age Class | Date | Length (mm) |                    |       | Weight (g) |                    |         | Condition Factor |                    |             | Percent <sup>1/</sup><br>Composition | n  |
|----------------|-----------|------|-------------|--------------------|-------|------------|--------------------|---------|------------------|--------------------|-------------|--------------------------------------|----|
|                |           |      | Mean        | Standard Deviation | Range | Mean       | Standard Deviation | Range   | Mean             | Standard Deviation | Range       |                                      |    |
| McKenzie Creek | 0.0       | 7/27 | 47.6        | 4.8                | 39-59 | 1.1        | 0.4                | 0.7-2.1 | 1.012            | 0.085              | 0.822-1.142 | 100                                  | 24 |
| Whiskers Creek | 0.0       | 8/5  | 45.0        | 4.0                | 41-49 | 1.1        | 0.3                | 0.8-1.4 | 1.175            | 0.014              | 1.161-1.190 | 100                                  | 2  |
|                | 0.0       | 9/24 | 53.0        | 3.7                | 49-59 | 1.9        | 0.3                | 1.5-2.5 | 1.246            | 0.033              | 1.209-1.282 | 100                                  | 4  |

<sup>1/</sup> Percent composition of each age class within sampling period.



Appendix I Table 6. Stomach content analysis of juvenile chinook, coho, and sockeye salmon collected in sloughs and clearwater tributaries of the Susitna River during summer and fall, Devils Canyon Project, 1977.

| Location              | Date | Number<br>Specimens | Estimated Percent of Combined Gut Contents |                     |                  |       | Predominate Organisms                               |
|-----------------------|------|---------------------|--|---------------------|------------------|-------|---|
|                       |      |                     | Crustacea                                  | Immature<br>Insecta | Adult<br>Insecta | Other |   |
| <u>Chinook-Summer</u> |      |                     |  |                     |                  |       |   |
| Susitna<br>#1         | 8/5  | 7                   | 0  | 10                  | 90               | 0     | Adult Diptera and Hymenoptera                       |
| Susitna<br>#2         | 8/5  | 6                   | <5   | 20                  | 75               | 0     | Adult Diptera; Diptera Chironomid larvae            |
| Susitna<br>#3         | 8/5  | 10                  | <5   | 80                  | >14              | <1    | Diptera Chironomid larvae and pupae                 |
| Susitna<br>#5         | 7/27 | 4                   | <1   | 10                  | >89              | 0     | Adult Diptera                                       |
| Susitna<br>#6         | 7/27 | 7                   | 50   | <5                  | >45              | 0     | Ostracoda and Calanoid Copepoda;<br>Adult Diptera   |
| Susitna<br>#6A        | 7/27 | 8                   | 0  | 4                   | >95              | <1    | Adult Diptera                                       |
| Slough<br>#8          | 7/27 | 5                   | 0  | 0                   | 100              | 0     | Adult Diptera                                       |
| Slough<br>#8A         | 8/3  | 2                   | 0  | 0                   | 0                | 0     | Guts empty  |
| Susitna<br>#8B        | 8/4  | 5                   | 0  | 5                   | 75               | 20    | Adult Diptera; unidentified capsules (plant seeds?) |
| Susitna<br>#8C        | 8/4  | 5                   | 0  | <5                  | >94              | <1    | Adult Diptera and Hymenoptera                       |

Appendix I Table 6. Stomach content analysis of juvenile chinook, coho, and sockeye salmon collected in sloughs and clearwater tributaries of the Susitna River during summer and fall, Devils Canyon Project, 1977, (continued).

| Location    | Date | Number Specimens | Estimated Percent of Combined Gut Contents |                  |               |       | Predominate Organisms   |
|-------------|------|------------------|--|------------------|---------------|-------|---|
|             |      |                  | Crustacea                                  | Immature Insecta | Adult Insecta | Other |   |
| Susitna #9  | 7/14 | 10               | 0  | <1               | >80           | 19    | Adult Diptera and Hymenoptera; unidentified structures (plant seeds?) |
| Susitna #9A | 7/27 | 11               | 0  | 5                | 95            | 0     | Adult Diptera   |
| Susitna #10 | 7/27 | 8                | 20   | 10               | 70            | 0     | Unidentified adult insect fragments; Cladocera Bosminidae             |
| Susitna #15 | 7/26 | 10               | <1   | 10               | >89           | 0     | Adult Diptera   |
| Susitna #16 | 7/26 | 20               | 0  | 70               | 30            | 0     | Diptera Chironomid larvae; Adult Diptera                              |
| Susitna #17 | 7/14 | 10               | 0  | 10               | 90            | 0     | Adult Diptera   |
| Susitna #17 | 7/26 | 9                | 0  | 40               | 60            | 0     | Adult Diptera and Homoptera; Diptera Chironomid larvae                |
| Susitna #18 | 7/26 | 10               | 20   | 20               | 60            | 0     | Adult Homoptera and Hymenoptera; Ostracoda                            |
| Susitna #19 | 8/2  | 2                | 0  | 50               | 50            | 0     | Diptera Chironomid larvae and pupae; Adult Diptera                    |
| Susitna #20 | 7/25 | 20               | 0  | <5               | >95           | 0     | Adult Diptera and Coleoptera  |

Appendix I Table 6. Stomach content analysis of juvenile chinook, coho, and sockeye salmon collected in sloughs and clearwater tributaries of the Susitna River during summer and fall, Devils Canyon Project, 1977, (continued).

| Location             | Date | Number Specimens | Estimated Percent of Combined Gut Contents |                  |               |       | Predominate Organisms   |
|----------------------|------|------------------|--|------------------|---------------|-------|---|
|                      |      |                  | Crustacea                                  | Immature Insecta | Adult Insecta | Other |   |
| Susitna #21          | 7/13 | 3                | 0  | 10               | 90            | 0     | Adult Diptera   |
| Whiskers Creek       | 8/5  | 2                | 10   | <5               | >85           | 0     | Adult Diptera   |
| Chase Creek          | 8/6  | 11               | 1  | 4                | 95            | 0     | Adult Homoptera and Hymenoptera   |
| McKenzie Creek       | 7/27 | 21               | 0  | 40               | 60            | 0     | Adult Diptera; Diptera Chironomid larvae and pupae  |
| Fourth of July Creek | 8/3  | 13               | 0  | 40               | 60            | 0     | Adult Diptera; adult Chironomid larvae and pupae  |
| <u>Chinook-Fall</u>  |      |                  |  |                  |               |       |   |
| Susitna #3           | 9/24 | 6                | 0  | 20               | 80            | 0     | Adult Hemiptera and unidentified adult Insecta; Diptera Chironomid larvae                                       |
| Susitna #4           | 9/24 | 1                | 0  | 10               | >85           | <5    | Unidentified adult insect fragments   |
| Slough #A            | 9/23 | 2                | 0  | 5                | 35            | 60    | Oligochaeta (?); Unidentified adult insect fragments  |
| Susitna #8           | 9/23 | 10               | 0  | >45              | 50            | <5    | Adult Diptera and Hymenoptera; Diptera Chironomid pupae and larvae; Trichoptera pupae; Diptera Tepulidae larvae |

Appendix I Table 6. Stomach content analysis of juvenile chinook, coho, and sockeye salmon collected in sloughs and clearwater tributaries of the Susitna River during summer and fall, Devils Canyon Project, 1977, (continued).

| Location    | Date | Number Specimens | Estimated Percent of Combined Gut Contents |                  |               |       | Predominate Organisms  |
|-------------|------|------------------|--|------------------|---------------|-------|--|
|             |      |                  | Crustacea                                  | Immature Insecta | Adult Insecta | Other |  |
| Susitna #8B | 9/23 | 3                | 0  | 10               | 90            | 0     | Adult Diptera, Hymenoptera and Lepidoptera   |
| Susitna #8C | 9/23 | 6                | 0  | 30               | 40            | 30    | Adult Homoptera and unidentified adult insect fragment; Oligochaeta (?); Diptera Chironomid larvae and pupae |
| Susitna #8D | 9/23 | 1                | 0  | 30               | 70            | 0     | Adult insect fragments; Diptera Chironomid larvae  |
| Susitna #9  | 9/22 | 10               | 0  | <5               | >95           | 0     | Adult Diptera, Hymenoptera, Homoptera and Lepidoptera  |
| Susitna #11 | 9/21 | 20               | 0  | 70               | 20            | 10    | Trichoptera and Diptera Chironomid pupae; adult Hemiptera and unidentified adult fragments                   |
| Susitna #13 | 9/22 | 4                | 0  | 30               | 70            | 0     | Adult Diptera and unidentified adult fragments; Diptera Chironomid larvae and pupae                          |
| Susitna #14 | 9/22 | 10               | 0  | 9                | 90            | 1     | Adult Diptera, Hymenoptera, Plecoptera   |
| Susitna #15 | 9/21 | 19               | 0  | 10               | 30            | 60    | Oligochaeta (?); Adult Diptera and Hemiptera   |
| Susitna #16 | 9/21 | 14               | 0  | 10               | 85            | 5     | Adult Diptera and Hemiptera  |
| Susitna #19 | 9/21 | 8                | 1  | 14               | 85            | 0     | Diptera Chironomid pupae   |

Appendix I Table 6. Stomach content analysis of juvenile chinook, coho, and sockeye salmon collected in sloughs and clearwater tributaries of the Susitna River during summer and fall, Devils Canyon Project, 1977, (continued).

| Location             | Date | Number Specimens | Estimated Percent of Combined Gut Contents |                  |               |       | Predominate Organisms  |
|----------------------|------|------------------|--|------------------|---------------|-------|--|
|                      |      |                  | Crustacea                                  | Immature Insecta | Adult Insecta | Other |  |
| Susitna #20          | 9/20 | 19               | 0  | 30               | 65            | 5     | Adult Hemiptera Diptera and Hymenoptera fragments; Diptera Chironomid larvae |
| Susitna #21          | 9/20 | 14               | 0  | <5               | >95           | 0     | Adult Diptera  |
| Whiskers Creek       | 9/24 | 4                | 0  | >95              | <5            | 0     | Trichoptera pupae  |
| Fourth of July Creek | 9/22 | 7                | 0  | 15               | 75            | 10    | Adult Diptera, Hemoptera, and Hymenoptera                                    |
| <u>Coho-Summer</u>   |      |                  |  |                  |               |       |  |
| Susitna #1           | 8/5  | 5                | 10   | 10               | 80            | 0     | Adult Lepidoptera and unidentified adult insect fragments                    |
| Susitna #6           | 7/27 | 1                | 70   | 0                | 30            | 0     | Calanoid Copepoda; Adult insect fragments                                    |
| Susitna #6A          | 7/27 | 2                | 0  | 80               | 20            | 0     | Diptera Chironomid larvae, unidentified adult insect fragments               |
| Susitna #10          | 7/27 | 2                | 0  | 20               | 80            | 0     | Unidentified adult insect fragments; Diptera Chironomid pupae                |
| Whiskers Creek       | 8/5  | 2                | 10   | 10               | 80            | 0     | Adult Coleoptera fragments   |
| Chase Creek          | 8/6  | 2                | <5   | 0                | >45           | 50    | Sand grains; adult Hymenoptera and Diptera                                   |

Appendix I Table 6. Stomach content analysis of juvenile chinook, coho, and sockeye salmon collected in sloughs and clearwater tributaries of the Susitna River during summer and fall, Devils Canyon Project, 1977, (continued).

| Location         | Date | Number Specimens | Estimated Percent of Combined Gut Contents |                  |               |       | Predominate Organisms  |
|------------------|------|------------------|--|------------------|---------------|-------|--|
|                  |      |                  | Crustacea                                  | Immature Insecta | Adult Insecta | Other |  |
| McKenzie Creek   | 7/27 | 3                | 0  | 10               | 90            | 0     | Adult Diptera  |
| <u>Coho-Fall</u> |      |                  |  |                  |               |       |  |
| Susitna #1       | 9/24 | 14               | <1   | 1                | >98           | 0     | Adult Hemiptera, Homoptera, Coleoptera, Lepidoptera and fragments                  |
| Susitna #4       | 9/24 | 13               | 0  | <5               | >35           | 60    | 3 salmonid juveniles; adult Coleoptera, Hemiptera, Homoptera, Hymenoptera, Diptera |
| Susitna #5       | 9/23 | 4                | 0  | 1                | 89            | 10    | Adult Diptera, Coleoptera, Hemiptera, and Homoptera                                |
| Slough #A        | 9/23 | 1                | 0  | 70               | 30            | 0     | Diptera Chironomid larvae; Ephemeroptera Plecoptera nymphs; adult Diptera          |
| Susitna #8C      | 9/23 | 1                | 0  | 40               | 60            | 0     | Unidentified adult Insecta; Diptera Chironomid larvae                              |
| Susitna #13      | 9/22 | 1                | <1   | 10               | >84           | 5     | Adult Diptera and Homoptera  |
| Susitna #16      | 9/21 | 2                | 0  | 20               | 0             | 80    | Algae; Diptera Chironomid larvae   |
| Susitna #19      | 9/21 | 2                | 0  | 20               | 80            | 0     | Adult Coleoptera and Diptera; Diptera Chironomid pupae                             |
| Susitna #21      | 9/20 | 1                | 0  | 10               | 90            | 0     | Adult Diptera  |

Appendix I Table 6. Stomach content analysis of juvenile chinook, coho, and sockeye salmon collected in sloughs and clearwater tributaries of the Susitna River during summer and fall, Devils Canyon Project, 1977, (continued).

| Location              | Date | Number Specimens | <u>Estimated Percent of Combined Gut Contents</u> |                  |               |       | Predominate Organisms  |
|-----------------------|------|------------------|---|------------------|---------------|-------|--|
|                       |      |                  | Crustacea   | Immature Insecta | Adult Insecta | Other |  |
| Whiskers Creek        | 9/24 | 6                | 0   | 20               | 80            | 0     | Adult Homoptera, Coleoptera, Hemiptera, Hymenoptera; Trechoptera pupae |
| <u>Sockeye-Summer</u> |      |                  |   |                  |               |       |  |
| Susitna #1            | 8/5  | 2                | 50  | 20               | 20            | 10    | Calanoid and Cyclopoid Copepoda; Diptera adults and Chironomid larvae  |
| Susitna #5            | 7/27 | 1                | 90  | 10               | 0             | 0     | Cladocera Bosminidae   |
| Susitna #6            | 7/27 | 3                | 50  | 50               | 0             | 0     | Ostracoda; Diptera Chironomid larvae                                   |
| Susitna #8            | 7/27 | 4                | 0   | 10               | 90            | 0     | Adult Diptera, Homoptera, Hymenoptera                                  |
| Susitna #12           | 7/29 | 10               | 0   | 90               | 10            | 0     | Diptera Chironomid larvae  |
| Susitna #17           | 7/26 | 1                | 20  | 80               | 0             | 0     | Diptera Chironomid larvae  |
| Susitna #19           | 7/14 | 3                | 20  | 40               | 40            | 0     | Adult Diptera; Diptera Chironomid larvae; Cladocera Bosminidae         |
| Susitna #19           | 7/26 | 11               | 50  | 50               | 0             | 0     | Cladocera Bosminidae; Diptera Chironomid larvae                        |

Appendix I Table 6. Stomach content analysis of juvenile chinook, coho, and sockeye salmon collected in sloughs and clearwater tributaries of the Susitna River during summer and fall, Devils Canyon Project, 1977, (continued).

| Location            | Date | Number<br>Specimens | Estimated Percent of Combined Gut Contents |                     |                  |       | Predominate Organisms   |
|---------------------|------|---------------------|--|---------------------|------------------|-------|---|
|                     |      |                     | Crustacea                                  | Immature<br>Insecta | Adult<br>Insecta | Other |   |
| <u>Sockeye-Fall</u> |      |                     |  |                     |                  |       |   |
| Susitna<br>#3       | 9/24 | 1                   | 5  | 10                  | 85               | 0     | Unidentified adult insect fragments;<br>Diptera Chironomid larvae |
| Susitna<br>#8       | 9/23 | 2                   | 1  | 9                   | 90               | 0     | Adult Diptera   |
| Susitna<br>#8B      | 9/23 | 12                  | <1   | 3                   | >95              | <1    | Adult Hymenoptera, Diptera, and<br>Lepidoptera                    |
| Susitna<br>#11      | 9/21 | 2                   | 5  | 95                  | 0                | 0     | Diptera Chironomid pupae and larvae                               |
| Susitna<br>#19      | 9/21 | 1                   | 10   | 50                  | 40               | 0     | Diptera Chironomid larvae; unidentified<br>adult insect fragments |



## APPENDIX II

Tables in the following appendix include data on water quality and quantity within the mainstem Susitna River and its clearwater sloughs and tributaries collected by ADF&G and USGS water quality data collected at established gaging stations.

## Appendix II

Table 1. Susitna River discharge at Cold Creek (USGS provisional data) 1977.

| Day | May          |           | June         |           | July         |           | August       |           | September    |           |
|-----|--------------|-----------|--------------|-----------|--------------|-----------|--------------|-----------|--------------|-----------|
|     | Gauge Height | Discharge | Gauge Height | Discharge | Gauge Height | Discharge | Gauge Height | Discharge | Gauge Height | Discharge |
| 1   |              |           | 11.49        | 30,900    |              | 35,000    | 10.60        | 24,200    | 8.00         | 10,600    |
| 2   |              |           | 12.19        | 36,700    |              | 32,000    | 10.89        | 26,200    | 8.03         | 10,700    |
| 3   |              |           | 12.44        | 39,000    |              | 32,000    | 10.49        | 23,400    | 8.02         | 10,700    |
| 4   |              |           | 12.52        | 39,700    |              | 30,000    | 10.17        | 21,500    | 7.97         | 10,500    |
| 5   |              |           | 12.34        | 38,100    |              | 30,000    | 10.20        | 21,700    | 7.81         | 9,840     |
| 6   |              |           | 11.77        | 33,200    |              | 28,000    | 10.49        | 23,400    | 7.72         | 9,520     |
| 7   |              |           | 11.54        | 31,300    |              | 27,000    | 10.48        | 23,400    | 7.82         | 9,880     |
| 8   |              |           | 11.56        | 31,500    |              | 27,000    | 10.12        | 21,200    | 8.20         | 11,400    |
| 9   |              |           | 11.61        | 31,900    |              | 27,000    | 9.80         | 19,300    | 8.48         | 12,500    |
| 10  |              |           | 11.93        | 34,400    |              | 28,000    | 9.90         | 19,900    | 8.05         | 10,800    |
| 11  |              |           | 12.39        | 38,500    |              | 28,000    | 9.98         | 20,400    | 8.81         | 14,000    |
| 12  |              |           | 13.02        | 44,200    |              | 30,000    | 10.12        | 21,200    | 9.34         | 16,700    |
| 13  |              |           | 13.67        | 51,400    |              | 30,000    | 9.74         | 18,900    | 9.38         | 16,900    |
| 14  |              |           | 13.78        | 52,600    |              | 32,000    | 9.58         | 18,000    | 9.37         | 16,800    |
| 15  |              |           | 13.78        | 52,600    |              | 32,000    | 9.94         | 20,100    | 9.01         | 15,000    |
| 16  |              |           | 13.58        | 50,400    |              | 30,000    | 10.01        | 20,600    | 9.13         | 15,600    |
| 17  | 8.72         | 13,600    | 13.07        | 44,800    |              | 28,000    | 9.78         | 19,200    | 8.90         | 14,500    |
| 18  | 9.06         | 15,300    | 12.82        | 42,400    |              | 26,000    | 9.69         | 18,600    | 8.87         | 14,400    |
| 19  | 10.37        | 22,700    | 12.67        | 41,000    |              | 24,000    | 9.66         | 18,500    | 8.59         | 13,000    |
| 20  | 10.92        | 26,400    | 12.22        | 37,000    | 10.31        | 22,400    | 9.67         | 18,500    | 8.41         | 12,200    |
| 21  | 10.19        | 21,600    | 11.93        | 34,400    | 10.28        | 22,200    | 9.76         | 19,100    | 8.84         | 14,200    |
| 22  | 9.56         | 17,900    | 11.75        | 33,000    | 10.22        | 21,800    | 9.93         | 20,100    | 8.93         | 14,600    |
| 23  | 9.33         | 16,600    |              | 33,000    | 10.41        | 23,000    | 10.18        | 21,600    | 8.95         | 14,800    |
| 24  | 9.38         | 16,900    |              | 34,000    | 10.39        | 22,800    | 10.16        | 21,500    | 8.60         | 13,000    |
| 25  | 9.36         | 16,800    |              | 36,000    | 10.14        | 21,300    | 9.72         | 18,800    | 8.21         | 11,400    |
| 26  | 9.61         | 18,200    |              | 38,000    | 10.00        | 20,500    | 9.19         | 16,000    | 7.95         | 10,400    |
| 27  | 10.22        | 21,800    |              | 40,000    | 9.84         | 19,500    | 8.88         | 14,400    | 7.81         | 9,840     |
| 28  | 10.54        | 23,800    |              | 42,000    | 9.87         | 19,700    | 8.68         | 13,400    | 8.10         | 11,000    |
| 29  | 11.18        | 28,400    |              | 40,000    | 9.90         | 19,900    | 8.41         | 12,200    | 8.31         | 11,800    |
| 30  | 11.76        | 33,100    |              | 38,000    | 9.85         | 19,600    | 8.09         | 11,000    | 8.48         | 12,500    |
| 31  | 11.34        | 29,700    |              |           | 10.11        | 21,200    | 7.85         | 10,000    |              |           |

Appendix II Table 2. Water quality data and juvenile salmon surveys in sloughs and clearwater tributaries of the Susitna River between the Chulitna River and Portage Creek, Devils Canyon Project, 1977.

| Date                      | Time | Weather<br>Conditions | Water<br>Conditions | Temperature °C |      | D.O.<br>(PPM) | pH  | Specific<br>Conductance<br>(uMHOS/CM) | Gage<br>Height<br>(M) | Number of<br>Fry<br>Observed |
|---------------------------|------|-----------------------|---------------------|----------------|------|---------------|-----|---------------------------------------|-----------------------|------------------------------|
| Slough #1 (26N 05W 11DAD) |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/18                      | 1250 | Sunny                 | Silty               | 22.0           | 15.0 | 8.1           | 5.9 | —                                     | .90                   | —                            |
| 7/30                      | 1425 | Sunny                 | Silty               | 22.0           | 11.0 | 8.4           | —   | 100                                   | .38                   | —                            |
| 8/5                       | 1340 | Sunny                 | Silty               | 23.5           | 13.0 | 9.7           | 6.5 | 100                                   | .76                   | —                            |
| 8/12                      | 1535 | Cloudy                | Silty               | 16.0           | 11.5 | 8.4           | 6.6 | 290                                   | .76                   | 190                          |
| 8/22                      | 1620 | Cloudy                | Silty               | 17.5           | 14.0 | 8.9           | —   | 100                                   | .70                   | 150                          |
| 9/24                      | 1530 | Sunny                 | Clear               | 16.0           | 6.0  | 9.4           | 6.0 | 125                                   | .31                   | 14                           |
| Slough #2 (26N 05W 02CDD) |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/18                      | 1330 | Sunny                 | —                   | 22.0           | 11.5 | 8.7           | 6.2 | —                                     | .26                   | 500                          |
| 7/30                      | 1450 | Sunny                 | —                   | 21.0           | 10.0 | 8.6           | —   | 150                                   | .16                   | 4                            |
| 8/5                       | 1440 | Sunny                 | —                   | 20.0           | 9.0  | 8.4           | 6.5 | 190                                   | .17                   | 100                          |
| 8/12                      | 1615 | Sunny                 | —                   | 18.5           | 10.0 | 6.7           | 7.1 | 130                                   | .18                   | 60                           |
| 8/22                      | 1520 | Cloudy                | —                   | 18.0           | 10.0 | 8.0           | —   | 130                                   | .16                   | 125                          |
| 9/10                      | 1130 | Rain                  | —                   | 10.0           | 8.0  | 10.1          | 6.0 | 102                                   | —                     | —                            |
| Slough #3 (27N 05W 35CCB) |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/17                      | 1800 | Sunny                 | Silty               | 26.5           | 19.5 | 8.3           | 5.9 | —                                     | —                     | —                            |
| 7/30                      | 1600 | Sunny                 | Clear               | 19.0           | 13.5 | 7.1           | —   | 125                                   | —                     | 2                            |
| 8/5                       | 1745 | Sunny                 | Clear               | 19.5           | 11.0 | 7.0           | 5.6 | 100                                   | —                     | 100                          |
| 8/12                      | 1800 | Sunny                 | Clear               | 20.5           | 10.0 | 5.4           | 6.2 | 110                                   | —                     | 465                          |
| 8/22                      | 1800 | Sunny                 | Clear               | 17.0           | 13.0 | 5.6           | —   | 100                                   | —                     | 300                          |
| 9/10                      | 1100 | Rain                  | Silty               | 10.2           | 6.8  | 6.6           | 5.5 | 72                                    | —                     | —                            |
| 9/24                      | 1245 | Sunny                 | Clear               | 10.0           | 5.3  | 8.2           | 5.5 | 85                                    | —                     | 350                          |
| Slough #4 (27N 05W 25CCC) |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/17                      | 1725 | Sunny                 | Clear               | 22.0           | 17.0 | 7.5           | 6.5 | —                                     | .88                   | 1,000+                       |
| 8/14                      | 1800 | Rain                  | Clear               | 17.5           | 15.0 | 9.1           | 6.0 | 100                                   | .82                   | 500                          |
| 9/10                      | 0900 | Rain                  | Silty               | 9.0            | 10.1 | 11.1          | 6.0 | 78                                    | .61                   | —                            |
| 9/24                      | 1120 | Sunny                 | Clear               | 7.0            | 6.2  | 10.9          | 5.0 | 85                                    | .82                   | 52                           |
| Slough #5 (27N 05W 01CCA) |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/16                      | 1050 | —                     | Silty               | 23.0           | 17.0 | 4.3           | 7.3 | —                                     | .58                   | —                            |
| 7/27                      | 1800 | Sunny                 | Rusty               | 22.0           | 18.5 | 5.4           | 6.3 | 120                                   | —                     | 5                            |
| 8/6                       | 1200 | Rain                  | —                   | 16.0           | 15.0 | 7.2           | 6.0 | 105                                   | .26                   | 10                           |
| 8/13                      | 0845 | Rain                  | —                   | 14.0           | 13.5 | 2.7           | 6.0 | 180                                   | .13                   | —                            |
| 8/21                      | 1330 | Rain                  | —                   | 21.0           | 14.0 | 1.2           | —   | 240                                   | .11                   | —                            |
| 8/29                      | 1730 | Partly<br>Sunny       | —                   | 17.0           | 15.0 | 7.6           | 6.5 | 100                                   | —                     | 7                            |
| 9/9                       | 1915 | Overcast              | Algae               | 11.0           | 12.5 | 10.8          | 6.0 | 88                                    | —                     | 90                           |
| 9/23                      | 1720 | Overcast              | —                   | 11.0           | 9.9  | 11.0          | 5.0 | 68                                    | —                     | 4                            |
| Slough #6 (27N 05W 01BAD) |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/16                      | 1115 | —                     | —                   | 23.5           | 14.0 | 9.2           | 7.0 | —                                     | .85                   | —                            |
| 7/27                      | 1715 | Sunny                 | Rusty               | 24.0           | 22.0 | 6.2           | 6.7 | 100                                   | .36                   | 100                          |
| 8/6                       | 1230 | Rain                  | Rusty               | 19.5           | 14.0 | 5.6           | 5.0 | 100                                   | .57                   | 5                            |
| 8/13                      | 0800 | Rain                  | Rusty               | 13.5           | 12.0 | 4.8           | 6.0 | 110                                   | .42                   | 12                           |
| 8/21                      | 1315 | Rain                  | Rusty               | 21.0           | 16.0 | 7.2           | —   | 100                                   | .39                   | 42                           |
| 8/29                      | 1700 | Partly<br>Sunny       | Rusty &<br>Algae    | 18.5           | 17.2 | 9.8           | 6.0 | 130                                   | .33                   | —                            |
| 9/9                       | 1850 | Overcast              | Rusty &<br>Algae    | 11.0           | 12.5 | 10.8          | 6.0 | 88                                    | —                     | —                            |
| 9/23                      | 1650 | Overcast              | —                   | 11.0           | 8.3  | 10.4          | 6.0 | 38                                    | .36                   | —                            |

Appendix II Table 2. Water quality data and juvenile salmon surveys in sloughs and clearwater tributaries of the Susitna River between the Chulitna River and Portage Creek, Devils Canyon Project, 1977 (continued).

| Date                                    | Time | Weather<br>Conditions | Water<br>Conditions | Temperature °C |      | D.O.<br>(PPM) | pH  | Specific<br>Conductance<br>(uMHOS/CM) | Gage<br>Height<br>(M) | Number of<br>Fry<br>Observed |
|---|------|-----------------------|---------------------|----------------|------|---------------|-----|---------------------------------------|-----------------------|------------------------------|
| Slough #7 (28N 05W 12DCA)               |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/13                                    | 1530 | Sunny                 | Clear               | 20.0           | 14.0 | 8.4           | 8.1 | —                                     | —                     | —                            |
| 8/4                                     | 1500 | Sunny                 | Clear               | 16.0           | 12.0 | 11.2          | —   | 100                                   | —                     | —                            |
| 8/11                                    | 2025 | Sunny                 | Clear               | 16.0           | 17.0 | 9.4           | 6.0 | 130                                   | —                     | 12                           |
| 8/19                                    | 1930 | Sunny                 | Clear               | 17.5           | 16.0 | 8.1           | —   | 100                                   | —                     | 10                           |
| 8/29                                    | 1530 | Partly<br>Sunny       | Clear               | 18.0           | 17.0 | 11.0          | 6.0 | 90                                    | —                     | 30                           |
| 9/9                                     | 1815 | Overcast              | Clear               | 11.0           | 14.5 | 10.0          | 5.0 | 100                                   | —                     | 80                           |
| 9/23                                    | 1550 | Overcast              | Clear               | 11.0           | 9.0  | 12.2          | 5.0 | 100                                   | —                     | —                            |
| Slough #8 (28N 04W 07BCB and 07BCC)     |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/13                                    | 1500 | Sunny                 | Clear               | 18.0           | 12.0 | 9.2           | 7.8 | —                                     | .36                   | 3,500                        |
| 7/27                                    | 1510 | Sunny                 | Clear               | 23.0           | 13.5 | 9.1           | 7.3 | 70                                    | .26                   | —                            |
| 8/4                                     | 1435 | Rain                  | —                   | 16.0           | 9.5  | 9.2           | —   | 65                                    | —                     | —                            |
| 8/11                                    | 1915 | Sunny                 | Clear               | 16.0           | 9.0  | 10.4          | 6.2 | 90                                    | .26                   | 670                          |
| 8/19                                    | 1850 | Sunny                 | Clear               | 18.0           | 11.0 | 9.2           | —   | 102                                   | .25                   | 400                          |
| 8/29                                    | 1500 | Sunny                 | Clear               | 18.5           | 11.5 | 10.3          | 6.0 | 70                                    | .24                   | —                            |
| 9/9                                     | 1800 | Overcast              | —                   | 12.5           | 8.8  | 10.7          | —   | 100                                   | .21                   | 1,200                        |
| 9/23                                    | 1500 | Overcast              | —                   | 12.0           | 7.8  | 9.8           | 5.0 | 100                                   | .20                   | 35                           |
| Slough #8A (30N 03W 20C, 29BBB and 30A) |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/12                                    | 1730 | Sunny                 | Clear               | 23.0           | 17.0 | 10.1          | 7.6 | —                                     | —                     | —                            |
| 8/3                                     | 1730 | Fair                  | Silty               | 19.0           | 16.0 | 8.4           | 6.8 | 140                                   | —                     | 1,500                        |
| 8/11                                    | 1400 | Sunny                 | Clear               | 17.5           | 17.0 | 7.1           | 6.5 | 175                                   | —                     | —                            |
| 8/19                                    | 1400 | Sunny                 | Clear               | 17.0           | 14.0 | 8.0           | —   | 45                                    | —                     | 2,000                        |
| 8/29                                    | 0825 | Partly<br>Cloudy      | Clear               | 5.0            | 10.0 | 10.3          | 5.8 | 118                                   | —                     | 90                           |
| 9/9                                     | 1350 | Partly<br>Cloudy      | Clear               | 12.5           | 12.5 | 10.1          | 6.0 | 145                                   | —                     | 135                          |
| 9/22                                    | 1700 | Overcast              | —                   | 9.0            | 7.1  | 10.3          | 5.6 | 75                                    | —                     | —                            |
| Slough #8B (29N 04W 02CBA)              |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/13                                    | 1105 | Sunny                 | Silty               | 17.0           | 11.0 | 8.1           | 7.9 | —                                     | —                     | 1,000                        |
| 8/4                                     | 1000 | Sunny                 | Silty               | 14.0           | 8.5  | 9.2           | 6.5 | 100                                   | —                     | —                            |
| 8/11                                    | 1515 | Sunny                 | Clear               | 19.5           | 13.0 | 8.4           | 6.7 | 170                                   | —                     | 560                          |
| 8/19                                    | 1510 | Sunny                 | Clear               | 17.0           | 12.0 | 7.7           | —   | 200                                   | —                     | 650                          |
| 8/29                                    | 0930 | Partly<br>Cloudy      | Clear               | 10.0           | 7.2  | 11.2          | 5.8 | 110                                   | —                     | 350                          |
| 9/9                                     | 1510 | Partly<br>Cloudy      | Clear               | 14.0           | 9.9  | 8.8           | 6.0 | 135                                   | —                     | —                            |
| 9/23                                    | 1100 | Sunny                 | Clear               | 6.5            | 5.0  | 10.8          | 5.6 | 68                                    | —                     | 25                           |
| Slough #8C (29N 04W 02CCC)              |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/13                                    | 1140 | Sunny                 | Clear               | 21.5           | 9.5  | 7.2           | 7.8 | —                                     | —                     | 500                          |
| 8/4                                     | 1100 | Sunny                 | Clear               | 16.0           | 7.5  | 7.5           | 6.0 | 80                                    | —                     | 30                           |
| 8/11                                    | 1625 | Sunny                 | Clear               | 16.5           | 11.0 | 7.2           | 6.9 | 70                                    | —                     | 34                           |
| 8/19                                    | 1540 | Sunny                 | Clear               | 16.0           | 11.0 | 6.8           | —   | 130                                   | —                     | 850                          |
| 8/29                                    | 1020 | Partly<br>Cloudy      | Clear               | 11.0           | 6.9  | 9.0           | 5.5 | 60                                    | —                     | —                            |
| 9/9                                     | 1540 | Partly<br>Cloudy      | Clear               | 14.0           | 8.8  | 7.5           | 5.5 | 60                                    | —                     | —                            |
| 9/23                                    | 1130 | Sunny                 | Clear               | 10.0           | 7.0  | 10.8          | 5.6 | 45                                    | —                     | 7                            |
| Slough #8D (29N 04W 11BBA)              |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/13                                    | 1200 | Sunny                 | Clear               | 23.0           | 11.0 | 11.0          | 7.2 | —                                     | —                     | —                            |
| 8/4                                     | 1130 | Sunny                 | Clear               | 17.0           | 8.0  | 9.2           | 5.0 | 50                                    | —                     | 4                            |
| 8/11                                    | 1640 | Sunny                 | Clear               | 18.0           | 14.0 | 8.0           | 6.9 | 90                                    | —                     | —                            |
| 8/19                                    | 1600 | Sunny                 | Clear               | 15.0           | 13.0 | 8.6           | —   | 130                                   | —                     | 40                           |
| 8/29                                    | 1045 | Partly<br>Cloudy      | Clear               | 13.2           | 8.0  | 10.8          | 5.8 | 58                                    | —                     | 50                           |
| 9/9                                     | 1600 | Partly<br>Cloudy      | Clear               | 13.0           | 9.9  | 9.8           | 5.5 | 72                                    | —                     | 750                          |
| 9/23                                    | 1210 | Sunny                 | Clear               | 12.2           | 6.5  | 10.3          | 5.6 | 55                                    | —                     | 1                            |

Appendix II Table 2. Water quality data and juvenile salmon surveys in sloughs and clearwater tributaries of the Susitna River between the Chulitna River and Portage Creek, Devils Canyon Project, 1977 (continued).

| Date  | Time | Weather<br>Conditions | Water<br>Conditions | Temperature °C |      | D.O.<br>(PPM) | pH  | Specific<br>Conductance<br>(µMHOS/CM) | Gage<br>Height<br>(M) | Number of<br>Fry<br>Observed |
|---|------|-----------------------|---------------------|----------------|------|---------------|-----|---------------------------------------|-----------------------|------------------------------|
| <u>Slough A (30N 04W 25DBB)</u>             |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/13  | 1020 | Sunny                 | Clear               | 17.5           | 9.0  | 9.9           | 7.7 | —                                     | —                     | —                            |
| 8/3   | 1830 | Sunny                 | Clear               | 24.0           | 12.0 | 7.0           | 6.3 | 110                                   | —                     | 20                           |
| 8/11  | 1415 | Sunny                 | Clear               | 15.5           | 14.0 | 7.1           | 6.9 | 110                                   | —                     | 27                           |
| 8/19  | 1415 | Sunny                 | Clear               | 20.0           | 13.0 | 7.1           | —   | 200                                   | —                     | 85                           |
| 8/29  | 0900 | Partly<br>Cloudy      | Clear               | 11.0           | 9.8  | 11.3          | 5.0 | 110                                   | —                     | —                            |
| 9/9   | 1400 | Partly<br>Cloudy      | Clear               | 12.2           | 9.9  | 12.4          | —   | 85                                    | —                     | —                            |
| 9/23  | 1010 | Sunny                 | Clear               | 5.2            | 6.0  | 9.8           | 5.6 | 58                                    | —                     | 3                            |
| <u>Slough #9 (30N 03W 16BD)</u>             |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/12  | 1600 | Clear                 | Silty               | 20.0           | 15.5 | 9.6           | 8.0 | —                                     | .39                   | —                            |
| 7/27  | 0850 | Cloudy                | Clear               | 15.0           | 8.0  | 8.9           | 6.7 | 190                                   | .38                   | 40                           |
| 8/3   | 1630 | Clear                 | Silty               | 17.0           | 13.0 | 8.8           | 7.0 | 115                                   | .39                   | —                            |
| 8/11  | 1200 | Clear                 | Clear               | 17.0           | 11.0 | 7.7           | 6.8 | 175                                   | .38                   | 140                          |
| 8/19  | 1150 | Clear                 | Clear               | 17.0           | 10.0 | 8.0           | —   | 210                                   | .38                   | 700                          |
| 8/29  | 1015 | Rain                  | Clear               | 15.0           | 12.0 | 7.0           | 5.4 | —                                     | .38                   | 600                          |
| 9/9   | 1230 | Overcast              | Clear               | 11.0           | 8.0  | 9.9           | 6.0 | 135                                   | .36                   | 250                          |
| 9/22  | 1500 | Clear                 | Clear               | 10.5           | 7.8  | 10.8          | 5.6 | 100                                   | .43                   | 78                           |
| <u>Slough #10 (31N 03W 36AAB)</u>           |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/8   | 1115 | Clear                 | Clear               | 22.0           | 7.0  | 11.0          | 6.8 | —                                     | .68                   | —                            |
| 7/12  | 1445 | Clear                 | Clear               | 22.0           | 11.5 | 9.1           | 7.8 | —                                     | .81                   | —                            |
| 7/27  | 0850 | Partly<br>Cloudy      | Clear               | 19.0           | 7.0  | 9.7           | 6.5 | 150                                   | .71                   | 1,000                        |
| 8/3   | 1135 | Clear                 | Clear               | 16.0           | 7.5  | 9.4           | 6.5 | 100                                   | .88                   | 1,000                        |
| 8/19  | 0900 | Clear                 | Clear               | 12.0           | 6.0  | 8.5           | 6.0 | 140                                   | .65                   | 1,200                        |
| 8/26  | 1415 | Rain                  | Clear               | 13.0           | 4.0  | 7.0           | 6.4 | —                                     | .52                   | 2,500                        |
| 9/9   | 1015 | Overcast              | Clear               | 9.0            | 5.1  | 9.6           | 6.0 | 145                                   | .30                   | 250                          |
| <u>Slough #11 (31N 02W 30AAB, 20B, 20C)</u> |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/1   | 1500 | Clear                 | Clear               | 23.0           | 10.0 | —             | —   | —                                     | 1.38                  | —                            |
| 7/12  | 1350 | Clear                 | Clear               | 23.0           | 11.0 | 10.6          | 7.8 | —                                     | 1.24                  | 3,000                        |
| 7/27  | 1745 | Partly<br>Cloudy      | Clear               | 27.0           | 12.0 | 11.0          | 7.9 | —                                     | <1.00                 | 8,000                        |
| 8/2   | 2010 | Clear                 | Silty               | 14.0           | 9.0  | 9.2           | 7.5 | —                                     | <1.00                 | —                            |
| 8/3   | 1710 | Mostly<br>Sunny       | Silty               | 28.0           | 12.0 | 11.0          | 7.3 | 180                                   | <1.00                 | 8,000                        |
| 8/10  | 1600 | Partly<br>Cloudy      | Silty               | 16.5           | 11.0 | 10.0          | 7.0 | 155                                   | <1.00                 | 8,000                        |
| 8/17  | 1530 | Clear                 | Silty               | 16.0           | 9.0  | 11.0          | —   | 150                                   | 1.08                  | 8,000                        |
| 8/28  | 1820 | Clear                 | Clear               | 13.0           | 8.5  | 9.8           | 5.0 | 170                                   | .92                   | 10,000                       |
| 9/8   | 1900 | Clear                 | Clear               | 10.5           | 5.2  | 10.8          | 5.5 | 190                                   | .89                   | 2,000                        |
| 9/22  | 1215 | Clear                 | Clear               | 8.5            | 6.2  | 11.5          | 5.6 | 105                                   | 1.00                  | 87                           |
| <u>Slough #12 (31N 02W 19DCD)</u>           |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/1   | 1400 | Mostly<br>Cloudy      | Silty               | 22.0           | 11.0 | —             | —   | —                                     | —                     | —                            |
| 7/12  | 1330 | Clear                 | Silty               | 23.0           | —    | 9.1           | 8.2 | —                                     | .95                   | —                            |
| 7/27  | 1545 | Partly<br>Cloudy      | Silty               | 20.0           | 9.0  | 8.0           | 7.6 | 150                                   | .85                   | 30                           |
| 8/2   | 1945 | Clear                 | Silty               | 14.0           | 9.5  | 9.2           | 6.8 | 200                                   | <1.00                 | —                            |
| 8/3   | 1540 | Mostly<br>Sunny       | Clear               | 20.0           | 11.0 | 9.0           | 7.5 | 200                                   | .94                   | 175                          |
| 8/10  | 1540 | Clear                 | Silty               | 17.5           | 13.0 | 8.9           | 7.4 | 170                                   | .85                   | 10                           |
| 8/17  | 1600 | Rain                  | Silty               | 15.0           | 10.0 | 10.0          | —   | 175                                   | .80                   | —                            |
| 8/23  | —    | —                     | —                   | —              | —    | —             | —   | —                                     | —                     | 10                           |
| 8/28  | 1750 | Clear                 | —                   | 15.0           | 10.0 | 10.0          | 6.0 | 158                                   | —                     | 10                           |
| 9/8   | 1830 | Rain                  | Silty               | 10.0           | 9.1  | 11.1          | 6.0 | 160                                   | .51                   | 9                            |
| 9/22  | 1200 | Clear                 | Clear               | 7.0            | 6.0  | 11.0          | 6.5 | 175                                   | .64                   | 0                            |

Appendix II Table 2. Water quality data and juvenile salmon surveys in sloughs and clearwater tributaries of the Susitna River between the Chulitna River and Portage Creek, Devils Canyon Project, 1977 (continued).

| Date                              | Time | Weather<br>Conditions | Water<br>Conditions | Temperature °C<br>Air      Water | D.O.<br>(PPM) | pH  | Specific<br>Conductance<br>(uMHOS/CM) | Gage<br>Height<br>(M) | Number of<br>Fry<br>Observed |
|-----------------------------------|------|-----------------------|---------------------|----------------------------------|---------------|-----|---------------------------------------|-----------------------|------------------------------|
| <u>Slough #13 (31N 02W 19DAB)</u> |      |                       |                     |                                  |               |     |                                       |                       |                              |
| 7/1                               | 1340 | Mostly<br>Cloudy      | Clear               | 24.0      7.0                    | —             | —   | —                                     | —                     | —                            |
| 7/12                              | 1310 | Clear                 | Clear               | 26.0      —                      | 11.1          | 7.7 | —                                     | .26                   | —                            |
| 7/28                              | 1210 | Clear                 | Clear               | 25.0      7.0                    | 10.0          | 7.4 | 180                                   | .25                   | 75                           |
| 8/2                               | 1900 | Clear                 | Clear               | 18.0      7.5                    | 9.6           | 6.5 | 185                                   | .27                   | 400                          |
| 8/3                               | 1800 | Partly<br>Cloudy      | Clear               | 25.0      7.0                    | 9.0           | 7.5 | 180                                   | .26                   | 75                           |
| 8/10                              | 1500 | Clear                 | Clear               | 20.0      9.0                    | 8.8           | 6.9 | 205                                   | .25                   | 310                          |
| 8/17                              | 1630 | Clear                 | Clear               | 15.0      7.0                    | 11.0          | —   | 170                                   | .25                   | 400                          |
| 8/28                              | 1725 | Clear                 | Clear               | 15.0      7.0                    | 9.5           | 6.0 | 160                                   | .25                   | 400                          |
| 9/8                               | 1800 | Clear                 | Clear               | 10.0      7.0                    | 11.7          | 6.0 | 130                                   | .29                   | 120                          |
| 9/22                              | 1130 | Clear                 | Clear               | 8.5      5.0                     | 11.4          | 6.0 | 105                                   | .25                   | 5                            |
| <u>Slough #14 (31N 02W 19AAA)</u> |      |                       |                     |                                  |               |     |                                       |                       |                              |
| 7/1                               | 1300 | Mostly<br>Cloudy      | Clear               | 26.0      15.0                   | —             | —   | —                                     | .58                   | —                            |
| 7/12                              | 1240 | Clear                 | Clear               | 26.0      —                      | 7.1           | 7.4 | —                                     | .43                   | —                            |
| 7/28                              | 1305 | Clear                 | Clear               | 26.0      15.0                   | 9.0           | 6.9 | 85                                    | .35                   | —                            |
| 7/29                              | 0950 | Clear                 | Clear               | —                                | —             | —   | —                                     | —                     | 500                          |
| 8/2                               | 1800 | Clear                 | Clear               | 18.0      13.5                   | 7.1           | 6.3 | 85                                    | —                     | 2,000                        |
| 8/3                               | 1900 | Partly<br>Cloudy      | Clear               | 18.0      11.0                   | 6.8           | 6.0 | 80                                    | .42                   | —                            |
| 8/10                              | 1435 | Clear                 | Clear               | 18.0      15.0                   | 7.8           | 6.0 | 95                                    | .35                   | 100                          |
| 8/19                              | 0830 | Clear                 | Clear               | 12.0      10.5                   | 7.8           | 6.0 | 78                                    | .28                   | 120                          |
| 8/26                              | 1530 | Rain                  | —                   | 10.0      14.0                   | —             | 6.5 | —                                     | .23                   | 500                          |
| 8/28                              | 1645 | Clear                 | Clear               | 13.0      12.0                   | 8.0           | 6.5 | 85                                    | .65                   | 100                          |
| 9/8                               | 1720 | Overcast              | Cloudy              | 12.0      8.9                    | 10.8          | 8.5 | 60                                    | 1.15                  | 20                           |
| 9/22                              | 1030 | Clear                 | Clear               | 7.5      6.5                     | 10.3          | 5.6 | 34                                    | .50                   | 10                           |
| <u>Slough #15 (31N 02W 17CAC)</u> |      |                       |                     |                                  |               |     |                                       |                       |                              |
| 7/1                               | 1235 | Mostly<br>Cloudy      | Clear               | 25.0      15.0                   | —             | —   | —                                     | .88                   | —                            |
| 7/12                              | 1215 | Clear                 | Silty               | 22.5      14.0                   | 8.3           | 8.4 | —                                     | .66                   | —                            |
| 7/26                              | 1745 | Cloudy                | Silty               | 18.5      13.5                   | 7.3           | 6.7 | 70                                    | .53                   | 1,500                        |
| 8/2                               | 1400 | Mostly<br>Cloudy      | Silty               | 17.0      12.5                   | 7.8           | 6.4 | 105                                   | .93                   | 2,000                        |
| 8/10                              | 1145 | Cloudy                | Silty               | 22.5      12.5                   | 7.2           | 6.4 | 105                                   | .55                   | —                            |
| 8/16                              | 2000 | Clear                 | Silty               | 16.5      14.0                   | 6.8           | 6.0 | 78                                    | .53                   | 1,000                        |
| 8/28                              | 1515 | Clear                 | Clear               | 16.0      8.8                    | 8.8           | 6.0 | 58                                    | .80                   | 155                          |
| 9/8                               | 1410 | Overcast              | Clear               | 7.0      10.6                    | 11.2          | 5.5 | 30                                    | —                     | 20                           |
| 9/21                              | 1630 | Rain                  | Clear               | 7.5      6.5                     | 10.8          | 5.6 | 18                                    | .29                   | 19                           |
| 10/5                              | —    | Clear                 | Clear               | —                                | —             | —   | —                                     | .09                   | 50                           |
| (Ice Cover)                       |      |                       |                     |                                  |               |     |                                       |                       |                              |
| <u>Slough #16 (31N 02W 17AAC)</u> |      |                       |                     |                                  |               |     |                                       |                       |                              |
| 7/1                               | 1210 | Partly<br>Cloudy      | Clear               | 21.5      9.0                    | —             | —   | —                                     | .68                   | —                            |
| 7/11                              | 1600 | Clear                 | Clear               | —                                | 9.0           | 7.2 | —                                     | .26                   | —                            |
| 7/26                              | 1710 | Mostly<br>Cloudy      | Clear               | 17.5      12.5                   | 9.5           | 6.7 | 50                                    | .17                   | 9,000                        |
| 8/2                               | 1248 | Cloudy                | Silty               | 16.5      11.5                   | 10.4          | 7.6 | 95                                    | .72                   | —                            |
| 8/3                               | 1200 | Partly<br>Cloudy      | Silty               | 21.0      11.5                   | —             | —   | —                                     | .48                   | 99                           |
| 8/10                              | 1100 | Cloudy                | Clear               | 21.0      11.0                   | 10.8          | 6.6 | 80                                    | .18                   | 600                          |
| 8/16                              | 1925 | Clear                 | Clear               | 16.0      11.5                   | 8.7           | 6.0 | 75                                    | .17                   | —                            |
| 8/28                              | 1500 | Clear                 | Clear               | 17.0      10.0                   | 9.5           | 6.0 | 75                                    | —                     | —                            |
| 9/8                               | 1340 | Overcast              | Clear               | 10.5      7.2                    | 11.6          | 6.0 | 50                                    | .08                   | 300                          |
| 9/21                              | 1500 | Rain                  | Clear               | 8.0      5.5                     | 10.2          | 5.6 | 10                                    | —                     | 18                           |
| 10/5                              | —    | Clear                 | Clear               | —                                | —             | —   | —                                     | .13                   | 150                          |
| (Ice Cover)                       |      |                       |                     |                                  |               |     |                                       |                       |                              |

Appendix II Table 2. Water quality data and juvenile salmon surveys in sloughs and clearwater tributaries of the Susitna River between the Chulitna River and Portage Creek, Devils Canyon Project, 1977 (continued).

| Date                       | Time | Weather<br>Conditions | Water<br>Conditions | Temperature °C |      | D.O.<br>(PPM) | pH  | Specific<br>Conductance<br>(uMHOS/CM) | Gage<br>Height<br>(M) | Number of<br>Fry<br>Observed |
|----------------------------|------|-----------------------|---------------------|----------------|------|---------------|-----|---------------------------------------|-----------------------|------------------------------|
| Slough #17 (31N 02W 09DBD) |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 6/14                       | 0030 | Clear                 | Silty               | —              | —    | —             | —   | —                                     | 1.03                  | —                            |
| 6/30                       | 2345 | Clear                 | Silty               | 9.0            | 6.0  | 9.0           | 6.9 | 140                                   | .84                   | —                            |
| 7/7                        | 1740 | Mostly<br>Sunny       | Silty               | 19.0           | 10.0 | —             | —   | —                                     | .71                   | —                            |
| 7/11                       | 1710 | Clear                 | Silty               | —              | —    | 10.8          | 7.5 | —                                     | .94                   | 50                           |
| 7/26                       | 1315 | Clear                 | Clear               | 20.0           | 8.5  | 9.8           | 6.8 | 90                                    | .84                   | 900                          |
| 8/2                        | 1145 | Clear                 | Silty               | 17.0           | 8.0  | 10.2          | 6.2 | 85                                    | 1.12                  | 1                            |
| 8/10                       | 1000 | Cloudy                | Clear               | 20.0           | 7.0  | 8.4           | 6.0 | 95                                    | .83                   | 230                          |
| 8/16                       | 1830 | Clear                 | Clear               | 16.0           | 9.5  | 9.9           | 6.3 | 100                                   | .82                   | 3                            |
| 8/28                       | 1100 | Clear                 | Clear               | 17.0           | 6.5  | 8.5           | 5.5 | 58                                    | .75                   | 15                           |
| 9/8                        | 1200 | Rain                  | Clear               | 8.0            | 4.5  | 10.8          | 6.5 | 50                                    | .17                   | 10                           |
| 9/21                       | 1400 | Rain                  | Clear               | 7.5            | 4.5  | 10.8          | 5.9 | 70                                    | .37                   | —                            |
| 10/5                       | —    | Clear                 | Clear               | —              | —    | —             | —   | —                                     | .12                   | 6                            |
| (Ice Cover)                |      |                       |                     |                |      |               |     |                                       |                       |                              |
| Slough #18 (31N 02W 10CBC) |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/7                        | 1725 | Mostly<br>Sunny       | Clear               | 19.0           | 15.0 | 9.7           | 7.2 | —                                     | .51                   | —                            |
| 7/11                       | 1720 | Clear                 | Clear               | 20.0           | 13.5 | 9.1           | 7.5 | —                                     | .65                   | —                            |
| 7/26                       | 1245 | Clear                 | Clear               | 25.0           | 12.0 | 7.8           | 7.3 | 140                                   | .63                   | 1,000                        |
| 8/2                        | 1125 | Clear                 | Clear               | 15.0           | 8.5  | 8.2           | 6.7 | 145                                   | .98                   | 125                          |
| 8/10                       | 0945 | Cloudy                | Clear               | 20.5           | 9.0  | 7.0           | 6.4 | 115                                   | .63                   | 60                           |
| 8/16                       | 1835 | Clear                 | Clear               | 15.0           | 12.0 | 8.4           | —   | 125                                   | .60                   | 100                          |
| 8/28                       | 1015 | Clear                 | Clear               | 13.0           | 7.0  | 5.4           | 5.5 | 80                                    | —                     | —                            |
| 9/8                        | 1230 | Rain                  | Clear               | 9.5            | 8.2  | 9.7           | 5.5 | 88                                    | .48                   | 12                           |
| 9/21                       | 1330 | Rain                  | Clear               | 7.5            | 6.0  | 10.4          | 6.0 | 100                                   | .52                   | 3                            |
| 10/5                       | —    | Clear                 | Clear               | —              | —    | —             | —   | —                                     | .47                   | 50                           |
| (Ice Cover)                |      |                       |                     |                |      |               |     |                                       |                       |                              |
| Slough #19 (31N 02W 10DBD) |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 6/30                       | 2235 | Mostly<br>Sunny       | Clear               | —              | —    | 10.0          | 7.2 | 125                                   | .74                   | —                            |
| 7/7                        | 1700 | —                     | Clear               | 20.0           | 12.0 | —             | —   | —                                     | .43                   | —                            |
| 7/11                       | 1745 | Clear                 | Clear               | 15.0           | 18.5 | 11.4          | 6.7 | —                                     | .54                   | —                            |
| 7/26                       | 1210 | Mostly<br>Cloudy      | Clear               | 20.0           | 8.5  | 9.5           | 7.7 | 150                                   | .54                   | 2,000                        |
| 8/2                        | 1000 | Partly<br>Sunny       | Clear               | 11.5           | 7.5  | 10.8          | 6.6 | 130                                   | .78                   | 2,000                        |
| 8/10                       | 0845 | Cloudy                | Clear               | 17.0           | 9.0  | 8.6           | 6.4 | 140                                   | .53                   | 200                          |
| 8/16                       | 1750 | Cloudy                | Clear               | 18.5           | 10.0 | 8.3           | 6.8 | 130                                   | .53                   | 800                          |
| 8/27                       | 2010 | Cloudy                | Clear               | 12.0           | 8.5  | 9.0           | 6.9 | 100                                   | .19                   | 100                          |
| 9/7                        | 1935 | Rain                  | Clear               | 9.0            | 7.0  | 8.9           | 6.5 | 100                                   | .17                   | —                            |
| 9/21                       | 1100 | Rain                  | Clear               | 6.5            | 5.0  | 10.8          | 5.5 | 100                                   | .17                   | 11                           |
| 10/5                       | —    | Clear                 | Clear               | —              | —    | —             | —   | —                                     | .11                   | 500                          |
| (Ice Cover)                |      |                       |                     |                |      |               |     |                                       |                       |                              |
| Slough #20 (31N 02W 11BBD) |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 6/30                       | 2130 | Mostly<br>Sunny       | Silty               | 14.0           | 9.0  | 10.0          | 7.8 | 70                                    | .33                   | —                            |
| 7/7                        | 1630 | —                     | Silty               | 24.0           | 12.0 | —             | —   | —                                     | .39                   | —                            |
| 7/11                       | 1817 | Clear                 | Clear               | 19.0           | 12.0 | 10.4          | 7.5 | —                                     | .37                   | —                            |
| 7/25                       | 2045 | Mostly<br>Cloudy      | Silty               | 13.0           | 10.0 | 10.0          | 7.9 | 75                                    | .34                   | 56                           |
| 8/1                        | 1855 | Mostly<br>Sunny       | Silty               | 15.5           | 11.5 | 10.4          | 7.0 | 125                                   | .56                   | —                            |
| 8/9                        | 1945 | Cloudy                | Clear               | 16.5           | 12.5 | 9.6           | 6.7 | 140                                   | .31                   | 700                          |
| 8/16                       | 1725 | Partly<br>Cloudy      | Silty               | 18.0           | 13.0 | 9.6           | 7.2 | 180                                   | —                     | 700                          |
| 8/27                       | 1945 | Mostly<br>Sunny       | Silty               | 12.5           | 11.5 | 10.3          | 6.0 | 90                                    | —                     | 1,000                        |
| 9/7                        | 1910 | Rain                  | Clear               | 9.5            | 7.2  | 9.8           | 6.0 | 60                                    | .38                   | —                            |
| 9/20                       | 1910 | Rain                  | Clear               | 6.0            | 5.1  | 12.2          | —   | 60                                    | .77                   | 19                           |
| 10/5                       | —    | Clear                 | Clear               | —              | —    | —             | —   | —                                     | .44                   | —                            |
| (Ice Cover)                |      |                       |                     |                |      |               |     |                                       |                       |                              |

Appendix II Table 2. Water quality data and juvenile salmon surveys in sloughs and clearwater tributaries of the Susitna River between the Chulitna River and Portage Creek, Devils Canyon Project, 1977 (continued).

| Date                                   | Time | Weather<br>Conditions | Water<br>Conditions  | Temperature °C<br>Air Water | D.O.<br>(PPM) | pH  | Specific<br>Conductance<br>(µMHOS/CM) | Gage<br>Height<br>(M) | Number of<br>Fry<br>Observed |
|--|------|-----------------------|----------------------|-----------------------------|---------------|-----|---------------------------------------|-----------------------|------------------------------|
| <u>Slough #21 (32N 02W 36CCC)</u>      |      |                       |                      |                             |               |     |                                       |                       |                              |
| 6/30                                   | 1940 | Mostly<br>Sunny       | Silty                | 19.5 7.0                    | 8.0           | 7.9 | 175                                   | .52                   | —                            |
| 7/7                                    | 1530 | —                     | Silty                | 14.0 24.0                   | —             | —   | —                                     | .25                   | —                            |
| 7/11                                   | 2010 | Clear                 | Clear                | 21.5 8.0                    | 10.2          | 7.9 | —                                     | .30                   | —                            |
| 7/25                                   | 1945 | Clear                 | Clear                | 19.5 8.0                    | 8.9           | 8.0 | 180                                   | .28                   | —                            |
| 8/1                                    | 1710 | Mostly<br>Cloudy      | Silty                | 17.0 10.0                   | 9.6           | 6.8 | 200                                   | .43                   | 33                           |
| 8/9                                    | 1800 | Mostly<br>Cloudy      | Silty                | — 9.0                       | 9.9           | 7.6 | 245                                   | .26                   | 385                          |
| 8/16                                   | 1635 | Clear                 | Silty                | 18.0 12.0                   | 8.5           | 7.7 | 210                                   | .29                   | 600                          |
| 8/27                                   | 1820 | Partly<br>Cloudy      | Clear                | 17.0 7.5                    | 10.2          | 6.7 | 170                                   | —                     | 180                          |
| 9/7                                    | 1730 | Rain                  | Clear                | 11.5 7.5                    | 10.4          | 6.0 | 135                                   | —                     | —                            |
| 9/20                                   | 1800 | Rain                  | Clear                | 7.0 3.9                     | 11.6          | 5.6 | 145                                   | —                     | 15                           |
| 10/5                                   | —    | Clear                 | Clear<br>(Ice Cover) | —                           | —             | —   | —                                     | .01                   | 350                          |
| <u>Whisker's Creek (26N 05W 03AAC)</u> |      |                       |                      |                             |               |     |                                       |                       |                              |
| <u>Downstream Gage</u>                 |      |                       |                      |                             |               |     |                                       |                       |                              |
| 7/17                                   | 1820 | —                     | Silty                | 22.0 15.5                   | 9.0           | 6.2 | —                                     | —                     | —                            |
| 7/30                                   | 1530 | —                     | Clear                | 21.5 17.0                   | 8.9           | —   | 95                                    | .44                   | —                            |
| 8/5                                    | 1525 | —                     | Silty                | 17.0 13.0                   | 10.8          | 6.0 | 100                                   | .53                   | —                            |
| 8/12                                   | 1655 | —                     | Silty                | 17.0 14.0                   | 9.8           | 7.0 | 80                                    | .52                   | —                            |
| 8/22                                   | 1700 | —                     | Silty                | 17.5 16.0                   | 9.0           | —   | 70                                    | .48                   | 200                          |
| 9/10                                   | 1000 | Rain                  | Clear                | 9.5 9.1                     | 10.8          | 5.5 | 30                                    | .13                   | —                            |
| 9/24                                   | —    | —                     | —                    | —                           | —             | —   | —                                     | .25                   | —                            |
| <u>Upstream Gage</u>                   |      |                       |                      |                             |               |     |                                       |                       |                              |
| 7/17                                   | 1825 | —                     | Clear                | 25.0 15.5                   | 9.3           | 5.3 | —                                     | —                     | —                            |
| 7/30                                   | 1540 | —                     | Clear                | 20.0 16.0                   | 10.8          | —   | 60                                    | .76                   | 1,000                        |
| 8/5                                    | 1535 | —                     | Clear                | 17.5 14.5                   | 9.5           | 5.6 | 90                                    | .78                   | —                            |
| 8/12                                   | 1700 | —                     | Clear                | 19.5 14.0                   | 9.5           | 6.3 | 60                                    | .78                   | 500                          |
| 8/22                                   | 1715 | —                     | Clear                | 17.0 15.5                   | 9.9           | —   | 35                                    | .76                   | 200                          |
| 9/10                                   | —    | Rain                  | Clear                | —                           | —             | —   | —                                     | .78                   | —                            |
| 9/24                                   | 1320 | —                     | Clear                | 8.0 7.0                     | 11.2          | 5.5 | 38                                    | .69                   | 10                           |
| <u>McKenzie Creek (29N 04W 32ABA)</u>  |      |                       |                      |                             |               |     |                                       |                       |                              |
| 7/13                                   | 1350 | Sunny                 | Clear                | 21.0 11.0                   | 11.2          | 8.0 | —                                     | —                     | 30,000                       |
| 7/27                                   | 1405 | Sunny                 | Clear                | 20.5 10.5                   | 10.8          | 7.7 | 105                                   | —                     | 12,500                       |
| 8/4                                    | 1310 | Sunny                 | Clear                | 16.0 8.5                    | 11.8          | 6.9 | 100                                   | —                     | 2,000                        |
| 8/11                                   | 1800 | Sunny                 | Clear                | 17.0 11.0                   | 9.8           | 5.9 | 125                                   | —                     | 1,800                        |
| 8/19                                   | 1800 | Sunny                 | Clear                | 16.0 10.0                   | 10.3          | —   | 105                                   | —                     | 1,300                        |
| 8/29                                   | 1200 | Overcast              | Clear                | 12.0 8.5                    | 13.8          | 5.2 | 130                                   | —                     | 3,500                        |
| 9/9                                    | 1650 | Overcast              | Clear                | 13.5 9.9                    | 10.8          | 5.5 | 78                                    | —                     | 2,500                        |
| 9/23                                   | 1340 | Overcast              | Clear                | 12.5 7.3                    | 10.8          | 5.6 | 70                                    | —                     | 20                           |
| <u>Chase Creek (27N 05W 12BCC)</u>     |      |                       |                      |                             |               |     |                                       |                       |                              |
| 7/16                                   | 1130 | Sunny                 | Clear                | 23.0 16.0                   | 12.8          | 7.0 | —                                     | —                     | —                            |
| 8/6                                    | 1330 | Sunny                 | Clear                | 21.0 17.0                   | 9.6           | 6.0 | 60                                    | —                     | 10,000                       |
| 8/13                                   | 0905 | Sunny                 | Clear                | 15.0 13.0                   | 8.6           | 6.0 | 78                                    | —                     | 5,000                        |
| 8/21                                   | 1400 | Sunny                 | Clear                | 21.0 18.0                   | 8.1           | —   | 50                                    | —                     | 5,000                        |
| 8/29                                   | 1800 | Overcast              | Clear                | 19.0 14.5                   | 8.9           | 6.5 | 48                                    | —                     | —                            |
| 9/9                                    | 1935 | Overcast              | Clear                | 11.6 11.2                   | 10.8          | 5.5 | 45                                    | —                     | —                            |
| 9/23                                   | 1800 | Overcast              | Clear                | 10.5 11.2                   | 7.5           | 5.0 | 52                                    | —                     | —                            |



Appendix II Table 2. Water quality data and juvenile salmon surveys in sloughs and clearwater tributaries of the Susitna River between the Chulitna River and Portage Creek, Devils Canyon Project, 1977 (continued).

| Date  | Time | Weather<br>Conditions | Water<br>Conditions | Temperature °C |      | D.O.<br>(PPM) | pH  | Specific<br>Conductance<br>(µMHOS/CM) | Gage<br>Height<br>(M) | Number of<br>Fry<br>Observed |
|---|------|-----------------------|---------------------|----------------|------|---------------|-----|---------------------------------------|-----------------------|------------------------------|
| <u>Lane Creek (28N 05W 12DAA)</u>           |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/13  | 1450 | Sunny                 | Clear               | 18.0           | 11.0 | 9.6           | 7.7 | —                                     | —                     | —                            |
| 7/27  | 1535 | Sunny                 | Clear               | 24.5           | 12.0 | 10.9          | 8.0 | 60                                    | —                     | —                            |
| 8/4   | 1420 | Sunny                 | Clear               | 17.0           | 11.0 | 10.4          | 5.4 | 60                                    | —                     | —                            |
| 8/11  | 2000 | Sunny                 | Clear               | 17.0           | 11.0 | 10.0          | 6.2 | 90                                    | —                     | —                            |
| 8/19  | 1900 | Sunny                 | Clear               | 18.0           | 12.0 | 9.0           | —   | 90                                    | —                     | —                            |
| 8/29  | 1430 | Overcast              | Clear               | 24.0           | 10.5 | 10.7          | 6.0 | 62                                    | —                     | —                            |
| 9/9   | 1730 | Overcast              | Clear               | 13.0           | 9.0  | 11.4          | —   | 99                                    | —                     | —                            |
| 9/23  | 1520 | Overcast              | Clear               | 14.0           | 6.2  | 10.6          | 5.0 | 75                                    | —                     | —                            |
| <u>Fourth of July Creek (30N 03W 03DAC)</u> |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/29  | 1140 | Clear                 | Clear               | 23.0           | 15.0 | 9.0           | 7.3 | 30                                    | —                     | —                            |
| 8/3   | 1300 | Clear                 | Clear               | 22.0           | 16.0 | 9.0           | 7.4 | 125                                   | —                     | 5,000                        |
| 8/11  | 0945 | Clear                 | Clear               | 14.0           | 13.0 | 9.5           | 7.1 | 50                                    | —                     | —                            |
| 8/19  | 1030 | Clear                 | Clear               | 15.5           | 14.0 | 9.2           | —   | 45                                    | —                     | —                            |
| 8/26  | 1230 | Rain                  | Clear               | 12.0           | 12.0 | 6.6           | 8.0 | —                                     | —                     | 18                           |
| 8/28  | 2010 | Partly<br>Cloudy      | Clear               | 12.5           | 11.0 | 9.8           | 5.5 | 24                                    | —                     | —                            |
| 9/9   | 1120 | Cloudy                | Turbid              | 10.0           | 9.1  | 11.6          | 5.5 | 46                                    | —                     | —                            |
| 9/22  | 1330 | Clear                 | Turbid              | 9.0            | 7.0  | 11.7          | 5.6 | 31                                    | —                     | 7                            |
| <u>Gold Creek (31N 02W 20BAD)</u>           |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 6/14  | 2100 | Rain                  | Turbid              | 11.0           | 4.0  | 12.0          | 7.8 | 60                                    | —                     | —                            |
| 7/21  | 1200 | Partly<br>Cloudy      | Clear               | 23.0           | 10.0 | 10.0          | 7.8 | 160                                   | —                     | —                            |
| 8/17  | 1400 | Rain                  | Clear               | 16.5           | 11.0 | 12.0          | —   | 200                                   | —                     | —                            |
| 9/22  | 1030 | Clear                 | Clear               | —              | —    | —             | —   | —                                     | —                     | 28                           |
| <u>Indian River (31N 02W 09CDA)</u>         |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/29  | 1140 | Clear                 | Clear               | 20.0           | 12.0 | 11.0          | 7.1 | 50                                    | —                     | —                            |
| 8/18  | 1530 | Partly<br>Cloudy      | Clear               | 17.0           | 12.0 | 11.0          | 7.5 | 40                                    | —                     | 581                          |
| 8/28  | 1430 | Partly<br>Cloudy      | Clear               | 17.0           | 12.0 | 11.2          | 6.0 | 43                                    | —                     | —                            |
| 9/8   | 1300 | Cloudy                | Clear               | 10.0           | 7.8  | 10.0          | 5.9 | 40                                    | —                     | —                            |
| <u>Portage Creek (32N 01W 26CDB)</u>        |      |                       |                     |                |      |               |     |                                       |                       |                              |
| 7/7   | 1200 | Clear                 | Clear               | 27.0           | 10.0 | 14.0          | 7.5 | —                                     | —                     | —                            |
| 7/28  | 1645 | Clear                 | Clear               | 23.0           | 13.0 | 10.0          | 7.8 | 80                                    | —                     | —                            |
| 8/25  | 1200 | —                     | Clear               | —              | 11.0 | —             | —   | —                                     | —                     | 6                            |

Table 3. Thermograph set in Rabideux Creek, upper sub-area; daily maximum and minimum water temperature, Devils Canyon Project, 1977.

| Temp. °C |      |      | Temp. °C |      |      | Temp. °C |      |      | Temp. °C |      |      | Temp. °C |      |      |
|----------|------|------|----------|------|------|----------|------|------|----------|------|------|----------|------|------|
| Date     | Max. | Min. | Date     | Max. | Min. | Date     | Max. | Min. | Date     | Max. | Min. | Date     | Max. | Min. |
| 5/25     | 8.2  | 8.2  | 6/27     | 14.7 | 14.3 | 7/30     | 16.0 | 15.6 | 9/1      | 11.0 | 10.9 | 10/5     | 4.0  | 3.0  |
| 26       | 10.0 | 8.2  | 28       | 14.3 | 14.2 | 31       | 15.6 | 15.3 | 2        | 10.9 | 10.9 | 6        | 3.9  | 3.5  |
| 27       | 10.0 | 9.8  | 29       | 15.5 | 14.2 | 8/1      | 15.2 | 14.8 | 3        | 10.8 | 10.7 | 7        | 4.0  | 3.5  |
| 28       | 11.7 | 10.0 | 30       | 15.2 | 14.1 | 2        | 14.8 | 14.7 | 4        | 10.7 | 10.0 | 8        | 4.0  | 3.9  |
| 29       | 10.8 | 9.2  | 7/1      | 15.6 | 14.1 | 3        | 16.0 | 14.8 | 5        | 10.0 | 10.0 | 9        | 4.1  | 4.0  |
| 30       | 10.1 | 9.2  | 2        | 14.8 | 13.7 | 4        | 15.8 | 15.4 | 6        | 10.0 | 10.0 | 10       | 4.5  | 4.1  |
| 31       | 11.7 | 10.1 | 3        | 14.4 | 13.0 | 5        | 15.4 | 14.6 | 7        | 10.0 | 10.0 | 11       | 5.5  | 4.5  |
| 6/1      | 12.6 | 11.3 | 4        | 13.1 | 13.0 | 6        | 15.3 | 14.6 | 8        | 10.0 | 10.0 | 12       | 5.7  | 5.5  |
| 2        | 14.1 | 12.6 | 5        | 13.2 | 13.0 | 7        | 15.2 | 15.0 | 9        | 10.0 | 10.0 | 13       | 5.7  | 3.7  |
| 3        | 14.8 | 13.0 | 6        | 13.9 | 13.0 | 8        | 15.1 | 14.9 | 10       | 10.0 | 10.0 | 14       | 4.0  | 3.9  |
| 4        | 13.3 | 10.8 | 7        | 15.3 | 13.2 | 9        | 16.0 | 15.1 | 11       | 10.0 | 10.0 | 15       | 4.0  | 3.9  |
| 5        | 10.8 | 10.4 | 8        | 16.3 | 13.2 | 10       | 15.9 | 15.0 | 12       | 10.0 | 10.0 | 16       | 4.0  | 3.9  |
| 6        | 10.7 | 10.4 | 9        | 17.2 | 14.2 | 11       | 15.0 | 14.0 | 13       | 9.9  | 9.9  | 17       | 4.0  | 3.8  |
| 7        | 11.0 | 10.8 | 10       | 17.9 | 14.5 | 12       | 14.0 | 13.7 | 14       | 9.9  | 9.9  | 18       | 3.8  | 3.2  |
| 8        | 12.3 | 10.0 | 11       | 18.8 | 15.1 | 13       | 14.8 | 13.8 | 15       | 9.9  | 8.8  | 19       | 3.2  | 2.5  |
| 9        | 12.8 | 12.3 | 12       | 18.8 | 15.0 | 14       | 14.7 | 14.7 | 16       | 8.8  | 8.3  | 20       | 2.5  | 2.2  |
| 10       | 13.6 | 12.8 | 13       | 16.0 | 15.0 | 15       | 14.7 | 14.7 | 17       | 8.3  | 8.3  | 21       | 2.2  | 1.8  |
| 11       | 13.6 | 13.6 | 14       | 15.5 | 15.0 | 16       | 14.7 | 14.7 | 18       | 8.3  | 8.3  | 22       | 1.8  | 1.7  |
| 12       | 13.6 | 13.6 | 15       | 15.0 | 14.0 | 17       | 14.7 | 14.5 | 19       | 8.3  | 8.3  | 23       | 1.8  | 1.7  |
| 13       | 13.6 | 13.6 | 16       | 17.0 | 14.0 | 18       | 14.5 | 14.4 | 20       | 8.3  | 7.7  |          |      |      |
| 14       | 14.4 | 13.6 | 17       | 16.8 | 14.0 | 19       | 14.6 | 14.4 | 21       | 7.7  | 7.7  |          |      |      |
| 15       | 14.7 | 14.5 | 18       | 16.5 | 14.0 | 20       | 15.5 | 14.4 | 22       | 7.7  | 7.5  |          |      |      |
| 16       | 14.8 | 14.8 | 19       | 15.8 | 13.9 | 21       | 15.5 | 14.3 | 23       | 7.5  | 7.2  |          |      |      |
| 17       | 14.8 | 14.5 | 20       | 14.8 | 13.9 | 22       | 15.5 | 14.5 | 24       | 7.2  | 6.7  |          |      |      |
| 18       | 14.5 | 13.3 | 21       | 14.9 | 14.7 | 23       | 14.8 | 13.5 | 25       | N/A  | N/A  |          |      |      |
| 19       | 13.3 | 13.2 | 22       | 15.2 | 13.7 | 24       | 14.0 | 13.9 | 26       | N/A  | N/A  |          |      |      |
| 20       | 14.2 | 13.5 | 23       | 15.3 | 13.0 | 25       | 13.9 | 13.7 | 27       | 7.5  | 7.5  |          |      |      |
| 21       | 14.2 | 14.0 | 24       | 16.0 | 13.0 | 26       | 13.7 | 13.0 | 28       | 7.5  | 7.2  |          |      |      |
| 22       | 14.0 | 13.6 | 25       | 15.3 | 14.4 | 27       | 13.0 | 12.4 | 29       | 7.2  | 6.8  |          |      |      |
| 23       | 13.3 | 13.0 | 26       | 15.3 | 14.4 | 28       | 12.4 | 11.5 | 30       | 6.8  | 5.5  |          |      |      |
| 24       | 14.4 | 13.0 | 27       | 15.3 | 14.3 | 29       | 11.3 | 11.3 | 10/1     | 5.5  | 3.0  |          |      |      |
| 25       | 13.9 | 13.8 | 28       | 16.4 | 14.3 | 30       | 11.3 | 11.0 | 2        | 3.0  | 2.4  |          |      |      |
| 26       | 14.9 | 13.8 | 29       | 16.0 | 15.6 | 31       | 11.0 | 11.0 | 3        | 2.5  | 2.4  |          |      |      |
|          |      |      |          |      |      |          |      |      | 4        | 3.0  | 2.5  |          |      |      |

## Appendix II

Table 4. Thermograph set in Montana Creek, upper sub-area; daily maximum and minimum water temperature, Devils Canyon Project, 1977.

| Temp. °C |      |      | Temp. °C |      |      | Temp. °C |      |      | Temp. °C   |      |      | Temp. °C |      |      |
|----------|------|------|----------|------|------|----------|------|------|------------|------|------|----------|------|------|
| Date     | Max. | Min. | Date     | Max. | Min. | Date     | Max. | Min. | Date       | Max. | Min. | Date     | Max. | Min. |
| 5/25     | 3.0  | 2.8  | 6/27     | 10.1 | 9.5  | 8/17     | 13.2 | 12.5 | 9/19       | 6.0  | 5.5  | 11/1     | 0.2  | 0.1  |
| 26       | 5.1  | 3.2  | 28       | 10.1 | 9.3  | 18       | 13.2 | 12.0 | 20         | 5.8  | 5.5  | 2        | 0.1  | 0.1  |
| 27       | 5.2  | 2.9  | 29       | 10.5 | 10.0 | 19       | 13.8 | 12.3 | 21         | 6.5  | 5.6  | 3        | 0.1  | 0.0  |
| 28       | 6.5  | 4.7  | 30       | 10.3 | 10.1 | 20       | 13.7 | 12.6 | 22         | 6.8  | 5.0  | 4        | 0.0  | 0.0  |
| 29       | 3.9  | 3.0  | 7/ 1     | 11.1 | 9.8  | 21       | 13.6 | 12.6 | 23         | 5.6  | 4.7  | 5        | 0.0  | 0.0  |
| 30       | 4.9  | 3.0  | 2        | 11.1 | 10.5 | 22       | 13.3 | 12.7 | 9/24-10/24 | N/A  | N/A  | 6        | 0.0  | 0.0  |
| 31       | 5.8  | 4.0  | 3        | 10.5 | 10.0 | 23       | 13.7 | 13.0 | 10/ 5      | 3.4  | 3.0  |          |      |      |
| 6/ 1     | 5.3  | 4.0  | 4        | 10.5 | 10.1 | 24       | 13.2 | 12.7 | 6          | 3.5  | 3.3  |          |      |      |
| 2        | 5.8  | 4.0  | 5        | 10.0 | 10.0 | 25       | 12.7 | 11.3 | 7          | 4.0  | 3.6  |          |      |      |
| 3        | 6.9  | 4.0  | 7/24     | N/A  | N/A  | 26       | 11.6 | 11.0 | 8          | 4.1  | 4.0  |          |      |      |
| 4        | 4.5  | 4.1  | 25       | 14.0 | 12.3 | 27       | 12.0 | 10.5 | 9          | 4.4  | 4.2  |          |      |      |
| 5        | 4.7  | 4.1  | 26       | 14.0 | 11.7 | 28       | 11.0 | 9.8  | 10         | 4.5  | 4.5  |          |      |      |
| 6        | 5.1  | 4.5  | 27       | 14.8 | 12.7 | 29       | 10.6 | 9.9  | 11         | 4.5  | 3.2  |          |      |      |
| 7        | 5.5  | 5.0  | 28       | 15.0 | 13.7 | 30       | 10.6 | 9.8  | 12         | 3.8  | 3.2  |          |      |      |
| 8        | 7.0  | 5.1  | 29       | 13.8 | 12.7 | 31       | 10.2 | 9.8  | 13         | 3.9  | 3.4  |          |      |      |
| 9        | 6.5  | 6.1  | 30       | 13.8 | 11.0 | 9/ 1     | 10.2 | 9.2  | 14         | 3.4  | 3.0  |          |      |      |
| 10       | 7.8  | 6.1  | 31       | 12.8 | 11.3 | 2        | 10.2 | 9.8  | 15         | 3.0  | 2.3  |          |      |      |
| 11       | 7.8  | 7.2  | 8/ 1     | 13.7 | 11.0 | 3        | 11.9 | 9.6  | 16         | 2.3  | 1.1  |          |      |      |
| 12       | 7.0  | 6.7  | 2        | 14.0 | 12.8 | 4        | 10.1 | 9.4  | 17         | 1.1  | 0.8  |          |      |      |
| 13       | 7.6  | 6.4  | 3        | 12.8 | 12.1 | 5        | 10.0 | 9.8  | 18         | 0.8  | 0.7  |          |      |      |
| 14       | 8.3  | 7.2  | 4        | 12.2 | 11.1 | 6        | 9.8  | 8.4  | 19         | 0.8  | 0.8  |          |      |      |
| 15       | 8.0  | 7.7  | 5        | 12.2 | 11.8 | 7        | 8.4  | 8.4  | 20         | 0.9  | 0.2  |          |      |      |
| 16       | 7.7  | 7.5  | 6        | 11.0 | 10.5 | 8        | 8.7  | 8.4  | 21         | 0.5  | 0.1  |          |      |      |
| 17       | 8.0  | 7.7  | 7        | 12.8 | 10.8 | 9        | 9.0  | 8.8  | 22         | 0.8  | 0.5  |          |      |      |
| 18       | 7.8  | 7.0  | 8        | 13.2 | 12.5 | 10       | 8.9  | 8.9  | 23         | 0.6  | 0.5  |          |      |      |
| 19       | 8.7  | 6.9  | 9        | 12.8 | 11.7 | 11       | 9.0  | 8.6  | 24         | 0.6  | 0.4  |          |      |      |
| 20       | 9.8  | 8.3  | 10       | 12.5 | 11.5 | 12       | 8.6  | 7.7  | 25         | 0.6  | 0.5  |          |      |      |
| 21       | 9.0  | 9.0  | 11       | 13.0 | 11.8 | 13       | 7.7  | 7.7  | 26         | 0.8  | 0.5  |          |      |      |
| 22       | 9.0  | 8.8  | 12       | 13.5 | 12.7 | 14       | 7.9  | 7.5  | 27         | 0.9  | 0.7  |          |      |      |
| 23       | 9.0  | 8.5  | 13       | 13.0 | 12.3 | 15       | 7.5  | 6.6  | 28         | 0.8  | 0.8  |          |      |      |
| 24       | 10.9 | 8.7  | 14       | 12.7 | 12.2 | 16       | 7.2  | 6.3  | 29         | 0.9  | 0.8  |          |      |      |
| 25       | 10.7 | 9.5  | 15       | 12.9 | 12.0 | 17       | 6.8  | 6.3  | 30         | 0.9  | 0.5  |          |      |      |
| 26       | 11.0 | 9.5  | 16       | 13.7 | 12.1 | 18       | 6.8  | 5.7  | 31         | 0.5  | 0.3  |          |      |      |

Appendix II Table 5.

Water chemistry data, Rabideux Creek, Devils Canyon Project, 1977.

| <u>Date</u>     | <u>D.O.</u><br><u>(mg/l)</u> | <u>pH</u> | <u>Hardness</u><br><u>(mg/l)</u> | <u>Alkalinity</u><br><u>(mg/l)</u> |
|-----------------|------------------------------|-----------|----------------------------------|------------------------------------|
| Upper Sub Area  |                              |           |                                  |                                    |
| 5/25            | 11                           | 6.6       | 17                               | 17                                 |
| 6/7             | 12                           | 7.3       | 34                               | 17                                 |
| 6/16            | 8                            | 7.0       | 34                               | 34                                 |
| 6/30            | 7                            | 7.3       | 51                               | 51                                 |
| 7/13            | 6                            | 6.5       | 51                               | 51                                 |
| 7/26            | 8                            | 7.0       | 51                               | 51                                 |
| 8/8             | 7                            | 7.0       | 51                               | 51                                 |
| 8/23            | 6                            | 6.8       | 51                               | 51                                 |
| 9/15            | 8                            | 6.8       | 34                               | 17                                 |
| 9/27            | 10                           | 6.8       | 34                               | 17                                 |
| 10/12           | 9                            | 6.8       | 34                               | 17                                 |
| 10/27           | 11                           | 7.2       | 34                               | 17                                 |
| Middle Sub Area |                              |           |                                  |                                    |
| 5/25            | 11                           | 7.0       | 34                               | 34                                 |
| 6/7             | 11                           | 7.3       | 34                               | 17                                 |
| 6/16            | 9                            | 7.3       | 51                               | 51                                 |
| 6/30            | 9                            | 7.3       | 51                               | 51                                 |
| 7/13            | 9                            | 7.7       | 51                               | 51                                 |
| 7/26            | 8                            | 7.5       | 68                               | 51                                 |
| 8/8             | 8                            | 7.3       | 68                               | 68                                 |
| 8/23            | 8                            | 7.3       | 68                               | 68                                 |
| 9/15            | 9                            | 7.3       | 34                               | 34                                 |
| 9/28            | 10                           | 7.3       | 34                               | 17                                 |
| 10/12           | 10                           | 7.2       | 51                               | 34                                 |
| 10/27           | 10                           | 7.2       | 51                               | 51                                 |
| Lower Sub Area  |                              |           |                                  |                                    |
| 5/25            | 11                           | 7.2       | 17                               | 17                                 |
| 6/8             | 10                           | 7.5       | 34                               | 34                                 |
| 6/30            | 9                            | 7.3       | 51                               | 51                                 |
| 7/13            | 10                           | 7.7       | 51                               | 68                                 |
| 7/26            | 9                            | 7.7       | 51                               | 68                                 |
| 8/8             | 10                           | 7.3       | 68                               | 68                                 |
| 8/23            | 9                            | 7.3       | 68                               | 68                                 |
| 9/15            | 9                            | 7.3       | 34                               | 34                                 |
| 9/29            | 9                            | 7.3       | 34                               | 34                                 |
| 10/12           | 11                           | 7.2       | 34                               | 34                                 |
| 10/27           | 11                           | 7.2       | 34                               | 34                                 |

Appendix II Table 6.

Water chemistry data, Montana Creek, Devils Canyon Project, 1977.

| <u>Date</u>     | <u>D.O.<br/>(mg/l)</u> | <u>pH</u> | <u>Hardness<br/>(mg /l)</u> | <u>Alkalinity<br/>(mg/l)</u> |
|-----------------|------------------------|-----------|-----------------------------|------------------------------|
| Upper Sub Area  |                        |           |                             |                              |
| 6/7             | 11                     | 7.3       | 34                          | 17                           |
| 6/29            | 9                      | 7.7       | 34                          | 34                           |
| 7/13            | 9                      | 7.7       | 34                          | 34                           |
| 7/26            | 9                      | 7.3       | 34                          | 17                           |
| 8/10            | 9                      | 7.3       | 34                          | 34                           |
| 8/22            | 9                      | 7.3       | 34                          | 34                           |
| 9/13            | 10                     | 7.3       | 34                          | 34                           |
| 9/28            | 10                     | 7.3       | 17                          | 17                           |
| 10/11           | 10                     | 7.3       | 34                          | 34                           |
| 10/26           | 13                     | 7.3       | 34                          | 34                           |
| Middle Sub Area |                        |           |                             |                              |
| 6/7             | 12                     | 7.3       | 34                          | 17                           |
| 6/29            | 9                      | 7.3       | 34                          | 34                           |
| 7/13            | 8                      | 6.8       | 34                          | 34                           |
| 8/10            | 9                      | 7.3       | 34                          | 34                           |
| 8/22            | 9                      | 7.3       | 34                          | 34                           |
| 9/13            | 10                     | 7.3       | 34                          | 34                           |
| 10/11           | 9                      | 7.3       | 34                          | 34                           |
| 10/28           | 10                     | 7.4       | 34                          | 34                           |
| 11/11           | 10                     | 7.3       | 34                          | 34                           |
| Lower Sub Area  |                        |           |                             |                              |
| 6/7             | 11                     | 7.3       | 34                          | 17                           |
| 6/29            | 10                     | 7.3       | 34                          | 34                           |
| 7/13            | 9                      | 7.7       | 34                          | 34                           |
| 7/26            | 9                      | 7.3       | 34                          | 17                           |
| 8/10            | 9                      | 7.3       | 34                          | 34                           |
| 8/22            | 9                      | 7.3       | 34                          | 34                           |
| 9/2             | 9                      | 7.6       | 34                          | 34                           |
| 9/13            | 10                     | 7.3       | 34                          | 34                           |
| 9/30            | 10                     | 7.3       | 34                          | 17                           |
| 10/11           | 10                     | 7.3       | 34                          | 34                           |
| 10/28           | 10                     | 7.4       | 34                          | 34                           |

Appendix II Table 7.

UNITED STATES DEPARTMENT OF INTERIOR - GEOLOGICAL SURVEY  
624941149221500 - SUSITNA R AB PORTAGE C NR GOLD CREEK AK

PROCESS DATE 02/11  
DISTRICT CODE 02

## WATER QUALITY DATA, WATER YEAR OCTOBER 1976 TO SEPTEMBER 1977

| DATE         | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>.004 MM<br>(70338) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>.008 MM<br>(70339) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>.016 MM<br>(70340) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>.031 MM<br>(70341) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>.062 MM<br>(70342) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>.125 MM<br>(70343) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>.250 MM<br>(70344) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>.500 MM<br>(70345) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>1.00 MM<br>(70346) | TOTAL<br>MERCURY<br>(HG)<br>(UG/L)<br>(71900) | SUS-<br>PENDEO<br>SEDI-<br>MENT<br>(MG/L)<br>(80154) |
|--------------|--|--|--|--|--|--|--|--|--|---|--|
| JUN<br>14... | 3  | 6  | 11   | 22   | 39   | 59   | 82   | 98   | 100  | .4  | 956  |

| DIS-<br>SOLVED<br>MAN-<br>GANESE<br>(MN)<br>(UG/L)<br>(01056) | TOTAL<br>MOLYB-<br>DENUM<br>(MO)<br>(UG/L)<br>(01062) | TOTAL<br>NICKEL<br>(NI)<br>(UG/L)<br>(01067) | TOTAL<br>SILVER<br>(AG)<br>(UG/L)<br>(01077) | TOTAL<br>ZINC<br>(ZN)<br>(UG/L)<br>(01092) | TOTAL<br>ALUM-<br>INUM<br>(AL)<br>(UG/L)<br>(01105) | TOTAL<br>SELE-<br>NIUM<br>(SE)<br>(UG/L)<br>(01147) | DIS-<br>SOLVED<br>SOLIDS<br>(RESI-<br>DUE AT<br>180 C)<br>(MG/L)<br>(70300) | DIS-<br>SOLVED<br>SOLIDS<br>(SUM OF<br>CONSTI-<br>TUENTS)<br>(MG/L)<br>(70301) | DIS-<br>SOLVED<br>SOLIDS<br>(TONS<br>PER<br>DAY)<br>(70302) | DIS-<br>SOLVED<br>SOLIDS<br>(TONS<br>PER<br>AC-FT)<br>(70303) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>.002 MM<br>(70337) |
|---|---|--|--|--|---|---|---|--|---|---|--|
| 20  | 1   | 100  | <10  | 110  | 9500  | 1   | 50  | 51   | 6750  | .07   | 2  |

Appendix II Table 7.

UNITED STATES DEPARTMENT OF INTERIOR - GEOLOGICAL SURVEY  
624941149221500 - SUSITNA R AB PORTAGE C NR GOLD CREEK AK

PROCESS DATE 02/10/70  
DISTRICT CODE 02

WATER QUALITY DATA, WATER YEAR OCTOBER 1976 TO SEPTEMBER 1977

| DATE         | TIME | TYPE | TEMPER-<br>ATURE<br>(DEG C)<br>(00010)                                   | INSTAN-<br>TANEOUS<br>DIS-<br>CHARGE<br>(CFS)<br>(00061)                | COLOR<br>(PLAT-<br>INUM-<br>COBALT<br>UNITS)<br>(00080)                 | SPE-<br>CIFIC<br>CON-<br>DUCT-<br>ANCE<br>(MICRO-<br>MHOS)<br>(00095) | DIS-<br>SOLVED<br>OXYGEN<br>(MG/L)<br>(00300)                | PH<br>(UNITS)<br>(00400)                                    | CARBON<br>DIOXIDE<br>(CO2)<br>(MG/L)<br>(00405)                    | ALKA-<br>LITY<br>AS<br>CACO3<br>(MG/L)<br>(00410)     | BICAR-<br>BONATE<br>(HCO3)<br>(MG/L)<br>(00440)    | CAR-<br>BONATE<br>(CO3)<br>(MG/L)<br>(00445)                      |  |  |
|--------------|------|------|--|---|---|---|--|---|--|---|--|---|--|--|
| JUN<br>14... | 1130 | 2    |  | 0.0E50000   | 55  | 80  | 12.8   | 7.2   | 3.7  | 30  | 37   | 0   |  |  |
|              |      |      | DIS-<br>SOLVED<br>NITRITE<br>PLUS<br>NITRATE<br>(N)<br>(MG/L)<br>(00631) | DIS-<br>SOLVED<br>ORTHO<br>PHOS-<br>PHATE<br>(PO4)<br>(MG/L)<br>(00660) | DIS-<br>SOLVED<br>ORTHO.<br>PHOS-<br>PHORUS<br>(P)<br>(MG/L)<br>(00671) | HARD-<br>NESS<br>(CA, MG)<br>(MG/L)<br>(00900)                        | NON-<br>CAR-<br>BONATE<br>HARD-<br>NESS<br>(MG/L)<br>(00902) | DIS-<br>SOLVED<br>CAL-<br>CIUM<br>(CA)<br>(MG/L)<br>(00915) | DIS-<br>SOLVED<br>MAG-<br>NE-<br>SIUM<br>(MG)<br>(MG/L)<br>(00925) | DIS-<br>SOLVED<br>SODIUM<br>(NA)<br>(MG/L)<br>(00930) | SODIUM<br>AD-<br>SORP-<br>TION<br>RATIO<br>(00931) | DIS-<br>SOLVED<br>PO-<br>TAS-<br>SIUM<br>(K)<br>(MG/L)<br>(00935) | DIS-<br>SOLVED<br>CHLO-<br>RIDE<br>(CL)<br>(MG/L)<br>(00940) |  |
|              |      |      | .06  | .06   | .02   | 36  | 5  | 12  | 1.4  | 2.6   | .2   | 13  | 1.2  | 4.9  |
|              |      |      | DIS-<br>SOLVED<br>SULFATE<br>(SO4)<br>(MG/L)<br>(00945)                  | DIS-<br>SOLVED<br>FLUO-<br>RIDE<br>(F)<br>(MG/L)<br>(00950)             | DIS-<br>SOLVED<br>SILICA<br>(SI02)<br>(MG/L)<br>(00955)                 | TOTAL<br>ARSENIC<br>(AS)<br>(UG/L)<br>(01002)                         | TOTAL<br>BARIUM<br>(BA)<br>(UG/L)<br>(01007)                 | TOTAL<br>CAD-<br>MIUM<br>(CD)<br>(UG/L)<br>(01027)          | TOTAL<br>CHRO-<br>MIUM<br>(CR)<br>(UG/L)<br>(01034)                | TOTAL<br>COPPER<br>(CU)<br>(UG/L)<br>(01042)          | TOTAL<br>IRON<br>(FE)<br>(UG/L)<br>(01045)         | DIS-<br>SOLVED<br>IRON<br>(FE)<br>(UG/L)<br>(01046)               | TOTAL<br>LEAD<br>(PB)<br>(UG/L)<br>(01051)                   | TOTAL<br>MAN-<br>GANESE<br>(MN)<br>(UG/L)<br>(01055) |
|              |      |      | 5.0  | .0  | 5.2   | 11  | 100  | <10   | 30   | 200   | 15000  | 170   | 1200   | 280  |

# Appendix II Table 7.

## UNITED STATES DEPARTMENT OF THE INTERIOR GEOLOGICAL SURVEY CENTRAL LABORATORY, DENVER, COLORADO

### WATER QUALITY ANALYSIS LAB ID # 291069 RECORD # 42798

SAMPLE LOCATION: SUSITNA R AB PORTAGE C NR GOLD CREEK AK  
STATION ID: 624941149221500 LAT.LONG.SEQ.: 624941 1492215 00  
DATE OF COLLECTION: BEGIN--771005 END-- TIME--1300  
STATE CODE: 02 COUNTY CODE: 170 PROJECT IDENTIFICATION: 470200350  
DATA TYPE: 2 SOURCE: SURFACE WATER GEOLOGIC UNIT:  
COMMENTS:

FIELD VALUE USED FOR BICARB & CARBONATE.

|                     |        |     |                       |        |      |
|---------------------|--------|-----|-----------------------|--------|------|
| AIR TEMP (DEG C)    |        | 8.0 | MOLYBDENUM TOTAL      | UG/L   | 4    |
| ALK.TOT (AS CaCO3)  | MG/L   | 45  | NICKEL TOTAL          | UG/L < | 50   |
| ALUMINUM TOTAL      | UG/L   | 410 | NO2+NO3 AS N DISS     | MG/L   | 0.07 |
| ARSENIC TOTAL       | UG/L   | 1   | OXYGEN DISSOLVED      | MG/L   | 13.9 |
| BARIUM TOTAL        | UG/L   | 200 | PH FIELD              |        | 7.2  |
| BICARBONATE         | MG/L   | 55  | PHOS ORTHO DIS AS P   | MG/L   | 0.00 |
| CAOMIUM TOTAL       | UG/L < | 10  | PHOSPHATE DIS ORTHO   | MG/L   | 0.00 |
| CALCIUM DISS        | MG/L   | 20  | POTASSIUM DISS        | MG/L   | 1.6  |
| CARBONATE           | MG/L   | 0   | RESIDUE DIS CALC SUM  | MG/L   | 98   |
| CHLORIDE DISS       | MG/L   | 17  | RESIDUE DIS TON/AFT   |        | 0.12 |
| CHROMIUM TOTAL      | UG/L   | 10  | RESIDUE DIS TON/DAY   |        | 1530 |
| COLOR               |        | 6   | RESIDUE DIS 180C      | MG/L   | 87   |
| COPPER TOTAL        | UG/L   | 20  | SAR                   |        | 0.4  |
| FLUORIDE DISS       | MG/L   | 0.1 | SELENIUM TOTAL        | UG/L   | 0    |
| HARDNESS NONCARB    | MG/L   | 17  | SILICA DISSOLVED      | MG/L   | 8.7  |
| HARDNESS TOTAL      | MG/L   | 62  | SILVER TOTAL          | UG/L < | 10   |
| IRON DISSOLVED      | UG/L   | 40  | SODIUM DISS           | MG/L   | 7.1  |
| IRON TOTAL          | UG/L   | 730 | SODIUM PERCENT        |        | 19   |
| LEAD TOTAL          | UG/L < | 100 | SP. CONDUCTANCE FLD   |        | 165  |
| MAGNESIUM DISS      | MG/L   | 3.0 | SP. CONDUCTANCE LAB   |        | 170  |
| MANGANESE DISSOLVED | UG/L   | 0   | STREAMFLOW (CFS)-INST | 557    | 6500 |
| MANGANESE TOTAL     | UG/L   | 40  | SULFATE DISS          | MG/L   | 13   |
| MERCURY TOTAL       | UG/L   | 0.0 | WATER TEMP (DEG C)    |        | 2.0  |
|                     |        |     | ZINC TOTAL            | UG/L   | 20   |

| CATIONS        |        |         | ANIONS         |        |         |
|----------------|--------|---------|----------------|--------|---------|
|                | (MG/L) | (MEQ/L) |                | (MG/L) | (MEQ/L) |
| CALCIUM DISS   | 20     | 0.996   | BICARBONATE    | 55     | 0.90    |
| MAGNESIUM DISS | 3.0    | 0.247   | CARBONATE      | 0      | 0.00    |
| POTASSIUM DISS | 1.6    | 0.041   | CHLORIDE DISS  | 17     | 0.48    |
| SODIUM DISS    | 7.1    | 0.309   | FLUORIDE DISS  | 0.1    | 0.00    |
|                |        |         | SULFATE DISS   | 13     | 0.27    |
|                |        |         | NO2-NO3 AS N D | 0.07   | 0.00    |
| TOTAL          |        | 1.545   | TOTAL          |        | 1.66    |

PERCENT DIFFERENCE = -2.07



Appendix II Table 7.

UNITED STATES DEPARTMENT OF INTERIOR - GEOLOGICAL SURVEY  
15292000 - SUSITNA RIVER AT GOLD C AK

PROCESS DATE 03  
DISTRICT CODE 02

## WATER QUALITY DATA, WATER YEAR OCTOBER 1976 TO SEPTEMBER 1977

| DATE         | TIME | TYPE | TEMPER-<br>ATURE<br>(DEG C)<br>(00010) | AIR<br>TEMPER-<br>ATURE<br>(DEG C)<br>(00020) | SURFACE<br>AREA<br>(SQUARE<br>MILES)<br>(00049) | INSTAN-<br>TANEOUS<br>DIS-<br>CHARGE<br>(CFS)<br>(00061) | COLOR<br>(PLAT-<br>NUM-<br>COBALT<br>UNITS)<br>(00080) | SPE-<br>CIFIC<br>CON-<br>DUCT-<br>ANCE<br>(MICRO-<br>MHOS)<br>(00095) | DIS-<br>SOLVED<br>OXYGEN<br>(MG/L)<br>(00300) | PH<br>(UNITS)<br>(00400) | CARBON<br>DIOXIDE<br>(CO2)<br>(MG/L)<br>(00405) | ALKA-<br>LITY<br>AS<br>CACO3<br>(MG/L)<br>(00410) |
|--------------|------|------|--|---|---|--|--|---|---|--------------------------|---|---|
| OCT<br>01... | 1400 | 2    | 3.5                                    | --  | 6160  | 5330   | --   | --  | --  | --                       | --  | --  |
| MAY<br>10... | 1830 | 2    | 1.0                                    | --  | 6160  | 3730   | --   | --  | --  | --                       | --  | --  |
| 18...        | 1000 | 2    | --                                     | --  | 6160  | 14200  | --   | --  | --  | --                       | --  | --  |
| JUN<br>14... | 1630 | 2    | 8.0                                    | 17.0  | 6160  | 52000  | 45   | 102   | 12.2  | 6.8                      | 7.1   | 23  |
| JUL<br>28... | 1730 | 2    | 14.0                                   | --  | 6160  | 21000  | --   | --  | --  | --                       | --  | --  |
| AUG<br>10... | 1430 | 2    | 12.0                                   | --  | 6160  | 20000  | 25   | 163   | 11.1  | 7.9                      | 1.1   | 45  |

| DATE         | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>.062 MM<br>(70342) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>.125 MM<br>(70343) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>.250 MM<br>(70344) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>.500 MM<br>(70345) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>1.00 MM<br>(70346) | TOTAL<br>MERCURY<br>(UG/L)<br>(71900) | ELEV.<br>OF LAND<br>SURFACE<br>DATUM<br>(FT.<br>ABOVE<br>HSL)<br>(72000) | SUS-<br>PENDE<br>SED-<br>MENT<br>CHARGE<br>(MG/L)<br>(80154) | SUS-<br>PENDE<br>SED-<br>MENT<br>DIS-<br>CHARGE<br>(T/DAY)<br>(80155) |
|--------------|--|--|--|--|--|---------------------------------------|--|--|---|
| OCT<br>01... | --   | --   | --   | --   | --   | --                                    | 677  | 10   | 144   |
| MAY<br>10... | --   | --   | --   | --   | --   | --                                    | 677  | 120  | 1210  |
| 18...        | --   | --   | --   | --   | --   | --                                    | 677  | 1110   | 42600   |
| JUN<br>14... | 40   | 62   | 84   | 97   | 100  | .2                                    | 677  | 915  | 128000  |
| JUL<br>28... | --   | --   | --   | --   | --   | --                                    | 677  | 394  | 22300   |
| AUG<br>10... | --   | --   | --   | --   | --   | .3                                    | 677  | 656  | 35400   |

Appendix II Table 7.

UNITED STATES DEPARTMENT OF INTERIOR - GEOLOGICAL SURVEY  
15292000 - SUSITNA RIVER AT GOLD C AK

PROCESS DATE 0  
DISTRICT CODE 02

## WATER QUALITY DATA, WATER YEAR OCTOBER 1976 TO SEPTEMBER 1977

| DATE         | BICAR-<br>BONATE<br>(HCO3)<br>(MG/L)<br>(00440) | CAR-<br>BONATE<br>(CO3)<br>(MG/L)<br>(00445) | DIS-<br>SOLVED<br>NITRITE<br>PLUS<br>NITRATE<br>(N)<br>(MG/L)<br>(00631) | DIS-<br>SOLVED<br>ORTHO<br>PHOS-<br>PHATE<br>(P04)<br>(MG/L)<br>(00660) | DIS-<br>SOLVED<br>ORTHO.<br>PHOS-<br>PHORUS<br>(P)<br>(MG/L)<br>(00671) | HARD-<br>NESS<br>(CA, MG)<br>(MG/L)<br>(00900) | NON-<br>CAR-<br>BONATE<br>HARD-<br>NESS<br>(MG/L)<br>(00902) | DIS-<br>SOLVED<br>CAL-<br>CIUM<br>(CA)<br>(MG/L)<br>(00915) | DIS-<br>SOLVED<br>MAG-<br>NE-<br>SIUM<br>(MG)<br>(MG/L)<br>(00925) | DIS-<br>SOLVED<br>SODIUM<br>(NA)<br>(MG/L)<br>(00930) | SODIUM<br>AD-<br>SORP-<br>TION<br>RATIO<br>(00931) | PERCENT<br>SODIUM<br>(00932) |
|--------------|---|--|--|---|---|--|--|---|--|---|--|------------------------------|
| OCT<br>01... | --  | --   | --   | --  | --  | --   | --   | --  | --   | --  | --   | --                           |
| MAY<br>10... | --  | --   | --   | --  | --  | --   | --   | --  | --   | --  | --   | --                           |
| 18...        | --  | --   | --   | --  | --  | --   | --   | --  | --   | --  | --   | --                           |
| JUN<br>14... | 28  | 0  | .06  | .06   | .02   | 36   | 13   | 12  | 1.4  | 2.4   | .2   | 12                           |
| JUL<br>28... | --  | --   | --   | --  | --  | --   | --   | --  | --   | --  | --   | --                           |
| AUG<br>10... | 55  | 0  |  | .06   | .02   | 75   | 30   | 23  | 4.3  | 3.6   | .2   | 9                            |

| DATE         | DIS-<br>SOLVED<br>PO-<br>TAS-<br>SIUM<br>(K)<br>(MG/L)<br>(00935) | DIS-<br>SOLVED<br>CHLO-<br>RIDE<br>(CL)<br>(MG/L)<br>(00940) | DIS-<br>SOLVED<br>SULFATE<br>(SO4)<br>(MG/L)<br>(00945) | DIS-<br>SOLVED<br>FLUO-<br>RIDE<br>(F)<br>(MG/L)<br>(00950) | DIS-<br>SOLVED<br>SILICA<br>(SI02)<br>(MG/L)<br>(00955) | TOTAL<br>ARSENIC<br>(AS)<br>(UG/L)<br>(01002) | TOTAL<br>BARIUM<br>(BA)<br>(UG/L)<br>(01007) | TOTAL<br>CAD-<br>MIUM<br>(CD)<br>(UG/L)<br>(01027) | TOTAL<br>CHRO-<br>MIUM<br>(CR)<br>(UG/L)<br>(01034) | TOTAL<br>COPPER<br>(CU)<br>(UG/L)<br>(01042) | TOTAL<br>IRON<br>(FE)<br>(UG/L)<br>(01045) | DIS-<br>SOLVED<br>IRON<br>(FE)<br>(UG/L)<br>(01046) |
|--------------|---|--|---|---|---|---|--|--|---|--|--|---|
| OCT<br>01... | --  | --   | --  | --  | --  | --  | --   | --   | --  | --   | --   | --  |
| MAY<br>10... | --  | --   | --  | --  | --  | --  | --   | --   | --  | --   | --   | --  |
| 18...        | --  | --   | --  | --  | --  | --  | --   | --   | --  | --   | --   | --  |
| JUN<br>14... | 1.1   | 15   | 4.7   | .1  | 5.2   | 5   | 0  | <10  | 30  | 50   | 20000                                      | 100   |
| JUL<br>28... | --  | --   | --  | --  | --  | --  | --   | --   | --  | --   | --   | --  |
| AUG<br>10... | 4.4   | 5.4  | 14  | .1  | 4.9   | 12  | 500  | <10  | 40  | 50   | 18000                                      | --  |

Appendix II Table 7.

UNITED STATES DEPARTMENT OF INTERIOR - GEOLOGICAL SURVEY  
15292000 - SUSITNA RIVER AT GOLD C AKPROCESS DATE 11-  
DISTRICT CODE 02

## WATER QUALITY DATA, WATER YEAR OCTOBER 1976 TO SEPTEMBER 1977

| DATE         | DIS-<br>SOLVED<br>SOLIDS<br>(TONS<br>PER<br>AO-FT)<br>(70303) | SUS.<br>SED.<br>SIEVE<br>DIAM.<br>% FINER<br>THAN<br>.062 MM<br>(70331) | SUS.<br>SED.<br>SIEVE<br>DIAM.<br>% FINER<br>THAN<br>.125 MM<br>(70332) | SUS.<br>SED.<br>SIEVE<br>DIAM.<br>% FINER<br>THAN<br>.250 MM<br>(70333) | SUS.<br>SED.<br>SIEVE<br>DIAM.<br>% FINER<br>THAN<br>.500 MM<br>(70334) | SUS.<br>SED.<br>SIEVE<br>DIAM.<br>% FINER<br>THAN<br>1.00 MM<br>(70335) | SUS.<br>SED.<br>SIEVE<br>DIAM.<br>% FINER<br>THAN<br>2.00 MM<br>(70336) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>.002 MM<br>(70337) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>.004 MM<br>(70338) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>.008 MM<br>(70339) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>.016 MM<br>(70340) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>.031 MM<br>(70341) |
|--------------|---|---|---|---|---|---|---|--|--|--|--|--|
| OCT<br>01... | --  | --  | --  | --  | --  | --  | --  | --   | --   | --   | --   | --   |
| MAY<br>10... | --  | 44  | 64  | 87  | 99  | 100   | --  | --   | --   | --   | --   | --   |
| 18...        | --  | 63  | 76  | 90  | 99  | 100   | --  | 7  | 9  | 17   | 27   | 44   |
| JUN<br>14... | .09   | --  | --  | --  | --  | --  | --  | 2  | 4  | 6  | 11   | 22   |
| JUL<br>28... | --  | 70  | 80  | 92  | 99  | 100   | --  | 14   | 19   | 29   | 44   | 54   |
| AUG<br>10... | .10   | 65  | 74  | 86  | 95  | 98  | 99  | 13   | 19   | 27   | 39   | 52   |

| DATE         | TOTAL<br>LEAD<br>(PB)<br>(UG/L)<br>(01051) | TOTAL<br>MAN-<br>GANESE<br>(MN)<br>(UG/L)<br>(01055) | DIS-<br>SOLVED<br>MAN-<br>GANESE<br>(MN)<br>(UG/L)<br>(01056) | TOTAL<br>MOLYB-<br>DENUM<br>(MO)<br>(UG/L)<br>(01062) | TOTAL<br>NICKEL<br>(NI)<br>(UG/L)<br>(01067) | TOTAL<br>SILVER<br>(AG)<br>(UG/L)<br>(01077) | TOTAL<br>ZINC<br>(ZN)<br>(UG/L)<br>(01092) | TOTAL<br>ALUM-<br>INUM<br>(AL)<br>(UG/L)<br>(01105) | TOTAL<br>SELE-<br>NIUM<br>(SE)<br>(UG/L)<br>(01147) | DIS-<br>SOLVED<br>SOLIDS<br>(RESI-<br>DUE AT<br>180 C)<br>(MG/L)<br>(70300) | DIS-<br>SOLVED<br>SOLIDS<br>(SUM OF<br>CONSTITUENTS)<br>(MG/L)<br>(70301) | DIS-<br>SOLVED<br>SOLIDS<br>(TONS<br>PER<br>DAY)<br>(70302) |
|--------------|--|--|---|---|--|--|--|---|---|---|---|---|
| OCT<br>01... | --   | --   | --  | --  | --   | --   | --   | --  | --  | --  | --  | --  |
| MAY<br>10... | --   | --   | --  | --  | --   | --   | --   | --  | --  | --  | --  | --  |
| 18...        | --   | --   | --  | --  | --   | --   | --   | --  | --  | --  | --  | --  |
| JUN<br>14... | 100  | 370  | 40  | 1   | 50   | <10  | 80   | 14000   | 0   | 63  | 56  | 8850  |
| JUL<br>28... | --   | --   | --  | --  | --   | --   | --   | --  | --  | --  | --  | --  |
| AUG<br>10... | <100                                       | 320  | 180   | 0   | <50  | <10  | 80   | 13000   | 1   | 76  | 130   | 4100  |

Appendix II Table 7.

UNITED STATES DEPARTMENT OF THE INTERIOR  
GEOLOGICAL SURVEY  
CENTRAL LABORATORY, DENVER, COLORADO

WATER QUALITY ANALYSIS  
LAB ID # 291068 RECORD # 42795

SAMPLE LOCATION: SUSITNA RIVER AT GOLD C AK  
STATION ID: 15292000 LAT.LONG.SEQ.: 524604 1494128 00  
DATE OF COLLECTION: BEGIN--771004 END-- TIME--1600  
STATE CODE: 02 COUNTY CODE: 170 PROJECT IDENTIFICATION: 470200350  
DATA TYPE: 2 SOURCE: SURFACE WATER GEOLOGIC UNIT:  
COMMENTS:  
FIELD VALUE USED FOR BICARB & CARBONATE.

|                     |        |     |                      |        |      |
|---------------------|--------|-----|----------------------|--------|------|
| AIR TEMP (DEG C)    |        | 9.0 | MOLYBDENUM TOTAL     | UG/L   | 7    |
| ALK+TOT (AS CaCO3)  | MG/L   | 37  | NICKEL TOTAL         | UG/L < | 50   |
| ALUMINUM TOTAL      | UG/L   | 500 | NO2+NO3 AS N DISS    | MG/L   | 0.11 |
| ARSENIC TOTAL       | UG/L   | 1   | OXYGEN DISSOLVED     | MG/L   | 12.6 |
| BARIUM TOTAL        | UG/L   | 200 | PH FIELD             |        | 7.4  |
| BICARBONATE         | MG/L   | 45  | PHOS ORTHO DIS AS P  | MG/L   | 0.00 |
| CADMIUM TOTAL       | UG/L < | 10  | PHOSPHATE DIS ORTHO  | MG/L   | 0.00 |
| CALCIUM DISS        | MG/L   | 18  | POTASSIUM DISS       | MG/L   | 1.4  |
| CARBONATE           | MG/L   | 0   | RESIDUE DIS CALC SUM | MG/L   | 85   |
| CHLORIDE DISS       | MG/L   | 11  | RESIDUE DIS TON/AFT  |        | 0.10 |
| CHROMIUM TOTAL      | UG/L   | 0   | RESIDUE DIS TON/DAY  |        | 1740 |
| COLOR               |        | 12  | RESIDUE DIS 180C     | MG/L   | 76   |
| COPPER TOTAL        | UG/L   | 50  | SAR                  |        | 0.4  |
| FLUORIDE DISS       | MG/L   | 0.1 | SELENIUM TOTAL       | UG/L   | 0    |
| HARDNESS NONCARB    | MG/L   | 20  | SILICA DISSOLVED     | MG/L   | 8.6  |
| HARDNESS TOTAL      | MG/L   | 57  | SILVER TOTAL         | UG/L < | 10   |
| IRON DISSOLVED      | UG/L   | 40  | SODIUM DISS          | MG/L   | 6.5  |
| IRON TOTAL          | UG/L   | 850 | SODIUM PERCENT       |        | 19   |
| LEAD TOTAL          | UG/L < | 100 | SP. CONDUCTANCE FLD  |        | 150  |
| MAGNESIUM DISS      | MG/L   | 3.0 | SP. CONDUCTANCE LAB  |        | 154  |
| MANGANESE DISSOLVED | UG/L   | 0   | STREAMFLOW(CFS)-INST |        | 8500 |
| MANGANESE TOTAL     | UG/L   | 20  | SULFATE DISS         | MG/L   | 14   |
| MERCURY TOTAL       | UG/L   | 0.2 | WATER TEMP (DEG C)   |        | 3.5  |
|                     |        |     | ZINC TOTAL           | UG/L   | 30   |

| CATIONS        |        |         | ANIONS         |        |         |
|----------------|--------|---------|----------------|--------|---------|
|                | (MG/L) | (MEQ/L) |                | (MG/L) | (MEQ/L) |
| CALCIUM DISS   | 18     | 0.899   | BICARBONATE    | 45     | 0.73    |
| MAGNESIUM DISS | 3.0    | 0.247   | CARBONATE      | 0      | 0.00    |
| POTASSIUM DISS | 1.4    | 0.036   | CHLORIDE DISS  | 11     | 0.31    |
| SODIUM DISS    | 6.5    | 0.283   | FLUORIDE DISS  | 0.1    | 0.00    |
|                |        |         | SULFATE DISS   | 14     | 0.29    |
|                |        |         | NO2+NO3 AS N O | 0.11   | 0.00    |
| TOTAL          |        | 1.464   | TOTAL          |        | 1.35    |

PERCENT DIFFERENCE = 3.94

Appendix II Table 7.

UNITED STATES DEPARTMENT OF INTERIOR - GEOLOGICAL SURVEY  
15292780 - SUSITNA R AT SUNSHINE AK

PROCESS DATE 03/16/78  
DISTRICT CODE 02

WATER QUALITY DATA, WATER YEAR OCTOBER 1976 TO SEPTEMBER 1977

| DATE         | TIME | TYPE | TEMPER-<br>ATURE<br>(DEG C)<br>(00010)                                   | INSTAN-<br>TANEOUS<br>DIS-<br>CHARGE<br>(CFS)<br>(00061)                 | COLOR<br>(PLAT-<br>INUM-<br>COBALT<br>UNITS)<br>(00080)                 | SPE-<br>CIFIC<br>CON-<br>DUCT-<br>ANCE<br>(MICRO-<br>MHOS)<br>(00095) | DIS-<br>SOLVED<br>OXYGEN<br>(MG/L)<br>(00300)                | PH<br>(UNITS)<br>(00400)                                    | CARBON<br>DIOXIDE<br>(CO2)<br>(MG/L)<br>(00405)                    | ALKA-<br>LITY<br>AS<br>CACO3<br>(MG/L)<br>(00410)     | BICAR-<br>BONATE<br>(HCO3)<br>(MG/L)<br>(00440)    | CAR-<br>BONATE<br>(CO3)<br>(MG/L)<br>(00445) |   |  |
|--------------|------|------|--|--|---|---|--|---|--|---|--|--|---|--|
| JUN<br>15... | 1630 | 2    | 8.0  | 115000   | 100   | 100   | 12.0   | 7.1   | 3.9  | 25  | 31   | 0  |   |  |
| AUG<br>10... | 2100 | 2    | 12.0   | 70000  | 25  | 112   | 10.6   | 7.6   | 2.1  | 43  | 52   | 0  |   |  |
| DATE         |      |      | DIS-<br>SOLVED<br>NITRITE<br>PLUS<br>NITRATE<br>(N)<br>(MG/L)<br>(00631) | DIS-<br>SOLVED<br>ORTHO-<br>PHOS-<br>PHATE<br>(PO4)<br>(MG/L)<br>(00660) | DIS-<br>SOLVED<br>ORTHO.<br>PHOS-<br>PHORUS<br>(P)<br>(MG/L)<br>(00671) | HARD-<br>NESS<br>(CA, MG)<br>(MG/L)<br>(00900)                        | NON-<br>CAR-<br>BONATE<br>HARD-<br>NESS<br>(MG/L)<br>(00902) | DIS-<br>SOLVED<br>CAL-<br>CIUM<br>(CA)<br>(MG/L)<br>(00915) | DIS-<br>SOLVED<br>MAG-<br>NE-<br>SIUM<br>(MG)<br>(MG/L)<br>(00925) | DIS-<br>SOLVED<br>SODIUM<br>(NA)<br>(MG/L)<br>(00930) | SODIUM<br>AD-<br>SORP-<br>TION<br>RATIO<br>(00931) | PERCENT<br>SODIUM<br>(00932)                 | DIS-<br>SOLVED<br>PO-<br>TAS-<br>SIUM<br>(K)<br>(MG/L)<br>(00935) | DIS-<br>SOLVED<br>CHLO-<br>RIDE<br>(CL)<br>(MG/L)<br>(00940) |
| JUN<br>15... |      |      | .13  | .12  | .04   | 37  | 11   | 12  | 1.6  | 1.9   | .1   | 10   | 1.1   | 7.3  |
| AUG<br>10... |      |      |  | .06  | .02   | 72  | 29   | 23  | 3.5  | 2.7   | .1   | 7  | 2.8   | 2.7  |

Appendix II Table 7.

UNITED STATES DEPARTMENT OF INTERIOR - GEOLOGICAL SURVEY  
15292780 - SUSITNA R AT SUNSHINE AK

PROCESS DATE 03/1  
DISTRICT CODE 02

## WATER QUALITY DATA, WATER YEAR OCTOBER 1976 TO SEPTEMBER 1977

| DATE      | SUS.<br>SED.<br>SIEVE<br>DIAM.<br>% FINER<br>THAN<br>(70332) | SUS.<br>SED.<br>SIEVE<br>DIAM.<br>% FINER<br>THAN<br>(70333) | SUS.<br>SED.<br>SIEVE<br>DIAM.<br>% FINER<br>THAN<br>(70334) | SUS.<br>SED.<br>SIEVE<br>DIAM.<br>% FINER<br>THAN<br>(70335) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>(70337) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>(70338) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>(70339) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>(70340) | SUS.<br>SED.<br>FALL<br>DIAM.<br>% FINER<br>THAN<br>(70341) | TOTAL<br>MERCURY<br>(UG/L)<br>(71900) | SUS-<br>PENDE<br>SED-<br>MENT<br>(MG/L)<br>(80154) |
|-----------|--|--|--|--|---|---|---|---|---|---------------------------------------|--|
| JUN 15... | 64   | 84   | 97   | 100  | 7   | 9   | 15  | 22  | 33  | .2                                    | 1630   |
| AUG 10... | 83   | 92   | 99   | 100  | 16  | 28  | 40  | 51  | 64  | .1                                    | 900  |

| DATE      | DIS-<br>SOLVED<br>MAN-<br>GANESE<br>(MN)<br>(UG/L)<br>(01056) | TOTAL<br>MOLYB-<br>DENUM<br>(MO)<br>(UG/L)<br>(01062) | TOTAL<br>NICKEL<br>(NI)<br>(UG/L)<br>(01067) | TOTAL<br>SILVER<br>(AG)<br>(UG/L)<br>(01077) | TOTAL<br>ZINC<br>(ZN)<br>(UG/L)<br>(01092) | TOTAL<br>ALUM-<br>INUM<br>(AL)<br>(UG/L)<br>(01105) | TOTAL<br>SELE-<br>NIUM<br>(SE)<br>(UG/L)<br>(01147) | DIS-<br>SOLVED<br>SOLIDS<br>(RESI-<br>DUE AT<br>180 C)<br>(MG/L)<br>(70300) | DIS-<br>SOLVED<br>SOLIDS<br>(SUM OF<br>CONSTI-<br>TUENTS)<br>(MG/L)<br>(70301) | DIS-<br>SOLVED<br>SOLIDS<br>(TONS<br>PER<br>DAY)<br>(70302) | DIS-<br>SOLVED<br>SOLIDS<br>(TONS<br>PER<br>AC-FT)<br>(70303) | SUS.<br>SED.<br>SIEVE<br>DIAM.<br>% FINER<br>THAN<br>(70331) |
|-----------|---|---|--|--|--|---|---|---|--|---|---|--|
| JUN 15... | 20  | 1   | 100  | <10  | 150  | 22000   | 1   | 56  | 51   | 17400   | .08   | 46   |
| AUG 10... | --  | 0   | <50  | <10  | 120  | 15000   | 0   | --  | 102  | --  | .09   | 76   |

| DATE      | DIS-<br>SOLVED<br>SULFATE<br>(SO4)<br>(MG/L)<br>(00945) | DIS-<br>SOLVED<br>FLUO-<br>RIDE<br>(F)<br>(MG/L)<br>(00950) | DIS-<br>SOLVED<br>SILICA<br>(SiO2)<br>(MG/L)<br>(00955) | TOTAL<br>ARSENIC<br>(AS)<br>(UG/L)<br>(01002) | TOTAL<br>BARIUM<br>(BA)<br>(UG/L)<br>(01007) | TOTAL<br>CAD-<br>MIUM<br>(CD)<br>(UG/L)<br>(01027) | TOTAL<br>CHRO-<br>MIUM<br>(CR)<br>(UG/L)<br>(01034) | TOTAL<br>COPPER<br>(CU)<br>(UG/L)<br>(01042) | TOTAL<br>IRON<br>(FE)<br>(UG/L)<br>(01045) | DIS-<br>SOLVED<br>IRON<br>(FE)<br>(UG/L)<br>(01046) | TOTAL<br>LEAD<br>(PB)<br>(UG/L)<br>(01051) | TOTAL<br>MAN-<br>GANESE<br>(MN)<br>(UG/L)<br>(01055) |
|-----------|---|---|---|---|--|--|---|--|--|---|--|--|
| JUN 15... | 5.7   | .1  | 4.9   | 25  | 200  | <10  | 60  | 200  | 37000                                      | 180   | 300  | 790  |
| AUG 10... | 11  | .1  | 4.0   | 24  | 500  | <10  | 40  | 40   | 24000                                      | --  | <100                                       | 540  |

Appendix II Table 7.

UNITED STATES DEPARTMENT OF THE INTERIOR  
GEOLOGICAL SURVEY  
CENTRAL LABORATORY, DENVER, COLORADO

WATER QUALITY ANALYSIS  
LAB ID # 291070 RECORD # 42801

SAMPLE LOCATION: SUSITNA R AT SUNSHINE AK  
STATION ID: 15292780 LAT.LONG.SED.: 621035 1501018 00  
DATE OF COLLECTION: BEGIN--771004 END-- TIME--0915  
STATE CODE: 02 COUNTY CODE: 170 PROJECT IDENTIFICATION: 470200350  
DATA TYPE: 2 SOURCE: SURFACE WATER GEOLOGIC UNIT:  
COMMENTS:  
FIELD VALUE USED FOR BICARB & CARBONATE.

|                     |        |      |                        |        |       |
|---------------------|--------|------|------------------------|--------|-------|
| AIR TEMP (DEG C)    |        | 6.8  | MOLYBDENUM TOTAL       | UG/L   | 3     |
| ALK.TOT (AS CaCO3)  | MG/L   | 43   | NICKEL TOTAL           | UG/L < | 50    |
| ALUMINUM TOTAL      | UG/L   | 2200 | NO2+NO3 AS N DISS      | MG/L   | 0.23  |
| ARSENIC TOTAL       | UG/L   | 3    | OXYGEN DISSOLVED       | MG/L   | 12.8  |
| BARIUM TOTAL        | UG/L   | 200  | PH FIELD               |        | 7.4   |
| BICARBONATE         | MG/L   | 52   | PHOS ORTHO DIS AS P    | MG/L   | 0.00  |
| CAESIUM TOTAL       | UG/L < | 10   | PHOSPHATE DIS ORTHO    | MG/L   | 0.00  |
| CALCIUM DISS        | MG/L   | 17   | POTASSIUM DISS         | MG/L   | 1.2   |
| CARBONATE           | MG/L   | 0    | RESIDUE DIS CALC SUM   | MG/L   | 78    |
| CHLORIDE DISS       | MG/L   | 6.0  | RESIDUE DIS ION/AFT    |        | 0.09  |
| CHROMIUM TOTAL      | UG/L   | 10   | RESIDUE DIS ION/DAY    |        | 4880  |
| COLOR               |        | 8    | RESIDUE DIS 180C       | MG/L   | 66    |
| COPPER TOTAL        | UG/L   | 20   | SAR                    |        | 0.3   |
| FLUORIDE DISS       | MG/L   | 0.1  | SELENIUM TOTAL         | UG/L   | 0     |
| HARDNESS NONCARB    | MG/L   | 12   | SILICA DISSOLVED       | MG/L   | 7.4   |
| HARDNESS TOTAL      | MG/L   | 55   | SILVER TOTAL           | UG/L < | 10    |
| IRON DISSOLVED      | UG/L   | 60   | SODIUM DISS            | MG/L   | 4.4   |
| IRON TOTAL          | UG/L   | 3700 | SODIUM PERCENT         |        | 15    |
| LEAD TOTAL          | UG/L < | 100  | SP. CONDUCTANCE FLD    |        | 135   |
| MAGNESIUM DISS      | MG/L   | 3.0  | SP. CONDUCTANCE LAB    |        | 133   |
| MANGANESE DISSOLVED | UG/L   | 0    | STREAMFLOW (CFS) -INST |        | 27400 |
| MANGANESE TOTAL     | UG/L   | 100  | SULFATE DISS           | MG/L   | 12    |
| MERCURY TOTAL       | UG/L   | 0.0  | WATER TEMP (DEG C)     |        | 4.0   |
|                     |        |      | ZINC TOTAL             | UG/L   | 30    |

| CATIONS        |        |         | ANIONS         |        |         |
|----------------|--------|---------|----------------|--------|---------|
|                | (MG/L) | (MEQ/L) |                | (MG/L) | (MEQ/L) |
| CALCIUM DISS   | 17     | 0.849   | BICARBONATE    | 52     | 0.853   |
| MAGNESIUM DISS | 3.0    | 0.247   | CARBONATE      | 0      | 0.000   |
| POTASSIUM DISS | 1.2    | 0.031   | CHLORIDE DISS  | 6.0    | 0.170   |
| SODIUM DISS    | 4.4    | 0.192   | FLUORIDE DISS  | 0.1    | 0.006   |
|                |        |         | SULFATE DISS   | 12     | 0.250   |
|                |        |         | NO2+NO3 AS N D | 0.23   | 0.017   |
| TOTAL          |        | 1.317   | TOTAL          |        | 1.293   |

PERCENT DIFFERENCE = 0.92

### APPENDIX III

The following appendix is a synopsis of ADF&G's recommended plan of study for the aquatic environment. Yearly objectives and cost estimates are included.



## AQUATIC BIOLOGY STUDIES

### Introduction

The proposed Susitna River hydroelectric project will have various impacts on both the indigenous organisms and the natural conditions within the aquatic environment. The fish populations are the most obvious aspects of the aquatic community where impacts will be evident due to their economic and recreational importance to the people of Alaska and the nation. However, studies cannot be limited to the fishery resource alone due to the complex interrelationships between all biological components of, and within, the aquatic community and the associated habitat. The majority of the impacts on fish species will likely result from changes in the natural regimes of the river rather than direct impacts on the fish in the vicinity. Primary areas of concern are reduction of stream flow, increased turbidity levels during winter months, and thermal and chemical pollution. Alterations of the habitat may adversely affect the existing fish populations and render portions of the drainage either nonproductive or unavailable in future years.

Baseline fisheries inventories were conducted by the Alaska Department of Fish and Game in the upper Susitna River during the 1974-1977 field seasons. The Susitna Basin is the major coho, pink, chum, and chinook salmon production area within the Cook Inlet area. Although total escapement estimates have not been derived for this system, it is probably the second or third largest sockeye salmon production area within Cook Inlet. Grayling, rainbow trout, Dolly Varden, lake trout, whitefish, and burbot are among the important resident fish species present.

The interrelationships within the biological communities and between their habitats must be clearly defined to protect the aquatic ecosystem from losses incurred by hydroelectric development. The effects on the anadromous and resident fish populations are of primary concern to the Alaska Department of Fish and Game fisheries divisions. Aquatic studies will, therefore, concentrate on the seasonal life histories and critical habitat requirements of fish species present.

Seasonal fluctuations in the physiochemical composition of the aquatic habitat are apparently the major factors influencing distribution of fish within the upper drainage. Any alterations resulting from hydroelectric project activities which restrict or reduce quality or quantity of required habitat will also reduce fish populations and associated members of the aquatic community.

Each aquatic community is dependent upon various river mechanics to provide the necessary habitat for its existence. Depth, width, and velocity of the stream flow determine the quality and quantity of habitat available to aquatic organisms. High water discharge associated with spring and summer run-off results in important physical habitat alterations. Unregulated flowing waters dilute and transport natural and man-generated pollutants. A flushing or scouring action occurs during periods of high flows and removes deposited sediments and fines, resulting in an annual cleansing of the river bottom. This is an important factor in rivers like the Susitna

which transport large amounts of glacial silt. Deposition of sediment without the annual scouring could change the overall productivity of the river, eventually suffocating some of the aquatic organisms.

Individual study proposals are designed to provide the necessary background information to enable proper evaluation of impacts. Six general objectives have been outlined:

- 1) Determine the relative abundance and distribution of anadromous fish populations within the drainage.
- 2) Determine the distribution and abundance of selected resident fish populations.
- 3) Determine the seasonal habitat requirements of anadromous and resident fish species during each stage of their life histories.
- 4) Determine the economic, recreational, social, and aesthetic values of the existing resident and anadromous fish stocks and habitat.
- 5) Determine the impact the Devils Canyon project will have on the aquatic ecosystems and any required mitigation prior to construction approval.
- 6) Determine a long term plan of study, if the project is authorized, to monitor the impacts during and after project completion.

Fisheries and physiochemical sampling techniques and equipment for large rivers similar to the Susitna are in the early stages of development. Research and development must accompany the study to modify equipment and techniques to the habitat conditions of the specific environment to be evaluated.

The large drainage areas encompassed by the project are divided and categorized by location and activity. The three major study areas are:

- 1) The Susitna River basin between Denali Highway and Cook Inlet.
- 2) The proposed transmission line corridor and construction road drainage areas.
- 3) The Cook Inlet estuarine area.

All proposed studies are interrelated and have been coordinated to produce specific results. The elimination of any segment of a project will require revision of study plans. Investigations have been arbitrarily divided into anadromous and resident species studies. To insure precise and adequate aquatic data are collected each study is limited to a specific geographic area. A sufficient number of personnel must therefore be distributed throughout the study areas to insure a cross-section of habitat conditions are examined and movements of fish populations are monitored.

Title: Impact of the Proposed Devils Canyon-Watana Hydropower Projects On Anadromous Fish Populations Within the Susitna River Drainage.

Objectives: Determine the abundance and distribution of anadromous fish populations.

Determine the seasonal freshwater habitat requirements of adult and juvenile salmon, including spawning, incubation, rearing, and migration.

Background: The salmon stocks of the Susitna River drainage are major contributors to the Cook Inlet area fishery. Determining total escapement into this system is greatly complicated by the glacial conditions of the major streams and the enormity of the area. Management of the northern Cook Inlet salmon stocks has been difficult due to the mixed stock commercial fishery in Cook Inlet and the lack of adequate tools to provide accurate in season escapement estimates for the drainage.

The major hydroelectric project impacts on the anadromous fish species are expected to be due to changes in habitat. Alteration of the normal flow regimes and the physical and chemical water characteristics will probably be the most critical impacts. It is difficult at this time to determine the distance downstream from the proposed dams that changes will occur. Studies conducted by Townsend (1975) in the Peace River demonstrate that effects were observed 730 miles downstream from the Bennett Dam.

The Alaska Department of Fish and Game has conducted fisheries investigations in the area of proposed dam construction downstream since 1974. Emphasis has been on the inventory of adult and juvenile salmon stocks and habitat assessment. Current research investigations have concentrated on determining total escapement of salmon species into the Susitna drainage and intrasystem migrations of fry. Successful tag and recovery projects were operated in the lower river during 1975 and 1977 and the feasibility of sonar operation was tested in the mainstem Susitna River approximately 25 miles upstream from Cook Inlet during 1976.

Only through complete stock assessment will it be possible to determine what portion of the Susitna River anadromous fish runs will be affected by the project and determine the level of mitigative measures which will ultimately be required. It is essential to know what portion the affected stocks contribute to the total Susitna River salmon escapement in order to determine potential losses of fish populations and numbers. Economic values and relative importance can be determined after establishing this. Pink, chum, and chinook salmon are the dominant species utilizing the upper reaches of the drainage although sockeye and coho salmon are also observed.

#### Adults

Population estimates of salmon species utilizing the Susitna River above the Chulitna River confluence were estimated during the 1974, 1975, and 1977 field seasons based on tagging and subsequent recovery of fish. These studies indicate a portion of the salmon tagged are not destined to spawn above the tagging site, but rather below it. The importance and extent

of this milling behavior in the upper river areas requires definition. The alterations in flow and water quality in the mainstem river after project completion could significantly affect this behavior and consequently spawning success. Behavior modifications and disorientation of fish due to tagging and handling may have been a contributing factor.

Observations of spawning areas between the Chulitna and Susitna river confluence upstream to Portage Creek during fall surveys indicate that a reduction in flow to proposed post-construction levels would prevent access to many important spawning areas.

The degree of impact of reduced flows will be dependent on the total area affected. The distance affected downstream would depend partially on the contribution of the natural Susitna River flow regimes to that of each major tributary and the drainage as a whole.

Studies conducted during the late 1950's indicate that Cook Inlet salmon stocks are unable to ascend the Susitna River beyond Devils Canyon, the latter being a natural water velocity barrier to migration (U.S. Department of the Interior, 1957). Reports from local residents of salmon observations above Devils Canyon indicate that this should be investigated further.

#### Juveniles

Previous studies have defined important clearwater streams and spring fed sloughs within the Susitna River drainage which support juvenile anadromous fish species. Investigations have, however, concentrated primarily on summer rearing areas. Surveys indicate these populations are not static, but vary in abundance and distribution. Studies conducted during the winter of 1974-1975 revealed that juvenile anadromous species also utilize the mainstem Susitna River.

Data collected since 1974 provide only baseline information. Generalizations may be made, but sufficient information is not available to determine specific impacts of dam construction and operation on incubating and rearing anadromous species.

#### Adults

Procedures: Emphasis should be on determining total salmon escapement into the drainage, stock separation, and habitat evaluation. Types of sampling gear which can be utilized in the upper area of the river and catchability of adult salmon migrating upstream greatly affect the success of a tag and recovery program. Recent developments and improvements in sonar salmon counters are a viable option. A sonar counting system suitable for operation in the upper Susitna River would have to be designed and tested. Installation of weirs or counting towers to determine escapements would be feasible on most clearwater tributaries.

Commercial Fisheries Division will operate side-scanning sonar salmon counters in the lower Susitna River during 1978 as part of their ongoing studies. A salmon tag and recovery program to provide an alternate escapement estimate could be funded through Devils Canyon studies to provide additional data and supplement sonar escapement information. The duration of this project is dependent on correlation of population estimates and sonar counts. Data obtained from these studies would be correlated with population estimates in the upper Susitna River. Through these studies the importance of the Susitna River salmon stocks to the Cook Inlet area as a whole could be determined.

Evaluation of milling behavior of adult salmon in the upper Susitna River will require new sampling techniques. Obtaining escapement samples and marking them to determine migrational characteristics without causing some modification of normal behavior is difficult. Internal sonic transmitters may be utilized to evaluate this. The effectiveness of this type of tag in heavily silt laden waters would have to be tested. Recently developed stock separation techniques based on salmon scale characteristics may eventually enable researchers to assign unknown stocks to specific areas. This technique is still in the developmental research stage, but preliminary data indicate that samples obtained from Cook Inlet can be assigned to one of the three major salmon producing systems with  $\pm$  14 percent confidence. A large data base of scale characteristics from tributary systems would have to be established before analysis could be made.

Surveys and escapement sampling should be conducted in the proposed impoundment areas between the Denali Highway and Devils Canyon during periods of peak adult salmon abundance. Initial observations would be conducted by aerial surveys to document the presence or absence of adult salmon. Surveys would be done in conjunction with resident fish investigations. Data obtained would be utilized to determine necessary mitigation measures.

Water quality, quantity, and biological studies to predict the effects on spawning and migration habitat are described in the habitat study section.

### Juveniles

Year-round studies are required to determine complete juvenile salmon distribution and habitat utilization data.

Surveys of all rearing areas defined in previous studies should be continued. The distribution, species composition, and growth characteristics of juvenile salmonids should be monitored. Additional sampling equipment should be employed to assure representative samples are being collected. These include seines, minnow traps, small fyke traps, and dip nets. Fore-gut sample analysis should be continued and related to invertebrate studies. Winter sampling should be initiated on selected sloughs and clearwater tributaries that support significant populations of rearing fish during the summer and are also accessible during the winter months. Physio-chemical parameters of the aquatic habitat will be monitored during each survey.

The timing of migration of juvenile fish from sloughs and tributaries to the mainstem river and the extent of mainstem utilization should be documented. Factors which trigger the outmigration will be determined through habitat monitoring. These will include water temperature, ice cover, relative water levels, dissolved oxygen, pH, and conductivity. Fish samples will be collected primarily by traps. Coded wire tags and/or pigment dye marking may be effective methods of determining intrasystem migrations after initial documentation of this phenomenon.

The quantity and quality of water within the mainstem Susitna River will be monitored year round. Data will be obtained from U.S.G.S. gauging stations and at additional sites by field crews monitoring fry distribution. (See Habitat Section).

Schedule: Following is a preliminary schedule of anadromous fish project activities. The initiation of some segments of the studies will be dependent on testing of sampling equipment and delivery time required for more complex equipment, i.e., sonar counters.

The fiscal years (FY) outlined encompass the period of July 1 through June 30.

FY 79      Determine total salmon escapement estimate for the  
Susitna River drainage.

Determine total escapement in selected streams in the upper  
drainage.

Monitor abundance, distribution, characteristics, and  
habitat requirements of adult and juvenile salmonids.

Monitor physical, chemical and hydrological parameters of  
the mainstem Susitna River, sloughs, and clearwater  
tributaries.

Evaluate the feasibility of operation of various types of  
sampling gear for use in the upper river areas.

Begin building data base for stock separation studies.

FY 80      Continue salmon escapement estimates.

Continue fry and habitat studies.

Evaluate milling behavior of adult salmon.

Continue water quantity and quality monitoring.

Continue impoundment surveys, if salmon are observed  
during FY 79.

Continue stock separation studies and begin detailed  
analysis.

FY 81 Continue all FY 80 studies and revise programs as necessary.  
FY 82 Continue ongoing field projects (FY 81) and begin final analysis of projects.  
FY 83 Continue field monitoring and prepare final report.

Cost:

FY 79 \$909,800  
FY 80 \$592,700  
FY 81 \$592,700  
FY 82 \$592,700  
FY 83 \$592,700

Literature Cited:

Townsend, G.H. 1975. Impact of the Bennett Dam on the Peace-Athabasca Delta. J. Fish. Res. Board Can. Vol. 32 (1). pp. 171-176.

U. S. Dept. of the Interior. 1957. (Unpublished). Progress Report 1956 field investigation Devils Canyon Dam Site, Susitna River Basin. 15 pp.

Title: Impact of the Susitna Hydroelectric Project on Resident Fish Species

Objectives: Determine species present and distribution.

Determine seasonal abundance of selected populations.

Determine seasonal habitat requirements necessary to sustain the species present.

Background: The Alaska Department of Fish and Game has conducted limited fisheries investigations in the Susitna River and its tributaries, both upstream and downstream of the proposed dam sites and in lakes near the impoundment area. The general distribution of resident species was monitored and basic seasonal life history and habitat observations were conducted during portions of the spring, summer, fall, and winter seasons. Some resident species make major migrations from lake and tributary systems into the mainstem Susitna for purposes of overwintering. The importance of this intrasystem migration and the role of the mainstem Susitna River is not understood at this time. Surveys conducted between 1974 and 1977 document that a high quality sport fishery is provided by the Susitna River, its tributaries, and nearby lakes.

Procedure: Seasonal life history, distribution, population abundance, and habitat requirement investigations of selected resident fish species will be continued and expanded. These studies will be closely coordinated with the anadromous fish studies. Special attention will be given to those areas important to resident fish which may not coincide with anadromous fish habitat. The study area for resident fish investigations may be considerably greater, extending along the Susitna River from the mouth of the Tyone River to Cook Inlet, including tributaries bisected by transmission and road corridors.

Of particular importance in this study will be the determination of winter distribution, migrational and habitat requirements within areas subject to project impact. Studies will be made of the tributaries where resident fish predominately spawn and reside during the summer months, and the mainstem Susitna River where many of these same fish may winter. Emphasis will also be given to streams impacted by inundation. Human utilization of resident species will also be determined.

This study will be conducted in two parts, with results of the first two years of effort being compiled and analyzed for use in related studies and as a basis for determining areas where efforts should be concentrated during the remaining years of the study.

Due to difficulty in capturing fish from the Susitna River through the winter ice cover, high velocities and turbid water conditions in the summer, considerable equipment and sampling technique adaptations will be necessary. Boom and backpack electrofishing, side scanning sonar, sonar, angling, radio tags, anchor tags, coded wire tags, fyke nets, seines, gill nets, fixed traps, fish wheels, weirs, and ground surveys will be among the techniques to be employed.



Those elements of the physiochemical and trophic makeup of the existing natural habitat which will be analyzed are discussed under the Habitat Studies Section.

Schedule:

- FY 79      Organize Susitna River Basin study team and coordinate work schedule with other study teams where necessary.
- Establish base camps and begin fisheries inventory, seasonal life history, and associated habitat investigations.
- FY 80      Continue field activities and relocate various personnel as dictated by data which are generated. Areas of investigation include impoundment, transmission and road corridors, and downstream of Devils Canyon to Cook Inlet.
- FY 81      Continue field activities and relocate various personnel as dictated by data which are generated.
- FY 82      Continue field activities and relocate various personnel as dictated by data which are generated.
- Initiate report writing process.
- FY 83      Continue field activities and relocate various personnel as dictated by data which are generated, and integrate and summarize all data collected into final report.

Cost:

|       |           |
|-------|-----------|
| FY 79 | \$462,900 |
| FY 80 | \$416,600 |
| FY 81 | \$416,600 |
| FY 82 | \$416,600 |
| FY 83 | \$416,600 |

Title: Investigations of the Cook Inlet Estuarine Area and Potential Effects of Hydroelectric Development.

Objectives: Identify the fisheries resources of the lower Susitna River and the Cook Inlet estuary.

Determine the existing water quality and biological productivity of the lower Susitna River and the Cook Inlet estuary.

Determine the contribution and importance of the Susitna River to the Cook Inlet estuary.

Background: Cook Inlet is approximately 170 miles long and 60 miles wide at its mouth, with a total volume of  $1.7 \times 10^{13}$  feet<sup>3</sup>. It can be divided into two natural regions, a northern and southern portion, by a natural topographic feature, the East and West Forelands. The Susitna River and the major streams and rivers entering Knik Arm represent about 70-80 percent of the total freshwater entering the Inlet (Rosenberg, 1967).

Estuaries generally have exceptional usefulness in support of fisheries as rearing areas. It is generally a high food production area for primary consumers such as clams and other filter feeding organisms and the secondary and tertiary level consumers, including finfish and shellfish species. Migratory fishes such as salmon must pass through the estuarine area to reach their spawning grounds.

The estuary is, in many ways, the most complicated and variable of the aquatic ecosystems. Current and salinity shape the life of the estuary where the environment is neither fresh nor salt water. Estuarine currents result from the interaction of one-direction flow which varies with seasonal run-off, oscillating tides and the winds. The unique assemblages of organisms utilizing the estuarine habitat have evolved to survive these rigorous conditions.

Oceanographic data from the Cook Inlet estuarine area is limited. The extent to which juvenile and adult salmon species utilize this estuarine area is unknown. If natural flow regimes and water quality are altered by the hydroelectric project, adverse effects would possibly be observed within the Inlet. Baseline studies to determine existing physiochemical habitat conditions and biological productivity should be conducted. Parameters which need to be evaluated include: temperature, salinity, pH, nutrients, sedimentation processes, water stage and velocity, and biological activities.

Investigations of estuarine areas are more difficult than for river systems and will require elaborate equipment and use of large vessels.

Procedures: Baseline aquatic biology, and habitat studies and a thorough investigation of existing data available on the Cook Inlet area will be conducted prior to initiation of any comprehensive field investigations. This environmental data will provide an adequate data base for determining the direction and level of future field studies necessary to project the effects of the hydroelectric project on the estuarine ecosystem.

Schedule:

- FY 79      Conduct field research and analyze the data collected.
- Review and evaluate existing environmental data of the Cook Inlet area.
- Develop comprehensive study plan.
- FY 80      Activities will depend on FY 79 findings. Ongoing monitoring and previous studies may provide sufficient data. If not, additional field investigations will have to be initiated.

Cost:

- FY 79              \$75,000
- FY 80-83          Open. Will depend on FY 79 results. Overall allocation may have to be amended.

Literature Cited:

Rosenberg, D.H., S.C. Burrell, K.V. Matarajan, and D.W. Hook, 1967. Oceanography of Cook Inlet with special reference to the effluent from the Collier Carbon and Chemical Plant. Institute of Marine Science, University of Alaska. Report No. R67-5. 80 pp.

Title: Susitna River Basin Habitat Investigations

Objectives: Identify seasonal habitat characteristics associated with the Susitna River Basin anadromous and resident fisheries.

Define the complex interrelationships between the various components of the habitat.

Determine which habitat components are critical to the sustenance of the existing fisheries, and why.

Background: Maintenance of anadromous and resident fish populations within the Susitna River Basin will require a thorough understanding of their life sustaining habitat. Impacts by the hydroelectric project which alter or reduce the quantity or quality of the critical spawning, incubation, rearing, and migration habitat of these species will reduce or eliminate their populations. Major changes may take place in the biotic community with only a subtle change in the habitat.

Baseline physiochemical and biological aquatic habitat data were collected between 1974 and 1977 by the Alaska Department of Fish and Game at selected sites within the Susitna River drainage. The United States Geological Survey and other agencies have also monitored physiochemical parameters of the drainage.

Literature on the physiochemical and biological composition of aquatic habitat in lotic and lentic environments and its relationships to aquatic communities is also available.

Procedure: Personnel conducting seasonal fisheries life history investigations within the Susitna River Basin will concurrently collect the majority of the associated physiochemical field habitat data. In situ water velocity, width, depth, gradient, temperature, conductivity, pH and dissolved oxygen measurements will be collected with sophisticated electronic and mechanical instrumentation. Water samples will also be collected for laboratory analyses of basic metals, dissolved solids, total suspended solids, alkalinity, hardness, pH, conductivity, and total recoverable solids. Additional investigations by fisheries personnel will include water surface and sedimentation profiles. The U.S.G.S. will be contracted to install stream gauging stations at selected sites.

Biological habitat investigations will include primary productivity, benthos species composition and diversity, forage fish, pathological, and bioassay studies. Benthos, forage fish and fish pathology investigations will be integrated with fisheries life history studies. The remaining three will be conducted as individual studies.

To define the complex interrelationships of the dynamic habitat conditions of the Susitna River Basin it will be necessary to collect data over an extended period of time. Because of the precise measurements required, equipment for this investigation will be costly.

Schedule:

|       |   |
|-------|---|
| FY 79 | Organize field staff and procure equipment. Establish field camps, install equipment, and initiate field and office research. |
| FY 80 | Continue field and office research.   |
| FY 81 | Continue field and office research.   |
| FY 82 | Continue field and office research.   |
| FY 83 | Continue field and office studies, analyze data, and write report.  |

Cost: Personnel and their associated expenses are included in the fisheries investigations.

|       |           |
|-------|-----------|
| FY 79 | \$191,000 |
| FY 80 | \$149,000 |
| FY 81 | \$149,000 |
| FY 82 | \$149,000 |
| FY 83 | \$149,000 |

Title: Transmission Corridors, Access Road Corridor, and Construction Pad Sites Fisheries Investigations

Objectives: Identify all fishery resources within the four proposed transmission corridors, the access road corridor, and the construction pad sites.

Identify species present in these waters and determine seasonal presence.

Identify the habitat associated with these species.

Background: Four transmission corridor routes, one access road corridor, gravel and fill sites, and numerous building site pads are under consideration. The corridors will provide human access to previously inaccessible areas. This access will concentrate sportsman efforts in certain areas which may result in adverse impacts to aquatic life. Uncontrolled removal of gravel and fill for construction activities will also adversely affect the aquatic habitat. No hydroelectric related fishery investigations of these areas have been conducted. Other sources of fisheries data in these drainages are insufficient.

Procedures: Fishery resources, their seasonal presence and associated habitat will be identified within these areas. Ground surveys, fish trapping, fish marking, benthic species collection and physiochemical water quality measurement techniques will be conducted. Backpack electro-fishing, nets, traps, anchor and radio tags, electrophoresis instrumentation, weirs, benthic samplers, sophisticated water quality measurement devices, water quantity measurement equipment, and survey equipment are among the equipment which will be utilized.

Schedule:

- |       |  |
|-------|--|
| FY 79 | Organize corridor and building site study teams, procure equipment, and coordinate schedules with other study teams where necessary.           |
|       | Establish base camps and initiate fisheries resource identification, species identification, and seasonal presence and habitat investigations. |
| FY 80 | Continue field activities.   |
| FY 81 | Continue field activities and relocate various personnel as dictated by data and overall study findings.                                       |
| FY 82 | Continue field activities and relocate various personnel as dictated by data and overall study findings.                                       |
| FY 83 | Conduct concentrated studies if necessary and integrate and summarize all data collected.  |

Cost:

FY 79      \$130,500

FY 80      \$125,500

FY 81      \$125,500

FY 82      \$125,500

FY 83      \$125,500

Title: Existing Economic, Recreational, Social and Aesthetic Evaluations of the Susitna River.

Objectives: Determine the economic values of the aquatic and terrestrial ecosystems.

Determine the recreational values of the aquatic and terrestrial ecosystems.

Determine the social values of the aquatic and terrestrial ecosystems.

Determine the aesthetic values of the aquatic and terrestrial ecosystems.

Background: Economic, recreational, social, and aesthetic values of the project drainages must be determined in order to project whether the project will enhance or diminish these values. The close proximity of municipalities containing half the human population of Alaska emphasizes the need to assess these values. The Susitna drainage is highly used and important to the sport and commercial fisherman, the recreational enthusiast, industry, and municipalities. The popularity of Denali State Park and nearby Mt. McKinley National Park further attests to the high social, recreational, and aesthetic qualities of the area. Specific data on these subjects in the hydroelectric project area watersheds are incomplete or lacking.

Procedure: The four objectives will be accomplished through statistical surveys and analyses. Some of the methods employed will be literature searches, mail surveys, creel surveys, personal interviews, and fish tag return data.

Schedule:

|       |  |
|-------|--|
| FY 79 | Organize personnel, procure equipment, and begin literature searches, and develop survey approaches. |
| FY 80 | Continue literature searches, analyze data, and begin surveys.                                       |
| FY 81 | Continue literature searches, analyze data, and continue surveys.                                    |
| FY 82 | Continue literature searches, analyze data, and continue surveys.                                    |
| FY 83 | Continue data collection and analyses and write report.  |

Costs:

|       |           |
|-------|-----------|
| FY 79 | \$200,000 |
| FY 80 | \$200,000 |
| FY 81 | \$100,000 |
| FY 82 | \$100,000 |
| FY 83 | \$100,000 |



Title: Predict Project Impacts

Objectives: Determine the direct, indirect, and magnitude of effects the Devils Canyon/Watana project will have on the Susitna River Basin fisheries and other drainages prior to construction approval.

Background: Susitna River Basin investigations to date have not generated sufficient data to predict the impacts of this project on the aquatic ecosystem. Scientific literature is available on the ecological effects of hydroelectric dams which have been constructed in other areas.

Procedure: This study culminates all previously outlined studies. An evaluation of data obtained from the proposed fisheries related biological, habitat, socio-economic, and recreational studies will be combined with other engineering and design studies. A predictive model of the aquatic ecosystem with and without the hydroelectric project will be constructed. Concerns will not be limited to fisheries; secondary effects and how humans will be affected will also be addressed. Information required in this analysis includes seasonal life history habitat requirements of the existing aquatic community, a thorough understanding of the interrelationships between physical, chemical, and biological components of the habitat, and recreational and socio-economic values. Project engineering and design models will also be required, especially those concerned with sedimentation, temperature, dissolved gasses, discharge, and other related physiochemical characteristics.

Literature searches and various project data will be continually analyzed to insure all sources of pertinent data are included.

Schedule:

|       |   |
|-------|---|
| FY 79 | Literature research.                                |
| FY 80 | Literature research, analyze data.                  |
| FY 81 | Literature research, analyze data.                  |
| FY 82 | Literature research, analyze data.                  |
| FY 83 | Literature research, analyze data, predict impacts. |

Cost:

|       |          |
|-------|----------|
| FY 79 | \$ 5,000 |
| FY 80 | \$ 5,000 |
| FY 81 | \$20,000 |
| FY 82 | \$60,000 |
| FY 83 | \$60,000 |

Title: Mitigative Measures for Lost Aquatic Habitat

Objective: To identify and evaluate the Devils Canyon/Watana Dam project fisheries mitigation requirements and implementation costs prior to construction approval.

Background: Critical habitat for various life history stages of aquatic species could be eliminated or reduced in quality and quantity by the Susitna hydropower project. For example, regulation will result in decreased flows downstream of the dams during the summer months which could eliminate critical rearing areas for salmonid fry. The proposed aquatic and related habitat studies should quantify the losses and resulting impact on the fisheries. This activity is designed to provide information to assess the feasibility of mitigation and to indicate long term studies which would direct actual mitigation efforts. Evaluation of these studies will go beyond phase I if the project is deemed feasible.

Procedure: Analyze all project data collected which relate to the fisheries and aquatic habitat of the Susitna River Basin and other impacted drainages. Conduct special studies where necessary and analyze. Conduct literature research to obtain aquatic impact data relating to existing and proposed hydroelectric projects.

Conduct preliminary site surveys which include reconnaissance and topographic analysis. Detailed site surveys and analysis will begin in the last two years of this study.

Schedule:

|       |  |
|-------|--|
| FY 79 | Preliminary site surveys.<br><br>Reconnaissance and topographic analysis<br>Conduct literature research and review.  |
| FY 80 | Continue preliminary site surveys.<br>Analyze data and identify potential areas for mitigation.<br>Continue literature search and review.<br>Report on findings. |
| FY 81 | Detailed site surveys.<br>Analyze surveys.<br>Continue literature search and review.   |
| FY 82 | Continue literature search and review.   |
| FY 83 | Continue detailed site surveys and literature search and review.<br><br>Report on findings.  |

Cost:

|        |          |
|--------|----------|
| FY 79  | \$26,000 |
| FY 80  | \$10,000 |
| FY 81* | \$60,000 |
| FY 82  | \$50,000 |
| FY 83  | \$60,000 |

\* Assumes \$10,000 per site survey.

Title: Plan of Study During and After Completion


Objective: Develop a plan of study to monitor the effects of the project to the aquatic ecosystems during and after completion.

Procedure: This ongoing activity will be dependent on the feasibility results. The data generated from all of the pre-authorization studies will provide the ground work for this plan. Flexibility must be built into this plan until the results of the biological and detailed feasibility studies are available.

Schedule: Complete plan within an additional 14 months after completion of the detailed feasibility studies.

Cost: \$50,000

Fisheries and Habitat Investigations  
of the Susitna River - A Preliminary Study of  
Potential Impacts of the Devils  
Canyon and Watana Hydroelectric Projects



PRELIMINARY ENVIRONMENTAL ASSESSMENT  
OF HYDROELECTRIC DEVELOPMENT  
ON THE SUSITNA RIVER,

TK  
1425  
.58  
A23  
no.75

Prepared for the  
United States Fish and Wildlife Service

by the  
Alaska Department of Fish and Game  
333 Raspberry Road  
Anchorage, Alaska

1978

**ARLIS**  
Alaska Resources  
Library & Information Services  
Anchorage, Alaska