



FINAL REPORT
20 September 2002

Aerial Surveys of Molting Long-tailed Ducks and Other Waterfowl in the Barrier Island-Lagoon Systems Between Spy Island and Brownlow Point, Alaska, 2001



Prepared for

BP EXPLORATION (ALASKA) INC.
Environmental Studies Group
P.O. Box 196612
Anchorage, Alaska 99519-6612



FINAL REPORT
20 September 2002
P598

Aerial Surveys of Molting Long-tailed Ducks and Other Waterfowl in the Barrier Island-Lagoon Systems Between Spy Island and Brownlow Point, Alaska, 2001

by

Lynn E. Noel¹
Stephen R. Johnson²
Gillian M. O'Doherty¹

¹LGL ALASKA RESEARCH ASSOCIATES, INC.
1101 East 76th Avenue, Suite B
Anchorage, Alaska 99518

²LGL LIMITED, environmental research associates
9768 Second Street
Sidney British Columbia, Canada
V8L 3Y8

**Aerial Surveys of Molting Long-tailed Ducks and Other Waterfowl
in the Barrier Island-Lagoon Systems
Between Spy Island and Brownlow Point, Alaska, 2001**

This report was prepared under contract to BP Exploration (Alaska) Inc.
Inquiries about this report may be addressed to:

BP EXPLORATION (ALASKA) INC.
Environmental Studies Group
P.O. Box 196612
Anchorage, Alaska 99519-6612

Cite report as:

Noel, L.E., S.R. Johnson, and G.M. O'Doherty. 2002. Aerial surveys of molting long-tailed ducks and other waterfowl in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska, 2001. Report for BP Exploration (Alaska) Inc. by LGL Alaska Research Associates, Inc., Anchorage, Alaska, USA. 122 p.

Cover photo by John Warden

ABSTRACT

There is concern about declines in sea duck populations in western North America and Alaska. We collected aerial survey data for molting sea ducks and other waterfowl in central Alaskan Beaufort Sea barrier island-lagoon systems collected during July-August 2001. These data add to the 15 years of long-term monitoring data collected on the same survey transects over the 25-year period since 1977. Our study objectives were to determine current distribution and abundance of molting sea ducks and other waterfowl, and to compare current and historical numbers and distributions of long-tailed ducks (*Clangula hyemalis*) in the barrier island-lagoon system between Spy Island and Brownlow Point, Alaska. Three low-level aerial strip-transect surveys were conducted between 23 July-11 August 2001, covering 1986 linear km and 790 km². Long-tailed ducks comprised 78% of the avian fauna in July-August 2001. This is slightly lower than results since 1977 (80-98%) across this same survey area. Eiders, primarily common eiders (*Somateria mollissima v-nigrum*), were more abundant in the eastern lagoon system (west of the Arctic National wildlife Refuge; mean density \pm standard error; 5.0 ± 2.13 eiders/km²); while scoters, primarily surf scoters (*Melanitta perspicillata*), were more abundant in the western lagoon system (west of Prudhoe Bay; 1.8 ± 0.43 scoters/km²). Geese and swans were most

abundant along the mainland shoreline in the western lagoon (6.0 ± 2.04 geese/km²) and on tundra transects south of the eastern lagoon (4.1 ± 2.28 geese/km²). Gulls, primarily glaucous gulls (*Larus hyperboreus*), were more abundant in the western lagoon system (1.5 ± 0.78 gulls/km²), compared to the eastern lagoon system (0.8 ± 0.61 gulls/km²). During July-August 2001, density of long-tailed ducks was highest in the eastern lagoon system (18.5 ± 15.26 ducks/km²), followed by Stefansson Sound (9.5 ± 6.93 ducks/km²), the western lagoon system (8.1 ± 7.75 ducks/km²), and tundra (0.1 ± 0.10 ducks/km²). Throughout the survey area the total number of long-tailed ducks increased from 23 July ($n = 8973$) to 8 August ($n = 14,736$) and then decreased on 11 August 2001 ($n = 3169$). Mean density of long-tailed ducks within the entire survey area (combined western and eastern lagoons) during July and August declined from 1978-2001 ($P = 0.018$). Mean areal density declined for 9 of 16 western lagoon transects, 1 offshore, 4 barrier island, 2 lagoon, and 2 mainland, through the 3 summary time periods (1978-1984, 1989-1991, 1998-2001) between 1978 and 2001. In the eastern lagoon, mean long-tailed duck density decreased on 3 barrier island transects and 1 lagoon transect from 1989-1991 to 1998-2000, while density on 2 mainland shoreline transects increased.

Key Words: central Alaska Beaufort Sea, *Clangula hyemalis*, eider, glaucous gull, *Larus hyperboreus*, long-tailed duck, marine waterbirds, *Melanitta*, scoter, *Somateria*

TABLE OF CONTENTS

| | |
|--|-----|
| ABSTRACT..... | i |
| TABLE OF CONTENTS..... | ii |
| LIST OF FIGURES | iii |
| LIST OF TABLES..... | vii |
| INTRODUCTION..... | 1 |
| Business Rationale | 1 |
| Study Objectives | 1 |
| METHODS..... | 2 |
| Survey Conditions..... | 2 |
| Data Summary and Analysis..... | 3 |
| RESULTS..... | 3 |
| Overview..... | 3 |
| Long-tailed Duck | 3 |
| 2001 Distribution and Abundance..... | 3 |
| Current and Historical Distribution and Abundance | 3 |
| Loons..... | 4 |
| Seabirds..... | 4 |
| Gulls..... | 4 |
| Arctic Terns..... | 4 |
| Miscellaneous Seabirds..... | 4 |
| Ducks..... | 4 |
| Eiders..... | 4 |
| Scoters..... | 5 |
| Miscellaneous Ducks..... | 5 |
| Geese and Swans..... | 5 |
| 2001 Human Activity..... | 5 |
| DISCUSSION..... | 5 |
| Long-tailed Duck Distribution and Abundance | 5 |
| ACKNOWLEDGEMENTS..... | 6 |
| LITERATURE CITED..... | 7 |
| APPENDIX A: JULY AND AUGUST 2001 WEST DOCK NOAA STATION DATA..... | 74 |
| APPENDIX B: 2001 LONG-TAILED DUCK DENSITY BY SURVEY AND HISTORICAL DATA..... | 81 |
| APPENDIX C: 2001 DISTRIBUTION MAPS FOR SELECTED SPECIES..... | 99 |

LIST OF FIGURES

| | | |
|------------|--|----|
| Figure 1. | Locations and numbers for aerial survey transects and survey regions in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska | 9 |
| Figure 2. | Avian species occurring within the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska..... | 10 |
| Figure 3. | Survey aircraft and examples of habitats covered in the barrier-island lagoon systems between Spy Island and Brownlow Point, Alaska..... | 11 |
| Figure 4. | Long-tailed duck sightings as percentage of all bird sightings for aerial surveys with annual mean densities showing standard error in nearshore waters of the central Alaska Beaufort Sea, 1977–2001 | 12 |
| Figure 5. | Mean with standard error bar and standard deviation line for areal density of avian groups among survey regions for 2 or 3 aerial survey replicates between Spy Island and Brownlow Point, Alaska, July–August 1999–2001 | 13 |
| Figure 6. | Areal density of long-tailed ducks for 3 aerial surveys during 23 July–11 August 2001, and mean areal density with standard error for all surveys by region between Spy Island and Brownlow Point, Alaska..... | 14 |
| Figure 7. | Total number of long-tailed ducks on- and off-transect during aerial surveys in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001..... | 15 |
| Figure 8. | Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July–11 August 2001 | 16 |
| Figure 9. | Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 1-24 August 2000..... | 17 |
| Figure 10. | Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 30 July–26 August 1999 | 18 |
| Figure 11. | Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 19 |
| Figure 12. | Summary of density for long-tailed ducks by 30-s segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 1-24 August 2000 | 20 |
| Figure 13. | Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 30 July–26 August 1999 | 21 |
| Figure 14. | Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 5 August–3 September 1998..... | 22 |
| Figure 15. | Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July–11 August 2001..... | 23 |
| Figure 16. | Summary of density for long-tailed ducks by 30-s time period segments on the barrier island-lagoon system between West Dock and Pole Island, Alaska, 1-24 August 2000..... | 24 |
| Figure 17. | Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 30 July–26 August 1999..... | 25 |

| | | |
|------------|---|----|
| Figure 18. | Mean areal long-tailed duck density with standard error by year for aerial surveys in the lagoon systems between Spy Island and Brownlow Point, Alaska, during July and August 1978-2001 | 26 |
| Figure 19. | Mean areal long-tailed duck density with standard error by transect for aerial surveys between Spy Island and West Dock, Alaska, during male long-tailed duck molt period from 15 July to 21 August 1978-1984, 1989-1991, and 1998-2001 | 27 |
| Figure 20. | Mean areal long-tailed duck density with standard error by transect for aerial surveys between Pole Island and Brownlow Point, Alaska, during male long-tailed duck molt period from 15 July to 21 August 1978-1984, 1989-1991, and 1998-2001 | 28 |
| Figure 21. | Total number of loons on- and off-transect during aerial surveys in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska, 23 July-11 August 2001 | 29 |
| Figure 22. | Areal density of loons for 3 aerial surveys by region between Spy Island and Brownlow Point, Alaska, 23 July-11 August 2001 | 30 |
| Figure 23. | Summary of density for loons by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July-11 August 2001 | 31 |
| Figure 24. | Summary of density for loons by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July-11 August 2001 | 32 |
| Figure 25. | Summary of density for loons by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July-11 August 2001 | 33 |
| Figure 26. | Total number of seabirds on- and off-transect during aerial surveys in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska, 23 July-11 August 2001 | 34 |
| Figure 27. | Areal density of gulls for 3 aerial surveys by region between Spy Island and Brownlow Point, Alaska, 23 July-11 August 2001 | 35 |
| Figure 28. | Summary of density for gulls by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July-11 August 2001 | 36 |
| Figure 29. | Summary of density for gulls by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July-11 August 2001 | 37 |
| Figure 30. | Summary of density for gulls by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July-11 August 2001 | 38 |
| Figure 31. | Areal density of terns for 3 aerial surveys by region between Spy Island and Brownlow Point, Alaska, 23 July-11 August 2001 | 39 |
| Figure 32. | Summary of density for Arctic terns by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July-11 August 2001 | 40 |
| Figure 33. | Summary of density for Arctic terns by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July-11 August 2001 | 41 |
| Figure 34. | Summary of density for Arctic terns by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July-11 August 2001 | 42 |
| Figure 35. | Total number of eiders on- and off-transect during aerial surveys in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska, 23 July-11 August 2001 | 43 |
| Figure 36. | Areal density of eiders for 3 aerial surveys by region between Spy Island and Brownlow Point, Alaska, 23 July-11 August 2001 | 44 |

| | | |
|------------|---|----|
| Figure 37. | Summary of density for eiders by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July–11 August 2001 | 45 |
| Figure 38. | Summary of density for eiders by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July–11 August 2001 | 46 |
| Figure 39. | Summary of density for eiders by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 47 |
| Figure 40. | Total number of scoters on- and off-transect during aerial surveys in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 48 |
| Figure 41. | Areal density of scoters for 3 aerial surveys by region between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 49 |
| Figure 42. | Summary of density for scoters by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July–11 August 2001 | 50 |
| Figure 43. | Summary of density for scoters by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 51 |
| Figure 44. | Total number of geese and swans observed both on- and off-transect during aerial surveys in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 52 |
| Figure 45. | Areal density of geese and swans for 3 aerial surveys by region between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 53 |
| Figure 46. | Summary of density for geese and swans by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July–11 August 2001 | 54 |
| Figure 47. | Summary of density for geese and swans by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July–11 August 2001..... | 55 |
| Figure 48. | Summary of density for geese and swans by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 56 |
| Figure 49. | Mean areal density with standard error for vessel traffic and human activity during aerial surveys by region between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001..... | 57 |
| Figure A1. | Daily mean temperature and wind speed recorded at the Prudhoe Bay NOAA Station at West Dock, Alaska, 15 July–31 August 2001 | 75 |
| Figure A2. | Hourly mean water temperature recorded at the Prudhoe Bay NOAA Station at West Dock, Alaska, 15 July–1 September 2001 | 76 |
| Figure A3. | Hourly mean wind speed and direction recorded at the Prudhoe Bay NOAA Station at West Dock, Alaska, 15 July–1 September 2001..... | 77 |
| Figure A4. | Hourly mean air temperature recorded at the Prudhoe Bay NOAA Station at West Dock, Alaska, 15 July–1 September 2001 | 78 |
| Figure A5. | Water level relative to mean low, low water level recorded at 6-min intervals at the Prudhoe Bay NOAA Station at West Dock, Alaska, 1–31 July 2001..... | 79 |
| Figure A6. | Water level relative to mean low, low water level recorded at 6-min intervals at the Prudhoe Bay NOAA Station at West Dock, Alaska, 1–31 August 2001..... | 80 |
| Figure B1. | Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July 2001 | 82 |

| | | |
|-------------|---|-----|
| Figure B2. | Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 8 August 2001 | 83 |
| Figure B3. | Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 11 August 2001 | 84 |
| Figure B4. | Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July 2001 | 85 |
| Figure B5. | Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 8 August 2001 | 86 |
| Figure B6. | Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 11 August 2001 | 87 |
| Figure B7. | Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July 2001..... | 88 |
| Figure B8. | Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 8 August 2001..... | 89 |
| Figure B9. | Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 11 August 2001..... | 90 |
| Figure C1. | Summary of density for Pacific loons by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July–11 August 2001 | 100 |
| Figure C2. | Summary of density for Pacific loons by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July–11 August 2001..... | 101 |
| Figure C3. | Summary of density for Pacific loons by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 102 |
| Figure C4. | Summary of density for red-throated loons by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July–11 August 2001 | 103 |
| Figure C5. | Summary of density for red-throated loons by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July–11 August 2001..... | 104 |
| Figure C6. | Summary of density for red-throated loons by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 105 |
| Figure C7. | Summary of density for yellow-billed loons by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July–11 August 2001..... | 106 |
| Figure C8. | Summary of density for yellow-billed loons by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July–11 August 2001..... | 107 |
| Figure C9. | Summary of density for yellow-billed loons by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July–11 August 2001..... | 108 |
| Figure C10. | Summary of density for jaegers by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July–11 August 2001 | 109 |
| Figure C11. | Summary of density for jaegers by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 110 |
| Figure C12. | Summary of density for black guillemots by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July–11 August 2001 | 111 |

| | |
|---|-----|
| Figure C13. Summary of density for black guillemots by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July–11 August 2001..... | 112 |
| Figure C14. Summary of density for black guillemots by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 113 |
| Figure C15. Summary of density for glaucous gulls by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July–11 August 2001 | 114 |
| Figure C16. Summary of density for glaucous gulls by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July–11 August 2001..... | 115 |
| Figure C17. Summary of density for glaucous gulls by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 116 |

LIST OF TABLES

| | |
|---|----|
| Table 1. Summary of weather and lagoon conditions during 3 aerial surveys in the barrier island-lagoon systems between Spy Island and Brownlow Point and on tundra transects from Brownlow Point to the Shaviovik River, Alaska, 23 July–11 August 2001 | 58 |
| Table 2. Summary of survey effort in the barrier island-lagoon systems between Spy Island and Brownlow Point and tundra between Brownlow Point and the Shaviovik River, Alaska, 23 July–11 August 2001..... | 59 |
| Table 3. Total number of bird sightings and individuals seen on- and off-transect for all aerial survey transects in the barrier island-lagoon system, offshore, and on tundra between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001..... | 60 |
| Table 4. Total number of bird sightings and individuals seen on- and off-transect for aerial survey transects in the barrier island-lagoon system and offshore between Spy Island and West Dock, Alaska, 23 July–11 August 2001 | 61 |
| Table 5. Total number of bird sightings and individuals seen on- and off-transect for aerial survey transects in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July–11 August 2001..... | 62 |
| Table 6. Total number of bird sightings and individuals seen on- and off-transect for aerial survey transects in the barrier island-lagoon system and offshore between Pole Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 63 |
| Table 7. Total number of bird sightings and individuals seen on- and off-transect for aerial survey transects on tundra between Brownlow Point and the Shaviovik River, Alaska, 8 and 11 August 2001 | 64 |
| Table 8. Long-tailed duck density by aerial survey transect in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July–11 August 2001 | 65 |
| Table 9. Long-tailed duck density by aerial survey transect in the barrier island-lagoon system and on tundra between Pole Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 66 |
| Table 10. Long-tailed duck density by aerial survey transect in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July–11 August 2001..... | 67 |
| Table 11. Species densities for all aerial survey transects in the barrier island-lagoon systems and on tundra between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001..... | 68 |

| | | |
|-----------|---|----|
| Table 12. | Habitat associations of loons during aerial surveys in the barrier island-lagoon systems and tundra transects between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 69 |
| Table 13. | Habitat associations of seabirds during aerial surveys in the barrier island-lagoon systems and tundra transects between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 70 |
| Table 14. | Habitat associations of eiders during aerial surveys in the barrier island-lagoon systems and tundra transects between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 71 |
| Table 15. | Habitat associations of scoters during aerial surveys in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 72 |
| Table 16. | Habitat associations of geese and swans during aerial surveys in the barrier island-lagoon systems and tundra transects between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 73 |
| Table B1. | Numbers and percentages of long-tailed ducks counted during aerial surveys in nearshore waters of the central Alaska Beaufort Sea, 1977-1984; 1989-1991; 1998-2001 | 91 |
| Table B2. | Individual on-transect long-tailed duck densities for transects between Spy Island and Brownlow Point, Beaufort Sea, Alaska, 5 June–23 September 1977-1984; 1989–1991; 1998-2001 | 92 |
| Table B3. | Database records for sightings of human activity during aerial surveys in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 95 |
| Table B4. | Habitat associations of long-tailed ducks during aerial surveys in the barrier island-lagoon systems and tundra transects between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 97 |
| Table B5. | Long-tailed duck behavior during aerial surveys in the barrier island-lagoon systems and tundra transects between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001 | 98 |

Aerial Surveys of Molting Long-tailed Ducks and Other Waterfowl in the Barrier Island-Lagoon Systems Between Spy Island and Brownlow Point, Alaska, 2001

INTRODUCTION

Tens of thousands of molting/flightless waterfowl aggregate along the mainland and barrier island coastlines in the central Beaufort Sea, Alaska (Figures 1 and 2). Based on several decades of study, we know that these aggregations of molting/flightless waterfowl, primarily sea ducks and geese, are highly susceptible to disturbance, and to oil or fuel spills (Johnson and Richardson 1981; Johnson 1985, 1990; Johnson and Gazey 1992). Recently, concern has been expressed over the apparent decline in 10 of the 15 species of North American sea ducks (Elliot 1997, USFWS 1999). These include species that occur within the Spy Island to Brownlow Point barrier island-lagoon systems (Figures 1 and 2): long-tailed duck (*Clangula hyemalis*), common eider (*Somateria mollissima v-nigrum*), king eider (*Somateria spectabilis*), black scoter (*Melanitta nigra americana*), surf scoter (*Melanitta perspicillata*), and white-winged scoter (*Melanitta fusca deglandi*).

Long-tailed ducks (formerly called oldsquaw ducks) are the predominant species in central Alaskan Beaufort Sea lagoon systems, and therefore they were the focus for the development of a monitoring program that was designed to evaluate changes in abundance and distribution of marine waterfowl in relation to industrial development (Figure 1, Johnson and Gazey 1992). Currently, there is concern over reported declines in long-tailed duck populations in western North America, Alaska, and northwestern Canada (Hodges et al. 1996, Conant et al. 1997, Dickson and Gilchrist 2002, Mallek et al. 2002). Trend data for the long-tailed duck population nesting on the Arctic Coastal Plain, however, are conflicting. Larned et al. (1999:15) reported a slightly increasing trend in long-tailed ducks on the Arctic Coastal Plain of Alaska from 1992-1999. Other authors have reported stable long-tailed duck populations during the same period (Conant et al. 1997, Elliot 1997, Larned and Balogh 1997, USFWS 1999), whereas Mallek and King (2000) and Mallek et al. (2002) reported recent declines in the population.

In 1998-2001, BP Exploration (Alaska) Inc. (BPXA) and the Point Thomson Unit Owners (PTUO) funded LGL Alaska Research Associates, Inc. (LGL) to collect molting waterfowl distribution and abundance data (Noel et al. 1999, 2000, 2001) using methods developed and tested for this

monitoring protocol (Johnson and Gazey 1992). BPXA and the PTUO have continued to fund LGL to monitor molting long-tailed ducks and other waterfowl using the Johnson and Gazey (1992) protocol to document activities that could disturb molting waterfowl and provide current distribution and abundance information for use in environmental assessments and environmental impact statements.

Business Rationale

Nesting populations of long-tailed ducks may be declining across the Arctic Coastal Plain of Alaska (Mallek et al. 2002), and in northwestern Canada (Dickson and Gilchrist 2002). This species is the most abundant molting waterfowl within the barrier island lagoons adjacent to oilfields in the Prudhoe Bay region and in areas of proposed coastal developments at Point Thomson. Waterfowl such as long-tailed ducks are susceptible to disturbance and petroleum spills during the molt when they are flightless. Development of new near shore (Northstar) and coastal oilfields (Badami and Point Thomson) increases the potential for disturbance from resource exploration and development activities and disturbance associated with wildlife research activities (Johnson et al. in prep.). Documenting and understanding how these activities affect distribution and abundance of long-tailed ducks is important for developing meaningful environmental assessments and impact statements.

Study Objectives

This report presents the results of molting long-tailed duck and other waterfowl monitoring surveys during July and August 2001 between Spy Island and Brownlow Point, Alaska. Aerial survey data for molting long-tailed ducks and other waterfowl in central Alaskan Beaufort Sea barrier island-lagoon systems collected during July-August 2001 add to the 15 years of long-term monitoring data collected over the 25-year period since 1977.

Objectives for the 2001 study were:

1. Document the distribution and abundance of molting long-tailed ducks and other waterfowl in the barrier island-lagoon systems between Spy Island and Brownlow Point, and on large lakes along an inland transect between Brownlow Point and the Shaviovik River, Alaska.

2. Compare current and historical numbers and distributions of long-tailed ducks in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska.
3. Document the level of human activity during surveys in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska.

METHODS

Survey methods and conditions adopted in this study were based on analyses that identified variables that influenced the numbers of long-tailed ducks recorded during low-level aerial surveys (Johnson 1990, Johnson and Gazey 1992). Three of five scheduled low-level aerial (fixed-wing aircraft) surveys were flown during the period 23 July – 11 August 2001 in the barrier island-lagoon systems from Spy Island in the west to Brownlow Point in the east (Figure 1, Table 1). Brownlow Point lies on the mainland just east of Flaxman Island (Figure 1). Sampling was balanced by surveying both the western and eastern lagoons on the same day. Four contiguous transects were surveyed within 4 habitat strata (offshore, barrier island, lagoon, mainland shoreline) in both the western and eastern lagoons (Figure 1). All habitats within a lagoon system were surveyed before continuing to the next lagoon system to reduce the time between sampling these transects, thereby minimizing the possibility of birds moving between transects. In addition to transects established by the monitoring protocol in the western and eastern lagoons (Johnson and Gazey 1992), barrier island and mainland shoreline habitats across Stefansson Sound, and large lakes on the tundra between the Staines River and the Shaviovik River were also surveyed (Figure 1).

The survey crew consisted of a pilot and two observers in a float equipped Cessna 206 fixed-wing aircraft (Figure 3); one observer sat in the right front seat and the other sat in the left rear seat. Survey altitude was approximately 45 m above ground level and ground speed approximately 180 km/h. Transect width was 400 m total: 200 m on each side of the aircraft. On-transect sightings were within 200 m from the aircraft, and off-transect sightings were more than 200 m from the aircraft. Observers were trained to estimate large numbers of birds in dense concentrations using the simulation program Counting Wildlife® v. 2.0 (1986, Wildlife Counts, Juneau, AK) and using poppy seed scatter trials (Johnson 1990, Johnson and Gazey 1992).

The survey period was focused on the peak of the male long-tailed duck flightless period from 15 July to 21 August, based on progression of the wing molt

(Johnson and Richardson 1981, Johnson and Gazey 1992). Six to 8 replicate surveys at even intervals during this period were recommended (Johnson and Gazey 1992). Surveys were scheduled as late in the day as practical; as late as 1700 h Alaska Daylight Savings Time, because long-tailed ducks have been found to concentrate along the barrier islands in the late afternoon (Johnson and Richardson 1981; Johnson 1982, 1985). Surveys on days with high winds (>37 km/h, 20 kts) were delayed until winds, wave height, and chop diminished, thereby improving sightability of birds on the water (Johnson 1990, Johnson and Gazey 1992). Marginal survey conditions were when winds approached 37 km/h.

Tape recorders were used to record information about observed birds, mammals, habitats, and environmental conditions. Continuous audio-tape recordings were made which included information on transects, sightings, and 30-s intervals (time period) marked by an audio-intervalometer (timer). Variables recorded include: *Transect* - start time (h-min-s), general location, transect number, temperature, wind speed, wind direction, Beaufort Sea state, wave height, cloud cover, habitat type, stop time; *Sighting* - species, number, behavior, habitat type, human activity, on- or off-transect; *Time period* - time at "beep" sound, ice cover, wave height, glare on the water surface, wind speed, wind direction, any changes in any particular variable. A notebook computer equipped with a Global Positioning System (GPS) receiver coupled with mapping software was used to record the flight line on a map of the area at 1-s intervals during surveys. Observers synchronized their watches with the GPS satellite time, and these times (recorded for transects and time periods) were used to geo-reference the survey data.

Survey Conditions

General weather conditions in the study area were recorded during each survey and as broadcast by radio operators at the Deadhorse airport or at the Badami weather office. Wind speed and direction were assessed by a combination of water surface conditions and pilot observations. Temperatures were either recorded at altitude from the aircraft or from Deadhorse or Badami air-radio broadcast. Wind speed and direction greatly influence sightability and behavior of molting waterfowl (Johnson and Richardson 1981, Johnson 1990, Johnson and Gazey 1992). We attempted to limit survey days to those with winds <37 km/h (Table 1). Weather data recorded over this survey period by the National Oceanic and Atmospheric Administration (NOAA) station at West Dock is presented in Appendix A.

Data Summary and Analysis

Our unit of replication for summaries and analyses is the survey. For visual comparisons among regions, transects within regions were summed for each survey and mean values for the 3 replicate surveys during 2001 were computed with standard errors and standard deviations. To compare changes in the molting long-tailed duck population over time, we summed the number of long tailed ducks for transects in both the western and eastern lagoons for each survey and computed the mean values based on survey replicates for each year. We then completed simple linear regression to evaluate data for any trend over time (Zar 1974). To compare changes in long-tailed duck density over time for individual transects, we computed the mean and standard errors for long-tailed duck density during 3 time periods 1978-1984 ($n = 6$ to 15 surveys), 1989-1991 ($n = 11$ to 20 surveys), and 1998-2001 ($n = 9$ to 12 surveys) for transects in the western and eastern lagoons.

Habitat and behavior summaries were based on the number of sightings of each species. Sightings are based on flocks rather than individuals. Individuals within a flock behave similarly and respond to others within the flock, and therefore are not independent. Flocks consisting of 1 or many individuals were considered independent from one another.

RESULTS

Total survey effort was 1986 linear km (790.4 km², Table 2). Results are presented below as a general overview, followed by sections for taxonomic groupings. We rely on graphs, maps, and tables to illustrate relative abundances, distributions, and habitat associations. Graphics and tables are presented in the order cited, following the body of the report. Areal density maps for taxonomic groups are presented as the number of individuals per km² for on-transect data, plotted at the mid-point of each 30-s time period, for each transect. Data for all 3 surveys are presented on a series of 3 maps covering the survey area. Maps for each survey date in 2001 for long-tailed ducks are presented in Appendix B. Maps for other selected species are presented in Appendix C.

Overview

Long-tailed ducks dominated the avian fauna in 2001, both in terms of numbers of individuals and numbers of sightings (Table 3), which is consistent with results since 1977 (Figure 4, Table B1, Johnson and Gazey 1992). Long-tailed ducks dominated in the western lagoon, Stefansson Sound, and the eastern lagoon during 2001, ranging from 75-81% of the total number of individuals recorded (Figure 1, Tables 4-6).

Relative abundance among other avian groups differed among the 4 survey regions (western lagoon, Stefansson Sound, eastern lagoon, and tundra, Figure 1). Loons dominated in terms of number of sightings, while geese and swans dominated the number of individuals on tundra transects (Table 7). Relative abundance of gulls was highest in the western lagoon, but was highly variable (Figure 5). Eiders were more abundant in the eastern lagoon; while scoters were more abundant in the western lagoon (Figure 5). Geese and swans were most abundant in the western lagoon and tundra (Figure 5).

Long-tailed Duck

2001 Distribution and Abundance

Mean areal long-tailed duck density was highest in the eastern lagoon (18.5 ± 15.26 ducks/km²), followed by Stefansson Sound (9.5 ± 6.93 ducks/km²), the western lagoon (8.1 ± 7.75 ducks/km²), and tundra (0.1 ± 0.10 ducks/km²) for surveys in July and August 2001; although variability (ducks/km² \pm standard error) was high (Figure 6). Throughout the survey area the total number of long-tailed ducks increased from 23 July-8 August (8973, 27.5 ± 3.02 individuals/flock to 14,736, 38.7 ± 3.56 individuals/flock) and decreased between 8 August-11 August 2001 (14,736, 38.7 ± 3.56 individuals/flock to 3,169, 21.2 ± 3.41 individuals/flock, Figure 7). In the western lagoon, long-tailed ducks were concentrated along the barrier islands and in the lagoon, with few long-tailed ducks along the mainland shoreline (Figure 8, Table 8).

Current and Historical Distribution and Abundance

The relatively low use of the mainland shoreline in the western lagoon in 2001 (Figure 8) is consistent with the distribution of long-tailed ducks observed in 2000 (Figure 9) and 1999 (Figure 10). In the eastern lagoon, long-tailed ducks were concentrated near Pole Island (Transect #133), Tigvariak Island (Transect #193), and along the mainland shoreline (Transect #190, Figure 11, Table 9). Long-tailed duck use of the mainland shoreline in the eastern lagoon in 2001 is consistent with distributions observed in 2000, 1999 and 1998 (Figures 12, 13, and 14). In Stefansson Sound, long-tailed ducks were concentrated in the lee of Cross Island and the McClure Islands in 2001 (Figure 15), similar to distributions in 2000 (Figure 16) and 1999 (Figure 17, Table 10). Distribution maps for each region by survey, showing long-tailed duck density by 30-s period and vessel traffic, are presented in Appendix B.

Long-tailed duck was the dominant species in the barrier island-lagoon systems between Spy Island and Brownlow Point during 2001 (Table 3), consistent with survey data since 1977 (Figure 4, Table B1). Mean areal density of long-tailed ducks within this survey area during July and August appears to have declined from 1978-2001, although time appears to explain only 38% of the variation in long-tailed duck density (Figure 18). This summary does not correct for factors which influence mean density such as the total number of km² surveyed, regions covered, or variability in survey conditions (Johnson and Gazey 1992, Johnson et al. in prep.).

The molting long-tailed duck population in Beaufort Sea lagoons from 15 July to 21 August is primarily composed of flightless males, as determined by wing measurements (Johnson and Richardson 1981, Johnson 1985). Mean long-tailed duck density among some transects in the western lagoon during this molt period shows a general decline from 1978-1984 to 1998-2001 (Figure 19). This trend is most pronounced for the barrier island transects at the western end of Simpson Lagoon (Figure 19). Mean areal density declined for 9 of 12 western lagoon transects (1 offshore, 4 barrier island, 2 lagoon, and 2 mainland) through the 3 summary time periods between 1978 and 2001 (Figure 19). In the eastern lagoon, mean long-tailed duck density decreased on 3 of 4 barrier island transects and 1 of 4 lagoon transects from 1989-1991 to 1998-2001, while mean long-tailed duck density on 2 of 4 mainland shoreline transects increased (Figure 20).

Loons

Although there was variability between individual surveys, loons were most abundant on tundra transects, followed by the western lagoon, Stefansson Sound, and eastern lagoon (Figure 5). The number of loons per survey by region ranged from >180 on 8 August (western lagoon) to <1 on 23 July (tundra) and 11 August (Stefansson Sound, Figure 21). Areal loon density was highest in the western lagoon on 8 August and was lowest in Stefansson Sound on 11 August (Figure 22). Pacific loons (*Gavia pacifica*) and red-throated loons (*Gavia stellata*) were the most common loon species, with the total density of Pacific loons more than twice that of red-throated loons in the survey area (Table 11). Loons were scattered throughout the survey area in low numbers (Figures 23-25). Pacific loons were the most widely distributed across habitat types with a higher proportion of sightings in lagoon, mainland shoreline, and barrier island shoreline habitats (Table 12). Red-throated loons occurred in the same habitats as Pacific loons, except the red-throated loons were not recorded in the tundra habitats surveyed (Table 12).

Yellow-billed loons (*Gavia adamsii*) were the least common loon species, and occurred primarily along mainland shorelines and in lagoons (Table 12).

Seabirds

Gulls, primarily glaucous gulls (*Larus hyperboreus*), were the most common seabird within the survey area (Figure 26, Table 3-7).

Gulls

Gull abundance (both total numbers and density) decreased in the western and eastern lagoons from 23 July-8 August (Figures 26 and 27). The mean density of gulls in the western lagoon was nearly twice that of gulls in the eastern lagoon (Figure 5). Most glaucous gull sightings were along the barrier islands (55%), followed by the mainland shoreline (33%, Table 13). The greatest concentrations of gulls in the western lagoon occurred near Stump Island and Bertocini Island (Figure 28). In the Stefansson Sound region, gulls were scattered throughout the barrier islands (Figure 29). In the eastern lagoon, gulls were concentrated on the shoreline of Mikkelson Bay (Figure 30). Three flocks with 49 Sabine's gulls (*Xema sabini*) were also recorded, with the largest flock (84% of individuals) offshore in the western lagoon.

Arctic Terns

Arctic terns (*Sterna paradisaea*) were not abundant in the survey area, but occurred most frequently in the Stefansson Sound and eastern lagoon (Figures 5 and 26, Tables 3-7). Arctic tern density was highest on 11 August in the Stefansson Sound region (Figure 31). Most sightings were adjacent to the barrier islands (Figures 32-34, Table 13).

Miscellaneous Seabirds

Two Pomarine Jaegers (*Stercorarius pomarinus*) were sighted on 8 August near the Shavirovik River (Figures C10 and C11). A common murre (*Uria aalge*) and a black guillemot (*Cepphus grylle*) were sighted on 11 August in the eastern lagoon and Stefansson Sound regions respectively (Figure C12, Tables 3-7).

Ducks

Eiders

Eiders were most abundant in the eastern lagoon and Stefansson Sound with highest numbers on 23 July in both areas (Figures 5, 35, and 36). Eider densities were higher in the eastern lagoon and Stefansson Sound than in the western lagoon or tundra regions (Figure 5). Density was highest in the eastern lagoon on 23 July and was lowest in the

tundra region on 11 August (Figure 36). Common eiders were the most abundant species in this group, comprising over 99% of eiders that were classified to species (Table 3). In the western lagoon, most eiders were scattered in small numbers along the barrier islands and in the lagoon with a few large groups offshore from Pingok and Bodfish islands and along the mainland on either side of Milne Point (Figure 37). In Stefansson Sound, eiders occurred south of Reindeer, Cross, Narwhal, and Jeanette islands (Figure 38). In the eastern lagoon large flocks of eiders occurred along the barrier islands and in the lagoons between the Stockton Islands and the Maguire Islands (Figure 39). Over half of common eider sightings (57%) were associated with the barrier islands (Table 14).

Scoters

Scoters, primarily surf scoters, were most abundant in the western lagoon, with the highest density on 8 August (Figures 5, 40, and 41; Tables 3-7). Over 75% of scoter sightings were on lagoon transects (Table 15). The largest flocks of scoters were in Simpson Lagoon (Figures 42 and 43).

Miscellaneous Ducks

Aside from scoters, eiders, and long-tailed ducks, only a few other duck species were recorded. Scaup (*Aythya spp.*) were recorded in the western lagoon, eastern lagoon, and tundra; northern pintails (*Anas acuta*) were recorded in the eastern lagoon; and a red-breasted merganser (*Mergus serrator*) was recorded in the eastern lagoon.

Geese and Swans

Within the survey area during 2001, geese and swans were most abundant along the mainland shoreline of the western lagoon and on tundra (Figure 5, 44, and 45; Table 16), with highest numbers on 8 August in the western lagoon (Figure 44). Goose and swan density was consistently higher in the western lagoon than in the eastern lagoon (Figures 5, 45-48, Table 4 and 6). Greater white-fronted goose (*Anser albifrons*) and black brant (*Branta bernicla*) were the most common species, occurring primarily along the coastline in the western lagoon (Figure 46, Tables 4 and 16). Lesser snow geese (*Chen caerulescens caerulescens*) and Canada geese (*Branta canadensis*) were the most common species in the eastern lagoon primarily on Tigvariak Island and in the Shavirovik River delta (Figure 48, Tables 6 and 16). A few tundra swans (*Cygnus columbianus*) were recorded on coastal tundra in the eastern and western lagoons, and on tundra transects (Figure 44, Table 16).

2001 Human Activity

Vessel traffic was highest on 8 August 2001 in the western lagoon region (Table B3). Vessel traffic was generally higher within the western lagoon than within the eastern lagoon, although vessel traffic was recorded throughout the survey area during 2001 (Figure 49, Table B3). Vessel sightings included small and medium sized boats, seismic boats, large ships, and small aircraft (Table B3). Human activity other than the established oilfield facilities included field camps at Cottle Island, Flaxman Island, and the Point Thomson #3 pad (Table B3). Other human activity sightings included fyke nets, bird capture nets, telemetry monitoring stations, people, and telemetry antennas (Table B3). On transect vessel traffic has remained generally higher in the western lagoon than in the eastern lagoon from 1999-2001, although there does not appear to be a similar pattern for human activity (Figure 49).

DISCUSSION

Long-tailed Duck Distribution and Abundance

Survey data for the central Beaufort Sea barrier island-lagoon systems collected during August 2001 supplement the 14 years of long-term monitoring data collected over the 25 year period since 1977. Based on the 2001 data, long-tailed ducks continue to be the most numerous birds in Beaufort Sea barrier island-lagoon systems, where they feed primarily on epibenthic organisms including mysids (*Mysis relicta* and *M. litoralis*) and amphipods (*Onisimus glacialis*) (Johnson 1982). During the flightless molt period from mid July to late August, male long-tailed ducks and some non-breeding females congregate in very large numbers in barrier island-lagoon systems such as Simpson Lagoon, Gwydyr Bay, and south of Flaxman Island. The highest densities of molting flocks generally concentrate immediately south of barrier island shorelines. Barrier islands provide protection from prevailing winds and rough water, provide easy access to roosting areas along leeward beaches, and are close to abundant prey resources in the lagoons (Johnson 1982).

Mean density of long-tailed ducks within the combined western and eastern lagoons during July and August appears to have declined from 1978-2001 (Figure 18, $y = -4.96x + 178$, $R^2 = 0.382$, $P = 0.018$), without correction for number of km² surveyed, regions covered, or variability in survey conditions. During periods of disturbance (aircraft, boat and humans) and rough water, Johnson (1982) found that long-tailed ducks responded by moving to a nearby

location that provided protection from wind and waves. Changes in distribution appeared to be primarily related to weather conditions rather than disturbance, but a decline in overall long-tailed duck numbers within the study area in 1981 (4000 to 2000) indicated that there was a general pattern of movement away from the sources of disturbance (Johnson 1982). This suggests that continuous vessel traffic and aircraft disturbance during molting may lead to displacement of long-tailed ducks. Development and expansion of the oilfields in the Prudhoe Bay region since 1977, with an assumed increase in nearshore traffic, may have contributed to the decreased density of long-tailed ducks within the western lagoon systems from 1977-1984 to 1998-2001 (Figure 19).

Since 1998, an assumed increase in vessel and aircraft traffic associated with development of the Badami oilfield, exploration in the Flaxman Island area, and intensive environmental studies may also have contributed to a decrease in density of long-tailed ducks on 3 of 4 barrier islands and an increase in density on 2 of 4 mainland shoreline transects from 1989-1991 to 1998-2001 (Figure 20). This may in part be a reflection of sampling intensity, survey timing, survey weather conditions, and the general decline in the Arctic Coastal Plain nesting population; but may also be related to increased disturbance within this lagoon system. A detailed analysis of these data, incorporating disturbance and other environmental variables known to influence the number of long-tailed ducks recorded during monitoring surveys, is beyond the scope of this presentation of our 2001 survey results. This analysis has recently been completed and the manuscript based on this analysis is in preparation (Johnson et al. in prep.).

Weather conditions influence the total number and distribution of long-tailed ducks recorded during our monitoring surveys (Johnson 1982, Johnson and Gazey 1992). Moderate to strong winds (20-37 km/h) directly influence sightability of birds on the water by increasing wave height and surface disturbance (Johnson and Gazey 1992). During periods of strong southwesterly winds, long-tailed ducks move toward the mainland coast and/or out of the lagoon through the inter-island passes to take shelter in the lee of the mainland coast or north of the barrier islands (Johnson and Richardson 1981).

Undisturbed molting long-tailed ducks typically cycle through a 24-h period of activity, with peak numbers resting and preening in the leeward nearshore and beach habitats during evening and early morning (Johnson 1982, 1985; Flint et al. 2001). During mid-day, long-tailed ducks typically

move farther from shore into mid-lagoon habitats to feed (Johnson 1982, Flint et al. 2001). Although molting sea ducks have been found to spend more time in open water when disturbed intermittently by aircraft; time spent swimming and feeding, and population levels appeared to be unaffected (Gollop et al. 1974). Flint et al. (2001) conducted experimental boat disturbances but could not identify changes in location, changes in movement patterns, or changes in feeding frequency associated with these experimental disturbances. Poor resolution for radio triangulated locations, and behavioral differences between experimental and control areas were cited as reasons for an inability to measure a response to experimental disturbances (Flint et al. 2001). Johnson (1982) documented a breakdown in the 24-h activity cycle with disturbance and increased wind, although cyclic movements continued in an adjacent undisturbed sheltered area (Johnson 1982).

ACKNOWLEDGEMENTS

This study was funded by BP Exploration (Alaska) Inc. (BP). We thank Dave Trudgen and Bill Streever, with the Environmental Studies Group at BP Exploration (Alaska) Inc. for their support and suggestions concerning this study. Bill Streever provided comments which improved this report. Peter Wainwright designed, programmed, and maintains the customized software used for data analysis. The survey aircraft was piloted by Jim Helmericks, who has piloted these surveys since 1977.

LITERATURE CITED

- Conant, B., D.J. Groves, C. Ferguson, and R.J. King. 1997. Long-tailed ducks – toward listing. Seventh Alaska Bird Conference, 1–5 December 1997, U.S. Fish and Wildlife Service, Migratory Bird Management, Anchorage, AK.
- Dickson, D.L., and H.G. Gilchrist. 2002. Status of marine birds in the southeastern Beaufort Sea. *Arctic* 55 Supplement 1:46-58.
- Elliot, R.D. (ed) 1997. Conservation issues for North American Sea Ducks. A concept paper for a Sea Duck Joint Venture under the North American Waterfowl Management Plan. Report by Canadian Wildlife Service, U.S. Fish & Wildlife Service and U.S. Geological Survey (Biological Resources Division). Canadian Wildlife Service – Atlantic Region, Sackville, New Brunswick, Canada. 35 p.
- Flint, P. L., R. B. Lanctot, J. C. Franson, T. Hollmen, J. Fischer, J. B. Grand, and M. Howell. 2001. Monitoring Beaufort Sea waterfowl and marine birds. Annual Progress Report by U.S. Geological Survey, Alaska Biological Science Center, Anchorage, AK. 43 p + appendices.
- Gollop, M.A., J.R. Goldsberry, and R.A. Davis. 1974. Aircraft disturbance to moulting sea ducks, Herschel Island, Yukon Territory, August 1972. *Arctic Gas Biol. Rep. Ser.* 14(5):202-231.
- Hodges, J.I., J.B. King, B. Conant, and H.A. Hanson. 1996. Aerial surveys of waterbirds in Alaska 1957–1994: Population trends and observer variability. National Biological Service Information and Technology Report 4. 24 p.
- Johnson, S.R. 1982. Continuing investigations of long-tailed ducks (*Clangula hyemalis* L.) during the molt period in the Alaskan Beaufort Sea. Pages 547-563 in *Envir. Assess. Alaskan Cont. Shelf, Final Rep. Princ. Invest.* Vol. 23. BLM/NOAA, OCSEAP, Juneau, AK.
- Johnson, S.R. 1985. Adaptations of the long-tailed duck (*Clangula hyemalis* L.) during the period of molt in Arctic Alaska. *Proc. Internat. Ornithol. Congress* 18:530-540.
- Johnson, S.R. 1990. Monitoring Beaufort Sea waterfowl and marine birds. Report OCS Study MMS 90-0048 by LGL Alaska Research Assoc., Inc. for U.S. Department of Interior, Alaska Outer Continental Shelf Region of the Minerals Management Service, Anchorage, Alaska. 121 p.
- Johnson, S.R. and W. Gazey. 1992. Design and testing of a monitoring program for Beaufort Sea waterfowl and marine birds. Report OCS Study MMS 92-0060 by LGL Alaska Research Associates, Inc. for U.S. Department of Interior, Alaska Outer Continental Shelf Region of the Minerals Management Service, Anchorage, Alaska. 114 p.
- Johnson, S.R. and D.R. Herter. 1989. The Birds of the Beaufort Sea. BP Exploration (Alaska) Inc., Anchorage, AK. 372 p.
- Johnson, S.R. and W.J. Richardson. 1981. Barrier island lagoon ecological process studies: Final Report, Simpson Lagoon. Part 3, Birds. Pages 109–383 in *Environmental Assessment of the Alaskan Continental Shelf, Final Reports of Principal Investigators. Volume 7. Biological Studies. Outer Continental Shelf Environmental Assessment Program.* U.S. Department of Commerce, National Oceanic & Atmospheric Administration and U.S. Department of the Interior, Bureau of Land Management. Boulder, Colorado. National Technical Information Service PB82-192113/AS.
- Johnson, S.R., L.E. Noel, W.J. Gazey, and V.C. Hawkes. In prep. Aerial monitoring of marine waterfowl in the central Alaskan Beaufort Sea: 1989-1991 and 1999-2001.
- Larned, W.W. and G.R. Balogh. 1997. Eider breeding population survey Arctic Coastal Plain, Alaska, 1992–1996. Unpublished Report, U.S. Fish and Wildlife Service, Migratory Bird Management Division, Anchorage, AK. 51 p.
- Larned, W., T. Tiplady, R. Platte, and R. Stehn. 1999. Eider breeding population survey, Arctic Coastal Plain, Alaska, 1997-1998. Unpub. report by U.S. Fish and Wildlife Service, Office of Migratory Bird Management, Anchorage, AK. 22 p.
- Mallek, E.J. and R. J. King. 2000. Aerial breeding pair surveys of the Arctic Coastal Plain of Alaska, 1999. Unpublished report by U.S. Fish and Wildlife Service, Waterfowl Management, Fairbanks, AK. 17 p.
- Mallek, E.J., R. Platte, and R. Stehn. 2002. Aerial breeding pair surveys of the Arctic Coastal Plain of Alaska-2001. Unpublished report by U.S. Fish and Wildlife Service, Waterfowl Management, Fairbanks, AK. 25 p. + Append.

- Noel, L.E., S.R. Johnson, and P.F. Wainwright. 1999. Aerial surveys of molting waterfowl in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska, 1998. Report by LGL Alaska Research Associates, Inc. for BP Exploration (Alaska) Inc., Anchorage, AK. 53 p.
- Noel, L.E., S.R. Johnson, and P.F. Wainwright. 2000. Aerial surveys of molting waterfowl in the barrier island-lagoon systems between the Stockton Islands and Flaxman Island, Alaska, 1999. Report by LGL Alaska Research Associates, Inc. for BP Exploration (Alaska) Inc., Anchorage, AK. 64 p. + Append.
- Noel, L.E., S.R. Johnson, and R. Rodrigues. 2001. Aerial surveys of molting waterfowl in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska, 2000. Report by LGL Alaska Research Associates, Inc. for BP Exploration (Alaska) Inc., Anchorage, AK. 69 p. + Append.
- U.S. Fish and Wildlife Service (USFWS). 1999. Population status and trends of sea ducks in Alaska. Unpublished Report. U.S. Fish and Wildlife Service, Migratory Bird Management, Anchorage, AK 137 p.
- Zar, J.H. 1974. Biostatistical Analysis. Prentice Hall, Englewood Cliffs, NJ. 620 p.






Figure 1. Locations and numbers for aerial survey transects and survey regions in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska. Transect numbering is consistent with Johnson and Gazey (1992).


Beaufort Sea Waterfowl, 2001



Male long-tailed duck (*Clangula hyemalis*).
Photo by John Warden



Pair of long-tailed ducks.
Photo BPXA file



Nesting female common eider (*Somateria mollissima*
v-nigrum). Photo by John Warden



Pair of common eiders.
Photo by John Warden



Glaucous gull (*Larus hyperboreas*).
Photo by John Warden



Black brant (*Branta bernicla nigricans*).
Photo by Lynn Noel

Figure 2. Avian species occurring within the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska.




Photo by Lynn Noel

Survey aircraft; float-equipped Cessna 206.



Photo by Steve Johnson


Long-tailed duck molting habitat.



Photo by Lynn Noel

Common eider nesting habitat-Pole Island.

Figure 3. Survey aircraft and examples of habitats covered in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska.



Beaufort Sea Waterfowl, 2001

Figure 4. Long-tailed duck sightings as percentage of all bird sightings for aerial surveys with annual mean densities showing standard error in nearshore waters of the central Alaska Beaufort Sea, 1977-2001 (Table B1. Johnson and Gazey 1992; Noel et al. 1999, 2000, 2001).




Figure 5. Mean with standard error bar and standard deviation line for areal density of avian groups among survey regions for 2 or 3 aerial survey replicates between Spy Island and Brownlow Point, Alaska, 23 July-11 August 2001. Western lagoon and Stefansson Sound regions were surveyed on 3 dates. Two mainland shoreline transects in the eastern lagoon and the tundra region were surveyed on 2 dates.






Figure 6. Areal density of long-tailed ducks for 3 aerial surveys during 23 July-11 August 2001, and mean areal density with standard error for all surveys by region between Spy Island and Brownlow Point, Alaska. On 23 July 2001, two mainland transects in the eastern lagoon and all tundra transects were not flown because of fog.


WESTERN LAGOON (Spy Island to West Dock)



EASTERN LAGOON (Pole Island to Brownlow Point)



STEFANSSON SOUND (West Dock to Pole Island)



TUNDRA (Brownlow Point to Shavirovik River)




Figure 7. Total number of long-tailed ducks on- and off-transect during aerial surveys in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska, 23 July – 11 August 2001. Surveys were flown on 23 July, 8 August and 11 August 2001.




Figure 8. Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July-11 August 2001.




Figure 9. Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 1-24 August 2000.




Figure 10. Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 30 July-26 August 1999.




Figure 11. Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July-11 August 2001.




Figure 12. Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 1-24 August 2000.

Beaufort Sea Waterfowl, 2001




Figure 13. Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 30 July-26 August 1999.




Figure 14. Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 5 August-3 September 1998.




Figure 16. Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 1-24 August 2000.

Beaufort Sea Waterfowl, 2001






Figure 25. Summary of density for loons by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July-11 August 2001.


WESTERN LAGOON (Spy Island to West Dock)



STEFANSSON SOUND (West Dock to Pole Island)



EASTERN LAGOON (Pole Island to Brownlow Point)



TUNDRA (Brownlow Point to Shaviovik River)




Figure 26. Total number of seabirds on- and off-transect during aerial surveys in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001. Surveys were flown on 23 July, 8 August and 11 August 2001.




Figure 28. Summary of density for gulls by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July-11 August 2001.




Figure 29. Summary of density for gulls by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July-11 August 2001.




Figure 30. Summary of density for gulls by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July-11 August 2001.




Figure 31. Areal density of terns for 3 aerial surveys by region between Spy Island and Brownlow Point, Alaska, 23 July-11 August 2001. On 23 July 2001, two mainland transects in the eastern lagoon and all tundra transects were not flown because of fog.




Figure 32. Summary of density for Arctic terns by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July-11 August 2001.




Figure 33. Summary of density for Arctic terns by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July-11 August 2001.




Figure 34. Summary of density for Arctic terns by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July-11 August 2001.

Beaufort Sea Waterfowl, 2001







Figure 35. Total number of eiders on- and off-transect during aerial surveys in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001. Surveys were flown on 23 July, 8 August and 11 August 2001.




Figure 36. Areal density of eiders for 3 aerial surveys by region between Spy Island and Brownlow Point, Alaska, 23 July-11 August 2001. On 23 July 2001, two mainland transects in the eastern lagoon and all tundra transects were not flown because of fog.




Figure 37. Summary of density for eiders by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July-11 August 2001.

EIDERS 2001 - STEFANSSON SOUND
(West Dock to Pole Island)




Figure 38. Summary of density for eiders by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July-11 August 2001.

Beaufort Sea Waterfowl, 2001




Figure 39. Summary of density for eiders by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July-11 August 2001.

Beaufort Sea Waterfowl, 2001





Figure 40. Total number of scoters on- and off-transect during aerial surveys in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001. Surveys were flown on 23 July, 8 August and 11 August 2001.




Figure 41. Areal density of scoters for 3 aerial surveys by region between Spy Island and Brownlow Point, Alaska, 23 July-11 August 2001. On 23 July 2001, two mainland transects in the eastern lagoon and all tundra transects were not flown because of fog.




Figure 42. Summary of density for scoters by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July-11 August 2001.




Figure 43. Summary of density for scoters by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July-11 August 2001.







Figure 44. Total number of geese and swans observed both on- and off-transect during aerial surveys in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska, 23 July–11 August 2001. Surveys were flown on 23 July, 8 August and 11 August 2001.




Figure 45. Areal density of geese and swans for 3 aerial surveys by region between Spy Island and Brownlow Point, Alaska, 23 July-11 August 2001. On 23 July 2001, two mainland transects in the eastern lagoon and all tundra transects were not flown because of fog.




Figure 46. Summary of density for geese and swans by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July-11 August 2001.




Figure 47. Summary of density for geese and swans by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July-11 August 2001.

Table 3. Total number of bird sightings and individuals seen on- and off-transect for all aerial survey transects (total length = 1986.4 km) in the barrier island-lagoon system, offshore, and on tundra between Spy Island and Brownlow Point, Alaska, 23 July - 11 August 2001.

| Species Code | Species Name | Number of Sightings | Percent of Sightings | Number of Individuals | Percent of Individuals | Percent of Classified for Group |
|---|---|---------------------|----------------------|-----------------------|------------------------|---------------------------------|
| PALO | Pacific Loon (<i>Gavia pacifica</i>) | 146 | 7.9 | 212 | 0.6 | 70.7 |
| RTLO | Red-throated Loon (<i>Gavia stellata</i>) | 59 | 3.2 | 80 | 0.2 | 26.7 |
| YBLO | Yellow-billed Loon (<i>Gavia adamsii</i>) | 8 | 0.4 | 8 | 0.0 | 2.7 |
| LOSP | Loon Species (<i>Gavia</i> spp.) | 7 | 0.4 | 8 | 0.0 | |
| Loons | | 220 | 11.9 | 308 | 0.9 | |
| POJA | Pomarine Jaeger (<i>Stercorarius pomarinus</i>) | 1 | 0.1 | 2 | 0.0 | 100.0 |
| Jaegers | | 1 | 0.1 | 2 | 0.0 | |
| GLGU | Glaucous Gull (<i>Larus hyperboreus</i>) | 347 | 18.7 | 856 | 2.5 | 94.6 |
| SAGU | Sabine's Gull (<i>Xema sabini</i>) | 3 | 0.2 | 49 | 0.1 | 5.4 |
| Gulls | | 350 | 18.9 | 905 | 2.6 | |
| ARTE | Arctic Tern (<i>Sterna paradisaea</i>) | 21 | 1.1 | 34 | 0.1 | |
| COMU | Common Murre (<i>Uria aalge</i>) | 1 | 0.1 | 1 | 0.0 | |
| BLGU | Black Guillemot (<i>Cephus grylle</i>) | 1 | 0.1 | 1 | 0.0 | |
| SEABIRDS (Jaegers, Gulls, and Terns) | | 374 | 20.1 | 942 | 2.7 | |
| COEI | Common (Pacific) Eider (<i>Somateria mollissima v-nigrum</i>) | 190 | 10.3 | 2240 | 6.4 | 99.5 |
| KIEI | King Eider (<i>Somateria spectabilis</i>) | 1 | 0.1 | 11 | 0.0 | 0.5 |
| EISP | Eider Species (<i>Somateria</i> spp.) | 6 | 0.3 | 104 | 0.3 | |
| Eiders | | 197 | 10.6 | 2355 | 6.7 | |
| SUSC | Surf Scoter (<i>Melanitta perspicillata</i>) | 71 | 3.8 | 698 | 2.0 | 99.6 |
| BLSC | Black Scoter (<i>Melanitta nigra</i>) | 1 | 0.1 | 1 | 0.0 | 0.1 |
| SCOT | Scoter Species (<i>Melanitta</i> spp.) | 2 | 0.1 | 2 | 0.0 | 0.3 |
| Scoters | | 74 | 4.0 | 701 | 2.0 | |
| NOPI | Northern Pintail (<i>Anas acuta</i>) | 2 | 0.1 | 24 | 0.1 | |
| OLDS | Long-tailed Duck (<i>Clangula hyemalis</i>) | 857 | 46.3 | 26878 | 77.0 | |
| RBME | Red-breasted Merganser (<i>Mergus serrator</i>) | 1 | 0.1 | 1 | 0.0 | |
| GRSC | Greater Scaup (<i>Aythya marila</i>) | 1 | 0.1 | 20 | 0.1 | |
| SCAUP | Scaup Species (<i>Aythya</i> spp.) | 3 | 0.2 | 10 | 0.0 | |
| DKSP | Duck Species | 22 | 1.2 | 201 | 0.6 | |
| Ducks | | 1157 | 62.5 | 30190 | 86.5 | |
| BRAN | Black Brant (<i>Branta bernicla</i>) | 21 | 1.1 | 1129 | 3.2 | 33.6 |
| CAGO | Canada Goose (<i>Branta canadensis</i>) | 17 | 0.9 | 461 | 1.3 | 13.7 |
| GWGO | Greater White-fronted Goose (<i>Anser albifrons</i>) | 31 | 1.7 | 1230 | 3.5 | 36.6 |
| LSGO | Lesser Snow Goose (<i>Chen caerulescens caerulescens</i>) | 8 | 0.4 | 520 | 1.5 | 15.5 |
| TUSW | Tundra Swan (<i>Cygnus columbianus</i>) | 7 | 0.4 | 19 | 0.1 | 0.6 |
| DAGO | Dark Goose | 2 | 0.1 | 100 | 0.3 | |
| Geese & Swans | | 84 | 4.5 | 3359 | 9.6 | |
| WATERFOWL (Ducks, Geese & Swans) | | 1241 | 67.0 | 33549 | 96.1 | |
| PHSP | Phalarope Species (<i>Phalaropus</i> spp.) | 3 | 0.2 | 81 | 0.2 | |
| SMSH | Small Shorebird | 5 | 0.3 | 11 | 0.0 | |
| MESH | Medium Shorebird | 2 | 0.1 | 5 | 0.0 | |
| SHOREBIRDS | | 10 | 0.5 | 97 | 0.3 | |
| SNOW | Snowy Owl (<i>Nyctea scandiaca</i>) | 1 | 0.1 | 1 | 0.0 | |
| SNBU | Snow Bunting (<i>Plectrophenax nivalis</i>) | 1 | 0.1 | 1 | 0.0 | |
| RLHA | Rough-legged Hawk (<i>Buteo lagopus</i>) | 1 | 0.1 | 1 | 0.0 | |
| BISP | Bird Species | 4 | 0.2 | 8 | 0.0 | |
| ALL BIRDS | | 1852 | 100 | 34907 | 100 | |

Table 4. Total number of bird sightings and individuals seen on- and off-transect for aerial survey transects (total length = 847.2 km) in the barrier island-lagoon system and offshore between Spy Island and West Dock, Alaska, 23 July-11 August 2001.

| Species Code | WESTERN LAGOON Species Name | Number of Sightings | Percent of Sightings | Number of Individuals | Percent of Individuals | Percent of Classified for Group |
|---|---|---------------------|----------------------|-----------------------|------------------------|---------------------------------|
| PALO | Pacific Loon (<i>Gavia pacifica</i>) | 93 | 9.1 | 129 | 0.8 | 62.6 |
| RTLO | Red-throated Loon (<i>Gavia stellata</i>) | 51 | 5.0 | 71 | 0.4 | 34.5 |
| YBLO | Yellow-billed Loon (<i>Gavia adamsii</i>) | 6 | 0.6 | 6 | 0.0 | 2.9 |
| LOSP | Loon Species (<i>Gavia</i> spp.) | 7 | 0.7 | 8 | 0.0 | |
| Loons | | 157 | 15.3 | 214 | 1.3 | |
| GLGU | Glaucous Gull (<i>Larus hyperboreus</i>) | 208 | 20.3 | 559 | 3.4 | 93.2 |
| SAGU | Sabine's Gull (<i>Xema sabini</i>) | 2 | 0.2 | 41 | 0.2 | 6.8 |
| Gulls | | 210 | 20.4 | 600 | 3.6 | |
| ARTE | Arctic Tern (<i>Sterna paradisaea</i>) | 5 | 0.5 | 6 | 0.0 | |
| SEABIRDS (Jaegers, Gulls, and Terns) | | 215 | 20.93 | 606 | 3.7 | |
| COEI | Common (Pacific) Eider (<i>Somateria mollissima v-nigrum</i>) | 30 | 2.9 | 219 | 1.3 | 100.0 |
| EISP | Eider Species (<i>Somateria</i> spp.) | 2 | 0.2 | 45 | 0.3 | |
| Eiders | | 32 | 3.1 | 264 | 1.6 | |
| SUSC | Surf Scoter (<i>Melanitta perspicillata</i>) | 65 | 6.3 | 683 | 4.2 | 100.0 |
| SCOT | Scoter Species (<i>Melanitta</i> spp.) | 2 | 0.2 | 2 | 0.0 | |
| Scoters | | 67 | 6.5 | 685 | 4.2 | |
| NOPI | Northern Pintail (<i>Anas acuta</i>) | 1 | 0.1 | 4 | 0.0 | |
| OLDS | Long-tailed Duck (<i>Clangula hyemalis</i>) | 479 | 46.6 | 12207 | 74.2 | |
| GRSC | Greater Scaup (<i>Aythya marila</i>) | 1 | 0.1 | 20 | 0.1 | |
| SCAUP | Scaup Species (<i>Aythya</i> spp.) | 1 | 0.1 | 2 | 0.0 | |
| DKSP | Duck Species | 14 | 1.4 | 147 | 0.9 | |
| Ducks | | 595 | 57.8 | 13329 | 81.0 | |
| BRAN | Black Brant (<i>Branta bernicla</i>) | 21 | 2.0 | 1129 | 6.9 | 49.5 |
| GWGO | Greater White-fronted Goose (<i>Anser albifrons</i>) | 25 | 2.4 | 1009 | 6.1 | 44.3 |
| CAGO | Canada Goose (<i>Branta canadensis</i>) | 6 | 0.6 | 140 | 0.9 | 6.1 |
| TUSW | Tundra Swan (<i>Cygnus columbianus</i>) | 1 | 0.1 | 2 | 0.0 | 0.1 |
| DAGO | Dark Goose | 1 | 0.1 | 20 | 0.1 | |
| Geese & Swans | | 53 | 5.2 | 2280 | 13.9 | |
| WATERFOWL (Ducks, Geese & Swans) | | 648 | 63.0 | 15609 | 94.9 | |
| MESH | Medium Shorebird | 1 | 0.1 | 2 | | |
| SMSH | Small Shorebird | 1 | 0.1 | 1 | 0.0 | |
| SHOREBIRDS | | 2 | 0.1 | 3 | 0.0 | |
| RLHA | Rough-legged Hawk (<i>Buteo lagopus</i>) | 1 | 0.1 | 1 | 0.0 | |
| RAPTORS | | 1 | 0.1 | 1 | 0.0 | |
| BISP | Bird Species | 4 | 0.4 | 8 | 0.0 | |
| ALL BIRDS | | 1027 | 100 | 16441 | 100 | |

Table 5 Total number of bird sightings and individuals seen on- and off-transect for aerial survey transects (total length = 202.9 km) in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July-11 August 2001.

| Species Code | STEFANSSON SOUND Species Name | Number of Sightings | Percent of Sightings | Number of Individuals | Percent of Individuals | Percent of Classified for Group |
|---|---|---------------------|----------------------|-----------------------|------------------------|---------------------------------|
| PALO | Pacific Loon (<i>Gavia pacifica</i>) | 6 | 3.9 | 11 | 0.6 | 78.6 |
| RTLO | Red-throated Loon (<i>Gavia stellata</i>) | 2 | 1.3 | 2 | 0.1 | 14.3 |
| YBLO | Yellow-billed Loon (<i>Gavia adamsii</i>) | 1 | 0.6 | 1 | 0.1 | 7.1 |
| Loons | | 9 | 5.8 | 14 | 0.8 | |
| GLGU | Glaucous Gull (<i>Larus hyperboreus</i>) | 27 | 17.4 | 46 | 2.6 | 100.0 |
| Gulls | | 27 | 17.4 | 46 | 2.6 | |
| ARTE | Arctic Tern (<i>Sterna paradisaea</i>) | 11 | 7.1 | 18 | 1.0 | |
| BLGU | Black Guillemot (<i>Cepphus grylle</i>) | 1 | 0.6 | 1 | 0.1 | |
| SEABIRDS (Jaegers, Gulls, and Terns) | | 39 | 25.2 | 65 | 3.7 | |
| COEI | Common (Pacific) Eider (<i>Somateria mollissima v-nigrum</i>) | 52 | 33.5 | 251 | 14.1 | 100.0 |
| Eiders | | 52 | 33.5 | 251 | 14.1 | |
| OLDS | Long-tailed Duck (<i>Clangula hyemalis</i>) | 52 | 33.5 | 1370 | 77.2 | |
| DKSP | Duck Species | 1 | 0.6 | 4 | 0.2 | |
| Ducks | | 105 | 67.7 | 1625 | 91.6 | |
| LSGO | Lesser Snow Goose (<i>Chen caerulescens caerulescens</i>) | 1 | 0.6 | 30 | 1.7 | 100.0 |
| Geese & Swans | | 1 | 0.6 | 30 | 1.7 | |
| WATERFOWL (Ducks, Geese & Swans) | | 106 | 68.4 | 1655 | 93.3 | |
| PHSP | Phalarope Species (<i>Phalaropus</i> spp.) | 1 | 0.6 | 40 | 2.3 | 100.0 |
| SHOREBIRDS | | 1 | 0.6 | 40 | 2.3 | |
| ALL BIRDS | | 155 | 100 | 1774 | 100 | |

Table 6. Total number of bird sightings and individuals seen on- and off-transect for aerial survey transects (total length = 813.8 km) in the barrier island-lagoon system and offshore between Pole Island and Brownlow Point, Alaska, 23 July-11 August 2001.

| Species Code | EASTERN LAGOON Species Name | Number of Sightings | Percent of Sightings | Number of Individuals | Percent of Individuals | Percent of Classified for Group |
|---|---|---------------------|----------------------|-----------------------|------------------------|---------------------------------|
| PALO | Pacific Loon (<i>Gavia pacifica</i>) | 30 | 5.8 | 44 | 0.3 | 88.0 |
| RTLO | Red-throated Loon (<i>Gavia stellata</i>) | 5 | 1.0 | 5 | 0.0 | 10.0 |
| YBLO | Yellow-billed Loon (<i>Gavia adamsii</i>) | 1 | 0.2 | 1 | 0.0 | 2.0 |
| Loons | | 36 | 7.0 | 50 | 0 | |
| GLGU | Glaucous Gull (<i>Larus hyperboreus</i>) | 109 | 21.2 | 248 | 1.7 | 96.9 |
| SAGU | Sabine's Gull (<i>Xema sabini</i>) | 1 | 0.2 | 8 | 0.1 | 3.1 |
| Gulls | | 110 | 21.4 | 256 | 1.8 | |
| ARTE | Arctic Tern (<i>Sterna paradisaea</i>) | 5 | 1.0 | 10 | 0.1 | |
| COMU | Common Murre (<i>Uria aalge</i>) | 1 | 0.2 | 1 | 0.0 | |
| SEABIRDS (Jaegers, Gulls, and Terns) | | 116 | 22.5 | 267 | 1.8 | |
| COEI | Common (Pacific) Eider (<i>Somateria mollissima v-nigrum</i>) | 108 | 21.0 | 1770 | 12.1 | 99.4 |
| KIEI | King Eider (<i>Somateria spectabilis</i>) | 1 | 0.2 | 11 | 0.1 | 0.6 |
| EISP | Eider Species (<i>Somateria</i> spp.) | 3 | 0.6 | 54 | 0.4 | |
| Eiders | | 4 | 0.8 | 65 | 0.4 | |
| SUSC | Surf Scoter (<i>Melanitta perspicillata</i>) | 6 | 1.2 | 15 | 0.1 | 93.8 |
| BLSC | Black Scoter (<i>Melanitta nigra</i>) | 1 | 0.2 | 1 | 0.0 | 6.3 |
| Scoters | | 7 | 1.4 | 16 | 0.1 | |
| OLDS | Long-tailed Duck (<i>Clangula hyemalis</i>) | 321 | 62.3 | 13249 | 90.9 | |
| SCAUP | Scaup Species (<i>Aythya</i> spp.) | 1 | 0.2 | 6 | 0.0 | |
| NOPI | Northern Pintail (<i>Anas acuta</i>) | 1 | 0.2 | 20 | 0.1 | |
| RBME | Red-breasted Merganser (<i>Mergus serrator</i>) | 1 | 0.2 | 1 | 0.0 | |
| DKSP | Duck Species | 7 | 1.4 | 50 | 0.3 | |
| Ducks | | 342 | 66.4 | 13407 | 92.0 | |
| CAGO | Canada Goose (<i>Branta canadensis</i>) | 5 | 1.0 | 210 | 1.4 | 24.8 |
| GWGO | Greater White-fronted Goose (<i>Anser albifrons</i>) | 4 | 0.8 | 141 | 1.0 | 16.7 |
| LSGO | Lesser Snow Goose (<i>Chen caerulescens caerulescens</i>) | 7 | 1.4 | 490 | 3.4 | 57.9 |
| TUSW | Tundra Swan (<i>Cygnus columbianus</i>) | 2 | 0.4 | 5 | 0.0 | 0.6 |
| DAGO | Dark Goose | 1 | 0.2 | 80 | 0.5 | |
| Geese & Swans | | 18 | 3.5 | 846 | 5.8 | |
| WATERFOWL (Ducks, Geese & Swans) | | 360 | 70 | 14253 | 98 | |
| SMSH | Small Shorebird | 1 | 0.2 | 1 | 0.0 | |
| MESH | Medium Shorebird | 1 | 0.2 | 3 | 0.0 | |
| SHOREBIRDS | | 2 | 0.4 | 4 | 0.0 | |
| SNOW | Snowy Owl (<i>Nyctea scandiaca</i>) | 1 | | 1 | | |
| ALL BIRDS | | 515 | 100 | 14575 | 100 | |

Table 7. Total number of bird sightings and individuals seen on- and off-transect for aerial survey transects (total length = 122.5 km) on tundra between Brownlow Point and the Shaviovik River, Alaska, 8 and 11 August 2001.

| Species Code | <i>TUNDRA</i> Species Name | Number of Sightings | Percent of Sightings | Number of Individuals | Percent of Individuals | Percent of Classified for Group |
|---|--|---------------------|----------------------|-----------------------|------------------------|---------------------------------|
| PALO | Pacific Loon (<i>Gavia pacifica</i>) | 17 | 36.2 | 28 | 8.0 | 93.3 |
| RTLO | Red-throated Loon (<i>Gavia stellata</i>) | 1 | 2.1 | 2 | 0.6 | 6.7 |
| Loons | | 18 | 38.3 | 30 | 8.6 | |
| POJA | Pomarine Jaeger (<i>Stercorarius pomarinus</i>) | 1 | 2.1 | 2 | 0.6 | 100.0 |
| Jaegers | | 1 | 2.1 | 2 | 0.6 | |
| GLGU | Glaucous Gull (<i>Larus hyperboreus</i>) | 3 | 6.4 | 3 | 0.9 | 100.0 |
| Gulls | | 3 | 6.4 | 3 | 0.9 | |
| SEABIRDS (Jaegers, Gulls, and Terns) | | 4 | 8.5 | 5 | 1.4 | |
| EISP | Eider Species (<i>Somateria</i> spp.) | 1 | 2.1 | 5 | 1.4 | |
| Eiders | | 1 | 2.1 | 5 | 1.4 | |
| OLDS | Long-tailed Duck (<i>Clangula hyemalis</i>) | 5 | 10.6 | 52 | 14.9 | |
| SCAUP | Scaup Species (<i>Aythya</i> spp.) | 1 | 2.1 | 2 | 0.6 | |
| Ducks | | 7 | 14.9 | 59 | 17.0 | |
| CAGO | Canada Goose (<i>Branta canadensis</i>) | 6 | 12.8 | 111 | 31.9 | 54.7 |
| GWGO | Greater White-fronted Goose (<i>Anser albifrons</i>) | 2 | 4.3 | 80 | 23.0 | 39.4 |
| TUSW | Tundra Swan (<i>Cygnus columbianus</i>) | 4 | 8.5 | 12 | 3.4 | 5.9 |
| Geese & Swans | | 12 | 25.5 | 203 | 58.3 | |
| WATERFOWL (Ducks, Geese & Swans) | | 19 | 40.4 | 262 | 75.3 | |
| PHSP | Phalarope Species (<i>Phalaropus</i> spp.) | 2 | 4.3 | 41 | 11.8 | |
| SMSH | Small Shorebirds | 3 | 6.4 | 9 | 2.6 | |
| SHOREBIRDS | | 5 | 10.6 | 50 | 14.4 | |
| SNBU | Snow Bunting (<i>Plectrophenax nivalis</i>) | 1 | 2.1 | 1 | 0.3 | |
| ALL BIRDS | | 47 | 100 | 348 | 100 | |

Table 8. Long-tailed duck density (number of individuals/km²) by aerial survey transect in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July - 11 August 2001.

| Date | Transect Numbers | | | | | | | | | | | | | | | | | | |
|----------------------------|------------------|--------|-------|-------|------------|-------|-------|--------|-------|--------|----------------|------|------|------|-----------|-------|------|------|------|
| | Barrier Islands | | | | Mid-Lagoon | | | | | | Mainland Shore | | | | Off Shore | | | | |
| | 23 | 31 | 202 | 201 | 24 | 32 | 301 | 302 | 601 | 602 | 603 | 25 | 33 | 401 | 402 | 22 | 30 | 101 | 102 |
| Molt (15 July - 21 August) | | | | | | | | | | | | | | | | | | | |
| 23 Jul 01 | 15.50 | 124.44 | 34.18 | 41.25 | 3.29 | 32.57 | 27.92 | 28.48 | 55.21 | 10.66 | 32.43 | 0.44 | 0.00 | 3.69 | 0.71 | 24.06 | 7.46 | 2.67 | 3.15 |
| 8 Aug 01 | 21.36 | 218.39 | 27.04 | 36.84 | 0.72 | 68.18 | 45.08 | 182.79 | 91.03 | 119.04 | 40.61 | 6.53 | 0.53 | 9.39 | 45.18 | 0.00 | 0.38 | 2.75 | 0.00 |
| 11 Aug 01 | 1.25 | 34.79 | 0.97 | 4.13 | 0.00 | 0.84 | 9.04 | 34.76 | 0.00 | 90.26 | 29.70 | 2.16 | 0.00 | 6.32 | 6.08 | 1.58 | 0.00 | 0.12 | 0.00 |
| Average Density | 12.70 | 125.87 | 20.73 | 27.41 | 1.34 | 33.86 | 27.35 | 82.01 | 48.75 | 73.32 | 34.25 | 3.04 | 0.18 | 6.47 | 17.32 | 8.55 | 2.61 | 1.85 | 1.05 |

Table 9. Long-tailed duck density (number of individuals/km²) by aerial survey transect in the barrier island-lagoon system and on tundra between Pole Island and Brownlow Point, Alaska, 23 July-11 August 2001.

| Date | Barrier Islands | | | | Transect Numbers | | | | | | | | Mainland Shore | | | |
|---------------------------------|-----------------|-------|-------|-------|------------------|------|------|------|------|------|-------|-------|----------------|--------|--------|--------|
| | 133 | 134 | 135 | 136 | 180 | 181 | 182 | 183 | 604 | 605 | 606 | 607 | 190 | 191 | 192 | 193 |
| MOLT 23 July - 11 August | | | | | | | | | | | | | | | | |
| 23 Jul 01 | 133.22 | 49.80 | 42.19 | 95.92 | 35.46 | 9.96 | 0.00 | 0.00 | 0.00 | 0.38 | 2.89 | 34.38 | - | - | 118.75 | 206.34 |
| 8 Aug 01 | 211.25 | 56.65 | 5.87 | 28.08 | 20.95 | 5.20 | 7.96 | 0.00 | 0.00 | 7.63 | 73.16 | 40.26 | 133.59 | 235.64 | 123.63 | 44.35 |
| 11 Aug 01 | 68.35 | 0.00 | 0.00 | 10.78 | 3.08 | 8.61 | 0.00 | 0.20 | 0.00 | 0.00 | 11.65 | 4.46 | 25.31 | 34.20 | 42.11 | 9.09 |
| Average Density | 137.61 | 35.48 | 16.02 | 44.93 | 19.83 | 7.92 | 2.65 | 0.07 | 0.00 | 2.67 | 29.23 | 26.37 | 79.45 | 134.92 | 94.83 | 86.59 |
| Date | Off Shore | | | | Tundra | | | | | | | | | | | |
| | 60 | 61 | 62 | 63 | 500 | 501 | 502 | 503 | | | | | | | | |
| MOLT 23 July - 11 August | | | | | | | | | | | | | | | | |
| 23 Jul 01 | 4.88 | 9.00 | 0.00 | 0.00 | - | - | - | - | | | | | | | | |
| 8 Aug 01 | 0.00 | 0.20 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.18 | | | | | | | | |
| 11 Aug 01 | 11.54 | 0.00 | 0.61 | 0.00 | 0.00 | 0.00 | 0.92 | 8.15 | | | | | | | | |
| Average Density | 5.47 | 3.07 | 0.20 | 0.00 | 0.00 | 0.00 | 0.46 | 4.17 | | | | | | | | |

Table 10. Long-tailed duck density (number of individuals/km²) by aerial survey transect in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July-11 August 2001.

| Date | Transect Numbers | | |
|-----------------------------------|------------------|-------|-------|
| | Barrier Islands | | |
| | 130 | 131 | 132 |
| Molt (15 July - 21 August) | | | |
| 23 Jul 01 | 9.32 | 43.01 | 11.31 |
| 8 Aug 01 | 2.05 | 12.66 | 15.53 |
| 11 Aug 01 | 4.27 | 3.11 | 32.61 |
| Average Density | 5.21 | 19.59 | 19.82 |

Table 11. Species densities for all aerial survey transects (total area = 790.38 km²) in the barrier island-lagoon systems and on tundra between Spy Island and Brownlow Point, Alaska, 23 July-11 August 2001.

| Species Code | Species Name | Total Number of Birds on Transect | Number of Sightings on Transect | Bird Density (Number/km ²) |
|---|---|---|---------------------------------------|---|
| LOONS | | | | |
| PALO | Pacific Loon (<i>Gavia pacifica</i>) | 160 | 114 | 0.202 |
| RTLO | Red-throated Loon (<i>Gavia stellata</i>) | 76 | 57 | 0.096 |
| YBLO | Yellow-billed Loon (<i>Gavia adamsii</i>) | 8 | 8 | 0.010 |
| LOSP | Loon Species (<i>Gavia</i> spp.) | 2 | 2 | 0.003 |
| SEABIRDS (Jaegers, Gulls, and Terns) | | | | |
| GLGU | Glaucous Gull (<i>Larus hyperboreus</i>) | 772 | 314 | 0.977 |
| SAGU | Sabine's Gull (<i>Xema sabini</i>) | 49 | 3 | 0.062 |
| ARTE | Arctic Tern (<i>Sterna paradisaea</i>) | 34 | 21 | 0.043 |
| BLGU | Black Guillemot (<i>Cephus grylle</i>) | 1 | 1 | 0.001 |
| COMU | Common Murre (<i>Uria aalge</i>) | 1 | 1 | 0.001 |
| WATERFOWL (Ducks, Geese & Swans) | | | | |
| COEI | Common (Pacific) Eider (<i>Somateria mollissima v-nigrum</i>) | 1894 | 172 | 2.396 |
| KIEI | King Eider (<i>Somateria spectabilis</i>) | 11 | 1 | 0.014 |
| EISP | Eider Species (<i>Somateria</i> spp.) | 99 | 5 | 0.125 |
| SUSC | Surf Scoter (<i>Melanitta perspicillata</i>) | 643 | 68 | 0.814 |
| BLSC | Black Scoter (<i>Melanitta nigra</i>) | 1 | 1 | 0.001 |
| SCOT | Scoter Species (<i>Melanitta</i> spp.) | 1 | 1 | 0.001 |
| OLDS | Long-tailed Duck (<i>Clangula hyemalis</i>) | 22660 | 784 | 28.670 |
| GRSC | Greater Scaup (<i>Aythya marila</i>) | 20 | 1 | 0.025 |
| SCAUP | Scaup Species (<i>Aythya</i> spp.) | 8 | 2 | 0.010 |
| NOPI | Northern Pintail (<i>Anas acuta</i>) | 4 | 1 | 0.005 |
| RBME | Red-breasted Merganser (<i>Mergus serrator</i>) | 1 | 1 | 0.001 |
| DKSP | Duck Species | 11 | 3 | 0.014 |
| DAGO | Dark Goose | 20 | 1 | 0.025 |
| BRAN | Black Brant (<i>Branta bernicla</i>) | 1069 | 20 | 1.353 |
| CAGO | Canada Goose (<i>Branta canadensis</i>) | 348 | 10 | 0.440 |
| GWGO | Greater White-fronted Goose (<i>Anser albifrons</i>) | 893 | 26 | 1.130 |
| LSGO | Lesser Snow Goose (<i>Chen caerulescens caerulescens</i>) | 220 | 6 | 0.278 |
| TUSW | Tundra Swan (<i>Cygnus columbianus</i>) | 5 | 2 | 0.006 |
| Other Species | | | | |
| PHSP | Phalarope Species (<i>Phalaropus</i> spp.) | 40 | 1 | 0.051 |
| RLHA | Rough-legged Hawk (<i>Buteo lagopus</i>) | 1 | 1 | 0.001 |
| SMSH | Small Shorebirds | 2 | 2 | 0.003 |
| MESH | Medium Shorebirds | 5 | 2 | 0.006 |
| BISP | Bird Species | 2 | 2 | 0.003 |

Table 12. Habitat associations of loons during aerial surveys in the barrier island-lagoon systems and tundra transects between Spy Island and Brownlow Point, Alaska, 23 July-11 August 2001

| General Habitat Type | Specific Habitat Type | Pacific Loon | | Red-throated Loon | | Yellow-billed Loon | |
|---------------------------|-----------------------------|---------------------|------------------|---------------------|------------------|---------------------|------------------|
| | | Number of Sightings | Percent of Total | Number of Sightings | Percent of Total | Number of Sightings | Percent of Total |
| Lagoon (9) | Lagoon (9) | 55 | 38 | 23 | 39 | 3 | 38 |
| Barrier Island (11) | Shoreline (water side; 15) | 27 | 18 | 11 | 19 | 4 | 50 |
| Nearshore Sea <3 mi. (13) | Ocean (8) | 13 | 9 | 9 | 15 | - | - |
| | Lagoon (9) | 3 | 2 | - | - | - | - |
| Wet Tundra (25) | Pond or Lake (18) | 1 | 1 | - | - | - | - |
| | Completely vegetated (25) | 1 | 1 | - | - | - | - |
| | Pond without Emergents (46) | 1 | 1 | - | - | - | - |
| | Pond (47) | 6 | 4 | - | - | - | - |
| | River Delta (54) | 1 | 1 | - | - | - | - |
| | Stream (55) | 1 | 1 | - | - | - | - |
| | Pond with Emergents (45) | 6 | 4 | - | - | - | - |
| Mainland Coast (27) | Shoreline (water side; 15) | 28 | 19 | 14 | 24 | 1 | 13 |
| | River Delta (16) | 1 | 1 | - | - | - | - |
| River Delta (92) | Tide Flat (13) | 2 | 1 | 2 | 3 | - | - |
| TOTAL | | 146 | 100 | 59 | 100 | 8 | 100 |

Table 13. Habitat associations of seabirds during aerial surveys in the barrier island-lagoon systems and tundra transects between Spy Island and Brownlow Point, Alaska, 23 July-11 August 2001.

| General Habitat Type | Specific Habitat Type | Arctic Tern | | Glaucous Gull | | Sabine's Gull | |
|---------------------------|----------------------------|---------------------|------------------|---------------------|------------------|---------------------|------------------|
| | | Number of Sightings | Percent of Total | Number of Sightings | Percent of Total | Number of Sightings | Percent of Total |
| Lagoon (9) | Lagoon (9) | - | - | 25 | 7.2 | 1 | 33.3 |
| | Smooth fast ice(1) | - | - | 1 | 0.3 | - | - |
| Barrier Island (11) | Spit (10) | - | - | 6 | 1.7 | - | - |
| | Island (11) | - | - | 16 | 4.6 | - | - |
| | Shoal (12) | 1 | 4.8 | 3 | 0.9 | - | - |
| | Shoreline (land side; 14) | - | - | 9 | 2.6 | - | - |
| | Shoreline (water side; 15) | 13 | 61.9 | 158 | 45.5 | 1 | 33.3 |
| Nearshore Sea <3 mi. (13) | Ocean (8) | 4 | 19.0 | 3 | 0.9 | 1 | 33.3 |
| | Lagoon (9) | 3 | 14.3 | 11 | 3.2 | - | - |
| Wet Tundra (25) | Pond (47) | - | - | 1 | 0.3 | - | - |
| | Lake Shore (44) | - | - | 1 | 0.3 | - | - |
| Mainland Coast (27) | Spit (10) | - | - | 3 | 0.9 | - | - |
| | Island (11) | - | - | 4 | 1.2 | - | - |
| | Shoal (12) | - | - | 1 | 0.3 | - | - |
| | Shoreline (land side; 14) | - | - | 6 | 1.7 | - | - |
| | Shoreline (water side; 15) | - | - | 85 | 24.5 | - | - |
| | River Delta (16) | - | - | 1 | 0.3 | - | - |
| | Mudflat (41) | - | - | 1 | 0.3 | - | - |
| River Delta (92) | Tide Flat (13) | - | - | 8 | 2.3 | - | - |
| | Mudflat (41) | - | - | 4 | 1.2 | - | - |
| TOTAL | | 21 | 100 | 347 | 100 | 3 | 100 |

Table 14. Habitat associations of eiders during aerial surveys in the barrier island-lagoon systems and tundra transects between Spy Island and Brownlow Point, Alaska, 23 July-11 August 2001.

| General Habitat Type | Specific Habitat Type | Common Eider | | King Eider | | Eider Species | | All Eiders | |
|---------------------------|----------------------------|---------------------|------------------|---------------------|------------------|---------------------|------------------|---------------------|------------------|
| | | Number of Sightings | Percent of Total | Number of Sightings | Percent of Total | Number of Sightings | Percent of Total | Number of Sightings | Percent of Total |
| Lagoon (9) | Lagoon (9) | 29 | 15.3 | 1 | 100 | - | - | 30 | 15.6 |
| | Smooth fast ice (1) | 2 | 1.1 | - | - | - | - | - | - |
| Barrier Island (11) | Lagoon (9) | 1 | 0.5 | - | - | - | - | - | - |
| | Smooth fast ice (1) | 1 | 0.5 | - | - | - | - | - | - |
| | Shoreline (land side; 14) | 2 | 1.1 | - | - | - | - | 2 | 1.0 |
| | Shoreline (water side; 15) | 102 | 53.7 | - | - | 1 | 16.7 | 103 | 53.6 |
| | Shoal (12) | 2 | 1.1 | - | - | - | - | 2 | 1.0 |
| Nearshore Sea <3 mi. (13) | Ocean (8) | 12 | 6.3 | - | - | - | - | 12 | 6.3 |
| | Lagoon (9) | 20 | 10.5 | - | - | - | - | 20 | 10.4 |
| Wet Tundra (25) | Pond (47) | - | - | - | - | 1 | 16.7 | 1 | 0.5 |
| Mainland Coast (27) | Spit (10) | 1 | 0.5 | - | - | - | - | - | - |
| | Shoreline (water side; 15) | 18 | 9.5 | - | - | 4 | 66.7 | 22 | 11.5 |
| TOTAL | | 190 | 100 | 1 | 100 | 6 | 100 | 192 | 100 |

Table 15. Habitat associations of scoters during aerial surveys in the barrier island-lagoon systems between Spy Island and Brownlow Point, Alaska, 23 July-11 August 2001.

| General Habitat Type | Specific Habitat Type | Surf Scoter | | Black Scoter | | Scoter Species | | All Scoters | |
|---------------------------|----------------------------|---------------------|------------------|---------------------|------------------|---------------------|------------------|---------------------|------------------|
| | | Number of Sightings | Percent of Total | Number of Sightings | Percent of Total | Number of Sightings | Percent of Total | Number of Sightings | Percent of Total |
| Lagoon (9) | Lagoon (9) | 55 | 77.5 | 1 | 100 | - | - | 56 | 75.7 |
| Barrier Island (11) | Shoreline (water side; 15) | 13 | 18.3 | - | - | 1 | 50 | 14 | 18.9 |
| Nearshore Sea <3 mi. (13) | Ocean (8) | 2 | 2.8 | - | - | - | - | 2 | 2.7 |
| Mainland Coast (27) | Shoreline (water side; 15) | 1 | 1.4 | - | - | 1 | 50 | 2 | 2.7 |
| TOTAL | | 71 | 100 | 1 | 100 | 2 | 100 | 74 | 100 |

Table 16. Habitat associations of geese and swans during aerial surveys in the barrier island-lagoon systems and tundra transects between Spy Island and Brownlow Point, Alaska, 23 July-11 August 2001.

| General Habitat Type | Specific Habitat Type | Black Brant | | Canada Goose | | Greater White-fronted Goose | | Tundra Swan | |
|----------------------|----------------------------|---------------------|------------------|---------------------|------------------|-----------------------------|------------------|---------------------|------------------|
| | | Number of Sightings | Percent of Total | Number of Sightings | Percent of Total | Number of Sightings | Percent of Total | Number of Sightings | Percent of Total |
| Barrier Island (11) | Shoreline (water side; 15) | 1 | 4.8 | 2 | 11.8 | 1 | 3.2 | - | - |
| Wet Tundra (25) | Pond or Lake (18) | - | - | 1 | 5.9 | - | - | - | - |
| | Lake with Emergents (42) | - | - | - | - | - | - | 1 | 14.3 |
| | Lake (50) | - | - | 1 | 5.9 | - | - | - | - |
| | Pond (47) | - | - | - | - | 1 | 3.2 | 3 | 42.9 |
| | River Delta (54) | - | - | 1 | 5.9 | - | - | - | - |
| | Lake Shore (44) | - | - | 1 | 5.9 | - | - | - | - |
| | Pond with Emergents (45) | - | - | 1 | 5.9 | - | - | - | - |
| | Large Lake (49) | - | - | - | - | 1 | 3.2 | - | - |
| Mainland Coast (27) | Shoreline (water side; 15) | 20 | 95.2 | 9 | 52.9 | 28 | 90.3 | 3 | 42.9 |
| River Delta (92) | Tide Flat (13) | - | - | 1 | 5.9 | - | - | - | - |
| TOTAL | | 21 | 100 | 17 | 100 | 31 | 100 | 7 | 100 |

APPENDIX A:
JULY AND AUGUST 2001
WEST DOCK NOAA STATION DATA




Figure A1. Daily mean temperature and wind speed recorded at the Prudhoe Bay NOAA Station at West Dock, Alaska, 15 July – 31 August 2001. Data available at <http://co-ops.nos.noaa.gov/co-ops.html>.

NOAA/NOS/CO-OPS
Water Temperature Plot
9497645 Prudhoe Bay, AK
From 07/15/2001 - 08/31/2001




Figure A2. Hourly mean water temperature recorded at the Prudhoe Bay NOAA Station at West Dock, Alaska, 15 July - 1 September 2001. Data available at <http://co-ops.nos.noaa.gov/co-ops.html>.

NOAA/NOS/CO-OPS
Wind Speed/Dir
9497645 Prudhoe Bay, AK
From 07/15/2001 - 08/31/2001




Figure A3. Hourly mean wind speed and direction recorded at the Prudhoe Bay NOAA Station at West Dock, Alaska, 15 July - 1 September 2001. Data available at <http://co-ops.nos.noaa.gov/co-ops.html>.

NOAA/NOS/CO-OPS
Air Temperature Plot
9497645 Prudhoe Bay, AK
From 07/15/2001 - 08/31/2001




Figure A4. Hourly mean air temperature recorded at the Prudhoe Bay NOAA Station at West Dock, Alaska, 15 July-1 September 2001. Data available at <http://co-ops.nos.noaa.gov/co-ops.html>.

NOAA/NOS/CO-OPS
 Preliminary 6 Minute Water Level (A1) vs Predictions Plot
 9497645 Prudhoe Bay, AK
 from 07/01/2001 - 07/31/2001




Figure A5. Water level relative to mean low, low water level recorded at 6-min intervals at the Prudhoe Bay NOAA Station at West Dock, Alaska, 1-31 July, 2001. Data downloaded at <http://co-ops.nos.noaa.gov/co-ops.html>.

NOAA/NOS/CO-OPS
Preliminary 6 Minute Water Level (A1) vs Predictions Plot
9497645 Prudhoe Bay, AK
From 08/01/2001 - 08/31/2001




Figure A6. Water level relative to mean low, low water level recorded at 6-min intervals at the Prudhoe Bay NOAA Station at West Dock, Alaska, 1-31 August 2001. Data downloaded at <http://co-ops.nos.noaa.gov/co-ops.html>.

**APPENDIX B:
2001 LONG-TAILED DUCK DENSITY BY SURVEY
AND HISTORICAL DATA**




Figure B1. Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July 2001.

Beaufort Sea Waterfowl 2001




Figure B2. Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 8 August 2001.

LONG-TAILED DUCK--11 AUGUST 2001
WESTERN LAGOON
(Spy Island to West Dock)




Figure B3. Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 11 August 2001.




Figure B4. Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July 2001.




Figure B5. Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 8 August 2001.




Figure B6. Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 11 August 2001.




Figure B7. Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July 2001.

Beaufort Sea Waterfowl, 2001




Figure B8. Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 8 August 2001.




Figure B9. Summary of density for long-tailed ducks by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 11 August 2001.

Table B1. Numbers and percentages of long-tailed ducks counted during aerial surveys in nearshore waters of the central Alaska Beaufort Sea, 1977–1984; 1989–1991; 1998–2001 (Johnson and Gazey 1992; Noel et al. 1999, 2002). Data for 1998 include only the eastern lagoon, 1999–2001 include western and eastern lagoons exclusive of 600 series mid-lagoon transects.

| Category* | Survey Year | | | | | | | | | | | | | | | All Years |
|-------------|-------------|---------|--------|--------|--------|--------|-------|--------|---------|---------|---------|--------|--------|--------|--------|-----------|
| | 1977 | 1978 | 1979 | 1980 | 1981 | 1982 | 1983 | 1984 | 1989 | 1990 | 1991 | 1998 | 1999 | 2000 | 2001 | |
| Numbers | | | | | | | | | | | | | | | | |
| 1 | 20,695 | 111,594 | 28,598 | 22,777 | 30,597 | 31,927 | - | 21,998 | 102,968 | 163,915 | 31,316 | 24,455 | 26,478 | 17,836 | 7,355 | 642,509 |
| 2 | 58,310 | 141,801 | 36,157 | 27,826 | 48,711 | 46,964 | 6,144 | 28,399 | 110,975 | 220,758 | 61,441 | 41,344 | 52,663 | 35,618 | 17,560 | 934,671 |
| 3 | 94,461 | 215,199 | 49,456 | 37,549 | 65,768 | 66,794 | - | 33,987 | 138,729 | 277,327 | 120,397 | 43,639 | 57,834 | 40,862 | 21,119 | 1,263,121 |
| 4 | 104,318 | 231,307 | 54,049 | 38,364 | 71,104 | 69,775 | - | 34,972 | 149,408 | 312,073 | 138,408 | 48,979 | 67,427 | 50,787 | 27,175 | 1,398,146 |
| Percentages | | | | | | | | | | | | | | | | |
| 5 | 90.55 | 93.04 | 91.50 | 97.88 | 92.50 | 95.73 | - | 97.18 | 92.85 | 88.87 | 86.99 | 89.10 | 85.77 | 80.46 | 77.71 | 91.83 |
| 6 | 55.90 | 61.30 | 66.90 | 72.53 | 68.51 | 67.31 | - | 81.20 | 74.28 | 70.74 | 44.39 | 84.41 | 78.10 | 70.13 | 64.62 | 68.80 |
| 7 | 21.91 | 51.86 | 57.83 | 60.66 | 46.52 | 47.80 | - | 64.72 | 74.22 | 59.11 | 26.01 | 56.04 | 45.78 | 43.65 | 34.83 | 51.04 |
| 8 | 19.84 | 48.24 | 52.91 | 59.37 | 43.03 | 45.76 | - | 62.90 | 68.92 | 52.52 | 22.63 | 49.93 | 39.27 | 35.12 | 27.07 | 47.11 |
| 9 | 61.73 | 65.89 | 73.11 | 74.11 | 74.06 | 70.31 | - | 83.56 | 79.99 | 79.60 | 51.03 | 94.74 | 91.06 | 87.17 | 83.15 | 74.93 |
| 10 | 35.49 | 78.70 | 79.09 | 81.86 | 62.81 | 67.98 | - | 77.46 | 92.78 | 74.25 | 50.97 | 59.15 | 50.28 | 50.08 | 41.88 | 67.57 |

- * 1 = No. of long-tailed ducks on-transect only on barrier island transects during all surveys
- 2 = No. of long-tailed ducks on-transect on all lagoon transects during all surveys
- 3 = No. of long-tailed ducks on- and off-transect on all lagoon transects during all surveys
- 4 = No. of all birds of all species on- and off-transect on all lagoon transects during all surveys
- 5 = Cat. 3/Cat. 4
- 6 = Cat. 2/Cat. 4
- 7 = Cat. 1/Cat. 3
- 8 = Cat. 1/Cat. 4
- 9 = Cat. 2/Cat. 3
- 10 = Cat. 1/Cat. 2

Table B2. Continued.

| Survey Date Transect Number Transect Area | Non-barrier Island Transect Nos. and Areas Sampled (km ²) | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|---|---|--------|--------|------|------|------|------|-------|-------|-------|-------|--------|--------|--------|---------|---------|---------|--------|---------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--|
| | 30 | 32 | 33 | 50 | 51 | 52 | 53 | 60 | 61 | 62 | 63 | 101 | 102 | 301 | 302 | 401 | 402 | 180 | 181 | 182 | 183 | 184 | 190 | 191 | 192 | 193 | 601 | | | |
| 5 Jun.77 | 5.00 | 5.72 | 7.52 | 4.72 | 4.24 | 4.28 | 5.64 | 5.76 | 4.44 | 4.52 | 5.36 | 8.44 | 6.00 | 6.92 | 5.24 | 7.28 | 5.64 | 5.68 | 4.24 | 5.16 | 5.64 | | | | | | | | | |
| 20 Jun.77 | | | | | | | | | | | | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 2.13 | | | | | | | | | | | | | |
| 5 Jul.77 | | | | | | | | | | | | 0.00 | 0.00 | 0.00 | 0.00 | 0.82 | 2.84 | | | | | | | | | | | | | |
| 28-29 Jul.77 | | | | | | | | | | | | 0.12 | 5.00 | 4.91 | 0.38 | 119.23 | 44.68 | | | | | | | | | | | | | |
| 15 Aug.77 | | | | | | | | | | | | 0.12 | 6.67 | 72.25 | 1505.73 | 1550.55 | 2741.41 | | | | | | | | | | | | | |
| 30 Aug.77 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| 22 Sep.77 | | | | | | | | | | | | 5.45 | 0.17 | 143.35 | 244.08 | 428.57 | 71.63 | | | | | | | | | | | | | |
| 23 Jun.78 | | 1.05 | | | | | | | | | | 5.21 | 0.83 | 626.45 | 1458.78 | 276.53 | 1334.04 | | | | | | | | | | | | | |
| 5 Jul.78 | | 11.71 | | | | | | | | | | 0.00 | 0.00 | 0.00 | 0.00 | 0.41 | 0.00 | | | | | | | | | | | | | |
| 15 Jul.78 | | 6.47 | | | | | | | | | | 0.24 | 0.00 | 3.90 | 7.44 | 3.43 | 0.53 | 24.47 | 107.78 | 0.00 | 0.00 | | | | | | | | | |
| 25 Jul.78 | | 3.50 | | | | | | | | | | 2.96 | 1.67 | 14.74 | 72.14 | 29.69 | 111.03 | | | | | | | | | | | | | |
| 5-6 Aug.78 | | 8.04 | | | | | | | | | | 18.26 | 2.17 | 82.80 | 62.02 | 28.43 | 7.09 | 4.40 | 11.79 | 0.00 | 0.00 | | | | | | | | | |
| 15 Aug.78 | | 14.86 | | | | | | | | | | 0.36 | 0.53 | 53.76 | 75.78 | 11.40 | 0.00 | 183.10 | 117.62 | 0.00 | 0.00 | | | | | | | | | |
| 25 Aug.78 | | 54.37 | | | | | | | | | | 0.00 | 0.00 | 4.77 | 10.88 | 0.00 | 0.00 | | | | | | | | | | | | | |
| 5-6 Sep.78 | | 38.64 | | | | | | | | | | 52.37 | 45.33 | 24.86 | 45.42 | 0.00 | 1.77 | 1.07 | 0.00 | 0.00 | 0.00 | | | | | | | | | |
| 15 Sep.78 | | | | | | | | | | | | 23.70 | 14.50 | 11.13 | 14.12 | 0.14 | 0.00 | 309.15 | 1354.01 | 206.59 | 18.97 | | | | | | | | | |
| 23 Sep.78 | | 29.10 | | | | | | | | | | 163.15 | 228.50 | 33.24 | 25.57 | 2.06 | 13.65 | | | | | | | | | | | | | |
| 22 Jun.79 | | 2.62 | | | | | | | | | | 2.61 | 0.00 | 369.36 | 56.30 | 157.69 | 114.54 | 438.03 | 79.01 | 0.00 | 152.30 | | | | | | | | | |
| 28 Jul.79 | | 4.55 | | | | | | | | | | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.53 | | | | | | | | | | | | | |
| 31 Aug.-1 Sep.79 | | | | | | | | | | | | 0.00 | 9.43 | 131.21 | 138.45 | 41.48 | 17.73 | 18.84 | 104.01 | 23.26 | 0.00 | | | | | | | | | |
| 22 Sep.79 | | 21.50 | | | | | | | | | | 6.57 | 45.83 | 98.70 | 640.29 | 0.00 | 0.00 | | | | | | | | | | | | | |
| 2 Aug.80 | | 71.68 | | | | | | | | | | 19.79 | 9.50 | 5.76 | 0.57 | 76.10 | 162.59 | | | | | | | | | | | | | |
| 18 Jul.81 | | | | | | | | | | | | 0.00 | 0.00 | 250.87 | 61.83 | 134.71 | 213.48 | 0.00 | 16.98 | 0.00 | 0.00 | | | | | | | | | |
| 29 Jul.81 | | | | | | | | | | | | 107.94 | 0.00 | 53.28 | 28.38 | 9.75 | 8.76 | | | | | | | | | | | | | |
| 2 Aug.81 | | 122.03 | | | | | | | | | | 0.00 | 0.17 | 129.48 | 177.87 | 0.00 | 0.35 | | | | | | | | | | | | | |
| 12 Aug.81 | | | | | | | | | | | | 0.00 | 0.00 | 51.16 | 233.59 | 37.23 | 27.66 | 32.22 | 20.05 | 0.00 | 0.00 | | | | | | | | | |
| 29 Aug.81 | | | | | | | | | | | | 0.00 | 0.00 | 7.37 | 8.87 | 0.27 | 4.79 | | | | | | | | | | | | | |
| 11 Sep.81 | | | | | | | | | | | | 8.06 | 0.33 | 15.32 | 8.02 | 0.00 | 0.00 | | | | | | | | | | | | | |
| 18 Jul.82 | 2.40 | 73.43 | 2.66 | | | | | | | | | 71.56 | 18.00 | 194.87 | 20.42 | 0.69 | 1.24 | | | | | | | | | | | | | |
| 31 Jul.82 | | | | | | | | | | | | 0.12 | 0.00 | 12.86 | 55.34 | 5.91 | 9.04 | | | | | | | | | | | | | |
| 14 Aug.82 | | | | | | | | | | | | 0.12 | 0.00 | 386.99 | 193.51 | 22.80 | 11.70 | | | | | | | | | | | | | |
| 28 Aug.82 | 8.00 | 7.69 | 19.95 | | | | | | | | | 1.18 | 0.00 | 28.12 | 0.38 | 0.00 | 0.00 | | | | | | | | | | | | | |
| 23 Sep.82 | 0.20 | 47.90 | 419.68 | | | | | | | | | 10.31 | 1.50 | 58.09 | 94.03 | 0.00 | 0.00 | | | | | | | | | | | | | |
| 29 Jul.83 | | | | | | | | | | | | 8.53 | 0.00 | 254.77 | 197.90 | 17.17 | 73.58 | | | | | | | | | | | | | |
| 8 Aug.84 | 12.00 | 70.28 | 0.00 | | | | | | | | | 7.94 | 0.00 | 209.54 | 387.80 | 0.00 | 0.00 | 101.76 | 227.19 | 113.37 | 7.80 | | | | | | | | | |
| 5 Aug.89 | 0.00 | 10.49 | | 3.18 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 275.01 | 318.70 | | 0.00 | 0.00 | 322.29 | 0.00 | | | | | | | | | | |
| 8 Aug.89 | 0.00 | 0.00 | | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 79.48 | 0.57 | | 4.40 | 23.58 | 23.45 | 0.00 | | | | | | | | | | |
| 9 Aug.89 | 1.41 | 0.00 | | 0.00 | 7.08 | 5.79 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.33 | 14.45 | 0.57 | | 0.00 | 0.00 | 211.24 | 39.01 | | | | | | | | | | |
| 18 Jul.90 | 1.81 | 24.65 | 0.00 | | | | | | | | | 36.89 | 41.52 | 0.00 | 0.00 | 8.29 | 0.00 | 227.75 | 28.72 | 7.28 | 0.00 | 62.68 | 195.52 | 34.88 | 0.00 | 490.03 | 372.32 | 165.29 | 63.62 | |
| 20 Jul.90 | 9.68 | 15.56 | 1.20 | | | | | | | | | 0.35 | 1.56 | 0.22 | 0.56 | 18.84 | 0.00 | 147.40 | 83.21 | 7.01 | 0.00 | 144.72 | 66.75 | 34.30 | 21.28 | 200.32 | 242.63 | 160.14 | 77.95 | |
| 23 Jul.90 | 0.00 | 15.03 | 0.00 | | | | | | | | | 0.59 | 0.00 | 59.39 | 17.75 | 2.47 | 0.35 | 22.18 | 2.36 | 0.19 | 2.66 | 387.64 | 452.46 | 216.57 | 53.25 | 387.64 | 452.46 | 216.57 | 53.25 | |
| 3 Aug.90 | 0.00 | 123.95 | 1.06 | | | | | | | | | 13.63 | 2.50 | 130.64 | 28.72 | 0.00 | 4.26 | 67.81 | 54.25 | 4.84 | 0.00 | 187.03 | 802.23 | 153.00 | 100.00 | 187.03 | 802.23 | 153.00 | 100.00 | |
| 4 Aug.90 | 0.00 | 156.99 | 0.00 | | | | | | | | | 0.00 | 0.00 | 112.86 | 34.35 | 0.96 | 0.00 | 13.20 | 29.25 | 26.74 | 2.66 | 319.30 | 277.01 | 150.57 | 32.01 | 319.30 | 277.01 | 150.57 | 32.01 | |
| 9 Aug.90 | 54.55 | 1.20 | | | | | | | | | | 59.97 | 58.30 | 0.14 | 0.18 | 0.18 | 31.16 | 41.04 | 14.15 | 0.00 | | 36.23 | 189.06 | 56.00 | 25.91 | 36.23 | 189.06 | 56.00 | 25.91 | |
| 16 Aug.90 | 0.17 | 0.27 | | | | | | | | | | 7.23 | 4.39 | 2.61 | 0.00 | | | | | | | | | | | | | | | |
| 20 Aug.90 | 3.23 | 14.16 | 2.93 | | | | | | | | | 0.00 | 0.00 | 2.21 | 0.00 | 4.27 | 0.00 | 19.51 | 222.14 | 4.95 | 1.06 | 0.35 | 0.00 | 27.52 | 0.00 | 45.09 | 406.92 | 234.43 | 2.13 | |
| 2 Sep.90 | 0.00 | 0.52 | 14.10 | | | | | | | | | 0.00 | 0.22 | 0.00 | 0.00 | 0.00 | 15.46 | 1.34 | 69.00 | 0.18 | 44.54 | 13.21 | 0.97 | 1.95 | 18.20 | 40.85 | 35.00 | 4.27 | | |
| 4 Sep.90 | | 7.52 | 5.98 | | | | | | | | | 19.94 | 8.11 | 0.00 | 0.35 | 37.32 | 8.02 | 237.21 | 3.72 | | | 24.21 | 71.65 | 31.00 | 5.18 | 24.21 | 71.65 | 31.00 | 5.18 | |
| 5 Sep.90 | 0.00 | 14.63 | | | | | | | | | | 1.59 | 3.24 | 1.10 | 0.53 | 0.00 | 0.47 | 62.02 | 192.73 | | | 3.01 | 140.40 | 48.86 | 20.83 | 3.01 | 140.40 | 48.86 | 20.83 | |
| 18 Jul.91 | 1.22 | 0.13 | | | | | | | | | | 12.43 | 5.73 | 5.22 | 2.84 | | | | | | | | | | | | | | | |
| 19 Jul.91 | 15.03 | 4.65 | | | | | | | | | | 22.40 | 12.80 | 7.42 | 0.18 | 23.06 | 7.55 | 6.78 | 0.00 | | | 45.70 | 108.04 | 23.29 | 4.57 | 45.70 | 108.04 | 23.29 | 4.57 | |
| 20 Jul.91 | 25.00 | 14.36 | | | | | | | | | | 30.35 | 15.46 | 11.40 | 4.96 | 5.11 | 7.78 | 4.84 | 0.35 | | | 11.78 | 70.76 | 33.00 | 8.84 | 11.78 | 70.76 | 33.00 | 8.84 | |
| 22 Jul.91 | 33.22 | 0.66 | | | | | | | | | | 27.89 | 10.89 | 19.23 | 0.18 | 66.90 | 63.44 | 116.67 | 0.00 | | | 67.20 | 101.34 | 30.71 | 50.81 | 67.20 | 101.34 | 30.71 | 50.81 | |
| 4 Aug.91 | 41.96 | 13.96 | | | | | | | | | | 76.59 | 88.28 | 16.48 | 13.48 | 36.44 | 34.67 | 61.24 | 0.35 | | | 49.52 | 78.13 | 83.14 | 44.31 | 49.52 | 78.13 | 83.14 | 44.31 | |
| 10 Aug.91 | 144.58 | 10.24 | | | | | | | | | | 69.94 | 10.31 | 13.32 | 1.77 | 30.11 | 16.04 | 67.44 | 0.53 | | | 39.65 | 94.64 | 38.43 | 17.28 | 39.65 | 94.64 | 38.43 | 17.28 | |
| 14 Aug.91 | 55.94 | 0.00 | | | | | | 4.50 | 0.00 | 0.00 | 0.59 | 0.00 | 41.47 | 16.41 | 2.34 | 14.18 | 0.00 | 28.30 | 75.39 | 0.00 | | 101.43 | 10.71 | 65.71 | 5.89 | 101.43 | 10.71 | 65.71 | 5.89 | |
| 16 Aug.91 | 0.60 | 36.89 | 1.33 | | | | | 23.44 | 65.99 | 23.23 | 1.31 | 0.47 | 1.67 | 138.87 | 147.52 | 1.92 | 0.00 | 35.21 | 23.35 | 83.72 | 0.18 | 19.90 | 9.60 | 9.71 | 8.94 | 19.90 | 9.60 | 9.71 | 8.94 | |
| 21 Aug.91 | 17.20 | 17.48 | 1.73 | | | | | 9.20 | 4.50 | 11.06 | 30.22 | 0.47 | 10.33 | 217.77 | 9.16 | 3.57 | 0.18 | 10.21 | 30.66 | 46.32 | 0.00 | 10.19 | 33.26 | 2.86 | 2.44 | 10.19 | 33.26 | 2.86 | 2.44 | |
| 5 Aug. 98 | | | | | | | | | | | | | | | | | | 8.09 | 9.11 | 0.00 | 0.00 | | | | | 122.97 | 87.41 | 85.03 | 0.57 | |
| 14 Aug. 98 | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |

Table B2. Continued.

| Survey Date | Tundra Transect Nos. and Areas Sampled (km ²) | | | | | | Tundra Transect Nos. and Areas Sampled (km ²) | | | |
|------------------|---|--------|-------|--------|-------|--------|---|------|------|------|
| Transect Number | 602 | 603 | 604 | 605 | 606 | 607 | 500 | 501 | 502 | 503 |
| Transect Area | 5.16 | 5.64 | 6.28 | 4.48 | 7.00 | 9.84 | 6.56 | 5.42 | 6.91 | 5.52 |
| 5 Jun.77 | | | | | | | | | | |
| 20 Jun.77 | | | | | | | | | | |
| 5 Jul.77 | | | | | | | | | | |
| 28-29 Jul.77 | | | | | | | | | | |
| 15 Aug.77 | | | | | | | | | | |
| 30 Aug.77 | | | | | | | | | | |
| 22 Sep.77 | | | | | | | | | | |
| 23 Jun.78 | | | | | | | | | | |
| 5 Jul.78 | | | | | | | | | | |
| 15 Jul.78 | | | | | | | | | | |
| 25 Jul.78 | | | | | | | | | | |
| 5-6 Aug.78 | | | | | | | | | | |
| 15 Aug.78 | | | | | | | | | | |
| 25 Aug.78 | | | | | | | | | | |
| 5-6 Sep.78 | | | | | | | | | | |
| 15 Sep.78 | | | | | | | | | | |
| 23 Sep.78 | | | | | | | | | | |
| 22 Jun.79 | | | | | | | | | | |
| 28 Jul.79 | | | | | | | | | | |
| 31 Aug.-1 Sep.79 | | | | | | | | | | |
| 22 Sep.79 | | | | | | | | | | |
| 2 Aug.80 | | | | | | | | | | |
| 18 Jul.81 | | | | | | | | | | |
| 29 Jul.81 | | | | | | | | | | |
| 2 Aug.81 | | | | | | | | | | |
| 12 Aug.81 | | | | | | | | | | |
| 29 Aug.81 | | | | | | | | | | |
| 11 Sep.81 | | | | | | | | | | |
| 18 Jul.82 | | | | | | | | | | |
| 31 Jul.82 | | | | | | | | | | |
| 14 Aug.82 | | | | | | | | | | |
| 28 Aug.82 | | | | | | | | | | |
| 23 Sep.82 | | | | | | | | | | |
| 29 Jul.83 | | | | | | | | | | |
| 8 Aug.84 | | | | | | | | | | |
| 6 Aug.89 | | | | | | | | | | |
| 8 Aug.89 | | | | | | | | | | |
| 9 Aug.89 | | | | | | | | | | |
| 18 Jul.90 | | | | | | | | | | |
| 20 Jul.90 | | | | | | | | | | |
| 23 Jul.90 | | | | | | | | | | |
| 3 Aug.90 | | | | | | | | | | |
| 4 Aug.90 | | | | | | | | | | |
| 9 Aug.90 | | | | | | | | | | |
| 16 Aug.90 | | | | | | | | | | |
| 20 Aug.90 | | | | | | | | | | |
| 2 Sep.90 | | | | | | | | | | |
| 4 Sep.90 | | | | | | | | | | |
| 5 Sep.90 | | | | | | | | | | |
| 18 Jul.91 | | | | | | | | | | |
| 19 Jul.91 | | | | | | | | | | |
| 20 Jul.91 | | | | | | | | | | |
| 22 Jul.91 | | | | | | | | | | |
| 4 Aug.91 | | | | | | | | | | |
| 10 Aug.91 | | | | | | | | | | |
| 14 Aug.91 | | | | | | | | | | |
| 16 Aug.91 | | | | | | | | | | |
| 21 Aug.91 | | | | | | | | | | |
| 5 Aug.98 | | | | | | | 0.00 | 0.00 | 0.00 | 0.00 |
| 14 Aug.98 | | | | | | | 0.00 | 0.00 | 0.00 | 0.00 |
| 16 Aug.98 | | | | | | | 0.00 | 0.00 | 0.00 | 0.00 |
| 31 Aug.98 | | | | | | | 0.00 | 5.79 | 0.00 | 0.00 |
| 3 Sept.98 | | | | | | | 0.00 | 9.11 | 0.00 | 0.00 |
| 30 Jul.99 | 19.51 | 35.70 | 78.96 | 65.98 | 21.23 | 188.73 | 0.00 | 5.76 | 0.76 | 6.16 |
| 1 Aug.99 | 116.21 | 75.14 | 0.00 | 16.60 | 5.17 | 151.63 | 0.00 | 0.88 | 0.00 | 0.00 |
| 4 Aug.99 | | | | | | | | | | |
| 5 Aug.99 | 30.66 | 100.54 | 0.75 | 9.23 | 8.33 | 31.99 | 0.00 | 0.00 | 0.00 | 0.00 |
| 11 Aug.99 | 33.71 | 109.57 | | | | | | | | |
| 26 Aug.99 | 16.60 | 13.83 | 47.20 | 111.07 | 1.87 | 31.08 | 0.00 | 0.00 | 0.00 | 0.00 |
| 1 Aug.00 | 8.54 | 53.16 | 38.67 | 39.42 | 25.41 | 34.12 | 0.00 | 6.41 | 0.00 | 2.82 |
| 8 Aug.00 | 75.99 | 122.22 | 14.66 | 31.35 | 5.28 | 96.08 | 0.00 | 0.00 | 0.00 | 1.03 |
| 12 Aug.00 | 41.86 | 26.12 | 3.70 | 17.42 | 41.60 | 21.17 | 0.00 | 4.17 | 0.00 | 2.41 |
| 14 Aug.00 | 60.63 | 9.19 | | | | | | | | |
| 24 Aug.00 | 0.19 | 50.29 | 0.00 | 5.58 | 19.01 | 29.41 | 0.00 | 3.40 | 0.00 | 4.43 |
| 23 Jul.01 | 10.66 | 32.43 | 0.00 | 0.38 | 2.89 | 34.38 | | | | |
| 8 Aug.01 | 119.04 | 40.61 | 0.00 | 7.63 | 73.16 | 40.26 | 0.00 | 0.00 | 0.00 | 0.18 |
| 11 Aug.01 | 67.31 | 29.70 | 0.00 | 0.00 | 11.65 | 4.46 | 0.00 | 0.00 | 0.92 | 8.15 |
| 3 | 13 | 13 | 11 | 11 | 11 | 11 | 15 | 15 | 15 | 15 |
| Mean = | 46.17 | 53.73 | 16.54 | 27.70 | 19.60 | 60.30 | 0.00 | 2.37 | 0.11 | 1.68 |
| s.d. = | 39.34 | 36.90 | 26.20 | 33.97 | 21.42 | 59.27 | 0.00 | 3.12 | 0.30 | 2.63 |
| c.v. = | 0.85 | 0.69 | 1.58 | 1.23 | 1.09 | 0.98 | | 1.32 | 2.65 | 1.58 |

Table B3. Continued.

| coll. code | sex | band | loc. no. | date | habitat | species | no. birds | type | behavior | habitat | no. birds | off. time | sub. birds | off. time | total | time | remarks |
|------------|------|------|----------|-----------|---------|------------|-----------|------|----------|---------|-----------|-----------|------------|-----------|-------|--------|---------------------------|
| P598 | 1106 | 27 | 134 | 7/23/2001 | 001 LN | 001 BOAT | 1 | 01 | 07 | 15 | 0 | 0 | 0 | 0 | 0 | 145119 | NET AND BOAT AT SEA |
| P598 | 1106 | 27 | 134 | 7/23/2001 | 001 LN | 001 BOAT | 1 | 01 | 07 | 15 | 0 | 0 | 0 | 0 | 0 | 145119 | BOAT ANCHORED ON SH |
| P598 | 2186 | 68 | 135 | 6/8/2001 | 003 LN | 001 BOAT | 1 | 01 | 07 | 15 | 0 | 0 | 0 | 0 | 0 | 145013 | ZODIAC AT MONTICORN |
| P598 | 3074 | 74 | 607 | 6/8/2001 | 003 RR | 001 BOAT | 1 | 01 | 07 | 15 | 0 | 0 | 0 | 0 | 0 | 172855 | SMALL BOAT WITH PROP |
| P598 | 3268 | 81 | 191 | 6/8/2001 | 003 LN | 001 BOAT | 1 | 01 | 07 | 15 | 0 | 0 | 0 | 0 | 0 | 172855 | ZODIACS ANCHORED |
| P598 | 4134 | 114 | 135 | 6/11/2001 | 003 LN | 001 BOAT | 1 | 01 | 07 | 15 | 0 | 0 | 0 | 0 | 0 | 165012 | TELEMETRY WITH BOAT |
| P598 | 4437 | 135 | 193 | 6/11/2001 | 003 LN | 001 BOAT | 1 | 01 | 07 | 15 | 0 | 0 | 0 | 0 | 0 | 175001 | TELEMETRY FORTWAT |
| P598 | 4710 | 125 | 193 | 6/11/2001 | 003 LN | 001 BOAT | 1 | 01 | 07 | 15 | 0 | 0 | 0 | 0 | 0 | 174808 | ZAMBONIA ALONG SH |
| P598 | 4813 | 127 | 191 | 6/11/2001 | 003 LN | 001 BOAT | 1 | 01 | 07 | 15 | 0 | 0 | 0 | 0 | 0 | 174808 | ZAMBONIA ALONG SH |
| P598 | 4824 | 127 | 191 | 6/11/2001 | 003 LN | 001 BOAT | 1 | 01 | 07 | 15 | 0 | 0 | 0 | 0 | 0 | 174808 | OFFSHORE AT PT THOM |
| P598 | 4838 | 127 | 191 | 6/11/2001 | 003 LN | 001 BOAT | 1 | 01 | 07 | 15 | 0 | 0 | 0 | 0 | 0 | 174808 | ZODIAC ON BEACH |
| P598 | 814 | 19 | 22 | 7/23/2001 | 001 LN | 001 BOAT | 1 | 01 | 07 | 15 | 0 | 0 | 0 | 0 | 0 | 145119 | ALASKA CLEAN SEAS |
| P598 | 815 | 19 | 22 | 7/23/2001 | 001 LN | 001 BOAT | 1 | 01 | 07 | 15 | 0 | 0 | 0 | 0 | 0 | 145119 | ALASKA CLEAN SEAS |
| P598 | 1006 | 45 | 24 | 6/8/2001 | 001 LN | 002 WBOAT | 1 | 04 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 151613 | 2 BOOM BARGES, 2 BA |
| P598 | 2180 | 59 | 22 | 6/8/2001 | 001 LN | 002 WBOAT | 1 | 04 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 151613 | 2 BOOM BARGES, 2 BA |
| P598 | 2189 | 59 | 22 | 6/8/2001 | 001 LN | 002 WBOAT | 1 | 04 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 151613 | 2 BOOM BARGES, 2 BA |
| P598 | 4634 | 122 | 161 | 6/11/2001 | 003 LN | 002 WBOAT | 1 | 04 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 175405 | 3 BOATS AT BARGE, B |
| P598 | 4761 | 126 | 192 | 6/11/2001 | 003 LN | 002 WBOAT | 1 | 01 | 07 | 15 | 27 | 0 | 0 | 0 | 0 | 175405 | HEADED OUT THROUGH |
| P598 | 4768 | 126 | 192 | 6/11/2001 | 003 LN | 002 WBOAT | 1 | 01 | 07 | 15 | 27 | 0 | 0 | 0 | 0 | 175508 | BOAT PULLING IN TO |
| P598 | 816 | 19 | 22 | 7/23/2001 | 001 LN | 001 BOAT | 1 | 01 | 07 | 15 | 0 | 0 | 0 | 0 | 0 | 145119 | BOAT ABOVE-CROSSED |
| P598 | 1911 | 45 | 24 | 6/8/2001 | 001 LN | 003 BEISMC | 1 | 04 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 145559 | 1 SEISMIC BARGE, 2 |
| P598 | 2189 | 59 | 22 | 6/8/2001 | 001 LN | 003 BEISMC | 1 | 04 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 145559 | 1 SEISMIC BARGE, 2 |
| P598 | 2401 | 56 | 101 | 6/8/2001 | 001 LN | 003 BEISMC | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 150210 | R OF ISLAND |
| P598 | 2403 | 56 | 101 | 6/8/2001 | 001 LN | 003 BEISMC | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 150210 | SEISMIC SOURCE BOAT |
| P598 | 3990 | 101 | 201 | 6/11/2001 | 001 LN | 003 BEISMC | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 145559 | MOVING WEST |
| P598 | 3996 | 101 | 201 | 6/11/2001 | 001 LN | 003 BEISMC | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 145559 | MOVING WEST |
| P598 | 4000 | 101 | 201 | 6/11/2001 | 001 LN | 003 BEISMC | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 145559 | WESTERN ENDEAVOR |
| P598 | 4011 | 101 | 201 | 6/11/2001 | 001 LN | 003 BEISMC | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 145559 | WESTERN ENDEAVOR |
| P598 | 4062 | 102 | 101 | 6/11/2001 | 001 LN | 003 BEISMC | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 145559 | WESTERN ENDEAVOR |
| P598 | 4072 | 102 | 101 | 6/11/2001 | 001 LN | 003 BEISMC | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 145559 | WESTERN ENDEAVOR |
| P598 | 4074 | 102 | 101 | 6/11/2001 | 001 LN | 003 BEISMC | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 145559 | WESTERN ENDEAVOR |
| P598 | 4151 | 105 | 22 | 7/23/2001 | 001 LN | 004 SHIP | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 181418 | 2 BOOM BARGES, NOT H |
| P598 | 3446 | 91 | 24 | 6/11/2001 | 001 LN | 004 SHIP | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 181418 | 2 BOOM BARGES, NOT H |
| P598 | 2468 | 60 | 130 | 6/8/2001 | 002 LN | 004 SHIP | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 181418 | 2 BOOM BARGES, NOT H |
| P598 | 2500 | 60 | 130 | 6/8/2001 | 002 LN | 004 SHIP | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 181418 | 2 BOOM BARGES, NOT H |
| P598 | 4165 | 106 | 130 | 6/11/2001 | 002 LN | 004 SHIP | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 144022 | BARGE AT WEST DOCK-3 BOOM |
| P598 | 4168 | 106 | 130 | 6/11/2001 | 002 LN | 004 SHIP | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 144022 | BARGE AT WEST DOCK-3 BOOM |
| P598 | 517 | 13 | 31 | 7/23/2001 | 002 LN | 006 PLANE | 1 | 01 | 11 | 15 | 11 | 1 | 0 | 0 | 0 | 150143 | LARGE BARGE |
| P598 | 1252 | 30 | 133 | 7/23/2001 | 003 LN | 006 PLANE | 1 | 01 | 11 | 15 | 11 | 1 | 0 | 0 | 0 | 150143 | LARGE BARGE |
| P598 | 1408 | 35 | 163 | 7/23/2001 | 003 LN | 006 PLANE | 1 | 01 | 11 | 15 | 11 | 1 | 0 | 0 | 0 | 150143 | LARGE BARGE |
| P598 | 1468 | 36 | 163 | 7/23/2001 | 003 LN | 006 PLANE | 1 | 01 | 11 | 15 | 11 | 1 | 0 | 0 | 0 | 150143 | LARGE BARGE |
| P598 | 2372 | 55 | 201 | 6/8/2001 | 001 RR | 007 BOUY | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 150327 | USPWS PLANE ON SHOR |
| P598 | 2381 | 56 | 101 | 6/8/2001 | 001 LN | 007 BOUY | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 150327 | USPWS PLANE ON SHOR |
| P598 | 2387 | 56 | 101 | 6/8/2001 | 001 LN | 007 BOUY | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 150327 | USPWS PLANE ON SHOR |
| P598 | 2392 | 56 | 101 | 6/8/2001 | 001 LN | 007 BOUY | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 150327 | USPWS PLANE ON SHOR |
| P598 | 2399 | 56 | 101 | 6/8/2001 | 001 RR | 007 BOUY | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 150327 | USPWS PLANE ON SHOR |
| P598 | 4659 | 102 | 101 | 6/11/2001 | 001 LN | 007 BOUY | 1 | 01 | 07 | 08 | 13 | 0 | 0 | 0 | 0 | 150327 | USPWS PLANE ON SHOR |

Table B4. Habitat associations of long-tailed ducks during aerial surveys in the barrier island-lagoon systems and tundra transects between Spy Island and Brownlow Point, Alaska, 23 July-11 August 2001.

| General Habitat Type | Specific Habitat Type | Number of Sightings 23 July 01 | Number of Sightings 8 Aug 01 | Number of Sightings 11 Aug 01 | Total All Surveys | Percent of Total |
|---------------------------|----------------------------|-----------------------------------|---------------------------------|----------------------------------|-------------------|------------------|
| Lagoon (9) | Smooth fast ice (1) | 1 | - | - | 1 | 0.1 |
| | Lagoon (9) | 91 | 145 | 69 | 305 | 35.6 |
| | Shoreline (land side) (14) | 1 | - | - | 1 | 0.1 |
| Barrier Island (11) | Lagoon (9) | 2 | - | - | 2 | 0.2 |
| | Smooth fast ice (1) | 3 | - | - | 3 | 0.4 |
| | Island (11) | 1 | - | - | 1 | 0.1 |
| | Shoreline (land side; 14) | 5 | - | - | 5 | 0.6 |
| | Shoreline (water side; 15) | 166 | 150 | 40 | 356 | 41.5 |
| Artificial Structure (79) | Shoreline (land side) (14) | 1 | - | - | 1 | 0.1 |
| | Shoreline (15) | 1 | - | - | 1 | 0.1 |
| Nearshore Sea <3 mi. (13) | Smooth fast ice (1) | 1 | - | - | 1 | 0.1 |
| | Ocean (8) | 17 | 7 | 6 | 30 | 3.5 |
| | Lagoon (9) | 12 | 4 | - | 16 | 1.9 |
| Unclassified Tundra (24) | Shoreline (waterside) (15) | 1 | - | - | 1 | 0.1 |
| Wet Tundra (25) | Completely vegetated (25) | - | 1 | - | 1 | 0.1 |
| | Pond (47) | - | - | 1 | 1 | 0.1 |
| | Medium lake (51) | - | - | 3 | 3 | 0.4 |
| Mainland Coast (27) | Spit (10) | 1 | - | - | 1 | 0.1 |
| | Island (11) | 1 | - | - | 1 | 0.1 |
| | Shoreline (water side; 15) | 21 | 73 | 31 | 125 | 14.6 |
| River Delta (92) | Tidal flat (13) | - | 1 | - | 1 | 0.1 |
| ALL HABITATS | | 326 | 381 | 150 | 857 | 100 |

Table B5. Long-tailed duck behavior during aerial surveys in the barrier island-lagoon systems and tundra transects between Spy Island and Browlow Point, Alaska, 23 July-11 August 2001.

| Specific Behaviors | Sightings 23 July 01 | Sightings 8 Aug 01 | Sightings 11 Aug 01 | Total All Surveys | Percent of Total |
|-----------------------------------|-------------------------|-----------------------|------------------------|----------------------|---------------------|
| Hauled out on ice (01) | - | - | 1 | 1 | 0.1 |
| Hauled out on land (02) | 1 | 3 | - | 4 | 0.5 |
| Hauled out (general) (03) | - | 12 | 4 | 16 | 1.9 |
| Hauled out on land-swim/dive (06) | - | - | - | 0 | 0.0 |
| Swimming (07) | 277 | 359 | 110 | 746 | 87.0 |
| Swimming then diving (08) | 12 | 2 | 4 | 18 | 2.1 |
| Swimming then flying (09) | - | 1 | - | 1 | 0.1 |
| Flying (11) | 2 | - | - | 2 | 0.2 |
| Standing (15) | 2 | 1 | 1 | 4 | 0.5 |
| Rest (23) | 9 | - | - | 9 | 1.1 |
| Incubate (30) | 1 | - | - | 1 | 0.1 |
| Unknown or unrecorded (99) | 22 | 3 | 30 | 55 | 6.4 |
| TOTAL | 326 | 381 | 150 | 857 | 100 |

**APPENDIX C:
2001 DISTRIBUTION MAPS FOR SELECTED SPECIES**




Figure C3. Summary of density for Pacific loons by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July-11 August 2001.

**RED-THROATED LOONS 2001;
WESTERN LAGOON
(Spy Island to West Dock)**




Figure C4. Summary of density for red-throated loons by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July-11 August 2001.




Figure C5. Summary of density for red-throated loons by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July-11 August 2001.




Figure C6. Summary of density for red-throated loons by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July-11 August 2001.

Beaufort Sea Waterfowl, 2001

YELLOW-BILLED LOONS 2001
WESTERN LAGOON
(Spy Island to West Dock)




Figure C7. Summary of density for yellow-billed loons by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July-11 August 2001.




Figure C8. Summary of density for yellow-billed loons by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July-11 August 2001.




Figure C9. Summary of density for yellow-billed loons by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July-11 August 2001.

Beaufort Sea Waterfowl, 2001

**JAEGERS 2001
WESTERN LAGOON
(Spy Island to West Dock)**






Figure C10. Summary of density for jaegers by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July-11 August 2001.

**JAEGERS 2001
EASTERN LAGOON (Pole Island to Brownlow Point)**



Projection: UTM Zone 6, NAD 27 for Alaska



Figure C11. Summary of density for jaegers by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July-11 August 2001.




Figure C12. Summary of density for black guillemots by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July-11 August 2001.




Figure C13. Summary of density for black guillemots by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July-11 August 2001.




Figure C14. Summary of density for black guillemots by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July-11 August 2001.

Beaufort Sea Waterfowl, 2001




Figure C15. Summary of density for glaucous gulls by 30-s time period segments in the barrier island-lagoon system between Spy Island and West Dock, Alaska, 23 July-11 August 2001.

GLAUCOUS GULLS 2001 - STEFANSSON SOUND
 (West Dock to Pole Island)




Figure C16. Summary of density for glaucous gulls by 30-s time period segments in the barrier island-lagoon system between West Dock and Pole Island, Alaska, 23 July-11 August 2001.




Figure C17. Summary of density for glaucous gulls by 30-s time period segments in the barrier island-lagoon system between Pole Island and Brownlow Point, Alaska, 23 July-11 August 2001.

Beaufort Sea Waterfowl, 2001